

## Review Paper

## Research progress on the contamination status and control policy of *Giardia* and *Cryptosporidium* in drinking water

Cuimin Feng<sup>a,b,\*</sup>, Zhen Xu<sup>a,b</sup>, Ying Li<sup>a,b</sup>, Na Zhu<sup>a,b</sup> and Zile Wang<sup>id a,b</sup><sup>a</sup> Key Laboratory of Urban Stormwater System and Water Environment, Ministry of Education, Beijing University of Civil Engineering and Architecture, Beijing 100044, China<sup>b</sup> National Demonstration Centre for Experimental Water Environment Education, Beijing University of Civil Engineering and Architecture, Beijing 100044, China

\*Corresponding author. E-mail: feng-cuimin@sohu.com

 ZW, 0000-0003-2481-6015

### ABSTRACT

*Cryptosporidium* and *Giardia* are globally recognized protozoa that directly cause human diarrhea. Their transmission route mainly involves drinking contaminated water, thus needing proper water treatment to avoid human infection. At present, there is a lack of review on the infection status and control measures of the two protozoa. Hence, this article summarizes and compares the infection status and the role of drinking water in transmitting the *Cryptosporidium* and *Giardia* in some key countries in Asia, Africa, Europe, and the Americas. With collected data, this review offers recommendations for sanitary control and provides theoretical support for the application of drinking water treatment projects.

**Key words:** *Cryptosporidium*, disinfection, drinking water, *Giardia*, water treatment

### HIGHLIGHTS

- Need to strengthen preventive measures of *Giardia* and *Cryptosporidium* in animals.
- Improve sanitation facilities, pay attention to non-core areas, and avoid concentrated outbreaks.
- Reasonable selection of testing methods according to the needs and conditions of each country.
- The water treatment process parameters need to be reasonably selected according to the natural water quality conditions.

### INTRODUCTION

*Giardia lamblia* and *Cryptosporidium* are two parasitic protozoa that directly cause human diarrhea (Smith *et al.* 2006). Giardiasis and cryptosporidiosis are common non-viral infectious diseases, ranking highest among diarrhea-causing parasites (Cai 2005). *Giardia* and *Cryptosporidium* are commonly found in natural water bodies, especially in areas polluted with agriculture and animal husbandry wastes (Bukhari *et al.* 1997). Although the global distribution of water resources is uneven, the common feature is that natural water bodies are used as the main source of domestic water. Therefore, the water treatment industry should emphasize biosafety risks caused by these two protozoa (Cui *et al.* 2006; Xiao *et al.* 2013; Zhang *et al.* 2010; Ma *et al.* 2014; Sun *et al.* 2014).

*Giardia* is a genus of the anaerobic flagellated protozoa and a common parasite that cause human intestinal infections. The tetranuclear cyst is the infective stage following a fecal-oral transmission. Once humans and animals ingest the infective cyst, it decapsulates through the action of the digestive juice and then develops into a trophozoite. Trophozoites parasitize the duodenum or the front end of the small intestine and multiply by longitudinal binary fission. Encystation occurs as the trophozoites move to the colon and then form the cysts excreted through the feces (Kofoid & Christiansen 1915). The *Giardia* cysts measure (8–12)  $\mu\text{m} \times$  (7–10)  $\mu\text{m}$  and become infective after two weeks at 25 °C or 11 weeks at 4 °C (Graczyk *et al.* 2008; Silva & Sabogal-Paz 2020; Singer *et al.* 2020).

*Cryptosporidium* is one of the apicomplexan parasitic alveolates, causing a zoonotic disease called cryptosporidiosis manifested clinically by watery diarrhea. Cryptosporidiosis is one of the six common diseases that cause diarrhea in the world

This is an Open Access article distributed under the terms of the Creative Commons Attribution Licence (CC BY-NC-ND 4.0), which permits copying and redistribution for non-commercial purposes with no derivatives, provided the original work is properly cited (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

(Cheng 2015). The life history of *Cryptosporidium* involves five developmental stages: the trophozoite, schizont, gametocyte, zygote, and oocyst. Oocyst is the infective stage of *Cryptosporidium*. Oocysts are round or elliptical with a diameter of 4–6  $\mu\text{m}$ . The wall is smooth and colorless. A mature oocyst contains four crescent-shaped sporozoites and one residual body. After humans and animals swallow the oocysts, the sporozoites escape through the digestive juice and invade the intestinal epithelial cells. Sporozoites develop into trophozoites and type I schizonts. After three nuclear divisions, type I schizonts will have eight merozoites. After being released, the merozoites invade the intestinal epithelial cells and develop into second-generation trophozoites, which develop into type II schizonts after two nuclear divisions. Mature type II merozoites contain 4 merozoites. After the merozoites are released, they invade epithelial cells and develop into female and male gametocytes, further developing into female and male gametes. After fertilization of gametes, it forms the zygote that develops into oocysts. Mature oocysts contain four naked sporozoites. There are two types of oocysts: thin-walled and thick-walled oocysts. Sporozoites of the thin-walled oocysts (~20%) invade the intestinal epithelial cells directly after escaping and proliferate to form an autologous infection. Meanwhile, thick-walled oocysts (~80%) are formed in the intestinal epithelial cells or intestinal cavity and are excreted with the host's feces. It takes 5–11 days to complete the entire life cycle of *Cryptosporidium* (DuPont *et al.* 1995; Xue 2006; Wang & Luo 2016).


## HARM AND CONTAMINATION STATUS OF *GIARDIA* AND *CRYPTOSPORIDIUM*

### Hazards and transmission of *Giardia* and *Cryptosporidium*

*Giardia* and *Cryptosporidium* have a high biosecurity risk due to their wide-spreading routes and difficulty inactivating their infective stages. These intestinal pathogenic microorganisms grow and reproduce in the host body and are finally excreted in the feces in the form of oocysts and cysts, respectively. They can infect other hosts through contaminated water or food (Figure 1).

The transmission routes of *Giardia* and *Cryptosporidium* are complex and diverse, including contact transmission, water transmission, food transmission, and respiratory transmission (Thompson 2008). Among them, water transmission is the most important route. Interestingly, a few studies have shown that unclean sex is also a transmission route (Escobedo *et al.* 2014). Even developed and developing countries with comprehensive water treatment technologies can also be vulnerable to *Giardia* and *Cryptosporidium* environmental contamination. Some of these reasons are described below.

First, *Giardia* and *Cryptosporidium* have a wide transmission route. Although countries with comprehensive water treatment technology can guarantee drinking water quality, other ways such as vegetable pollution and contact with livestock and pets still pose greater risks (Fayer *et al.* 2000; Chen *et al.* 2002; Nahhas & Aboualchamat 2020). Some studies have shown that raw vegetables play a role in spreading parasitic food-borne diseases (Blackburn & McClure 2002; Eraky *et al.*



**Figure 1** | The transmission route of *Giardia* and *Cryptosporidium*.

2014). Several studies have also shown that people are more susceptible to intestinal worms and protozoa by eating fruits or vegetables (Al-Shawa & Mwafy 2007; Adanir & Tasci 2013; Ismail 2016). Researches have revealed that the reason for this phenomenon may be that vegetables are polluted by wastewater during irrigation or directly contaminated by animals and humans during harvesting, packaging, transportation, processing, distribution, and sales (Amoah *et al.* 2007; Gabre & Shakir 2016). The increase in the number of pets raised has also increased the infection rate of humans with *Giardia* and *Cryptosporidium*. In particular, common pets in most areas of China are likely to be infected with *Giardia* and *Cryptosporidium*, which increases the risk of infection by human contact. A summary of reports on pet infections in China and some foreign countries is shown in Table 1.

Second, *Giardia* and *Cryptosporidium* are highly contagious. A study by Hunter *et al.* (2009) has shown that when there is a problem with water safety, individuals will substitute emergency water (untreated or improperly treated water), which are of great risks to *Giardia* cysts and *Cryptosporidium* oocysts contaminations. Many studies have pointed out that the pathogenic doses of *Giardia* cysts and *Cryptosporidium* oocysts to humans are 10–100 live cysts and 1–10 live oocysts, respectively (Xu & Hu 2007). Under *Giardia* infection, patients can excrete as many as  $1 \times 10^{10}$  cysts per day, thus increasing the spread of the parasite in the environment and the risk of transmission.

Third, humans are highly susceptible to *Giardia* and *Cryptosporidium* infections. Except for children, the high-risk group includes the elderly, immunodeficient individuals, hospitalized individuals, travelers, gay men, people in post-disaster areas, and homeless people in the outbreak area (Escobedo *et al.* 2010).

Fourth, giardiasis and cryptosporidiosis are hard to cure. At present, there is no specific medicine for giardiasis and cryptosporidiosis (Chen *et al.* 2018). In immunodeficient patients, the death rate after protozoan infection is extremely high (Kotloff *et al.* 2013; Utami *et al.* 2020). A few studies have shown that children infected with cryptosporidiosis will have a short-term developmental delay with varying degrees of cognitive development impairment (Bushen *et al.* 2007).

**Table 1** | Statistics of *Giardia* and *Cryptosporidium* in pets in some areas at home and abroad

Parasite category	Country/region	Species	Infection rate	Human insect species can be infected in positive samples	References	
Cryptosporidium	China	Henan	Totoro	10.00%	<i>C. parvum</i>	Qi <i>et al.</i> (2015)
			Birds	8.10%	<i>C. meleagridis</i>	Qi <i>et al.</i> (2011)
		Hubei	Birds	20.20%	<i>C. meleagridis</i>	Liao (2019)
		Guangzhou	Dogs	3.20–6.90%	<i>C. parvum</i>	Liao <i>et al.</i> (2020), Zheng <i>et al.</i> (2019)
			Cats	6.20%	<i>C. felis</i> , <i>C. parvum</i>	Zheng <i>et al.</i> (2019)
		Sichuan	Dogs	4.30%	<i>C. canis</i>	Hu (2011)
		Heilongjiang	Dogs	2.20%	<i>C. canis</i>	Yang (2015)
			Cats	3.80%	<i>C. felis</i> , <i>C. parvum</i>	
		Anhui	Dogs	1.50%	<i>C. canis</i>	Gu <i>et al.</i> (2015)
		Zhejiang	Dogs	1.50%	<i>C. canis</i>	
		Shanghai	Dogs	8.00%	<i>C. canis</i>	Xu (2016)
			Cats	3.80%	<i>C. felis</i>	
		Xinjiang	Dogs	6.80%	<i>C. canis</i> , <i>C. parvum</i>	Zhang <i>et al.</i> (2017)
		Ethiopia	Cattle	7.80%	–	Wegayehu <i>et al.</i> (2013)
Giardia	China	Guangdong	Dogs	3.10–9.40%	<i>G. lamblia</i> (Aggregate A)	Xiao <i>et al.</i> (2013), Zheng <i>et al.</i> (2019)
				Cats	3.60%	<i>G. lamblia</i> (Aggregate A)
		Henan	Totoro	37.50%	–	Lu (2009)
		Heilongjiang	Cats	1.90%	–	Yang (2015)
			Dogs	4.50%	<i>G. lamblia</i> (Aggregate C)	
		Anhui	Dogs	3.20%	<i>G. lamblia</i> (Aggregate B & D)	Gu <i>et al.</i> (2015)
		Zhejiang	Dogs	3.20%	<i>G. lamblia</i> (Aggregate B & D)	
		Shanghai	Dogs	6.00%	<i>G. lamblia</i> (Aggregate A & B)	Xu (2016)
			Cats	5.60%	<i>G. lamblia</i> (Aggregate A & B)	
		Russia	Dogs	4.60%	–	Kurnosova <i>et al.</i> (2019)
			Cats	9.80%	–	
			Totoro	47.40%	–	
		Colombia	Dogs	47.00%	–	Hernández <i>et al.</i> (2021)
		Ethiopia	Cattle	2.30%	–	Wegayehu <i>et al.</i> (2013)

## The contamination status of *Giardia* and *Cryptosporidium*

### Contamination status of *Giardia* and *Cryptosporidium* in China

China first detected a case of *Cryptosporidium* in Nanjing in 1987. After that, *Giardia* and *Cryptosporidium* infections investigations have been carried out in many areas, specifically in Xuzhou, Anhui, Inner Mongolia, Fujian, Shandong, Hunan, and other provinces and cities (Cai 1995; Fu *et al.* 1998; Gao 2008). In 2005, an epidemiological survey of *Cryptosporidium* was conducted for the first time in Chinese rural populations in Jiangshan, Zhejiang Province. The results showed that the infection rate was as high as 56.72% (Shan 2007). In 2010, the survey of the contamination of *Giardia* and *Cryptosporidium* in drinking water and environmental water in Shanghai showed that oocysts were not detected in 156 water samples collected from 16 districts in Shanghai, including factory water, pipe network water, and community drinking water. Among the 70 water samples collected from the water supply plant, water source, Huangpu river, animal feed farm surrounding a river, sewage treatment plant effluent, and domestic sewage collected in 5 districts, the total detection rate of *Cryptosporidium* oocysts was 17.1%. The total detection rate of *Giardia* cysts was 20.0% (Zhang *et al.* 2010). In 2014, survey results showed that *Cryptosporidium* oocysts and *Giardia* cysts in the water source of the centralized water supply source in rural areas in southern China had a prevalence of 23.33% and 33.33%, respectively (Sun *et al.* 2014). In the same year, the survey results of *Giardia* and *Cryptosporidium* contamination in source and drinking water in typical areas of Jiangsu Province showed that 7 out of 222 samples were positive, of which the detection rates of *Cryptosporidium* and *Giardia* were 0.5% and 2.7%, respectively (Zheng *et al.* 2014). In 2019, Wang Lu *et al.* summarized *Cryptosporidium* studies and analyzed the map of *Cryptosporidium* infection in China using GIS technology. Results showed that infected people were concentrated in Jiangsu, Anhui, Shandong, Henan, and other places (Wang *et al.* 2019a, 2019b). In addition, a large survey indicated that the water environment of some provinces in China (Liaoning, Northeast, Qinghai, Shaanxi, and Inner Mongolia) is seriously contaminated by *Giardia* cysts and *Cryptosporidium* oocysts. Even the water environment of first-tier cities and key projects (Shanghai, Three Gorges Reservoir) had exceeded the standard contamination rate of the two protozoa (Hou *et al.* 2011; Liu *et al.* 2011; Wang *et al.* 2012, 2019a, 2019b; Wang 2013; Xiao *et al.* 2013).

### Contamination status of *Giardia* and *Cryptosporidium* in other countries

Most of the research on *Giardia* and *Cryptosporidium* infections in countries other than China is concentrated in Asia and Africa. The backward health system of most countries in Asia and Africa is the main reason for the serious situation of *Giardia* and *Cryptosporidium* infections. Several studies have shown that children under five years of age in developing countries with poor basic health conditions account for cryptosporidiosis to 30–50% of overall child mortality (Checkley *et al.* 2015; Platts-Mills *et al.* 2015). Because children and immunodeficient patients are high-risk and susceptible groups, most of the researches centered on these two groups. A summary of related studies on children and immunodeficient patients in Asian and African countries infected with *Giardia* and *Cryptosporidium* is shown in Table 2.

Infections with *Giardia* and *Cryptosporidium* are common in children in Southeast Asia. For instance, Thailand's large water systems provide a good environment for transmission. Recent research reports showed that the infection rate of *Cryptosporidium* in children in Thailand was as high as 15% (Choekphaibulkit *et al.* 2001), while the infection rate of *Giardia* was as high as 10.2% (Sagnuankiat *et al.* 2014). At the same time, the study also pointed out that the infection rate of *Cryptosporidium* in children with AIDS was 33%, twice that of normal children and 11.5% higher than that of normal AIDS patients. Children in Cambodia, Malaysia, Myanmar, the Philippines, Laos, and Indonesia are also at risk of being infected by the two parasitic protozoa. In South Asia, India and Bangladesh have the most serious infections in children, especially in terms of *Cryptosporidium* infection in which the prevalence was recorded at 27.4 and 64%, respectively (Mahmoudi *et al.* 2017; Steiner *et al.* 2018). In Bangladesh, the reported *Giardia* infection rate was as high as 40% (Berendes *et al.* 2020). West Asia is the hardest-hit area in Asia, with most infections occurring in children. For example, in a study in Kuwait, the infection rate among the surveyed groups was as high as 94%.

In Africa, the child infection rates in Ethiopia, Kenya, and Egypt were higher than 20%, far higher than that of other countries, with both ordinary AIDS and child AIDS patients showing the highest infection rates. However, there was no obvious difference in infection rate between the two AIDS patient types, indicating that these patients could be both infected by *Giardia* and *Cryptosporidium*. It is worth noting that the infection rate of *Giardia* and *Cryptosporidium* is closely related to the sanitary conditions and water environment of these countries.

Compared with Asian and African regions, other countries have fewer studies on the infection status of *Giardia* and *Cryptosporidium*. Results of the epidemiological data of some European and American countries are shown in Table 3. There are

**Table 2** | Epidemiological statistics of children and immunodeficiency patients infected with the *Giardia* and *Cryptosporidium* in some Asian and African countries

Country	Infected people	Infection rate	Infected species	References	
Southeast Asia	Cambodia	Child	7.40%	<i>Cryptosporidium</i> ( <i>C. hominis</i> , <i>C. parvum</i> )	Arthur <i>et al.</i> (1992)
		AIDS patient	45%	<i>Cryptosporidium</i> ( <i>C. hominis</i> , <i>C. parvum</i> )	Chhin <i>et al.</i> (2006)
	Malaysia	Child	0.40–10.60%/2.60%	<i>Cryptosporidium</i> / <i>Giardia</i> (uncategorized)	Lai (1992), Mahmoudi <i>et al.</i> (2017), Lim <i>et al.</i> (2008)
		Child(CWD)	4.60%	<i>Cryptosporidium</i> (uncategorized)	Latif & Rossle (2015)
		AIDS patient	3–64%	<i>Cryptosporidium</i> (uncategorized)	Lim <i>et al.</i> (2005), Zaidah <i>et al.</i> (2008), Lim <i>et al.</i> (2011)
		Child	3.40%	<i>Cryptosporidium</i> (uncategorized)	Aye <i>et al.</i> (1994)
	Philippines	Child	2.50–2.90%	<i>Cryptosporidium</i> (uncategorized)	Mahmoudi <i>et al.</i> (2017)
	Thailand	Cancer patient	28.30%	<i>Cryptosporidium</i> ( <i>C. hominis</i> , <i>C. parvum</i> )	Rivera <i>et al.</i> (2005)
		Child	15%/0.80–10.20%	<i>Cryptosporidium</i> / <i>Giardia</i> (uncategorized)	Chokephaibulkit <i>et al.</i> (2001), Sagnuankiat <i>et al.</i> (2014), Assavapongpaiboon <i>et al.</i> (2018), Sanprasert <i>et al.</i> (2016)
		Child(HIV)	33%	<i>Cryptosporidium</i> (uncategorized)	Chokephaibulkit <i>et al.</i> (2001)
AIDS patient		11.50%	<i>Cryptosporidium</i> ( <i>C. hominis</i> , <i>C. meleagridis</i> , <i>C. parvum</i> , <i>C. felis</i> , <i>C. canis</i> )	Pinlaor <i>et al.</i> (2005)	
Laos	AIDS patient	13.90%	<i>Cryptosporidium</i> (uncategorized)	Paboriboune <i>et al.</i> (2014)	
Indonesia	AIDS patient	4.90%	<i>Cryptosporidium</i> (uncategorized)	Kurniawan <i>et al.</i> (2009)	
South Asia	India	Child	1.30–27.40%	<i>Cryptosporidium</i> ( <i>C. hominis</i> , <i>C. parvum</i> )	Mahmoudi <i>et al.</i> (2017)
		AIDS patient	2–77%	<i>Cryptosporidium</i> ( <i>C. parvum</i> )	
	Sri Lanka	Child(CWD)	5.70%	<i>Cryptosporidium</i> (uncategorized)	Sirisena <i>et al.</i> (2014)
	Bangladesh	Child	44–64%/40%	<i>Cryptosporidium</i> ( <i>C. hominis</i> , <i>C. parvum</i> , <i>C. meleagridis</i> )/ <i>Giardia</i> (uncategorized)	Steiner <i>et al.</i> (2018), Berendes <i>et al.</i> (2020)
		AIDS patient	47.10%	<i>Cryptosporidium</i> (uncategorized)	Mahmoudi <i>et al.</i> (2017)
	Nepal	Child	1–16%	<i>Cryptosporidium</i> (uncategorized)	Mahmoudi <i>et al.</i> (2017), Paudyal <i>et al.</i> (2013)
		AIDS patient	11–35.70%	<i>Cryptosporidium</i> (uncategorized)	
Pakistan	Cancer, Diabetes, Dialysis	40%	<i>Cryptosporidium</i> (uncategorized)	Baqai <i>et al.</i> (2005)	
West Asia	Iran	Child	3.30–10.30%	<i>Cryptosporidium</i> (uncategorized)	Mahmoudi <i>et al.</i> (2017)
		AIDS patient	2–7%	<i>Cryptosporidium</i> ( <i>C. hominis</i> , <i>C. parvum</i> )	
	Iraq	Child(CWD)	1.50–7%	<i>Cryptosporidium</i> ( <i>C. hominis</i> , <i>C. parvum</i> )	Meamar <i>et al.</i> (2006)
		Child	8.56%	<i>Cryptosporidium</i> (uncategorized)	Latif & Rossle (2015)
	Israel	Child	6–33.83%	<i>Cryptosporidium</i> ( <i>C. parvum</i> )	Azeez & Alsakee (2017), Rahi <i>et al.</i> (2013)
		Child	1.30–31.90%	<i>Cryptosporidium</i> (uncategorized)	Mahmoudi <i>et al.</i> (2017)
	Jordan	Child	42.60%	<i>Giardia</i> (uncategorized)	Ammoura (2010)
		Child(CWD)	37.30%	<i>Cryptosporidium</i> (uncategorized)	Latif & Rossle (2015)
		AIDS patient	6%	<i>Cryptosporidium</i> (uncategorized)	
	Kuwait	Hemodialysis patients	11%	<i>Cryptosporidium</i> (uncategorized)	Zueter <i>et al.</i> (2019)
Child		3.40–94%	<i>Cryptosporidium</i> ( <i>C. hominis</i> , <i>C. parvum</i> )	Ahmed & Karanis (2020)	
Lebanon		Child	10.40%	<i>Cryptosporidium</i> (uncategorized)	Osman <i>et al.</i> (2018)
Palestine		Child	11.60%	<i>Cryptosporidium</i> (uncategorized)	

(Continued.)

**Table 2** | Continued

Country	Infected people	Infection rate	Infected species	References	
Yemen	Child(CWD)	1%	<i>Giardia</i> (uncategorized)	Abu-Elamreen <i>et al.</i> (2008)	
	Child	43.70%	<i>Cryptosporidium</i> (uncategorized)	Mahmoudi <i>et al.</i> (2017)	
	Cancer patient	30.1%/18%	<i>Cryptosporidium/Giardia</i> (uncategorized)	Al-Qobati <i>et al.</i> (2012)	
Saudi Arabia	Child	1.70–11%	<i>Cryptosporidium</i> ( <i>C. hominis</i> , <i>C. parvum</i> )	El-Malky <i>et al.</i> (2018), Shalaby <i>et al.</i> (2014)	
	AIDS patient	8.10–69.70%	<i>Cryptosporidium</i> ( <i>C. hominis</i> , <i>C. parvum</i> , <i>C. meleagridis</i> , <i>C. muris</i> )	Al-Megrin 2010, Al-Brikan <i>et al.</i> (2008)	
Africa	Nigeria	Child	19.40%	<i>Cryptosporidium</i> ( <i>C. hominis</i> , <i>C. parvum</i> )	Molloy <i>et al.</i> (2010)
	Egypt	Child	35%	<i>Cryptosporidium</i> ( <i>C. hominis</i> , <i>C. parvum</i> )	Gharieb <i>et al.</i> (2018)
	Ethiopia	Child	4.60%/55%	<i>Cryptosporidium/Giardia</i> (uncategorized)	Wegayehu <i>et al.</i> (2016)
		Child(HIV)	9.60%	<i>Cryptosporidium</i> (uncategorized)	Gebre <i>et al.</i> (2019)
	Botswana	Child	10%/7%	<i>Cryptosporidium/Giardia</i> (uncategorized)	Alexander <i>et al.</i> (2012)
	Kenya	Child	45.20%	<i>Cryptosporidium</i> (uncategorized)	Mutai <i>et al.</i> (2020)
	Tanzania	Child	6%	<i>Cryptosporidium</i> (uncategorized)	Korpe <i>et al.</i> (2018)
	Gabon	Child	13.30%/15.60%	<i>Cryptosporidium/Giardia</i> (uncategorized)	Bouyou-Akotet <i>et al.</i> (2015)

Note: Child(CWD): Children with diarrhea; Child(HIV): Children with AIDS.

**Table 3** | Epidemiological statistics of population infections with *Giardia* and *Cryptosporidium* in some European and American countries

Country	Infected people	Infection rate	Infected species	References
United States	Patients with diarrhea	2.99/10 <sup>5</sup>	<i>Cryptosporidium</i> (uncategorized)	Alleyne <i>et al.</i> (2020)
Croatia	Food industry personnel	0.07%	<i>Giardia</i> (uncategorized)	Plutzer <i>et al.</i> (2018)
	Symptoms of bowel disease	0.24%	<i>Giardia</i> (uncategorized)	
Czech Republic	Symptoms of bowel disease	0.52%	<i>Giardia</i> (uncategorized)	
Estonia	Patients with diarrhea	(0.05/10 <sup>5</sup> )/(18.28/10 <sup>5</sup> )	<i>Cryptosporidium</i> / <i>Giardia</i> (uncategorized)	
Hungary	–	0.03%/1.2%	<i>Cryptosporidium</i> / <i>Giardia</i> (uncategorized)	
Latvia	Patients with diarrhea	(0.29/10 <sup>5</sup> )/ (2.48/10 <sup>5</sup> )	<i>Cryptosporidium</i> / <i>Giardia</i> (uncategorized)	
Poland	–	(0.006/10 <sup>5</sup> )/(5.43/10 <sup>5</sup> )	<i>Cryptosporidium</i> / <i>Giardia</i> (uncategorized)	
Romania	–	(0.01/10 <sup>5</sup> )	<i>Cryptosporidium</i> (uncategorized)	
Slovenia	Patients with diarrhea	1.53%	<i>Cryptosporidium</i> (uncategorized)	
Bosnia and Herzegovina	Symptoms of bowel disease	0.96%	<i>Giardia</i> (uncategorized)	
	Patients with diarrhea	9.09%	<i>Giardia</i> (uncategorized)	
Serbia	Food industry personnel	0.28%	<i>Giardia</i> (uncategorized)	
Slovakia	Child	6.30%	<i>Giardia</i> (Aggregate A II, B, F)	Pipiková <i>et al.</i> (2018)
Austria	Child	1.50%	<i>Cryptosporidium</i> (uncategorized)	Joachim (2004)
Italy	Child(CWD)	7.20%	<i>Cryptosporidium</i> (uncategorized)	
Switzerland	Symptoms of bowel disease	0.20%	<i>Cryptosporidium</i> (uncategorized)	
	Child(CWD)	5.50%	<i>Cryptosporidium</i> (uncategorized)	
New Zealand	–	12.90/10 <sup>5</sup>	<i>Cryptosporidium</i> ( <i>C.hominis</i> , <i>C.parvum</i> )	Pipiková <i>et al.</i> (2018)
Brazil	Child	13.70–18%	<i>Giardia</i> (uncategorized)	Prado <i>et al.</i> (2003), Teixeira <i>et al.</i> (2007)

Note: Child(CWD): Children with diarrhea; Child(HIV): Children with AIDS; -: The survey population is not clearly indicated.

little data on human infections, and most of the relevant research directions are concentrated in agriculture, with research focusing mainly on animal or livestock scenario. In the case of human infection, the infection rate of *Giardia* and *Cryptosporidium* in European and American countries is extremely low. In Slovakia, the relatively higher infection rate (6.3%) can be explained by the survey participants in poor areas in Northeast Slovakia. This place lacks water supply and sewage treatment systems and has low sanitation, education, and health awareness.

In South America, Brazil has the highest infection rate (13.7–18%). The main reason is that the survey was performed in 29 slum areas in the city of Zfifola in southeastern Brazil. Secondly, Brazil has rich water resources, with freshwater resources accounting for about 13% of the world's total. In addition, the number of rivers entering the sea accounts for about 20% of the total rivers in the world. Thus, the large water system provides unique environmental conditions for the spread of *Giardia* and *Cryptosporidium*.

Tables 2 and 3 show that research data from Asia and Africa are relatively rich. Data suggest that *Giardia* and *Cryptosporidium* contamination improvements have been slow across time, and there has been a sharp increase in recent years. In some regions, such as India, Bangladesh, Iraq, and Pakistan, the infection rate in children is relatively high, directly related to the country's health system conditions. Although few relevant studies in Europe and America, the infection rate is far lower than

that of Asia and Africa. However, the data of Europe and America have not shown a steady decline. In New Zealand, human *Cryptosporidium* infection rates in 2001 and 2002 revealed a total of 33.2 cases per 100,000 and 26.1 cases per 100,000, respectively. In addition, infection rates in 2013 and 2014 were 30.3 cases per 100,000 and 12.9 cases per 100,000, respectively (Garcia-R *et al.* 2020). In Stockholm, Sweden, there were more than 300 cases of cryptosporidiosis between October and November 2019. In France, the water supply system was polluted by heavy rain, causing 92 cases of cryptosporidiosis in October 2019. Therefore, European and American countries maintain a low infection rate while steadily reducing the infection rate.

## DETECTION METHOD OF *GIARDIA* AND *CRYPTOSPORIDIUM* IN WATER

The earliest monitoring method for *Giardia* and *Cryptosporidium* was the Information Collection Rule (ICR) protozoan method proposed by U.S. Environmental Protection Agency (USEPA). However, this method is prone to sample loss in the elution, concentration, and purification stages. Also, it needs experienced staff to operate the fluorescence microscope, which could lead to relatively large subjective detection results. Moreover, this method cannot evaluate the activity of *Giardia* and *Cryptosporidium* and cannot further distinguish the protozoan species (Zong *et al.* 2005).

Since 1996, USEPA began to use immunomagnetic separation technology to optimize the detection methods of *Giardia* and *Cryptosporidium*. The first release of the EPA1622 method for the detection of *Cryptosporidium* was in 1999. Then, the EPA1623 method that can detect two types of protozoa was released. This method benefits from filter cartridge filtration, immunomagnetic bead separation, and immunofluorescence microscopy detection, improving this method's recovery rate and accuracy. In addition, with the help of DAPI staining and differential interference microscope, the internal structure can now be observed to confirm the presence of oocysts and cysts (USEPA, Method 1623 2001; Sun 2007). However, this method has several drawbacks. First, the binding site of the immunofluorescence antibody is located on the outer surface of oocysts and cysts, and the cyst wall after decapsulation will still be detected, which may increase the detection rate. Second, chlorine disinfection, increasing age of parasites, and environmental changes are likely to cause inactivation of binding sites and reduce the detection rate. Third, the method still cannot clearly distinguish between activity and protozoan species. However, some researches showing the optimization of the EPA1623 process could improve the system. For example, Ye *et al.* optimized the immunomagnetic separation stage, which replaced the *Filta-Max Xpress* filter element with a filter membrane for rapid filtration. It also replaced the air compressor with an ultrasonic elution filter membrane to quickly elute the filter element. After optimization, the recovery rate of *Giardia* and *Cryptosporidium* standard addition is much higher than the recovery rate obtained by operating following the national standard (Ye *et al.* 2017).

Enzyme-linked immunosorbent assay (ELISA) is an immunological detection technique. Because of its good specificity and sensitivity, ELISA has been used often in epidemiological studies for *Giardia* and *Cryptosporidium* detection (Li 2015). Although it can be qualitatively or quantitatively measured, the operation is complicated in quantitative measurement, and many factors affect the reaction. Moreover, each group of experiments needs to measure multiple concentrations to draw a positive standard curve.

Due to the limitations of immunological technology, molecular technology has been rapidly developed and showed great potential in detecting pathogenic protozoa. Polymerase chain reaction (PCR) is a molecular biology detection method developed in the mid-1980s (Lu 2016). However, this method has strict requirements on the test environment and is often sensitive to DNA contamination. Therefore, it can be combined with immunomagnetic separation technology (IMS-PCR) to improve PCR specificity (Giovanni *et al.* 1999; Rimhanen-Finne *et al.* 2002). Other molecular biology techniques include the Nested PCR, reverse transcription-polymerase chain reaction (RT-PCR), Quantitative Real-time PCR (qRT-PCR), reverse transcription PCR (RT-PCR), multiplex fluorescent PCR, loop-mediated isothermal amplification (LAMP), and PCR-restriction fragment polymorphism analysis (PCR-PFLP).

The Nested PCR uses two pairs of PCR primers based on traditional PCR, making it highly specific and sensitive. This technique is widely used in the detection of pathogenic microorganisms (Wang *et al.* 2014). For example, Meng *et al.* (2011) used nested PCR to detect water-borne *Cryptosporidium* in Xinjiang, China. Li *et al.* (2010) also established a nested PCR method to detect cryptosporidiosis.

Quantitative Real-time PCR (qRT-PCR) is the real-time monitoring of the entire PCR process through fluorescent signals in the traditional PCR amplification process. Kumar *et al.* (2016) used Real-time PCR to detect *Cryptosporidium* in the water environment in Malaysia, the Philippines, Thailand, and Vietnam in Southeast Asia. Results showed that Real-time PCR is



sensitive and specific in the quantitative detection of *Cryptosporidium*. But this method still cannot distinguish and detect the activity of live oocysts. Reverse transcription PCR (RT-PCR) is based on mRNA for detection. The above PCR-derived methods are based on DNA for detection. DNA can still be stored intact for a long time after the oocysts or cysts die, so the detection rate is relatively high. Only living cells can produce mRNA. This method overcomes the shortcomings of other DNA-based molecular biology detection methods.

Traditional PCR technology can only detect one kind of pathogenic microorganism in each PCR reaction tube, which has low efficiency and high cost. To address such shortcomings, Chamberlain *et al.* (1988) first proposed the concept of multiplex PCR (multiplex-PCR) in their research in 1988. This method can detect multiple pathogenic microorganisms at the same time, saving workforce and material resources. For example, Moniot *et al.* (2020) established a simultaneous detection method for intestinal microsporidia and *Cryptosporidium* using multiplex PCR technology. PCR-restriction fragment polymorphism analysis (PCR-RFLP) is a technique that combines PCR amplification and restriction enzyme digestion to detect polymorphisms. It uses a specific restriction enzyme to cut the amplified product and run on gel electrophoresis. The technique is simple and can analyze samples in a short time. Therefore, it is suitable for biotyping and identifying multiple protozoan species, such as *Cryptosporidium* and *Giardia*. Rafiei *et al.* (2014) used PCR-RFLP technology to identify the types of *Cryptosporidium* infecting Iranians. The study identified three genotypes *C. parvum*, *C. hominis*, and *C. meleagridis*, of which *C. meleagridis* was the first case in Iran.

Loop-mediated isothermal amplification technology (LAMP) is a relatively new technology. In 2000, a Japanese scholar Notomi proposed a constant temperature accounting amplification technology suitable for genetic diagnosis. It uses strand displacement DNA polymerase to amplify a large amount in a short time under a constant temperature environment, and the product produces a large amount of magnesium pyrophosphate white precipitate. As a result, the presence of the target gene can be observed by naked eyes. Karanis *et al.* (2007) applied this technology to detect *Cryptosporidium* for the first time due to its high specificity, simple operation, and low requirements for equipment (PCR requires expensive equipment, LAMP only needs a water bath or incubator). Thus, this method is of great significance at the practical application level.

The above technology is the basic detection technology for *Giardia* and *Cryptosporidium* (as shown in Table 4). According to the characteristics of each detection method, combined with the actual conditions and requirements of the test, researchers from various countries often use a combination of multiple detection methods, thus proposing a variety of effective combinations. For example, Hallier-Soulier & Guillot (2000) combined immunomagnetic separation and PCR technology to detect the number of *Cryptosporidium* oocysts in rivers. Fontaine & Guillot (2003) used immunomagnetic separation technology combined with real-time PCR technology to detect *Cryptosporidium* oocysts in tap water and Seine water in France. Liao *et al.* (2014) combined real-time fluorescent PCR and multiplex PCR to establish a dual real-time fluorescent PCR detection method for *Cryptosporidium* and *Giardia*, creating a much faster and more accurate system.

## CONTROL OF GIARDIA AND CRYPTOSPORIDIUM BY DRINKING WATER TREATMENT PROCESS

Compared with *Giardia* cysts, *Cryptosporidium* oocysts are smaller, have a lower pathogenic dose, and are more resistant to disinfectants (Betancourt & Rose 2004). Researchers believe that when *Cryptosporidium* oocysts are removed from the water, it could also lead to the removal of *Giardia* cysts. Hence, *Cryptosporidium* oocysts are generally used as control targets, which is why there are more studies on *Cryptosporidium* than *Giardia*. The removal rate of *Giardia* and *Cryptosporidium* is often expressed in logarithm. For example, the logarithmic removal rate of 2.0 log corresponds to an inactivation rate of 99%. These two types of protozoa are ubiquitous in the natural water environment. Thus, certain control indicators need to be reached in each stage of water treatment to ensure that subsequent processes can run well and ensure that the final effluent reaches the standard.

### Coagulation-sedimentation-filtration

The outer surfaces of *Cryptosporidium* oocysts and *Giardia* cysts are negatively charged, similar to other low-density negatively charged colloids. As a result, a coagulation-sedimentation process can remove them to a certain extent. However, at the same time, because the impurity particles have a protective effect on pathogenic microorganisms, it affects the mechanism of disinfectants and the removal of pathogens (Yan & Chen 2004). Therefore, what kind of coagulant to choose, what type of filtration method, and how much effluent turbidity is controlled have become the main research directions.

The USEPA promulgated the 'Interim Enhanced Surface Water Treatment Rule' (IESWTR) on December 6, 1998. The regulation's core is that the maximum pollutant level indicator (MCLG) of *Cryptosporidium* must be set to zero. Furthermore,

**Table 4** | List of *Cryptosporidium* and *Giardia* detection methods

Detection method	Introduction	Advantage	Disadvantage
ICR	The earliest method used to monitor <i>Giardia</i> and <i>Cryptosporidium</i> in the water environment	Qualitatively analysis	<ol style="list-style-type: none"> <li>1. Large losses in the elution, concentration and purification stages can easily lead to low detection rates</li> <li>2. High professional requirements and strong subjectivity</li> <li>3. Unable to detect activity and distinguish types</li> </ol>
EPA1623	The globally recognized detection method for <i>Giardia</i> and <i>Cryptosporidium</i> , which has confirmed the existence of oocysts and spore cysts, and has now been widely used	<ol style="list-style-type: none"> <li>1. Higher detection rate</li> <li>2. Simple operation</li> <li>3. Observe the internal structure</li> </ol>	<ol style="list-style-type: none"> <li>1. More unstable factors</li> <li>2. Unable to detect activity and distinguish protozoan species</li> </ol>
ELISA	An immunological technique widely used in epidemiological testing	<ol style="list-style-type: none"> <li>1. Both quantitatively and qualitatively analysis</li> <li>2. High specificity and sensitivity</li> </ol>	<ol style="list-style-type: none"> <li>1. The operation is complicated in quantitative determination, and there are many influencing factors</li> </ol>
PCR	The first molecular biology detection method applied to the detection of <i>Giardia</i> and <i>Cryptosporidium</i>	Detectable genotype	<ol style="list-style-type: none"> <li>1. Unable to determine the activity of oocysts and cysts</li> <li>2. Poor sensitivity and high DNA purity requirements</li> </ol>
Nested PCR	PCR-derived detection method is one of the most effective molecular biology detections	<ol style="list-style-type: none"> <li>1. High specificity and sensitivity</li> <li>2. Detectable genotype</li> </ol>	<ol style="list-style-type: none"> <li>1. Difficult to quantify</li> <li>2. Cross-infection may occur during secondary amplification</li> <li>3. Unable to determine the activity of oocysts and cysts</li> </ol>
qRT-PCR	A detection method for adding fluorescent chemicals in DNA amplification to monitor the total amount of products in each PCR process	<ol style="list-style-type: none"> <li>1. Obvious data</li> <li>2. Quantitative analysis</li> </ol>	<ol style="list-style-type: none"> <li>1. Expensive</li> <li>2. Only suitable for detecting specific targets</li> <li>3. Unable to determine the activity of oocysts and cysts</li> </ol>
RT-PCR	MRNA-based detection method	<ol style="list-style-type: none"> <li>1. The activity of oocysts and cysts can be determined</li> <li>2. High sensitivity</li> </ol>	<ol style="list-style-type: none"> <li>1. A single reverse transcription can only amplify one gene</li> <li>2. Strict operating environment requirements</li> <li>3. RNA preservation is difficult</li> </ol>
multiplex PCR	One of the molecular biology detection methods in which multiple primers can be amplified in the same reaction system	<ol style="list-style-type: none"> <li>1. High efficiency</li> <li>2. Simple operation</li> <li>3. Low experiment cost</li> </ol>	<ol style="list-style-type: none"> <li>1. The pairing and competitive amplification among multiple primers affect the multiplex PCR amplification</li> <li>2. High quality requirements for DNA extraction</li> <li>3. Unable to determine the activity of oocysts and cysts</li> </ol>

(Continued.)

Table 4 | Continued

Detection method	Introduction	Advantage	Disadvantage
LAMP	The detection technology proposed by Japanese scholars has been applied to the detection of SARS, avian influenza, HIV and other diseases	<ol style="list-style-type: none"> <li>1. Simple operation</li> <li>2. High specificity and sensitivity</li> <li>3. Low requirements for experimental environment and testing costs</li> </ol>	<ol style="list-style-type: none"> <li>1. Unable to quantitatively study</li> <li>2. Unable to distinguish sample species</li> </ol>
PCR-RFLP	A method that can accurately identify the genotypes and species of <i>Giardia</i> and <i>Cryptosporidium</i>	<ol style="list-style-type: none"> <li>1. The operation is simple, fast and highly automated</li> <li>2. Low amount of DNA required</li> <li>3. Accurately perform typing research on samples</li> </ol>	<ol style="list-style-type: none"> <li>1. High requirements for digestion conditions</li> <li>2. Inability to distinguish heterozygotes</li> <li>3. Difficulty in distinguishing alleles</li> </ol>

it is required to use only the filtration process without disinfection process, and the removal rate of *Cryptosporidium* oocysts needs to reach 2.0 log. When the content of *Cryptosporidium* oocysts in the raw water ( $C_{(Cry)}$ ) is greater than 0.075 oocysts per liter (oocysts/L), the oocyst removal rate must reach 3.0 log. Meanwhile, when  $C_{(Cry)}$  is greater than one oocysts/L, the removal rate must reach 4.0 log. When  $C_{(Cry)}$  is greater than three oocysts/L, the removal rate needs to reach 4.5 log (USEPA IESWTR).

States *et al.* (2002) pointed out that the three coagulants of ferric chloride, alum, and polyaluminum chloride can effectively remove *Cryptosporidium* oocysts based on the effect of pH on the removal of *Cryptosporidium* oocysts and Total Organic Carbon (TOC). Here, the removal rate is up to 4.3 log, and pH does not affect the removal of *Cryptosporidium*. Alum is the most commonly used water treatment coagulant in Australia, with a few studies showing its good removal effect on oocysts. The removal rate of *Cryptosporidium* oocysts is greater than 1.0 log when alum dosage is at 40–100 mg/L (Keegan *et al.* 2008). Cornwell *et al.* (2003) and Logsdon & Johnson (2010) showed that the use of the lime softening method in water treatment could greatly reduce the content of *Giardia* and *Cryptosporidium* in water (2.5–3.5 log). Ongerth & Pecoraro (1995) used a designed filter (anthracite + silica sand + garnet) to explore the ability of the probability pool to treat the two protozoa. They found that when the influent turbidity is 0.38 NTU and the effluent turbidity is 0.03 NTU at room temperature, the removal rates of *Cryptosporidium* and *Giardia* can reach 3.1 log and 3.6 log, respectively.

Based on many studies, the three conventional water treatment processes of coagulation-sedimentation-filtration can effectively remove *Giardia* and *Cryptosporidium*. In contrast to the disinfection method, these two types of protozoa are removed as suspended particles. However, filtered water effluent needs that the removal rate reaches at least 2.0 log. Therefore, as the concentration of oocysts and cysts in the raw water increases, the process needs to be changed to achieve a higher removal rate.

## Disinfection

Disinfection is the most critical link in the water treatment process. Choosing the appropriate disinfectant and working conditions is key to ensuring the safety of the effluent. *Giardia* and *Cryptosporidium*, as a special type of protozoa in the water environment, are important indices to evaluate the disinfection effect and the quality of the effluent. Researchers have carried out many experimental studies on removing *Giardia* and *Cryptosporidium* to provide theoretical support for engineering practice, particularly on the impact of different disinfection methods.

## Chlorine disinfection

In 1981, Rice *et al.* (1982) first applied chlorine disinfection to the inactivation of *Giardia* cysts. The research object was the cysts excreted by *Giardia* cyst carriers, and the chlorine ( $Cl_2$ ) concentration used was 2.5 mg/L. Although this is medical research, it is a milestone for applying disinfection technology to inactivate *Giardia* and *Cryptosporidium*. The  $Cl_2$  has a

slight killing effect on *Giardia* and *Cryptosporidium*. The concentration of Cl<sub>2</sub> added to the water plant cannot completely kill the protozoa cysts and oocysts. If the inactivation rate reaches 99%, the CT value needs to be 7,200 mg.min/L. Moreover, because Cl<sub>2</sub> is used for drinking water disinfection, there are many hidden health and safety hazards, so there is less research on inactivating *Giardia* and *Cryptosporidium* using Cl<sub>2</sub> disinfectant. Compared to Cl<sub>2</sub>, chlorine dioxide (ClO<sub>2</sub>) has a stronger killing effect. [Ran et al. \(2011\)](#) explored the influencing factors of ClO<sub>2</sub> in the inactivation of *Cryptosporidium* and found that the best disinfection effect (inactivation rate is greater than 99%) is pH = 7, T = 25 °C, NTU = 1, C<sub>(ClO<sub>2</sub>)</sub> = 3 mg/L, and t (contact time) = 120 min. Moreover, turbidity was revealed as the main influencing factor, of which the higher the turbidity, the worse the inactivation effect. [Clark et al. \(2003\)](#) optimized the CT value equation for ClO<sub>2</sub> inactivation of *Cryptosporidium* and proposed a CT value equation to ensure drinking water safety effectively. The data given by the current specifications are also instructive for engineering practice. The USEPA LT2ESWTR: Toolbox Guidance Manual clearly shows the relationship between the inactivation rate of *Cryptosporidium*, temperature, and the CT value of ClO<sub>2</sub>, as shown in [Table 5](#).

### Ozone disinfection

[Yu et al. \(2008\)](#) pointed out that when the ozone dosage is 1 mg/L and the contact time is 5–10 min, the inactivation rate of *Cryptosporidium* oocysts can reach 1.0 log. [Cho & Yoon \(2007\)](#) showed that under the conditions of pH 5.6, 7.1, and 8.2, the inactivation rate of *Cryptosporidium* oocysts was 3.25, 3.10, and 2.78 mg/(L.min). [Sivaganesan & Mariñas \(2005\)](#) established a model based on the first-order reaction kinetics that can predict and calculate the minimum CT value with a confidence interval under the conditions of a given temperature and target survival rate. This model can provide technical and theoretical support for engineering practice. The USEPA LT2ESWTR: Toolbox Guidance Manual has developed a comparison table of CT values for inactivating *Cryptosporidium* oocysts with ozone, as shown in [Table 6](#). The CT value calculation formulas suitable for inactivating *Cryptosporidium* (1) and *Giardia* (2) in water treatment are also given.

$$\text{Cryptosporidium Log Credit} = 0.0397 \times (1.09757)^{\text{Temp}} \times \text{CT} \quad (1)$$

$$\text{Giardia Log Credit} = 1.0380 \times (1.0741)^{\text{Temp}} \times \text{CT} \quad (2)$$

### UV disinfection

The chemical reagents used in the disinfection stage of water treatment often produce disinfection by-products and bring potential harm to human health. Ultraviolet disinfection is gradually applied to the water treatment industry because of its broad-spectrum antibacterial properties and no by-product formation. Many scholars began to study the effect of ultraviolet rays on the removal of *Giardia* and *Cryptosporidium*. [King et al. \(2008\)](#) studied the impact of tap water and environmental water on *C. parvum* after being irradiated by the sun. Results showed that solar radiation could reduce the infectivity of *C. parvum* oocysts. [Soliman et al. \(2018\)](#) studied the inactivation of *Cryptosporidium* oocysts by solar ultraviolet and artificial ultraviolet radiation. The experimental results indicate that *Cryptosporidium* does not have the ability to infect mice again after 4 hours of artificial ultraviolet radiation of 10 mJ/cm<sup>2</sup> or natural sunlight of 8 hours ultraviolet radiation. This conclusion provides a simple, convenient, and economical method to inactivate *Giardia* cysts and *Cryptosporidium* oocysts. [Entrala et al.](#)

**Table 5** | The CT value of inactivated *cryptosporidium* by ClO<sub>2</sub> (mg-min/L)

Inactivation rate Log	Water temperature, (°C)										
	<=0.5	1	2	3	5	7	10	15	20	25	30
0.25	159	153	140	128	107	90	69	45	29	19	12
0.5	319	305	279	256	214	180	138	89	58	38	24
1.0	637	610	558	511	429	360	277	179	116	75	49
1.5	956	915	838	767	643	539	415	268	174	113	73
2.0	1,275	1,220	1,117	1,023	858	719	553	357	232	150	98
2.5	1,594	1,525	1,396	1,278	1,072	899	691	447	289	188	122
3.0	1,912	1,830	1,675	1,534	1,286	1,079	830	536	347	226	147

**Table 6** | The CT value of inactivated *cryptosporidium* by ozone (mg-min/L)

Inactivation rate Log	Water temperature, (°C)										
	<=0.5	1	2	3	5	7	10	15	20	25	>30
0.25	6.0	5.8	5.2	4.8	4.0	3.3	2.5	1.6	1.0	0.6	0.39
0.5	12	12	10	9.5	7.9	6.5	4.9	3.1	2.0	1.2	0.78
1.0	24	23	21	19	16	13	9.9	6.2	3.9	2.5	1.6
1.5	36	35	31	29	24	20	15	9.3	5.9	3.7	2.4
2.0	48	46	42	38	32	26	20	12	7.8	4.9	3.1
2.5	60	58	52	48	40	33	25	16	9.8	6.2	3.9
3.0	72	69	63	57	47	39	30	19	12	7.4	4.7

(2007) studied the effect of ultraviolet disinfection on the removal of *Cryptosporidium* oocysts by designing a medium-pressure ultraviolet reactor and a low-pressure reactor. Results showed that the flow rates in the two types of reactors are 15 m<sup>3</sup>/h and 42 m<sup>3</sup>/h, respectively. The inactivation rate of *Cryptosporidium* exposed to an effective UV dose of 400 J/m<sup>2</sup> can reach 4.92 log. The U.S. and German guidelines for UV disinfection are widely recognized internationally (Ru & Pan 2011). The U.S. Environmental Protection Agency's UV Disinfection Manual (USEPA-UVDGM 2006) provides the corresponding relationship between the inactivation rate of *Giardia*, *Cryptosporidium*, and viruses with the irradiation measurement, as shown in Table 7. Ru & Pan (2011) combined USEPA-UVDGM and German standard DVGM to apply UV disinfection to a water plant in Shanghai. The designed UV reactor has a maximum water volume of 6,627 m<sup>3</sup>/h and a minimum water volume of 3,313 m<sup>3</sup>/h. The ultraviolet dose in the early stage of the operation is 270 J/m<sup>2</sup>. The long-term ultraviolet measurement is 400 J/m<sup>2</sup>, making the removal rates of *Giardia* and *Cryptosporidium* reach 2.5–3.0 log. However, the data provided by USEPA-UVDGM suggests that, although ultraviolet rays can effectively kill the *Giardia* cysts and *Cryptosporidium* oocysts, the amount of exposure required for viruses in the water is extremely large.

## CONCLUSIONS

1. As intestinal pathogenic microorganisms, *Giardia* and *Cryptosporidium* have a wide range of transmission routes and high biosafety risks. Studies have found that pets are commonly infected with *Giardia* and *Cryptosporidium*. Therefore, the health department should initiate investigations and prevention of pets infected with zoonotic protozoa. At the same time, water source transmission is their main way of transmission, and attention should be paid to their removal and monitoring in drinking water treatment. The pollution of *Giardia* and *Cryptosporidium* is more serious in the water environment of many places in China, and there are more reports of infections in humans. The number of reports on *Giardia* and *Cryptosporidium* in other countries is uneven. Countries in Asia and Africa have more serious pollution problems. Thailand, India, Bangladesh, Jordan, and Egypt have high infection rates among children and immunodeficient patients. The high infection rate is closely related to the country's sanitary conditions and water environment. Therefore, it is necessary to improve the construction of sanitation facilities and close attention to high-risk susceptible groups.
2. The detection technology for *Giardia* and *Cryptosporidium* has matured through decades of diagnostic developments. The detection methods are abundant, and many research contents are combining multiple detection methods. Researchers should choose suitable detection methods according to their experimental conditions. International organizations should promote the LAMP method in Asia and Africa as the core detection method for such high-infection poverty

**Table 7** | Comparison table of inactivation rate and radiation dose of *Giardia* and *Cryptosporidium* by ultraviolet

Inactivation rate/log	0.5	1	1.5	2	2.5	3	3.5	4
<i>Giardia</i> /J/m <sup>2</sup>	16	25	39	58	85	120	150	220
<i>Cryptosporidium</i> /J/m <sup>2</sup>	15	21	30	52	77	110	150	220
Virus/J/m <sup>2</sup>	390	580	790	1,000	1,210	1,430	1,630	1,860

areas and apply it to water environment detection and medical emergency detection in poor areas. All stages of drinking water treatment have a certain removal effect on *Giardia* and *Cryptosporidium*. The coagulation-sedimentation-filtration method is to remove them as suspended particles. The removal rate is related to the selection of coagulant, filtration method, turbidity control, among other factors. Commonly used coagulants, such as ferric chloride, alum, and polyaluminum chloride, can effectively remove *Giardia* and *Cryptosporidium* and are not affected by environmental pH. It is recommended to use the combination of anthracite + silica sand + garnet to remove the cysts and oocysts of the two protozoa. The minimum limit of the removal rate of *Giardia* and *Cryptosporidium* in the filtered water is 2.0 log, taking into account that the removal rate needs to be increased according to the content of the raw water. The different disinfection processes have different killing intensities for *Giardia* and *Cryptosporidium*. Compared with Cl<sub>2</sub>, ClO<sub>2</sub> has a stronger killing effect, while ozone and ultraviolet disinfection have better killing effects among them. However, it is not recommended to use a single disinfectant. Moreover, the operating parameters should be implemented according to the USEPA standard.

- Improved drinking water treatment process is a key link to prevent *Giardia* and *Cryptosporidium* from spreading through water sources and causing harm to human health. However, this method requires strict treatment management and close monitoring. In the absence of a water treatment facility, protozoa infections still occur frequently in some regions of various countries. As an emerging water treatment technology, membrane separation technology can achieve excellent control for *Giardia* and *Cryptosporidium*. The most widely used microfiltration membrane can reach a removal rate of up to 6.0 log. Therefore, it is recommended that the conventional water treatment methods for *Giardia* and *Cryptosporidium* involve an appropriate amount of ozone or ultraviolet based on controlling the effluent of the filtered water. It should also use a microfiltration membrane to intercept the protozoa cysts and oocysts to ensure safety. In the emergency water source treatment, the treatment method of ultraviolet + microfiltration can be directly adapted to provide a higher water treatment per unit time under the premise of ensuring water quality safety.

## ACKNOWLEDGEMENTS

This research is funded by National Natural Science Foundation of China (51678026) and Beijing University of Civil Engineering Postgraduate Innovation Project (PG2021047). We would like to express our gratitude to EditSprings (<https://www.editsprings.com/>) for the expert linguistic services provided.

## CONFLICT OF INTEREST

We declare that we have no financial and personal conflicts of interest to this work.

## DATA AVAILABILITY STATEMENT

All relevant data are included in the paper or its Supplementary Information.

## REFERENCES

- Abu-Elamreen, F. H., Abed, A. A. & Sharif, F. 2008 Viral, bacterial and parasitic etiology of pediatric diarrhea in Gaza, Palestine. *Medical Principles and Practice* **17**, 296–301.
- Adanir, R. & Tasci, F. 2013 Prevalence of helminth eggs in raw vegetables consumed in Burdur, Turkey. *Food Control* **31**, 482–484.
- Ahmed, S. A. & Karanis, P. 2020 *Cryptosporidium* and cryptosporidiosis: the perspective from the Gulf countries. *International Journal of Environmental Research and Public Health* **17** (18), 1–34.
- Al-Brikan, F. A., Salem, H., Beeching, N. & Hilal, N. 2008 Multilocus genetic analysis of *Cryptosporidium* isolates from Saudi Arabia. *Journal of the Egyptian Society of Parasitology* **38** (2), 645–658.
- Alexander, K., Herbein, J. & Zajac, A. 2012 The occurrence of *cryptosporidium* and giardia infections among patients reporting diarrheal disease in Chobe District, Botswana. *Advances in Infectious Diseases* **2**, 143–147.
- Alleyne, L., Fitzhenry, R., Mergen, K., Espina, N., Amoroso, E., Cimini, D., Balter, S., Fireteanu, A. M., Seeley, A., Janus, L., Gutelius, B., Madison-Antenucci, S. & Thompson, C. 2020 *Epidemiology of Cryptosporidiosis, New York City, New York, USA, 1995–20181*. *Emerging Infectious Diseases* **26**, 409–419.
- Al-Megrin, W. 2010 Intestinal parasites infection among immunocompromised patients in Riyadh, Saudi Arabia. *Pakistan Journal of Biological Sciences: PJBS* **13** (8), 390–394.
- Al-Qobati, S. A., Al-Maktari, M. T., Al-Zoa, A. M. & Derhim, M. 2012 Intestinal parasitosis among Yemeni patients with cancer, Sana'a, Yemen. *Journal of the Egyptian Society of Parasitology* **42** (3), 727–734.

- Al-Shawa, R. M. & Mwafy, S. N. 2007 *The Enteroparasitic Contamination of Commercial Vegetables in Gaza Governorates*.
- Ammoura, A. 2010 *Impact of hygienic level on parasite infection*. *Asian Pacific Journal of Tropical Medicine* **3**, 148–149.
- Amoah, P., Drechsel, P., Abaidoo, R. & Klutse, A. 2007 *Effectiveness of common and improved sanitary washing methods in selected cities of West Africa for the reduction of coliform bacteria and helminth eggs on vegetables*. *Tropical Medicine & International Health* **12**, 40–50.
- Arthur, J., Bodhidatta, L., Echeverria, P., Phuphaisan, S. & Paul, S. 1992 *Diarrheal disease in Cambodian children at a camp in Thailand*. *American Journal of Epidemiology* **135** (5), 541–551.
- Assavapongpaiboon, B., Bunkasem, U., Sanprasert, V. & Nuchprayoon, S. 2018 *A cross-sectional study on intestinal parasitic infections in children in suburban public primary schools, saraburi, the central region of Thailand*. *The American Journal of Tropical Medicine and Hygiene* **98** (3), 763–767.
- Aye, T., Moe, K., Nyein, M. & Swe, T. 1994 *Cryptosporidiosis in Myanmar infants with acute diarrhea*. *The Southeast Asian Journal of Tropical Medicine and Public Health* **25** (4), 654–656.
- Azeez, S. S. & Alsakee, H. M. 2017 *Cryptosporidium spp. and rotavirus gastroenteritis and change of incidence after rotavirus vaccination among children in Raparin Pediatrics Hospital, Erbil, Iraq*. *Medical Journal of Indonesia* **26**, 190.
- Baqai, R., Anwar, S. & Kazmi, S. 2005 *Detection of cryptosporidium in immunosuppressed patients*. *Journal of Ayub Medical College, Abbottabad: JAMC* **17** (3), 38–40.
- Berendes, D., Capone, D., Knee, J., Holcomb, D., Sultana, S., Pickering, A. & Brown, J. 2020 *Associations between enteric pathogen carriage and height-for-age, weight-for-age and weight-for-height in children under 5 years old in urban Dhaka, Bangladesh*. *Epidemiology and Infection* **148**, e39.
- Betancourt, W. & Rose, J. 2004 *Drinking water treatment processes for removal of Cryptosporidium and Giardia*. *Veterinary Parasitology* **126** (1–2), 219–234.
- Blackburn, C. & McClure, P. 2002 *Foodborne Pathogens: Hazards, Risk Analysis and Control*.
- Bouyou-Akotet, M., Owono-Medang, M., Moussavou-Boussougou, M., Mamfoumbi, M. M., Mintsu-Nguéma, R., Mawili-Mboumba, D. P. & Kombila, M. 2015 *Low sensitivity of the ImmunocardSTAT® Crypto/Giardia rapid assay test for the detection of giardia and Cryptosporidium in fecal samples from children living in Libreville, Central Africa*. *Journal of Parasitic Diseases* **40**, 1179–1183.
- Bukhari, Z., Smith, H., Sykes, N., Humphreys, S., Paton, C., Girdwood, R. & Fricker, C. 1997 *Occurrence of cryptosporidium spp oocysts and Giardia spp cysts in sewage influents and effluents from treatment plants in England*. *Water Science and Technology* **35**, 385–390.
- Bushen, O., Kohli, A., Pinkerton, R., Dupnik, K., Newman, R., Sears, C., Fayer, R., Lima, A. & Guerrant, R. 2007 *Heavy cryptosporidial infections in children in northeast Brazil: comparison of Cryptosporidium hominis and Cryptosporidium parvum*. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **101** (4), 378–384.
- Cai, H. 1995 *Modern Environmental Hygiene*. People's Medical Publishing House, Beijing, China.
- Cai, J. 2005 *The prevalence and control of infections of Giardia lamblia and Cryptosporidium*. *Chinese Journal of Health Inspection* (11). doi: CNKI: SUN: ZWJZ.0.2005-11-068.
- Chamberlain, J., Gibbs, R., Ranier, J., Nguyen, P. & Caskey, C. 1988 *Deletion screening of the Duchenne muscular dystrophy locus via multiplex DNA amplification*. *Nucleic Acids Research* **16** (23), 11141–11156.
- Checkley, W., White, A., Jaganath, D., Arrowood, M., Chalmers, R., Chen, X., Fayer, R., Griffiths, J., Guerrant, R., Hedstrom, L., Huston, C., Kotloff, K., Kang, G., Mead, J., Miller, M. A., Petri, W., Priest, J., Roos, D., Striepen, B., Thompson, R. C., Ward, H., Voorhis, W. V., Xiao, L., Zhu, G. & Houpt, E. 2015 *A review of the global burden, novel diagnostics, therapeutics, and vaccine targets for cryptosporidium*. *The Lancet Infectious Diseases* **15** (1), 85–94.
- Chen, X. M., Keithly, J. S., Paya, C. V. & Larusso, N. F. 2002 *Cryptosporidiosis*. *New England Journal of Medicine* **346** (22), 1723.
- Chen, M., Li, H., Tang, L., Kan, Z., Li, W. & Gu, Y. 2018 *Molecular epidemiology of Cryptosporidium parvum*. *Science and Technology Information of Animal Husbandry and Veterinary Medicine* **9**, 24–26. doi:CNKI:SUN:XMKX. 0.2018-09-017.
- Cheng, X. 2015 *Human Parasitology*. Fudan University Press, Shanghai, China.
- Chhin, S., Harwell, J., Bell, J. D., Rózycki, G., Ellman, T., Barnett, J., Ward, H., Reinert, S. & Pugatch, D. 2006 *Etiology of chronic diarrhea in antiretroviral-naïve patients with HIV infection admitted to Norodom Sihanouk Hospital, Phnom Penh, Cambodia*. *Clinical Infectious Diseases: an Official Publication of the Infectious Diseases Society of America* **43** (7), 925–932.
- Cho, M. & Yoon, J. 2007 *Quantitative evaluation and application of Cryptosporidium parvum inactivation with ozone treatment*. *Water Science and Technology: A Journal of the International Association on Water Pollution Research* **55** (1–2), 241–250.
- Chokephaibulkit, K., Wanachiwanawin, D., Tosasuk, K., Pavitpok, J., Vanprapar, N. & Chearskul, S. 2001 *Intestinal parasitic infections among human immunodeficiency virus-infected and -uninfected children hospitalized with diarrhea in Bangkok, Thailand*. *The Southeast Asian Journal of Tropical Medicine and Public Health* **32** (4), 770–775.
- Clark, R., Sivaganesan, M., Rice, E. & Chen, J. 2003 *Development of a Ct equation for the inactivation of Cryptosporidium oocysts with chlorine dioxide*. *Water Research* **37** (11), 2773–2783.
- Cornwell, D., Macphée, M., Brown, R. & Via, S. 2003 *Demonstrating Cryptosporidium Removal using spore monitoring at lime-softening plants*. *Journal American Water Works Association* **95**, 124–133.
- Cui, F., Zuo, J., Zhao, Z. & Fei, X. 2006 *Research progress on Giardia and Cryptosporidium in drinking water*. *Journal of Harbin Institute of Technology* **9**, 1487–1491.
- DuPont, H., Chappell, C., Sterling, C., Okhuysen, P., Rose, J. & Jakubowski, W. 1995 *The infectivity of Cryptosporidium parvum in healthy volunteers*. *The New England Journal of Medicine* **332** (13), 855–859.

- El-Malky, M., Mowafy, N., Zaghloul, D., Al-Harathi, S., El-Bali, M., Mohamed, R. T., Bakri, R., Mohamed, A. A. & Elmedany, S. 2018 Prevalence of *Cryptosporidium* species isolated from diarrheic children in Makkah, Saudi Arabia. *Tropical Biomedicine* **35** (1), 76–81.
- Entrala, E., Garin, Y. J., Meneceur, P., Hayat, M., Scherpereel, G., Savin, C., Féliers, C. & Derouin, F. 2007 Pilot-scale evaluation of UV reactors' efficacy against *in vitro* infectivity of *Cryptosporidium parvum* oocysts. *FEMS Immunology and Medical Microbiology* **51** (3), 555–561.
- Eraky, M., Rashed, S., Nasr, M., El-Hamshary, A. M. & El-Ghannam, A. S. 2014 Parasitic contamination of commonly consumed fresh leafy vegetables in Benha, Egypt. *Journal of Parasitology Research* **2014**, 1–7.
- Escobedo, A., Almirall, P., Robertson, L., Franco, R. M., Hanevik, K., Mørch, K. & Cimerman, S. 2010 Giardiasis: the ever-present threat of a neglected disease. *Infectious Disorders Drug Targets* **10** (5), 329–348.
- Escobedo, A., Almirall, P., Alfonso, M., Cimerman, S. & Chacín-Bonilla, L. 2014 Sexual transmission of giardiasis: a neglected route of spread? *Acta Tropica* **132**, 106–111.
- Fayer, R., Morgan, U. & Upton, S. J. 2000 Epidemiology of *Cryptosporidium*: transmission, detection and identification. *International Journal for Parasitology* **30** (12–13), 1305–1322.
- Fontaine, M. & Guillot, E. 2003 An immunomagnetic separation-real-time PCR method for quantification of *Cryptosporidium parvum* in water samples. *Journal of Microbiological Methods* **54**1, 29–36.
- Fu, J., Jin, C. & Zhao, Y. 1998 Study on the secondary pollution of drinking water in residential areas and its prevention and control countermeasures. *Water Supply and Drainage* **7**, 55–59.
- Gabre, R. & Shakir, A. 2016 Prevalence of some human enteroparasites in commonly consumed raw vegetables in Tabuk, Saudi Arabia. *Journal of Food Protection* **79** (4), 655–658.
- Gao, Y. 2008 Preliminary Study on the Killing Effect and Mechanism of Chlorine Dioxide on *Cryptosporidium* Oocysts in Water. Master's Thesis, Harbin Institute of Technology, Harbin, China.
- Garcia-R, J. C., Pita, A., Velathanthiri, N., French, N. & Hayman, D. 2020 Species and genotypes causing human cryptosporidiosis in New Zealand. *Parasitology Research* **119**, 2317–2326.
- Gebre, B., Alemayehu, T., Girma, M., Ayalew, F., Tadesse, B. T. & Shemelis, T. 2019 Cryptosporidiosis and other intestinal parasitic infections and concomitant threats among HIV-Infected children In southern Ethiopia receiving first-line antiretroviral therapy. *HIV/AIDS (Auckland, N.Z.)* **11**, 299–306.
- Gharieb, R., Merwad, A., Saleh, A. A. & El-Ghany, A. M. 2018 Molecular screening and genotyping of cryptosporidium species in household dogs and in-contact children in Egypt: risk factor analysis and zoonotic importance. *Vector Borne and Zoonotic Diseases* **18** (8), 424–432.
- Giovanni, G. D., Hashemi, F., Shaw, N. J., Abrams, F., Lechevallier, M. & Abbaszadegan, M. 1999 Detection of infectious cryptosporidium parvum oocysts in surface and filter backwash water samples by immunomagnetic separation and integrated cell culture-PCR. *Applied and Environmental Microbiology* **65**, 3427–3432.
- Graczyk, T., Majewska, A. & Schwab, K. 2008 The role of birds in dissemination of human waterborne enteropathogens. *Trends in Parasitology* **24** (2), 55–59.
- Gu, Y., Wang, K., Liu, D., Mei, N., Chen, C., Chen, T., Han, M., Zhou, L., Cao, J., Zhang, H., Zhang, X., Fan, Z. & Li, W. 2015 Pet dogs Giardia lamblia and Cryptosporidium molecular detection of infection. *Chinese Journal of Parasitology and Parasitic Diseases* **5**, 362–367.
- Hallier-Soulier, S. & Guillot, E. 2000 Detection of cryptosporidia and *Cryptosporidium parvum* oocysts in environmental water samples by immunomagnetic separation–polymerase chain reaction. *Journal of Applied Microbiology* **89** (1), 5–10.
- Hernández, P. C., Pava, L. M., Chaparro-Olaya, J., López-Osorio, S., López-Arias, Á. & Chaparro-Gutierrez, J. 2021 Multilocus genotyping of *Giardia intestinalis* in pet dogs of Medellín Colombia. *Veterinary Parasitology, Regional Studies and Reports* **23**, 100520.
- Hou, Q., Chen, J. & Liu, W. 2011 Cryptosporidium infection and epidemic characteristics of children in Xi'an Children's Hospital. *Contemporary Medicine* **24**, 53.
- Hu, L. 2011 Research on the Molecular Genetic Characteristics and Phylogenetic Development of *Giardia Canis* and *Cryptosporidium* in Parts of Sichuan Province. Master's Thesis, Sichuan Agricultural University, Sichuan, China.
- Hunter, P., Zmirou-Navier, D. & Hartemann, P. 2009 Estimating the impact on health of poor reliability of drinking water interventions in developing countries. *The Science of the Total Environment* **407** (8), 2621–2624.
- Ismail, Y. 2016 Prevalence of parasitic contamination in salad vegetables collected from supermarkets and street vendors in Amman and Baqa'a – Jordan. *Polish Journal of Microbiology* **65** (2), 201–207.
- Joachim, A. 2004 Human cryptosporidiosis: an update with special emphasis on the situation in Europe. *Journal of Veterinary Medicine, B, Infectious Diseases and Veterinary Public Health* **51** (6), 251–259.
- Karanis, P., Thekisoe, O., Kiouptsi, K., Ongerth, J., Igarashi, I. & Inoue, N. 2007 Development and preliminary evaluation of a loop-mediated isothermal amplification procedure for sensitive detection of cryptosporidium oocysts in fecal and water samples. *Applied and Environmental Microbiology* **73**, 5660–5662.
- Keegan, A., Daminato, D. P., Saint, C. & Monis, P. 2008 Effect of water treatment processes on *Cryptosporidium* infectivity. *Water Research* **42** (6–7), 1805–1811.
- King, B., Hoefel, D., Daminato, D. P., Fanok, S. & Monis, P. 2008 Solar UV reduces *Cryptosporidium parvum* oocyst infectivity in environmental waters. *Journal of Applied Microbiology* **104**, 1311–1323.



- Kofoid, C. A. & Christiansen, E. B. 1915 [On the life-history of Giardia](#). *Proceedings of the National Academy of Sciences of the United States of America* **1** (11), 547–552.
- Korpe, P., Valencia, C., Haque, R., Mahfuz, M., McGrath, M., Houpt, E., Kosek, M., McCormick, B., Yori, P. P., Babji, S., Kang, G., Lang, D. R., Gottlieb, M., Samie, A., Bessong, P., Faruque, A., Mduma, E., Nshama, R., Havt, A., Lima, I., Lima, A., Bodhidatta, L., Shreshtha, A., Petri, W., Ahmed, T. & Duggal, P. 2018 [Epidemiology and risk factors for cryptosporidiosis in children from 8 low-income sites: results from the MAL-ED study](#). *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America* **67**, 1660–1669.
- Kotloff, K., Nataro, J., Blackwelder, W., Nasrin, D., Farag, T., Panchalingam, S., Wu, Y., Sow, S., Sur, D., Breiman, R., Faruque, A., Zaidi, A., Saha, D., Alonso, P., Tamboura, B., Sanogo, D., Onwuchekwa, U., Manna, B., Ramamurthy, T., Kanungo, S., Ochieng, J. B., Omoro, R., Oundo, J., Hossain, A., Das, S., Ahmed, S., Qureshi, S., Quadri, F., Adegbola, R., Antonio, M., Hossain, M., Akinsola, A., Mandomando, I., Nhampossa, T., Acácio, S., Biswas, K., O'Reilly, C., Mintz, E., Berkeley, L. Y., Muhsen, K., Sommerfelt, H., Robins-Browne, R. & Levine, M. 2013 [Burden and aetiology of diarrhoeal disease in infants and young children in developing countries \(the Global Enteric Multicenter Study, GEMS\): a prospective, case-control study](#). *The Lancet* **382**, 209–222.
- Kumar, T., Majid, M. A., Onichandran, S., Jaturas, N., Andiappan, H., Salibay, C., Tabo, H. A., Tabo, N., Dungca, J. Z., Tangpong, J., Phiriyasamith, S., Yuttayong, B., Polseela, R., Do, B. N., Sawangiaroen, N., Tan, T., Lim, Y. & Nissapatorn, V. 2016 [Presence of cryptosporidium parvum and Giardia lamblia in water samples from Southeast Asia: towards an integrated water detection system](#). *Infectious Diseases of Poverty* **5**, 1–12.
- Kurniawan, A., Karyadi, T., Dwintasari, S. W., Sari, I. P., Yunihastuti, E., Djauzi, S. & Smith, H. 2009 [Intestinal parasitic infections in HIV/AIDS patients presenting with diarrhoea in Jakarta, Indonesia](#). *Transactions of the Royal Society of Tropical Medicine and Hygiene* **103** (9), 892–898.
- Kurnosova, O. P., Arisov, M. & Odoyevskaya, I. M. 2019 [Intestinal parasites of pets and other house-kept animals in Moscow](#). *Helminthologia* **56**, 108–117.
- Lai, K. 1992 [Intestinal protozoan infections in Malaysia](#). *The Southeast Asian Journal of Tropical Medicine and Public Health* **23** (4), 578–586.
- Latif, B. & Rossle, N. F. 2015 [Cryptosporidiosis among children with diarrhoea in three Asian countries: a review](#). *Asian Pacific Journal of Tropical Biomedicine* **5**, 885–888.
- Li, X. 2015 [Study on the Detection and Removal of Cryptosporidium and Giardia in tap Water](#). Master's Thesis, Chang'an University, Shaanxi, China.
- Li, W., Wu, K., Zhang, X., Li, J., Zhang, G., Li, S., Gong, P. & Yang, J. 2010 [Detection of Cryptosporidium by nested PCR](#). *Chinese Journal of Veterinary Medicine* **6**, 779–783.
- Liao, C. 2019 [Molecular Epidemiological Survey of Bird Cryptosporidium in Some Areas of Hubei Province](#). Master's Thesis, Huazhong Agricultural University, Hubei, China.
- Liao, L., Xiao, N., Sun, J., Ruan, Z., Lin, Q., Lu, J. & Qin, Z. 2014 [Establishment of a dual real-time fluorescent PCR detection method for Cryptosporidium and Giardia](#). *Shanghai Animal Husbandry and Veterinary News* **3**, 16–18.
- Liao, S., Lin, X., Sun, Y., Qi, N., Lv, M., Wu, C., Li, J., Hu, J., Yu, L., Cai, H., Xiao, W., Sun, M. & Li, G. 2020 [Occurrence and genotypes of Cryptosporidium spp., Giardia duodenalis, and Blastocystis sp. in household, shelter, breeding, and pet market dogs in Guangzhou, southern China](#). *Scientific Reports* **10**.
- Lim, Y., Rohela, M., Sim, B., Jamaiah, I. & Nurbayah, M. 2005 [Prevalence of cryptosporidiosis in HIV-infected patients in Kajang Hospital, Selangor](#). *The Southeast Asian Journal of Tropical Medicine and Public Health* **36** (Suppl 4), 30–33.
- Lim, Y., Ahmad, R. A. & Smith, H. 2008 [Current status and future trends in Cryptosporidium and Giardia epidemiology in Malaysia](#). *Journal of Water and Health* **6** (2), 239–254.
- Lim, Y., Iqbal, A., Surin, J., Sim, B., Jex, A., Nolan, M. J., Smith, H. & Gasser, R. 2011 [First genetic classification of Cryptosporidium and Giardia from HIV/AIDS patients in Malaysia](#). *Infection, Genetics and Evolution: Journal of Molecular Epidemiology and Evolutionary Genetics in Infectious Diseases* **11** (5), 968–974.
- Liu, A., Zhang, X., Yang, F., Dong, Y., Han, S., Wen, J. & Zhang, W. 2011 [Molecular identification of Cryptosporidium andersonii in raw water of Harbin sewage treatment plant](#). *Chinese Journal of Pathogen Biology* **6**, 440–442.
- Logsdon, G. & Johnson, S. 2010 [The Removal and Disinfection Efficiency of Lime Softening Processes for Giardia and Viruses](#).
- Lu, C. 2009 [Investigation of Intestinal Parasite Infections in Certain Rodents and Analysis of Cryptosporidium Genetic Characteristics](#). Master's degree Thesis, Henan Agricultural University, Henan, China.
- Lu, J. 2016 [Establishment of Detection Methods for Giardia and Cryptosporidium in Water Source Water and Investigation of Pollution in Some Areas of my Country](#). Master's Thesis, Chinese Center for Disease Control and Prevention, Beijing, China.
- Ma, L., Sotiriadou, I., Cai, Q., Karanis, G., Wang, G., Wang, G., Lu, Y., Li, X. & Karanis, P. 2014 [Detection of Cryptosporidium and Giardia in agricultural and water environments in the Qinghai area of China by IFT and PCR](#). *Parasitology Research* **113**, 3177–3184.
- Mahmoudi, M., Ongerth, J. & Karanis, P. 2017 [Cryptosporidium and cryptosporidiosis: the Asian perspective](#). *International Journal of Hygiene and Environmental Health* **220** (7), 1098–1109.
- Meamar, A., Guyot, K., Certad, G., Dei-Cas, E., Mohraz, M., Mohebbi, M., Mohammad, K., Mehbod, A., Rezaie, S. & Rezaian, M. 2006 [Molecular Characterization of Cryptosporidium Isolates from Humans and Animals in Iran](#). *Applied and Environmental Microbiology* **73**, 1033–1035.
- Meng, J., Zhao, Y. & Asya, B. 2011 [Detection of water-borne Cryptosporidium in Xinjiang](#). *Chinese Journal of Pathogen Biology* **1**, 42–44 + 28.

- Method 1623: Cryptosporidium in Water by Filtration/IMS/FA EPA-821-r-01-025 2001 *United States Environmental Protection Agency*. Office of Water, Washington, DC, USA.
- Molloy, S., Tanner, C., Kirwan, P., Asaolu, S., Smith, H., Nichols, R., Connelly, L. & Holland, C. 2010 Sporadic Cryptosporidium infection in Nigerian children: risk factors with species identification. *Epidemiology and Infection* **139**, 946–954.
- Moniot, M., Nourrisson, C., Faure, C., Delbac, F., Favennec, L., Dalle, F., Garrouste, C. & Poirier, P. 2020 Assessment of a multiplex PCR for the simultaneous diagnosis of intestinal cryptosporidiosis and microsporidiosis: epidemiological report from a French prospective study. *The Journal of Molecular Diagnostics: JMD* **23** (4), 417–423.
- Mutai, D. C., Owili, P. & Muga, M. 2020 Trend of cryptosporidium infection among children below 24 months in an informal urban settlement, Kenya. *Open Journal of Medical Microbiology* **10**, 153–161.
- Nahas, S. A. & Aboualchamat, G. 2020 Investigation of parasitic contamination of salad vegetables sold by street vendors in city markets in Damascus, Syria. *Food and Waterborne Parasitology* **21**, 1–6.
- Ongerth, J. & Pecoraro, J. 1995 Removing Cryptosporidium using multimedia filters. *Journal American Water Works Association* **87**, 83–89.
- Osman, M., Benamrouz, S., Guyot, K., Safadi, D. E., Mallat, H., Dabboussi, F., Hamze, M., Viscogliosi, E. & Certad, G. 2018 Molecular epidemiology of Cryptosporidium spp. in North Lebanon. *Journal of Infection in Developing Countries* **12** 2.1, 34S.
- Paboriboune, P., Phoumindr, N., Borel, E., Sourinphoumy, K., Phaxayaseng, S., Luangkhot, E., Sengphilom, B., Vansilalom, Y., Odermatt, P., Delaporte, E., Etard, J. & Rabodonirina, M. 2014 Intestinal parasitic infections in HIV-infected patients, Lao People's Democratic Republic. *PLoS ONE* **9** (3), e91452.
- Paudyal, S., Shrestha, S. & Mahato, N. 2013 Zoonotic aspects of cryptosporidiosis in Nepal. *International Journal of Applied Sciences and Biotechnology* **1**, 21–26.
- Pinlaor, S., Mootsikapun, P., Pinlaor, P., Pipitgool, V. & Tuangnadee, R. 2005 Detection of opportunistic and non-opportunistic intestinal parasites and liver flukes in HIV-positive and HIV-negative subjects. *The Southeast Asian Journal of Tropical Medicine and Public Health* **36** (4), 841–845.
- Pipiková, J., Papajová, I., Majláthová, V., Šoltys, J., Bystrianska, J., Schusterová, I. & Vargová, V. 2018 First report on Giardia duodenalis assemblage F in Slovakian children living in poor environmental conditions. *Journal of Microbiology, Immunology, and Infection* **53**, 148–156.
- Platts-Mills, J., Babji, S., Bodhidatta, L., Gratz, J., Haque, R., Havt, A., McCormick, B., McGrath, M., Olórtegui, M., Samie, A., Shakoob, S., Mondal, D., Lima, I., Hariraju, D., Rayamajhi, B., Qureshi, S., Kabir, F., Yori, P., Mufamadi, B., Amour, C., Carreon, J., Richard, S., Lang, D. R., Bessong, P., Mduma, E., Ahmed, T., Lima, A., Mason, C., Zaidi, A., Bhutta, Z., Kosek, M., Guerrant, R., Gottlieb, M., Miller, M. A., Kang, G. & Houghton, E. 2015 Pathogen-specific burdens of community diarrhoea in developing countries: a multisite birth cohort study (MAL-ED). *The Lancet Global Health* **3** (9), e564–e575.
- Plutzer, J., Lassen, B., Jokelainen, P., Djurković-Djaković, O., Kucsera, I., Dorbek-Kolin, E., Šoba, B., Sréter, T., Imre, K., Omeragić, J., Nikolic, A., Bobić, B., Živičnjak, T., Lučinger, S., Stefanović, L. L., Kučinar, J., Sroka, J., Deksné, G., Keidāne, D., Kváč, M., Hůzová, Z. & Karanis, P. 2018 Review of Cryptosporidium and Giardia in the eastern part of Europe, 2016. *Eurosurveillance* **23**, 1–23.
- Prado, M. S., Strina, A., Barreto, M., Oliveira-Assis, A. M., Paz, L. M. & Cairncross, S. 2003 Risk factors for infection with Giardia duodenalis in pre-school children in the city of Salvador, Brazil. *Epidemiology and Infection* **131**, 899–906.
- Qi, M., Wang, R., Ning, C., Jian, F. & Zhang, L. 2011 Epidemiological investigation and molecular genetic analysis of cryptosporidiosis in pet birds in Zhengzhou. In: *Chinese Society of Animal Husbandry and Veterinary Medicine Proceedings of the Sixth Congress of the Society of Domestic Animal Parasitology and the Eleventh Academic Symposium*. p. 279.
- Qi, M., Luo, N., Wang, H., Yu, F., Wang, R., Huang, J. & Zhang, L. 2015 Investigation on the infection of Cryptosporidium and Enterozoites in pet chinchillas. In: *Animals in China The 15th National Academic Conference of the Society of Parasitology Professional Committee and the 6th International Parasitology Symposium Abstracts*. p. 140.
- Rafei, A., Rashno, Z., Samarbazadeh, A. & Khademvatan, S. 2014 Molecular characterization of Cryptosporidium spp. isolated from immunocompromised patients and children. *Jundishapur Journal of Microbiology* **7**, e9183.
- Rahi, A. A., Ali, M. A. & Al-Charrakh, A. H. 2013 Prevalence of Cryptosporidium parvum among children in Iraq. *American Journal of Life Sciences* **1**, 256.
- Ran, Z., Li, S., Zhu, J., Cui, C. & Yuan, Y. 2011 Study on the influencing factors and mechanism of chlorine dioxide inactivation of Cryptosporidium in water. *Chinese Environmental Science* **6**, 904–909.
- Rice, E., Hoff, J. C. & Schaefer, F. 1982 Inactivation of Giardia cysts by chlorine. *Applied and Environmental Microbiology* **43** (1), 250–251.
- Rimhanen-Finne, R., Hörman, A., Ronkainen, P. & Hänninen, M. 2002 An IC-PCR method for detection of Cryptosporidium and Giardia in natural surface waters in Finland. *Journal of Microbiological Methods* **50** (3), 299–303.
- Rivera, W., Yason, J. A. & Rivera, P. 2005 Serological detection of cryptosporidiosis among Filipino cancer patients. *Parasitology Research* **98**, 75–76.
- Ru, L. & Pan, X. 2011 Engineering application of low-pressure and high-intensity ultraviolet disinfection system in Shanghai Linjiang Water Plant. *Water Supply and Drainage* **8**, 17–21.
- Sagnuankiat, S., Wanichsuwan, M., Bhunnachet, E., Jungarat, N., Panraksa, K., Komalamisra, C., Maipanich, W., Yoonuan, T., Pubampen, S., Adisakwattana, P. & Wathanakulpanich, D. 2014 Health status of immigrant children and environmental survey of child daycare centers in Samut Sakhon Province, Thailand. *Journal of Immigrant and Minority Health* **18**, 21–27.

- Sanprasert, V., Srichaipon, N., Bunkasem, U., Srirungruang, S. & Nuchprayoon, S. 2016 Prevalence of intestinal protozoan infections among children in Thailand: a large-scale screening and comparative study of three standard detection methods. *The Southeast Asian Journal of Tropical Medicine and Public Health* **47** (6), 1123–1133.
- Shalaby, I., Gherbawy, Y., Jamjoom, M. & Banaja, A. 2014 Prevalence and genotyping of *Cryptosporidium* in stool samples collected from children in Taif City (Saudi Arabia). *Tropical Biomedicine* **31** (2), 215–224.
- Shan, H. 2007 *Seroepidemiological Study on Cryptosporidium Infection in Chinese Community Population Using Recombinant Cryptosporidium Specific Antigen*. Master's Thesis, Nanjing Medical University, Nanjing, China.
- Silva, K. J. & Sabogal-Paz, L. P. 2020 *Cryptosporidium spp. and Giardia spp. (oo)cysts as target-organisms in sanitation and environmental monitoring: a review in microscopy-based viability assays*. *Water Research* **189**, 116590.
- Singer, S., Angelova, V. V., Deleon, H. & Miskovsky, E. 2020 *What's eating you? an update on Giardia, the microbiome and the immune response*. *Current Opinion in Microbiology* **58**, 87–92.
- Sirisena, U., Iddawela, W., Noordeen, F. & Wickramasinghe, S. 2014 *Prevalence and identification of cryptosporidium species in paediatric patients with diarrhoea*. *The Ceylon Medical Journal* **59** (3), 75–78.
- Sivaganesan, M. & Mariñas, B. 2005 *Development of a Ct equation taking into consideration the effect of lot variability on the inactivation of Cryptosporidium parvum oocysts with ozone*. *Water Research* **39** (11), 2429–2437.
- Smith, H. V., Caccio, S. M., Tait, A., McLaughlin, J. & Thompson, R. C. 2006 *Tools for investigating the environmental transmission of Cryptosporidium and Giardia infections in humans*. *Trends Parasitol* **22** (4), 160–167.
- Soliman, A., El-Adawy, A., El-Aal, A. A., Elmallawany, M. A., Nahnoush, R. K., Eiaghni, A. R., Negm, M. & Mohsen, A. 2018 *Usefulness of sunlight and artificial UV radiation versus chlorine for the inactivation of Cryptosporidium Oocysts: an in vivo animal study*. *Open Access Macedonian Journal of Medical Sciences* **6**, 975–981.
- States, S., Tomko, R. F., Scheuring, M. & Casson, L. W. 2002 *Enhanced coagulation and removal of cryptosporidium*. *Journal American Water Works Association* **94**, 67–77.
- Steiner, K. L., Ahmed, S., Gilchrist, C., Burkey, C., Cook, H., Ma, J., Korpe, P., Ahmed, E., Alam, M., Kabir, M., Tofail, F., Ahmed, T., Haque, R., Petri, W. & Faruque, A. 2018 *Species of cryptosporidia causing subclinical infection associated with growth faltering in rural and urban Bangladesh: a birth cohort study*. *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America* **67**, 1347–1355.
- Sun, L. 2007 *Research on the Detection and Removal of Giardia and Cryptosporidium in Shanghai Tap Water*. Master's Thesis, Shanghai Jiaotong University, Shanghai, China.
- Sun, B., Xia, Y., Chen, X., Lai, W., He, C., Ding, Z. & Zhang, R. 2014 *Investigation of cryptosporidium and giardia in the centralized water supply of rural rural areas in Southern China*. *Public Health and Preventive Medicine* **4**, 11–13.
- Teixeira, J., Heller, L. & Barreto, M. 2007 *Giardia duodenalis infection: risk factors for children living in sub-standard settlements in Brazil*. *Cadernos de Saude Publica* **23** (6), 1489–1493.
- Thompson, R. 2008 Fayer R, Xiao L, (eds): *Cryptosporidium and Cryptosporidiosis*. *Parasites & Vectors* **1**, 1.
- USEPA *Ultraviolet Disinfection Guidance Manual for the Final Long Term Enhanced Surface Water Treatment Rule, EPA 815-D-03-007 2006*. Office of Water, Washington, DC, USA.
- Utami, W. S., Murhandarwati, E. H., Artama, W. & Kusnanto, H. 2020 *Cryptosporidium infection increases the risk for chronic diarrhea among people living with HIV in Southeast Asia: a systematic review and meta-analysis*. *Asia-Pacific Journal of Public Health* **32**, 8–18.
- Wang, G., Ma, L., Cai, Q., Lu, Y., Wang, G., Li, X., Ye, C. & Niu, X. 2012 *Detection of cryptosporidium and giardia in drinking water sources in sanjiangyuan region*. *Qinghai Journal of Animal Husbandry and Veterinary* **4**, 10–11.
- Wang, L. 2013 *The Distribution, Spread and Traceability of Cryptosporidium in the Water Environment and Susceptible Populations in Some Areas of my Country*. PhD Thesis, East China University of Science and Technology, Shanghai Science and Technology Publishing House, Shanghai, China.
- Wang, H., Ge, C., Zhao, J., Yu, M., Rong, J. & Wang, H. 2014 *Experimental study on the detection of Escherichia coli o157:H7 by immunopcr*. *Journal of Qiqihar Medical College* **3**, 327.
- Wang, J. & Luo, E. 2016 *Pathogenic Biology*, 2nd edn. Shanghai Science and Technology Publishing House, Shanghai, China.
- Wang, H., Yang, Z., Sun, D., Wu, H., Yang, D., Jin, M. & Yin, J. 2019a *Investigation of Cryptosporidium and Giardia in the pastoral area of Baotou City*. *Journal of Environment and Health* **6**, 522–524.
- Wang, L., Chen, Y., Huang, J. & Zhang, L. 2019b *Spatial analysis of cryptosporidium epidemiological data in China based on GIS technology*. *Chinese Journal of Zoonoses* **7**, 633–638.
- Wegayehu, T., Adamu, H. & Petros, B. 2013 *Prevalence of Giardia duodenalis and Cryptosporidium species infections among children and cattle in North Shewa Zone, Ethiopia*. *BMC Infectious Diseases* **13**, 419–419.
- Wegayehu, T., Karim, R., Anberber, M., Adamu, H., Erko, B., Zhang, L. & Tilahun, G. 2016 *Prevalence and genetic characterization of cryptosporidium species in dairy calves in central Ethiopia*. *PLoS ONE* **11**, 22.
- Xiao, G., Qiu, Z., Qi, J., Chen, J., Liu, F., Liu, W., Luo, J. & Shu, W. 2013 *Occurrence and potential health risk of cryptosporidium and giardia in the three gorges reservoir, China*. *Water Research* **47** (7), 2431–2445.
- Xu, H. 2016 *Infection Status and Genotype Distribution of Cryptosporidium, Giardia and Microsporidia in Shanghai Companion Animals*. Master's Thesis, East China University of Science and Technology, Shanghai, China.
- Xu, H. & Hu, M. 2007 *Research status of Cryptosporidium and Giardia in drinking water*. *Urban Water Supply* (03), 79–80. doi:10.14143/j.cnki.czgs.2007.03.007.

- Xue, Y. 2006 Emerging parasitic disease-cryptosporidiosis. *Journal of Clinical and Experimental Medicine* **9**, 1427–1429.
- Yan, M. & Chen, H. 2004 Pathogenic microorganisms and water treatment. *Journal of Zhejiang University of Technology* **3**, 96–100 + 111.
- Yang, J. 2015 *Molecular Epidemiology of Three Important Intestinal Protozoa in Children and Companion Animals in Heilongjiang Province*. Master's Thesis, Northeast Agricultural University, Heilongjiang, China.
- Ye, X., Li, C., Li, X., Zhang, Y., Yang, P., Wen, L., Liu, F. & Yang, G. 2017 Optimize the EPA 1623 method to detect *Cryptosporidium* and *Giardia* in drinking water. *Journal of Food Safety and Quality Inspection* **5**, 1838–1841.
- Yu, Y., Yin, J., Zhang, X., Zhou, X. & Wu, L. 2008 Harmfulness and removal of *Cryptosporidium* and *Giardia* in drinking water. *Journal of Jilin Institute of Civil Engineering and Architecture* **2**, 15–17 + 44.
- Zaidah, A., Chan, Y., Asma, H., Abdullah, S., Nurhaslindawati, A. R., Salleh, M., Zeehaida, M., Lalitha, P., Mustafa, M. & Ravichandran, M. 2008 Detection of *cryptosporidium parvum* in HIV-infected patients in Malaysia using a molecular approach. *The Southeast Asian Journal of Tropical Medicine and Public Health* **39** (3), 511–516.
- Zhang, X., He, Y., Zhu, Q., Ma, X. & Cai, L. 2010 Investigation on the pollution of *Cryptosporidium* and *Giardia lamblia* in drinking water and environmental water in Shanghai. *Chinese Journal of Parasitology and Parasitic Diseases* **6**, 435–438.
- Zhang, Y., Wang, Z., Lu, P., Xu, M., Wang, W. & Bo, X. 2017 The infection situation of tapeworm, *cryptosporidium* and *microsporidium* in pet dogs in Shihezi City. In: No. *The Sixteenth National Academic Conference and the Seventh International Symposium on Parasitology Abstracts*. p. 197.
- Zheng, H., Yu, Y., Fei, J., Wang, L., Zhang, R., Ding, Z. & Chen, X. 2014 Pollution status and analysis of *Cryptosporidium* and *Giardia* in source water and drinking water in typical areas of Jiangsu Province. *Journal of Environmental Hygiene* **6**, 535–537.
- Zheng, Z., Li, J., Dan, X., Guo, Y., Li, N., Feng, Y. & Xiao, L. 2019 Genotypes and zoonotic characteristics of *Cryptosporidium*, *Giardia* and *Microsporidia* in dogs and cats in Guangdong. In: *Abstracts of the First Academic Forum of Young Scientists of the Veterinary Parasitology Branch of the Chinese Society of Animal Husbandry and Veterinary Medicine*. p. 38.
- Zong, Z., Hu, H. & Lu, Y. 2005 Introduction to the analysis method of *cryptosporidium* and *Giardia*. *Water Supply and Drainage* **6**, 8–11.
- Zueter, A. M., Hijjawi, N., Hamadeneh, K. N., Al-Sheyab, M. M. & Hatamleh, A. M. 2019 *Cryptosporidiosis among hemodialysis patients in Jordan: first preliminary screening surveillance*. *Tropical Medicine and Infectious Disease* **4**, 1–7.

First received 13 August 2021; accepted in revised form 4 October 2021. Available online 20 October 2021