

Review Paper

Effectiveness of behaviour change interventions to reduce the risk of faecal contamination in urban irrigated vegetable value chains – applying the COM-B behavioural framework

David Galibourg ^{a,*}, Rebecca E. Scott ^a, Katherine V. Gough ^b, Pay Drechsel ^c and Barbara E. Evans ^d

^a Water Engineering and Development Centre (WEDC), School of Civil Engineering, Loughborough University, Loughborough, United Kingdom

^b Department of Geography and Environment, Loughborough University, United Kingdom

^c International Water Management Institute, Sri Lanka

^d School of Civil Engineering, University of Leeds, United Kingdom

*Corresponding author. E-mail: d.galibourg@lboro.ac.uk

 DG, 0000-0003-2676-3551; RES, 0000-0002-0069-065X; KVG, 0000-0002-9638-9879; PD, 0000-0002-2592-8812; BEE, 0000-0001-9815-3141

ABSTRACT

In low- and middle-income countries, inadequate sanitation results in faecal contamination of the water used by urban farmers for irrigation. Consumers of raw contaminated vegetables run the risk of developing diarrhoeal diseases and helminth infections, which are a leading cause of under-five mortality and impact the well-being and productivity of millions of adults. This review identifies the evidence base for assessing which factors determine the success and/or failure of interventions that aim to manage the risk of faecal contamination in the urban irrigated vegetable value chain. We carried out a systematic search of the literature from the perspective of the COM-B behaviour framework (Capability + Opportunity + Motivation = Behaviour). Our results reveal that most interventions address stakeholders' opportunity or capability to adopt safe practices without adequately considering their motivation. Interventions often focus on one sector rather than on the whole value chain (sanitation, agriculture, trade, consumption). To effectively change hygiene and food safety practices in the urban irrigated vegetable value chain, stakeholders' intrinsic motivations need to be identified. Where WHO's multi-barrier approach is the best option, we recommend building on local multistakeholder platforms and adopting a behaviour change framework to support the largely technical change from farm to fork.

Key words: behaviour change, food safety, informal sector, motivation, participatory governance

HIGHLIGHTS

- Adopting safe practices from farm to fork can reduce the risk of faecal contamination of urban irrigated vegetables.
- Adopting safe practices depends on stakeholders' opportunity, capability, and motivation to do so.
- Stakeholders' motivations are often neglected.
- Persuasion (i.e., leverage stakeholders' values and norms), environmental restructuring, and role modelling can stimulate stakeholders' motivations.

INTRODUCTION

Most leafy vegetables consumed in cities of low- and middle-income countries (LMICs) are grown locally (Orsini *et al.* 2013). Urban and peri-urban agriculture enhances urban livelihoods, food sovereignty, recycling of organic waste, greening cities, and limiting floods (FAO 2012). However, the urban water systems most often used to irrigate these vegetables are contaminated with faecal, chemical, and emerging contaminants due to lagging sanitation. It is estimated that urban farmers practise *de facto* indirect wastewater reuse on 30 million hectares of farmland, supplying fresh vegetables to 800 million people globally (Thebo *et al.* 2017), and these figures are likely to increase (Raschid-Sally & Jayakody 2008; Figuié 2019).

This is an Open Access article distributed under the terms of the Creative Commons Attribution Licence (CC BY 4.0), which permits copying, adaptation and redistribution, provided the original work is properly cited (<http://creativecommons.org/licenses/by/4.0/>).

Exposure to contaminated water and consuming raw contaminated vegetables are the two major pathways to contracting enteric, acute respiratory, and parasitic diseases (Hoffmann *et al.* 2017; Raj *et al.* 2020). Faecal pathogens (bacteria, protozoa, and viruses) can provoke diarrhoeal diseases (including cholera), acute respiratory infections (e.g., tuberculosis from breathing sprinklers droplets), as well as other water- and food-borne invasive infectious (e.g., hepatitis, typhoid fever) and parasitic diseases (e.g., helminth infections such as *Taenia* and *Ascaris*). Exposure to contaminated water caused about 1.4 million deaths and the loss of 74 million disability-adjusted life years (DALYs) in LMICs in 2019 (Wolf *et al.* 2023). The consumption of contaminated food caused over 220,000 deaths and the loss of 17 million DALYs in LMICs in 2010, mainly in Africa and Southeast Asia (WHO 2015). Even though these mostly preventable diseases primarily affect children under 5 years, they also impact the well-being and productivity of all age groups and impose a heavy economic burden (Hutton & Haller 2004; WHO 2015).

The World Health Organization (WHO) proposes a multiple-barrier approach adapted to LMICs to reduce health risks from using wastewater (WHO 2006, 2022). It suggests public authorities set appropriate health-based targets, working with primary stakeholders (farmers, vegetable traders, street food vendors, and consumers) to combine safe practices from 'farm to fork' to protect workers' and consumers' health. Where possible, sanitation is counted as the first barrier. Amoah *et al.* (2011) demonstrated the cost-effectiveness of adopting low-cost irrigation practices on farms, safe vegetable handling, and washing methods in markets (including street food restaurants) and households. Increasing uptake of these practices remains challenging (Drechsel *et al.* 2022) and should be considered a priority (Prain *et al.* 2022).

To date, there has been no systematic analysis of the factors that result in the success or failure of interventions to change stakeholder hygiene and food safety practices in the urban irrigated vegetable value chains of LMICs. This research responds to that gap and proposes modifications for more successful interventions. The COM-B behavioural model (Capability + Opportunity + Motivation = Behaviour) developed by Michie *et al.* (2011), explained in the following section, was used to structure a review of the range of approaches used to address the risk of faecal contamination in the value chain. This provides a basis for explaining why safe practices are not more widely adopted.

METHODS

This review systematically appraises the impact of interventions on stakeholders' adoption of good hygiene and food safety practices in the urban irrigated value chains of LMICs.

Search strategy

The systematic search of the literature ensured comprehensive conceptual coverage by querying the titles, abstracts, and keywords in the Scopus, Web of Science, and ProQuest-Social Science Premium Collection against the Boolean combination of five queries as follows:

#1 (the presence of human excreta through 16 terms including faeces, faecal sludge or wastewater)

AND

#2 (the occurrence of oro-faecal diseases through 25 terms including water-borne, diarrhoea or helminth)

AND

[#3 (irrigated vegetable production through 11 terms including farmer, horticulture or crop)

OR

#4 (vegetable trading through 21 terms including produce, trader and consumer)]

AND

#5 (low- and middle-income countries through terms such as Global South or developing countries and a list of LMICs countries).

Query #5 was adapted from Liverpool-Tasie *et al.* (2020) and is based on the World Bank classification of economies from 2021. The detailed terms of the search are available in Table 3 in the Supplementary Material.

Inclusion and exclusion criteria

The search was limited to articles focusing on LMICs published in English between 1 January 2000 and 1 August 2022. Furthermore, the Context-Intervention-Mechanism-Outcome (CIMO) logic proposed by Denyer & Tranfield (2009) was adopted to develop the inclusion and exclusion criteria as follows:

- Context: the study relates to the urban irrigated vegetable value chain in LMICs.

Excluded: high-income countries, rural settings, health facilities, schools, focus on drinking water or groundwater, and food products other than vegetables or non-faecal contaminants.

- Intervention: the study reports an initiative aiming to manage the risk of faecal contamination rather than assess the risk.

Excluded: risk assessments.

- Mechanisms: the study reports a risk management mechanism other than conventional effluent treatment.

Excluded: conventional onsite or centralised wastewater or sludge treatment.

- Outcomes: the study reports a reduction of contamination or a change of practices.

Excluded: absent results.

Records were screened using Rayyan (<https://rayyan.ai/>), and those not meeting eligibility and inclusion criteria were excluded. The remaining articles were included for qualitative analysis. Relevant information was systematically identified: authors, publication year, and country of implementation; sector of intervention (farm and/or market and/or household-consumer); mechanism of intervention; and whether the intervention targets stakeholders' opportunities, capabilities, and/or motivations.

Conceptual framework

We hypothesised that many WASH interventions follow top-down designs that primarily focus on providing stakeholders with an innovation (such as a technology, service, or technique), sometimes supported by education or training. They often, however, fail to account for stakeholders' perceptions and willingness to integrate the innovation into their practices. Assessing interventions against a robust theoretical behavioural framework helps identify which factors promote or hinder the success of the cases reviewed. The COM-B behavioural framework (Michie *et al.* 2011) was selected for its accessibility and robust theoretical foundation, unifying several theories and models. Numerous studies build on this model to explain behaviours or promote changes in contexts ranging from informing the British government's behaviour change policies (Michie & West 2013) to improving hand and food hygiene behaviours in LMICs (Wodnik *et al.* 2018).

The COM-B model posits that adopting a behaviour depends on stakeholders having the Capability (C), Opportunity (O), and Motivation (M) to perform this Behaviour (B), hence the COM-B acronym (see Figure 1). The model defines the 'Capability' component as the physical and psychological ability within an individual (e.g., through training and education) to perform a behaviour. Opportunity is the physical and socio-economic possibility of performing a behaviour (e.g., access to tangible and intangible resources, including regulations, technology, or services). Motivation refers to the 'brain processes that energise and direct behaviour', (Michie *et al.* 2011, p. 4) whether automatic (e.g., feelings) or reflexive (e.g., conscious planning).

Deficits in Capability, Opportunity, and/or Motivation can be addressed through nine functions (see Table 1): restricting opportunities for competing behaviours, restructuring the physical or social environment, enabling, educating, training, incentivising, persuading, coercing, and modelling (Michie *et al.* 2011). One intervention function can address more than one behavioural component and can incorporate functions. Enablement is understood broadly as providing the means to make it easier to enact a behaviour innovation to reduce barriers or improve the opportunity (beyond environmental restructuring) or the capability (beyond education and training). Examples include providing physical or social support and material




Figure 1 | The COM-B framework. Source: adapted from Michie *et al.* (2011).

Table 1 | Intervention functions, definitions, and the behaviour change component(s) addressed

Intervention functions	Definitions	Opportunity (physical)	Opportunity (social)	Capability (physical)	Capability (psychological)	Motivation (reflexive)	Motivation (automatic)
Restriction	Using rules to reduce the opportunity to engage in the target behaviour or increase the target behaviour by reducing the opportunity to engage in competing behaviours	✓	✓				
Environmental restructuring	Changing the physical/social context	✓	✓				✓
Enablement	Increasing means/reducing barriers to increase capability or opportunity beyond training, education, or environmental restructuring	✓	✓	✓	✓		✓
Education	Increasing knowledge or understanding				✓	✓	
Training	Imparting skills			✓	✓		
Incentivisation	Creating expectation of reward					✓	✓
Persuasion	Using communication to induce positive or negative feelings or stimulate action					✓	✓
Coercion	Creating expectation of punishment or cost					✓	✓
Modelling	Providing an example for people to aspire to or imitate						✓

Source: Adapted from Michie *et al.* (2011), pp. 7–8.

or financial resources. These interventions are promoted by different types of policy: legislation, service provision, communication, or marketing campaigns. Michie *et al.* (2011, 2014) recommend diagnosing what people need to adopt recommended behaviour by identifying intervention functions and policy categories adapted to foster change, then selecting the behaviour change technique and mode of delivery that best serves the intervention function.

The results presented and discussed below are structured around the intervention functions described in the records under review and clustered around the related Opportunity, Capability, and Motivation components of the COM-B model (see Table 1).

RESULTS AND DISCUSSION

Searching the published literature for ‘farm to fork’ experiences aimed at improving hygiene and safety in urban irrigated vegetable value chains in LMICs yielded 630 unique records. Of these, 597 were excluded during screening and eligibility assessment. The remaining 33 studies were included in the qualitative analysis (see Figure 2). They mention a range of approaches with the potential to complement conventional treatment systems in reducing contamination risk. In many cases, they focus on one sector rather than the whole value chain and overlook stakeholders capabilities and motivations (see Table 2). Their outcomes, often qualitative, are presented below, structured around the COM-B model, clustered by the mechanism (i.e. intervention function) they build on, and grouped under the main need they aim to address (i.e., Opportunity, Capability, or Motivation). These outcomes are discussed in relation to the broader literature. As this relatively small sample primarily represents interventions in sub-Saharan Africa, this may make our findings more representative of this area.

Opportunities

The COM-B framework suggests that physical (e.g., equipment and infrastructure) or social (e.g. services) opportunity needs are best addressed by restricting the opportunities to perform competing behaviours, restructuring the physical and/or social environment, and enablement interventions.




Figure 2 | Flow diagram of the search strategy.

Restriction: bans and regulations

Regulations on wastewater reuse in LMICs reflect different political priorities. To address water scarcity, Mexico and several countries in the Middle East/North Africa and other arid/semi-arid regions promote formal irrigation schemes using treated wastewater (Shoushtarian & Negahban-Azar 2020). Countries including Ghana, India, Jordan, Mexico, Morocco, Tunisia, and Vietnam, however, have officially banned the use of untreated wastewater with mixed success. Regulations around wastewater reuse for irrigation are inadequate or missing in many LMICs, including Southeast Asia (Winkler *et al.* 2017). Regulations that aim to protect public health from faecal pathogens in other sectors can also be unclear or inconsistent (see Moya *et al.* (2019) on the use of excreta-derived fertilisers in Kenya and Okoruwa & Onuigbo-Chatta (2021) on food safety regulations in Nigeria).

The lack of enforcement of regulations to effectively control stakeholders' practices limits the effectiveness of restriction (and coercion) interventions. Examples include Cuenca-Adame *et al.* (2001) for wastewater irrigation in Mexico, Musa & Akande (2002) for routine medical examination of food vendors in Nigeria, and Salamandane *et al.* (2020) for food markets in Mozambique. The lack of coordination among public agencies, the resulting poor regulatory coherence, and weak enforcement capability also exacerbate the problem (Okoruwa & Onuigbo-Chatta 2021). As largely informal food value chains, characterised by a multitude of small-scale farmers and mobile vendors, are challenging to control, most administrations focus on the formal/export sectors in capital cities (Antwi-Agyei *et al.* 2016; Nordhagen *et al.* 2022). The lack of understanding of microbial risk among urban farmers and vegetable and street food vendors, and their limited access to water, sanitation, or cold storage, make compliance with regulations particularly difficult (Keraita *et al.* 2014; Salamandane *et al.* 2020). When non-compliance is detected, sanctions are not always proportionate or dissuasive. Sanctions that are absent or too light entice stakeholders to ignore good practices (Seleman *et al.* 2020).

Ad hoc regulations and adequate enforcement can lead to more sustainable changes than temporary educational or promotional interventions (Idowu & Rowland 2006; Keraita *et al.* 2014). Mobilising the political will to develop adequate policies and secure funding sufficient to enforce resulting regulations is challenging, as local data on the prevalence of food-borne diseases are scarce and public authorities rarely prioritise the informal sector (Jaffee *et al.* 2019). Regulations are more likely to change behaviours if they avoid mimicking international standards they do not have the capability to

Table 2 | Included articles and their association with sectors of the value chain and components of behaviour change

Authors	Country of study	Focus	Integration across sectors of the value chain				Integration across the three sources of behaviour		
			Sanitation	Farmers	Food vendors	Households	Opportunity	Capability	Motivation
1 <i>Amoah et al. (2007)</i>	West Africa	Sanitary vegetable washing methods			X	X	X		
2 <i>Amoah et al. (2011)</i>	West Africa	Low-cost options to reduce contamination from farm to fork		X	X	X	X		
3 <i>Asirifi et al. (2021)</i>	Ghana	Biochar		X			X		
4 <i>Borghini et al. (2002)</i>	Burkina Faso	Hygiene promotion				X	X		
5 <i>Cuenca-Adame et al. (2001)</i>	Mexico	Wastewater irrigation of onion and decontamination by disinfectant		X			X		
6 <i>Davis et al. (2011)</i>	Tanzania	Informational interventions				X		X	
7 <i>De Buck et al. (2017)</i>	LMICs	Promote handwashing and sanitation behaviour change				X			X
8 <i>Donkor (2009)</i>	Ghana	Food handling			X			X	
9 <i>Googoolie et al. (2020)</i>	Mauritius	Water cress cultivation and washing practices		X			X		
10 <i>Hirai et al. (2016)</i>	Indonesia	Handwashing with soap				X			X
11 <i>Idowu & Rowland (2006)</i>	Nigeria	Personal hygiene			X				
12 <i>Keraiya et al. (2007a)</i>	Ghana	Effect of low-cost wastewater irrigation methods on lettuce contamination		X			X		
13 <i>Keraiya et al. (2007b)</i>	Ghana	Effect of wastewater irrigation cessation on lettuce contamination		X			X		
14 <i>Knopp et al. (2010)</i>	Tanzania	Patterns and risk factors of helminthiasis				X			
15 <i>Kragić Kok et al. (2020)</i>	Ethiopia	Serious games to promote water recycling and hygiene practices				X			X
16 <i>Luchesi et al. (2016)</i>	Brazil	Washing of leafy vegetables with chlorinated water			X ^a		X		
17 <i>Mihrshahi et al. (2009)</i>	Viet Nam	Deworming				X ^b			
18 <i>Morse et al. (2019)</i>	Malawi	Changing WASH and food hygiene practices (using the Risk, Attitude, Norm, Ability, and Self-Regulation (RANAS) method)				X			X

(Continued.)

Table 2 | Continued

Authors	Country of study	Focus	Integration across sectors of the value chain				Integration across the three sources of behaviour		
			Sanitation	Farmers	Food vendors	Households	Opportunity	Capability	Motivation
19 Moya <i>et al.</i> (2019)	Kenya	Barriers to the use of human excreta-derived fertiliser by horticultural export farmers		X			X		X
20 Musa & Akande (2002)	Nigeria	Routine medical examination of school food vendors			X				
21 Ngowi <i>et al.</i> (2017)	Burkina Faso	Health education (using the PHAST method) to control taeniasis and cysticercosis among pig growers		X				X	
22 Nondlazi <i>et al.</i> (2017)	South Africa	Onsite greywater treatment		X ^c			X		
23 Reddi <i>et al.</i> (2016)	India	Hand hygiene				X			
24 Salamandane <i>et al.</i> (2020)	Mozambique	Fresh vegetable handling			X		X		
25 Samadi <i>et al.</i> (2009)	Iran	Fresh vegetable washing with disinfectants			X ^d		X		
26 Sarter & Sarter (2012)	Madagascar	Promoting a culture of food safety (using disgust, social norms and accountability)			X				X
27 Silverman <i>et al.</i> (2014)	Ghana	On-farm wastewater treatment for vegetable irrigation		X			X		
28 Traoré <i>et al.</i> (2020)	Mali	Lettuce washing		X			X		
29 Tripathi <i>et al.</i> (2019)	India	Micro-irrigation with wastewater		X			X		
30 Udert <i>et al.</i> (2016)	South Africa	Nutrient recovery from urine	X				X		
31 Winkler <i>et al.</i> (2017)	Multiple	Sanitation safety planning	X	X			X		
32 Woldetsadik <i>et al.</i> (2017)	Ethiopia	Lettuce washing		X ^e			X		
33 Yihenew <i>et al.</i> (2014)	Ethiopia	Cooperative social organization to control malaria and parasitosis				X		X	

^aSupermarkets/lab.

^bWomen of reproductive age/lab.

^cGardens/lab.

^dMarkets and retail stores/lab.

^eFarms/lab.

implement (Mdee & Harrison 2019) and, instead, adapt them into national standards that account for the local context (Keraita & Drechsel 2015).

Environmental restructuring: certification and access to premium markets

Certifying vegetable safety to give access to premium channels should motivate urban farmers to adopt safe practices (Keraita & Drechsel 2015). Most domestic food safety management systems, however, consist primarily of controls in the formal sector, overlooking the informal sector despite it serving the largest proportion of the population and accounting for the majority of the food disease burden in LMICs (Figuíé 2019; Henson *et al.* 2023).

Effectively developing standards and enforcing controls seems to be a feature of the export-oriented private sector rather than governments (Moya *et al.* 2019). Certified farmers adhere to standards developed with little or no participation from their end (González & Nigh 2005), including record-keeping and controlling the quality of the water used for irrigation and washing. Most smallholder farmers, however, are unable to sustain the high certification costs or emulate these practices, and the food safety of domestic markets remains unaffected (Henson *et al.* 2023).

Third-party certification schemes have recently started to serve domestic markets, adopting requirements more accessible to smallholder farmers. The Ghanaian Ministry of Food and Agriculture and Standards Authority and cooperation partners instigated the Ghana Green Label in 2014 (GhanaGreenLabel.Org 2015). High costs and low domestic demand for certified vegetables limit the supply to a handful of packing houses and high-end outlets in Accra. As of 2022, about 700 farmers are certified nationally, including a few dozen urban and peri-urban farmers. Similarly, since 2016, the supermarket chain SPAR-South Africa has developed Localg.a.p. (for Local Good Agricultural Practices), a programme adapted from the European Farm Assurance Programme GlobalG.A.P. (GLOBALG.A.P./FoodPLUS GmbH 2023), which builds smallholder farmers' capability to comply with good hygiene and safety practices and supply its local supermarkets (SPAR 2021). Both these programmes allow farmers to use a potentially polluted source of irrigation water, provided they supply a water analysis demonstrating they took the necessary steps to ensure the water quality complies with 'accepted [water] standards' (Ghana Green Label) or the WHO (2006) Guidelines (SPAR/Localg.a.p.). Unfortunately, the principles of health-based targets and the multi-barrier approach are overlooked in both cases.

Participatory guarantee systems (PGSs) are an alternative to private third-party certification, consisting of farmers, consumers, and other interested stakeholders agreeing on good practices and working conditions adapted to socio-ecological contexts. Networks of local farmers and consumers assess compliance (Nelson *et al.* 2010), which can help address food safety issues in value chains where quality management is limited (Moustier *et al.* 2023) and distrust exists between stakeholders (Antwi-Agyei *et al.* 2016). More than 1.2 million producers are involved in over 240 PGSs cultivating 900,000 hectares globally, mostly in the Global South (Anselmi & Moura e Castro 2022). PGSs also provide platforms to reduce farmers' costs through collective buying, marketing, and logistics, exchanging good practices and locally adapted seeds (Home *et al.* 2017). PGSs, however, often suffer from a lack of recognition by institutions and struggle to convince consumers to allocate time and resources to maintain the certification process. Consequently, they are subject to tension among stakeholders (Kaufmann *et al.* 2023).

Enablement: developing and testing alternatives to conventional wastewater treatment

Several on-farm techniques can reduce the transmission of pathogens to the crop. Biochar proved ineffective in eliminating pathogens from the soil of farms in northern Ghana (Asirifi *et al.* 2021). On-farm fly ash/lime filters reduce faecal coliforms by 0.5 log units in domestic greywater, which can then be used for irrigation (Nondlazi *et al.* 2017). The focus on technical performance, however, often leaves aside cost efficiency, sustainability, and user acceptability, which shape stakeholders' capability and motivation to use them. The project 'Valorisation of Urine Nutrients in Africa' included a multi-disciplinary team to better account for economic viability, seek authorities' support, and explore users' attitudes towards the reuse of urine and dehydrated faeces, recognising that adoption is the users' decision (Udert *et al.* 2016). Sedimentation ponds and filtration techniques (Silverman *et al.* 2014; Tripathi *et al.* 2019) can help farmers reduce pathogen concentration in irrigation water, especially helminth eggs. Some Ghanaian urban farmers use small ponds but are not interested in maintaining their sedimentation properties (Silverman *et al.* 2014). Moreover, the risk of these ponds becoming reservoirs for malaria-transmitting mosquitoes needs to be considered, particularly with the expansion of *Anopheles stephensi*, which can develop in urban polluted waters (Drechsel *et al.* 2006; Sinka *et al.* 2020).

Various irrigation practices impact pathogen exposure differently. The most effective are drip irrigation systems (especially subsurface ones; see [Tripathi et al. 2019](#)), but most urban farmers are not willing to accommodate the filtration, pressurisation, and distribution equipment, and reduced cropping density they require ([Keraita et al. 2007a](#)). Contrary to the overhead flow and splashing on watering cans, flood and furrow irrigation limit the pathogen transfer to the aerial parts of vegetables but further expose farmers who work barefoot. Sprinklers and spray irrigation produce droplets that transfer pathogens to crops, farmers, passers-by, and farm neighbours ([Keraita et al. 2007a](#)). Stopping irrigation a few days before harvest for pathogens to die off naturally is acceptable under temperate climates where crops are only irrigated every few days ([Drechsel & Keraita 2014](#)), but not in hot climates such as in Ghana, where it results in wilting and revenue loss ([Keraita et al. 2007b](#)). Optimising irrigation to reduce soil moisture during and after the growing season also reduces pathogen concentration ([Cuenca-Adame et al. 2001](#); [Tripathi et al. 2019](#)) but is beyond most farmers' capability. Selecting crops that require cooking increases safety but is unattractive due to their lower market value. Switching to safer water sources is often constrained by piped water being too expensive or landlords forbidding farmers to drill boreholes on their land ([Amoah et al. 2011](#)).

Sanitary vegetable washing methods can also reduce the level of pathogenic contamination. Washing vegetables in basins, with or without the addition of salt or vinegar, is common in markets and households but less effective at removing pathogens than washing under running water or in water with added chlorine, permanganate potassium, or another disinfectant ([Amoah et al. 2007](#); [Googoollee et al. 2020](#); [Traoré et al. 2020](#)). As washing vegetables does not prevent the occupational exposure of farmers and vendors ([Woldetsadik et al. 2017](#)), nor is sufficient to protect consumers ([Amoah et al. 2007](#); [Samadi et al. 2009](#); [Luchesi et al. 2016](#); [Googoollee et al. 2020](#)), it is essential to address both on-farm and post-harvest contamination.

Medical examinations and deworming (chemotherapy) of street food vendors can limit contamination. The effectiveness of medical examination to prevent transmission of helminth eggs depends on whether the drug used targets all relevant parasites ([Idowu & Rowland 2006](#)) and the enforcement is sufficient to ensure results are reported and post-convalescence examinations performed ([Musa & Akande 2002](#)). Regular deworming left high levels of contamination in Nigeria as the limited campaign coverage and insufficient access to safe water and toilets did not prevent re-infection ([Idowu & Rowland 2006](#); [Knopp et al. 2010](#)). In contrast, Sri Lanka has almost eliminated morbidity due to worm infections thanks to mass deworming ([Montresor & Mupfasoni, 2019](#)) and higher access to at least basic urban water and sanitation services ([JMP 2022](#)). Education activities should complement deworming campaigns to address low-risk awareness, which is exacerbated by the infection remaining asymptomatic for extended periods ([Idowu & Rowland 2006](#); [Mihirshahi et al. 2009](#); [Knopp et al. 2010](#)).

Capabilities

The COM-B framework suggests that a physical, psychological (emotional or cognitive) capability deficit within an individual's realm is best addressed by education, training, and enablement interventions.

Education and training: hygiene promotion and training for good practices

Interventions often consider education and training together or even interchangeably. From the perspective of the COM-B framework, however, they serve different purposes and should be differentiated. Education increases knowledge or understanding by providing information, thus improving psychological capability or reflexive motivation. Training imparts skills to increase physical or psychological capabilities ([Michie et al. 2011](#)).

Providing opportunities to adopt safe practices needs to be complemented with education and guidance to increase stakeholders' awareness of microbial risks ([Luchesi et al. 2016](#); [Reddi et al. 2016](#); [Woldetsadik et al. 2017](#); [Salamandane et al. 2020](#)). Education intervention also aims to change their attitudes ([Moya et al. 2019](#)) and motivate them to act ([Kragić Kok et al. 2020](#)).

Indeed, throughout Africa, many urban residents do not link food-borne diseases to a lack of hygiene ([Salamandane et al. 2020](#)), with some claiming that 'stomach aches result from consuming food that the worms do not like' ([Idowu & Rowland 2006](#), p. 163) or that speaking of contaminated vegetables only aims to hurt others' businesses ([Abass et al. 2017](#)).

Education is a major health determinant ([Mihirshahi et al. 2009](#); [Ngowi et al. 2017](#)). Methods, such as PRECEDE (Predisposing, Reinforcing, and Enabling Constructs in Educational Diagnosis and Evaluation) or PHAST (Participatory Hygiene and Sanitation Transformation), help practitioners structure interventions to more effectively increase participants' knowledge appropriation ([Ngowi et al. 2017](#)). However, Information–Education–Communication (IEC) interventions alone offer

mixed results, especially if they misunderstand the context in which the behaviour is to be performed or the recipient's perception of the cost and benefits of the behaviour (De Buck *et al.* 2017). Individuals' choices are not purely rational and based on knowledge (Scott *et al.* 2007). Rather, they rely on shortcuts and cue associations, such as the credibility of the person conveying the information or the attractiveness of the medium (Glanz *et al.* 2008).

The profile and legitimacy of the educator, their interpersonal communication skills, and their attention to the gender sensitivity of some behaviours (Salamandane *et al.* 2020) are as important as their knowledge of the topic (Borghi *et al.* 2002; Winkler *et al.* 2017). Equally important is the quality and availability of educational material, as well as the duration and frequency of the interventions (De Buck *et al.* 2017). Co-designing the content with intended recipients and fieldworkers helps ensure it is appropriate (Amoah *et al.* 2009). Knowledge can be conveyed through plays, world cafés, and road shows that follow the pathogen pathway from farm to fork (Amoah *et al.* 2009), radio spots (Borghi *et al.* 2002), movies, and comic booklets (Ngowi *et al.* 2017) or educational games (Kragić Kok *et al.* 2020). Such community-based education involving experiential, empowering exchanges are more effective than one-way approaches, such as sharing health messaging, especially in the long-term (De Buck *et al.* 2017), but is more challenging to upscale.

Training often completes education interventions (Musa & Akande 2002; Donkor 2009; Salamandane *et al.* 2020). Training strengthens recipients' self-efficacy, i.e., 'I know I can do it' (Ngowi *et al.* 2017) by allowing them to observe and practise in a controlled environment the behaviour to be performed. Training food handlers is a widely applied strategy to increase food safety (FAO 2003). Urban farmers, however, receive little training to address the health risks of wastewater irrigation (Amoah *et al.* 2011). In many cases, extension officers can only offer urban farmers training developed for rural farmers, even though the crops and practices differ, reflecting the lack of recognition and marginality of urban agriculture (Keraita & Drechsel 2015). Even when tailored to recipients' needs, training requires refresher interventions (Salamandane *et al.* 2020) and is challenging to upscale (Grace 2015).

Motivation

The COM-B framework suggests that stakeholder automatic (e.g., wants, needs, emotions) and reflexive (i.e., conscious planning, beliefs about good and bad) motivation is best stimulated by persuasion, incentivisation, or coercion interventions. In addition, education can stimulate reflexive motivation, while automatic motivation responds to environmental restructuring, modelling, and enablement.

Incentivisation and coercion: market incentives

Incentivisation is one of the functions that intervention designers rely on the most to stimulate stakeholders' motivation to improve sanitation services (Moya *et al.* 2019) or increase the adoption of safe practices among farmers and traders (Amoah *et al.* 2011). Designers mostly think of market incentives, such as premium prices for safer vegetables, cash transfers, governmental subsidies, access to microcredit, or other forms of monetary incentives (Karg *et al.* 2010; De Buck *et al.* 2017).

Implementing viable market incentives by stimulating demand is challenging (Amoah *et al.* 2011). Studies suggest that stakeholders' returns need to increase by at least 30% for them to adopt a new practice (Karg *et al.* 2010). This accounts for often underestimated intangible costs, such as cultural resistance, perceived control, opportunity cost, and other psychological costs (Keraita & Drechsel 2015), but not necessarily for quality monitoring costs (Karg *et al.* 2010). Most customers of informal markets are unaware of the risk (Keraita & Drechsel 2015), and not interested in knowing (Nordhagen *et al.* 2022) or paying for safer vegetables. The demand from hotel restaurants, supermarkets, and other premium outlets that serve customers willing to pay such premiums is limited. Decision-makers could stimulate the offer of safe vegetables through coercion through fines, fees, taxes, or other disincentives (Karg *et al.* 2010). Funding the required monitoring of farmers' and food handlers' practices would require policymakers to have a clearer understanding of the return on investment (Keraita & Drechsel 2015).

Other monetary and explicit incentives do not always reach their goal (De Buck *et al.* 2017), as decision-makers sometimes misunderstand stakeholders' values and preferences. The existence, nature, size, and framing of the incentive convey either positive or negative signals to stakeholders. Too small an incentive may suggest that policymakers think the behaviour is unimportant, whereas if too high it can suggest policymakers consider the behaviour too complicated for stakeholders or that they are only self-interested (Gneezy *et al.* 2011). Thus, poorly designed incentives can reduce stakeholders' willingness to adopt a safe practice, even once it is removed. Incentives are more effective when framed in a way that preserves

stakeholders' feelings of trust and autonomy rather than through control or challenging social norms, image concerns, and other social motivations (Gneezy *et al.* 2011) as illustrated by Bahraseman *et al.* (2024).

Non-monetary incentives are also a possibility, including dedicated support from extension officers, 'best farmer' awards or compliance certificates, good media publicity, and new marketing channels (Karg *et al.* 2010; Keraita *et al.* 2014). The study by Morse *et al.* (2019) is one of the few to actually report leveraging implicit incentives that mobilise stakeholders' desire for social recognition to align their practices with their personal values and ethics, autonomy, and control. Experts often underestimate farmers' sensitivity to jeopardising consumer health and assume farmers are more interested in explicit incentives, such as being provided with free or subsidised seeds, fertilisers, or other inputs (Keraita *et al.* 2014).

Environmental restructuring, modelling, and persuasion

Interventions that cater to stakeholders' motivations are seldom articulated in the screened literature. Morse *et al.* (2019) suggests restructuring the social environment by advertising that many households are already performing the recommended hygiene-related behaviours, which are becoming the social norm. Improving farmers' land-tenure security would increase their willingness to adopt safe practices (Amponsah *et al.* 2016). Others leverage traditional beliefs such as 'today' (similar to Buddhist and Hinduist karma) to convince Malagasy food vendors to avoid making their customers sick (Sarter & Sarter 2012). Role models motivating the uptake of good health practices by those around them have also been highlighted (Yihenew *et al.* 2014). Opinion leaders or other individuals respected by their community can champion behaviour change (De Buck *et al.* 2017; Winkler *et al.* 2017). Modelling through serious games can also stimulate the adoption of good practices (Kragić Kok *et al.* 2020). Davis *et al.* (2011) generated personalised information by testing the water quality in homes to persuade households to adopt risk-reducing strategies; Hirai *et al.* (2016) identified the desire to smell nice as a powerful intrinsic motivation to wash hands with soap, while Kragić Kok *et al.* (2020) leveraged social learning through games.

Further reflections

Adopting a robust behavioural framework, such as the COM-B framework, helps intervention designers consider all relevant factors and establish a thorough diagnosis and theory of change instead of relying on assumptions and familiar go-to functions. The framework also helps articulate the functions (e.g., sharing information is not training) and objectives of the intervention (e.g., whether education is to strengthen capabilities and/or reinforce reflexive motivation) and align its content and delivery. It is helpful to consider which practices present the best combination of Acceptability, Practicability, Effectiveness, Affordability, Spill-over or Side-effects, and Equity (the APEASE criteria, see Michie *et al.* 2014) when selecting the ones to promote.

Changing the practices and attitudes of all stakeholders involved in the value chain has coordination and governance implications. Food safety and public health are public goods whose production is one of the sovereign duties of governments. A challenge governments face is to recognise and enable informal businesses to produce and distribute safer vegetables by providing them with information, support, and incentives, rather than just control and sanctions, to comply with minimum food safety standards (Karg *et al.* 2010; Henson *et al.* 2023). Henson *et al.* (2023) also recommended that governments integrate food safety in urban planning and other relevant policies and guide local multisectoral collective action at the municipal level.

LIMITATIONS

This review focuses on LMICs where the multi-barrier approach, as recommended by WHO (2006), remains the accepted procedure. Each country can adopt its own trajectory, especially in relation to enforcing water quality standards and food safety regulations (Drechsel *et al.* 2023). This review examines the contamination of vegetables primarily through irrigation with water contaminated with faecal matter. Faecal contamination introduced through food handlers' poor personal hygiene and food practices was also accounted for. Other contaminants, such as heavy metals and contaminants of emerging concerns that might require source control or other solutions not mentioned here, might also need to be considered. Experiences of adoption of organic farming practices, which also depend on premiums for 'safer' food, have not been included since these topics are extensively covered in the literature.

CONCLUSIONS

This review uses the COM-B behavioural framework to synthesise experiences with different approaches adopted to reduce the health risks associated with the use of wastewater contaminated with faecal pathogens in informal vegetable value chains in LMICs. It shows how most interventions focus on enabling the adoption of good practices by providing stakeholders with opportunities or strengthening their capabilities. Fewer interventions aim to stimulate stakeholder motivation to adopt safe practices, and those that do often simply convey their designers' rationales or struggle to identify sustainable monetary or other explicit incentives. Technical innovation, regulations, education, training, and explicit incentives are helpful and may be necessary but are often not sufficient. This review suggests that intervention designers should consider less-used strategies, including implicit incentivisation, promoting good examples to emulate, and persuasion, which can reinforce and complement technical and management controls and increase their impact. It is equally important not to consider behaviour change approaches as a shortcut to transferring responsibilities to primary stakeholders without investing in the required infrastructure or services. To improve their impact, hygiene and food safety interventions should address the opportunity, capability, and motivation needs of both primary stakeholders (farmers, vendors, and/or consumers) and secondary stakeholders, such as supporting institutions. More participatory approaches that include stakeholders early in the design process would likely increase their buy-in and the sustainability of the intervention. This aligns with the WHO recommendation that governments recognise that public agencies, farmers, food business associations, consumers, and community representatives are interdependent (WHO 2018). We suggest that where the multi-barrier approach remains the best option, urban farmers, vegetable traders, street food vendors, and consumer representatives should be encouraged to contribute to existing multistakeholder platforms to develop a shared vision of the value chain that ensures all their opportunities, capabilities, and motivations are catered for.

ACKNOWLEDGEMENT

The authors would also like to thank Kate Medicott for comments on the manuscript draft.

AUTHOR CONTRIBUTIONS

DG conceptualised the study; carried out the search, analysis, and interpretation; and drafted the review. RES and KVG supervised the study, reviewed the search, and edited the manuscript. PD and BEE reviewed and critiqued the manuscript for clarity and coherence.

FUNDING

This work was supported by the UKRI Engineering and Physical Science Research Council (EPSRC) through a PhD studentship received by the first author as part of the EPSRC Centre for Doctoral Training in Water and Waste Infrastructure and Services Engineered for Resilience (Water-WISER). EPSRC Grant No.: EP/S022066/1.

DATA AVAILABILITY STATEMENT

All relevant data are included in the paper or its Supplementary Information.

CONFLICT OF INTEREST

The authors declare there is no conflict.

REFERENCES

- Abass, K., Ganle, J. K. & Afriyie, K. 2017 'The germs are not harmful': Health risk perceptions among consumers of peri-urban grown vegetables in Kumasi, Ghana. *GeoJournal* **82** (6), 1213–1227. <https://doi.org/10.1007/s10708-016-9747-6>.
- Amoah, P., Drechsel, P., Abaidoo, R. C. & Klutse, A. 2007 Effectiveness of common and improved sanitary washing methods in selected cities of West Africa for the reduction of coliform bacteria and helminth eggs on vegetables. *Tropical Medicine & International Health* **12** (S2), 40–50. <https://doi.org/10.1111/j.1365-3156.2007.01940.x>.
- Amoah, P., Drechsel, P., Schuetz, T., Kranjac-Berisavjevic, G. & Manning-Thomas, N. 2009 From world cafés to road shows: Using a mix of knowledge sharing approaches to improve wastewater use in urban agriculture. *Knowledge Management for Development Journal* **5** (3), 246–262. <https://doi.org/10/bqgsbh>.

- Amoah, P., Keraita, B., Akple, M., Drechsel, P., Abaidoo, R. C. & Konradsen, F. 2011 *Low-Cost Options for Reducing Consumer Health Risks from Farm to Fork Where Crops are Irrigated with Polluted Water in West Africa*. Research Report 141; IWMI Research Report, p. 45. International Water Management Institute (IWMI). <https://doi.org/10.5337/2011.201>.
- Amponsah, O., Vigre, H., Braimah, I., Schou, T. W. & Abaidoo, R. C. 2016 *The policy implications of urban open space commercial vegetable farmers' willingness and ability to pay for reclaimed water for irrigation in Kumasi, Ghana*. *Heliyon* 2 (3), e00078. <https://doi.org/10.1016/j.heliyon.2016.e00078>.
- Anselmi, S. & Moura e Castro, F. 2022 Participatory guarantee systems in 2021. In: *The World of Organic Agriculture. Statistics and Emerging Trends 2022*, pp. 170–177. Research Institute of Organic Agriculture FiBL, Frick, and IFOAM – Organics International. Available from: <https://www.fibl.org/fileadmin/documents/shop/1344-organic-world-2022.pdf>.
- Antwi-Agyei, P., Peasey, A., Biran, A., Bruce, J. & Ensink, J. 2016 *Risk perceptions of wastewater use for urban agriculture in Accra, Ghana*. *PLoS One* 11 (3), e0150603. <https://doi.org/10.1371/journal.pone.0150603>.
- Asirifi, I., Kaetzl, K., Werner, S., Saba, C. K. S., Abagale, F. K., Amoah, P. & Marschner, B. 2021 *Pathogen and heavy metal contamination in urban agroecosystems of northern Ghana: Influence of biochar application and wastewater irrigation*. *Journal of Environmental Quality* 50 (5), 1097–1109. <https://doi.org/10.1002/jeq2.20260>.
- Bahraseman, S. E., Firoozzare, A., Jaghdani, T. J. & Dourandish, A. 2024 *Intervention strategies for the safe use of semi-treated wastewater by Iranian farmers: An approach for safe food production in the circular economy*. *NJAS: Impact in Agricultural and Life Sciences* 96 (1), 2335376. <https://doi.org/10.1080/27685241.2024.2335376>.
- Borghi, J., Guinness, L., Ouedraogo, J. & Curtis, V. 2002 *Is hygiene promotion cost-effective? A case study in Burkina Faso*. *Tropical Medicine and International Health* 7 (11), 960–969. <https://doi.org/10.1046/j.1365-3156.2002.00954.x>.
- Cuenca-Adame, E., Riestra-Díaz, D., Pérez-Mangas, J. M. & Echeagaray-Alemán, A. 2001 *Use of sewage water and control of pathogenic organisms in the production of onions*. *Agrociencia* 35 (3), 255–265.
- Davis, J., Pickering, A. J., Rogers, K., Mamuya, S. & Boehm, A. B. 2011 *The effects of informational interventions on household water management, hygiene behaviors, stored drinking water quality, and hand contamination in peri-urban Tanzania*. *American Journal of Tropical Medicine and Hygiene* 84 (2), 184–191. <https://doi.org/10.4269/ajtmh.2011.10-0126>.
- De Buck, E., Hans Van, R., Hannes, K., Govender, T., Naidoo, S., Avau, B., Axel Vande, V., Musekiwa, A., Lutje, V., Cargo, M., Hans-Joachim, M., Vandekerckhove, P. & Young, T. 2017 *Approaches to promote handwashing and sanitation behaviour change in low- and middle-income countries: A mixed method systematic review*. *Campbell Systematic Reviews* 13 (1), 1–447. Social Science Premium Collection. <https://doi.org/10.4073/csr.2017.7>.
- Denyer, D. & Tranfield, D. 2009 *Producing a systematic review*. In: *The Sage Handbook of Organizational Research Methods* (Buchanan, D. A. & Bryman, A., eds.). Sage Publications Ltd., Thousand Oaks, CA, pp. 671–689.
- Donkor, E. S. 2009 *Application of the WHO five keys of food safety to improve food handling practices of food vendors in a poor resource community in Ghana*. *East African Journal of Public Health* 6 (2), 148–151.
- Drechsel, P. & Keraita, B. 2014 *Irrigated Urban Vegetable Production in Ghana: Characteristics, Benefits and Risk Mitigation*, 2nd ed. International Water Management Institute (IWMI). Available from: <http://csirspace.csirgh.com:80/handle/123456789/217>.
- Drechsel, P., Graefe, S., Sonou, M. & Cofie, O. O. 2006 *Informal Irrigation in Urban West Africa: An Overview*. Research Report 102; IWMI Research Report, Issue 44572, p. 34. International Water Management Institute. <https://doi.org/10.22004/ag.econ.44572>.
- Drechsel, P., Qadir, M. & Galibourg, D. 2022 *The WHO guidelines for safe wastewater use in agriculture: A review of implementation challenges and possible solutions in the Global South*. *Water* 14 (6), 864. <https://doi.org/10.3390/w14060864>.
- Drechsel, P., Marjani Zadeh, S. & Salcedo, F. P. 2023 *Water Quality in Agriculture: Risks and Risk Mitigation*. FAO & IWMI. <https://doi.org/10.4060/cc7340en>.
- FAO 2003 *The Informal Food Sector. Municipal Support Policies for Operators: A Briefing Guide for Mayors, City Executive and Urban Planners in Developing Countries and Countries in Transition*. FAO. Available from: <https://www.fao.org/documents/card/en?details=15578940-48bc-5b7e-962e-1ba70feef66/>.
- FAO 2012 *Growing Greener Cities in Africa: First Status Report on Urban and Peri-Urban Horticulture in Africa*. Food and Agriculture Organization of the United Nations, Rome, Italy, p. 116.
- Figuié, M. 2019 *Food safety risks*. In: *Food Systems at Risk. New Trends and Challenges* (Dury, S., Bendjebbar, P., Hainzelin, E., Giordano, T. & Bricas, N., eds.). FAO, CIRAD, pp. 115–118. <https://doi.org/10.19182/agritrop/00080>.
- GhanaGreenLabel.org 2015 *Ghana green label*. Available from: <https://www.ghanagreenlabel.org/about/history/>.
- Glanz, K., Rimer, B. K. & Viswanath, K. 2008 *Health Behavior and Health Education: Theory, Research, and Practice*, 4th ed. Jossey-Bass, San Francisco, CA.
- GLOBALG.A.P./FoodPLUS GmbH 2023 *Localg.a.p. PFA*. Globalgap.Org. Available from: https://www.globalgap.org/uk_en/what-we-do/globalg.a.p.-certification/localg.a.p.
- Gneezy, U., Meier, S. & Rey-Biel, P. 2011 *When and why incentives (don't) work to modify behavior*. *Journal of Economic Perspectives* 25 (4), 191–210. <https://doi.org/10.1257/jep.25.4.191>.
- González, A. A. & Nigh, R. 2005 *Smallholder participation and certification of organic farm products in Mexico*. *Journal of Rural Studies* 21 (4), 449–460. <https://doi.org/10.1016/j.jrurstud.2005.08.004>.
- Googoollee, A. M., Takooree, S. D., Goburdhun, D. & Neetoo, H. (2020). *Characterizing the cultivation practices and microbiological quality of watercress*. *Journal of Agriculture and Food Research* 2. <https://doi.org/10.1016/j.jafr.2020.100057>.

- Grace, D. 2015 Food safety in low and middle income countries. *International Journal of Environmental Research and Public Health* **12** (9), 10490–10507. <https://doi.org/10.3390/ijerph120910490>.
- Henson, S., Jaffee, S. & Wang, S. 2023 *New Directions for Tackling Food Safety Risks in the Informal Sector of Developing Countries (Report)*, p. 96. ILRI. Available from: <https://cgspace.cgiar.org/handle/10568/130652>.
- Hirai, M., Graham, J. P., Mattson, K. D., Kelsey, A., Mukherji, S. & Cronin, A. A. 2016 Exploring determinants of handwashing with soap in Indonesia: A quantitative analysis. *International Journal of Environmental Research and Public Health* **13** (9). <https://doi.org/10.3390/ijerph13090868>.
- Hoffmann, S., Devleeschauwer, B., Aspinall, W., Cooke, R., Corrigan, T., Havelaar, A., Angulo, F., Gibb, H., Kirk, M., Lake, R., Speybroeck, N., Torgerson, P. & Hald, T. 2017 Attribution of global foodborne disease to specific foods: Findings from a World Health Organization structured expert elicitation. *PLoS One* **12** (9), e0183641. <https://doi.org/10.1371/journal.pone.0183641>.
- Home, R., Bouagnimbeck, H., Ugas, R., Arbenz, M. & Stolze, M. 2017 Participatory guarantee systems: Organic certification to empower farmers and strengthen communities. *Agroecology and Sustainable Food Systems* **41** (5), 526–545. <https://doi.org/10.1080/21683565.2017.1279702>.
- Hutton, G. & Haller, L. 2004 *Evaluation of the Costs and Benefits of Water and Sanitation Improvements at the Global Level*. Technical Report WHO/SDE/WSH/04.04; Technical Documents, p. 87. World Health Organization.
- Idowu, O. A. & Rowland, S. A. 2006 Oral fecal parasites and personal hygiene of food handlers in Abeokuta, Nigeria. *African Health Sciences* **6** (3), 160–164. <https://doi.org/10.4314/ahs.v6i3.6945>.
- Jaffee, S., Henson, S., Unnevehr, L., Grace, D. & Cassou, E. 2019 *The Safe Food Imperative: Accelerating Progress in Low- and Middle-Income Countries*. World Bank, Washington, DC. <https://doi.org/10.1596/978-1-4648-1345-0>.
- JMP 2022 Household Data – 2022 – Service Levels. WHO & UNICEF Joint Monitoring Program. Available from: <https://washdata.org/>.
- Karg, H., Drechsel, P., Amoah, P., Jeitler, R., 2010 Facilitating the adoption of food-safety interventions in the street-food sector and on farms. In: *Wastewater Irrigation and Health: Assessing and Mitigating Risk in Low-Income Countries*, (Drechsel, P., Scott, C. A., Raschid-Sally, L., Redwood, M. & Bahri, A., eds.) IWMI, IDRC, Earthscan, Colombo, Sri Lanka, pp. 319–335.
- Kaufmann, S., Hruschka, N., Vildoza, L. & Vogl, C. R. 2023 Alternative food networks in Latin America – Exploring PGS (participatory guarantee systems) markets and their consumers: A cross-country comparison. *Agriculture and Human Values* **40** (1), 193–216. <https://doi.org/10.1007/s10460-022-10347-w>.
- Keraita, B. & Drechsel, P. 2015 *Consumer Perceptions of Fruit and Vegetable Quality: Certification and Other Options for Safeguarding Public Health in West Africa (Working Paper 164; IWMI Working Paper, p. 32)*. International Water Management Institute (IWMI). <https://doi.org/10.5337/2015.215>.
- Keraita, B., Konradsen, F., Drechsel, P. & Abaidoo, R. C. 2007a Effect of low-cost irrigation methods on microbial contamination of lettuce irrigated with untreated wastewater. *Tropical Medicine and International Health* **12** (Suppl. 2), 15–22. <https://doi.org/10.1111/j.1365-3156.2007.01937.x>.
- Keraita, B., Konradsen, F., Drechsel, P. & Abaidoo, R. C. 2007b Reducing microbial contamination on wastewater-irrigated lettuce by cessation of irrigation before harvesting. *Tropical Medicine & International Health* **12**, 8–14. <https://doi.org/10.1111/j.1365-3156.2007.01936.x>.
- Keraita, B., Amoah, P., Drechsel, P. & Akple, M. 2014 Enhancing adoption of food safety measures in urban vegetable production and marketing systems. In: *International Symposium on Urban and Peri-Urban Horticulture in the Century of Cities: Lessons, Challenges, Opportunities* 1021, pp. 391–399.
- Knopp, S., Mohammed, K. A., Stothard, J. R., Khamis, I. S., Rollinson, D., Marti, H. & Utzinger, J. 2010 Patterns and risk factors of helminthiasis and anemia in a rural and a peri-urban community in Zanzibar, in the context of helminth control programs. *PLoS Neglected Tropical Diseases* **4** (5), e681. <https://doi.org/10.1371/journal.pntd.0000681>.
- Kragić Kok, D., Bisschops, I., Knoop, L., Tulu, L., Kujawa-Roeleveld, K., Masresha, N. & Houtkamp, J. 2020 Game over or play again? Deploying games for promoting water recycling and hygienic practices at schools in Ethiopia. *Environmental Science and Policy* **111**, 83–90. <https://doi.org/10.1016/j.envsci.2020.05.016>.
- Liverpool-Tasie, L. S. O., Wineman, A., Young, S., Tambo, J., Vargas, C., Reardon, T., Adjognon, G. S., Porciello, J., Gathoni, N., Bizikova, L., Galië, A. & Celestin, A. 2020 A scoping review of market links between value chain actors and small-scale producers in developing regions. *Nature Sustainability* **3** (10), 799–808. <https://doi.org/10.1038/s41893-020-00621-2>.
- Luchesi, C., Dambrós, B. P., Gelinski, J. M. L. N., Frighetto, M. & Baratto, C. M. 2016 Leafy vegetables free of coliforms and parasites after washing in chlorine treated water. *Revista de Ciências Farmacéuticas Básica e Aplicada* **37** (1). Available from: <https://rcfba.fcfar.unesp.br/index.php/ojs/article/view/5>.
- Mdee, A. & Harrison, E. 2019 Critical governance problems for farmer-led irrigation: Isomorphic mimicry and capability traps. *Water Alternatives* **12** (1), 30–45.
- Michie, S. & West, R. 2013 Behaviour change theory and evidence: A presentation to government. *Health Psychology Review* **7** (1), 1–22. <https://doi.org/10.1080/17437199.2011.649445>.
- Michie, S., van Stralen, M. M. & West, R. 2011 The behaviour change wheel: A new method for characterising and designing behaviour change interventions. *Implementation Science* **6** (1), 42. <https://doi.org/10.1186/1748-5908-6-42>.
- Michie, S., Atkins, L. & West, R. 2014 *The Behaviour Change Wheel: A Guide to Designing Interventions*. Available from: <http://www.vlebooks.com/vleweb/product/openreader?id=Exeter&isbn=9781912141081>.

- Mihrshahi, S., Casey, G. J., Montresor, A., Phuc, T. Q., Thach, D. T. C., Tien, N. T. & Biggs, B. A. 2009 The effectiveness of 4 monthly albendazole treatment in the reduction of soil-transmitted helminth infections in women of reproductive age in Viet Nam. *International Journal for Parasitology* **39** (9), 1037–1043. <https://doi.org/10.1016/j.ijpara.2009.01.013>.
- Montresor, A. & Mupfasoni, D. 2019 Achievements of the deworming programme in Sri Lanka. *The Lancet Global Health* **7** (9). e1156–e1157. [https://doi.org/10.1016/S2214-109X\(19\)30309-2](https://doi.org/10.1016/S2214-109X(19)30309-2).
- Morse, T., Chidziwisano, K., Tilley, E., Malolo, R., Kumwenda, S., Musaya, J. & Cairncross, S. 2019 Developing a contextually appropriate integrated hygiene intervention to achieve sustained reductions in diarrheal diseases. *Sustainability* **11** (17). <https://doi.org/10.3390/su11174656>.
- Moustier, P., Holdsworth, M., Anh, D. T., Seck, P. A., Renting, H., Caron, P. & Bricas, N. 2023 The diverse and complementary components of urban food systems in the Global South: Characterization and policy implications. *Global Food Security* **36**, 100663. <https://doi.org/10.1016/j.gfs.2022.100663>.
- Moya, B., Parker, A. & Sakrabani, R. 2019 Challenges to the use of fertilisers derived from human excreta: The case of vegetable exports from Kenya to Europe and influence of certification systems. *Food Policy* **85**, 72–78. <https://doi.org/10.1016/j.foodpol.2019.05.001>.
- Musa, O. I. & Akande, T. M. 2002 Routine medical examination of food vendors in secondary schools in Ilorin. *Nigerian Journal of Medicine: Journal of the National Association of Resident Doctors of Nigeria* **11** (1), 9–12.
- Nelson, E., Gómez Tovar, L., Schwentesius Rindermann, R. & Gómez Cruz, M. Á. 2010 Participatory organic certification in Mexico: An alternative approach to maintaining the integrity of the organic label. *Agriculture and Human Values* **27** (2), 227–237. <https://doi.org/10.1007/s10460-009-9205-x>.
- Ngowi, H., Ozbolt, I., Millogo, A., Dermauw, V., Some, T., Spicer, P., Jervis, L. L., Ganaba, R., Gabriel, S., Dorny, P. & Carabin, H. 2017 Development of a health education intervention strategy using an implementation research method to control taeniasis and cysticercosis in Burkina Faso. *Infectious Diseases of Poverty* **6**. <https://doi.org/10.1186/s40249-017-0308-0>.
- Nondlazi, S., Ngqwala, N. P., Zuma, B. & Tandlich, R. 2017 Investigating the viability and performance of the pilot scale fly ash/lime filter tower for onsite greywater treatment. *Desalination and Water Treatment* **91**, 349–364. <https://doi.org/10.5004/dwt.2017.21461>.
- Nordhagen, S., Lee, J., Onuigbo-Chatta, N., Okoruwa, A., Monterrosa, E., Lambertini, E. & Pelto, G. H. 2022 ‘Sometimes you get good ones, and sometimes you get not-so-good ones’: Vendors’ and consumers’ strategies to identify and mitigate food safety risks in urban Nigeria. *Foods* **11** (2). <https://doi.org/10.3390/foods11020201>.
- Okoruwa, A. & Onuigbo-Chatta, N. 2021 Review of food safety policy in Nigeria. *Journal of Law, Policy and Globalization* **110** (0), 57. <https://doi.org/10.7176/JLPG/110-07>.
- Orsini, F., Kahane, R., Nono-Womdim, R. & Gianquinto, G. 2013 Urban agriculture in the developing world: A review. *Agronomy for Sustainable Development* **33** (4), 695–720. <https://doi.org/10/f5csvj>.
- Prain, G., Simon, D., Halliday, J. & Drechsel, P. 2022 Investment priorities for research and innovation in urban agri-food systems: Toward more resilient cities in the Global South. *Frontiers in Sustainable Food Systems* **6**, 965011. <https://doi.org/10.3389/fsufs.2022.965011>.
- Raj, S. J., Wang, Y., Yakubu, H., Robb, K., Siesel, C., Green, J., Kirby, A., Mairinger, W., Michiel, J., Null, C., Perez, E., Roguski, K. & Moe, C. L. 2020 The SaniPath exposure assessment tool: A quantitative approach for assessing exposure to fecal contamination through multiple pathways in low resource urban settlements. *PLoS One* **15** (6), e0234364. <https://doi.org/10.1371/journal.pone.0234364>.
- Raschid-Sally, L. & Jayakody, P. 2008 *Drivers and Characteristics of Wastewater Agriculture in Developing Countries: Results From A Global Assessment*. Research Report 127, p. 39. IWMI.
- Reddi, S., Kumar, R. N., SubbaRao, G. M., Rao, M. V. V. & Sudershan, R. V. 2016 Bacteriological quality of drinking water at point of use and hand hygiene of primary food preparers: Implications for household food safety. *Journal of Water Sanitation and Hygiene for Development* **6** (2), 224–230. <https://doi.org/10.2166/washdev.2016.184>.
- Salamandane, C., Fonseca, F., Afonso, S., Lobo, M. L., Antunes, F. & Matos, O. 2020 Handling of fresh vegetables: Knowledge, hygienic behavior of vendors, public health in Maputo markets, Mozambique. *International Journal of Environmental Research and Public Health* **17** (17). <https://doi.org/10.3390/ijerph17176302>.
- Samadi, N., Abadian, N., Bakhtiari, D., Fazeli, M. R. & Jamalifar, H. 2009 Efficacy of detergents and fresh produce disinfectants against microorganisms associated with mixed raw vegetables. *Journal of Food Protection* **72** (7), 1486–1490. <https://doi.org/10.4315/0362-028X-72.7.1486>.
- Sarter, G. & Sarter, S. 2012 Promoting a culture of food safety to improve hygiene in small restaurants in Madagascar. *Food Control* **25** (1), 165–171. <https://doi.org/10.1016/j.foodcont.2011.10.023>.
- Scott, B. E., Lawson, D. W. & Curtis, V. 2007 Hard to handle: Understanding mothers’ handwashing behaviour in Ghana. *Health Policy and Planning* **22** (4), 216–224. <https://doi.org/10.1093/heapol/czm014>.
- Seleman, A., Gabriellson, S., Mbwette, T. S. A. & Kimwaga, R. 2020 Drivers of unhygienic desludging practices in unplanned settlements of Dar es Salaam, Tanzania. *Journal of Water Sanitation and Hygiene for Development* **10** (3), 512–526. <https://doi.org/10.2166/washdev.2020.179>.
- Shoushtarian, F. & Negahban-Azar, M. 2020 Worldwide regulations and guidelines for agricultural water reuse: A critical review. *Water* **12** (4), 971. <https://doi.org/10.3390/w12040971>.
- Silverman, A. I., Akrong, M. O., Drechsel, P. & Nelson, K. L. 2014 On-farm treatment of wastewater used for vegetable irrigation: Bacteria and virus removal in small ponds in Accra, Ghana. *Journal of Water Reuse and Desalination* **4** (4), 276–286. <https://doi.org/10.2166/wrd.2014.010>.

- Sinka, M. E., Pironon, S., Massey, N. C., Longbottom, J., Hemingway, J., Moyes, C. L. & Willis, K. J. 2020 A new malaria vector in Africa: Predicting the expansion range of *Anopheles stephensi* and identifying the urban populations at risk. *Proceedings of the National Academy of Sciences* **117** (40), 24900–24908. <https://doi.org/10.1073/pnas.2003976117>.
- SPAR 2021 SPAR – Small Scale Farmers get Food Safety Accreditation. SPAR. Available from: <https://www.spar.co.za/Lifestyle/View/SPAR-Cares/Small-scale-farmers-get-food-safety-accreditation>.
- Thebo, A. L., Drechsel, P., Lambin, E. F. & Nelson, K. L. 2017 A global, spatially-explicit assessment of irrigated croplands influenced by urban wastewater flows. *Environmental Research Letters* **12** (7), 074008. <https://doi.org/10.1088/1748-9326/aa75d1>.
- Traoré, S., Cornelius, E. W., Samaké, F., Essilfie, G., Babana, A. H., Bengaly, M., Koné, A., Cissé, H. & Acheampong, M. A. 2020 Efficiency of common washing treatments in reducing microbial levels on lettuce in Mali. *West African Journal of Applied Ecology* **28** (2), 64–74. <https://doi.org/10.4314/wajae.v28i2>.
- Tripathi, V. K., Rajput, T. B. S., Patel, N. & Nain, L. 2019 Impact of municipal wastewater reuse through micro-irrigation system on the incidence of coliforms in selected vegetable crops. *Journal of Environmental Management* **251**. <https://doi.org/10.1016/j.jenvman.2019.109532>.
- Udert, K. M., Etter, B. & Gounden, T. 2016 Promoting sanitation in South Africa through nutrient recovery from urine. *Gaia* **25** (3), 194–196. Scopus. <https://doi.org/10.14512/gaia.25.3.12>.
- WHO 2006 *Guidelines for the Safe Use of Wastewater Excreta and Greywater*, Vol. 1–4, 3rd edn. World Health Organization, Geneva, Switzerland.
- WHO 2015 *WHO Estimates of the Global Burden of Foodborne Diseases: Foodborne Disease Burden Epidemiology Reference Group 2007–2015*. World Health Organization, Geneva, Switzerland. Available from: <https://apps.who.int/iris/handle/10665/199350>.
- WHO 2018 *Regional Framework for Action on Food Safety in the Western Pacific*. WHO Regional Office for the Western Pacific; WHO IRIS. Available from: <http://iris.wpro.who.int/handle/10665.1/14084>.
- WHO 2022 *Sanitation Safety Planning: Step-by-Step Risk Management for Safely Managed Sanitation Systems*, 1st edn. World Health Organization. Available from: <https://apps.who.int/iris/bitstream/handle/10665/364586/9789240062887-eng.pdf?sequence=1>.
- Winkler, M. S., Jackson, D., Sutherland, D., Payden, Lim, J. M. U., Srikantaiah, V., Fuhrmann, S. & Medlicott, K. 2017 Sanitation safety planning as a tool for achieving safely managed sanitation systems and safe use of wastewater. *WHO South-East Asia Journal of Public Health* **6** (2), 34–40. <https://doi.org/10.4103/2224-3151.213790>.
- Wodnik, B. K., Freeman, M. C., Ellis, A. S., Awino Ogutu, E., Webb Girard, A. & Caruso, B. A. 2018 Development and application of novel caregiver hygiene behavior measures relating to food preparation, handwashing, and play environments in rural Kenya. *International Journal of Environmental Research and Public Health* **15** (9). Article 9. <https://doi.org/10.3390/ijerph15091994>.
- Woldetsadik, D., Drechsel, P., Keraita, B., Itanna, F., Erko, B. & Gebrekidan, H. 2017 Microbiological quality of lettuce (*Lactuca sativa*) irrigated with wastewater in Addis Ababa, Ethiopia and effect of green salads washing methods. *International Journal of Food Contamination* **4** (1). <https://doi.org/10.1186/s40550-017-0048-8>.
- Wolf, J., Johnston, R. B., Ambelu, A., Arnold, B. F., Bain, R., Brauer, M., Brown, J., Caruso, B. A., Clasen, T., Colford, J. M., Mills, J. E., Evans, B., Freeman, M. C., Gordon, B., Kang, G., Lanata, C. F., Medlicott, K. O., Prüss-Ustün, A., Troeger, C., Boisson, S. & Cumming, O. 2023 Burden of disease attributable to unsafe drinking water, sanitation, and hygiene in domestic settings: A global analysis for selected adverse health outcomes. *The Lancet* **401** (10393), 2060–2071. [https://doi.org/10.1016/S0140-6736\(23\)00458-0](https://doi.org/10.1016/S0140-6736(23)00458-0).
- Yihenew, G., Adamu, H. & Petros, B. 2014 The impact of cooperative social organization on reducing the prevalence of malaria and intestinal parasite infections in Awramba, a rural community in South Gondar, Ethiopia. *Interdisciplinary Perspectives on Infectious Diseases* **2014**. <https://doi.org/10.1155/2014/378780>.

First received 10 January 2024; accepted in revised form 14 July 2024. Available online 25 July 2024