

## Spatial and temporal assessment of groundwater quality and hydrogeochemical processes in Urmia Lake Basin, Iran

Mohammadreza Pashaeifar<sup>a</sup>, Reza Dehghanzadeh<sup>b</sup>, Mohammad Ebrahim Ramazani<sup>a,\*</sup>,  
Omid Rafieyan<sup>a</sup> and Arezoo Nejaei<sup>a</sup>

<sup>a</sup> Department of Environmental Engineering, Tabriz Branch, Islamic Azad University, Tabriz, Iran

<sup>b</sup> Health and Environment Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

\*Corresponding author. E-mail: ramazani@iaut.ac.ir

### ABSTRACT

The evaluation of groundwater quality and geochemistry have an important role in the management of water resources in arid and semi-arid regions. In the present study, the spatio-temporal assessment of groundwater quality and hydrogeochemical processes, as well as, statistical analyses in the Azarshahr and Ajabshir planes located in the Urmia Lake basin were investigated. The results revealed that within six years (from 2014 to 2019), the value of total hardness was higher than the permissible level and the quality of groundwater for drinking was very hard and fresh in both planes. In 2019, 84 and 67% of the samples fell within the range of good to poor groundwater quality in the Azarshahr and Ajabshir planes, respectively. The temporal assessment with the help of water quality index values for both planes revealed a good groundwater quality for the Azarshahr plane and a good to poor groundwater quality for the Ajabshir plane. However, deterioration of water quality was observed in both planes from 2014 to 2019. The level of water quality for irrigation was better in the Azarshahr plane than in the Ajabshir plane, due to the presence of fewer salty sites. In addition, Ca–Mg–HCO<sub>3</sub> and rock dominance were identified as hydrochemical facies and the controlling factor in the groundwater of both planes, respectively. Multivariate statistical analyses indicated both natural and anthropogenic sources (such as weathering, fertilizers, and wastewater) for hydrochemical parameters. It was suggested to develop a comprehensive regulation to control the entry of pollutants into the groundwater of the study area.

**Key words:** drinking and irrigation, groundwater quality, hydrogeochemistry, Urmia Lake

### HIGHLIGHTS

- The level of water quality for drinking and irrigation was better in the Azarshahr plane than in the Ajabshir plane.
- Ca–Mg–HCO<sub>3</sub> and rock dominance were identified as hydrochemical facies and the controlling factors in the groundwater of the study area.
- The groundwater resources in the Urmia Lake Basin were found to be threatened by different sources of pollution.

### INTRODUCTION

In the last decades, water demand has been significantly increased due to water shortage caused by global warming and climate changes (Fallahati *et al.* 2020). The groundwater is considered as one of the most valuable water resources for drinking, agricultural, and industrial purposes (Raju *et al.* 2015; Jalili *et al.* 2018; Li *et al.* 2019a, 2019b). According to Qasemi *et al.* (2018, 2019), agricultural purposes rank first and account for more than 80% of groundwater consumption in Iran. In arid and semi-arid areas, the high demand for groundwater resources leads to reduction in the water table of groundwater aquifers (Abbasnia *et al.* 2019). Moreover, pollution of water resources is rapidly increasing due to urbanization, industrialization, and agricultural activities (Rao *et al.* 2017; Haghazadeh & Saneie 2019; Rahman *et al.* 2020). The anthropogenic sources including industrial effluents, municipal sewage, overuse of pesticides and fertilizers, and mining operations along with the natural sources such as weathering of parent materials are the main sources of water quality reduction (Taghipour *et al.* 2012; Brindha *et al.* 2017; Ezugwu *et al.* 2019; Onwuka *et al.* 2019; Haghazadeh *et al.* 2020). Groundwater quality is mainly affected by hydrological properties, geological structures, fertilizers, industrial pollution, sewage, and sand river mining (Mirzabeygi *et al.* 2017; Yousefi *et al.* 2017; Abboud 2018; Rezaei & Hassani 2018; Adimalla & Qian 2019). Therefore, control and assessment of groundwater quality are considered to be essential.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Licence (CC BY-NC-ND 4.0), which permits copying and redistribution for non-commercial purposes with no derivatives, provided the original work is properly cited (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Various studies have focused on the hydrochemical monitoring, geochemical processing, sources of pollution, and applicability of water for agricultural and drinking purposes (Alaya *et al.* 2014; Li *et al.* 2016; Abd El-Aziz 2017; Rezaei *et al.* 2018; Zhang *et al.* 2018; Berhe 2020). The sources of contamination can be determined by multivariate statistical analyses such as correlation matrix and principal component analysis (PCA). These approaches are useful in finding the complicated relationships among the physicochemical parameters of water resources (Wang *et al.* 2014; Ustaoglu *et al.* 2020). Control of groundwater quality can be done using the water quality index (WQI), which has been widely used by numerous researchers (Sahu & Sikdar 2008; Bhuiyan *et al.* 2016; Sethy *et al.* 2016; Rawat & Singh 2018; Sharmin *et al.* 2020). One of the advantages of WQI is its capability of overall assessment of physicochemical parameters (Abtahi *et al.* 2015). So far, different types of methods, such as Wilcox, Piper, and Gibbs diagrams, have been used for groundwater quality assessment in previous studies (Choramin *et al.* 2015; Li *et al.* 2016; Narsimha & Sudarshan 2017; Kawo & Karuppanan 2018; Sahoo & Khaoash 2020). The suitability for agricultural purposes can be assessed using Wilcox diagrams (Khoramabadi Shams *et al.* 2014). Furthermore, hydrochemical facies and factors for controlling the groundwater chemistry are evaluated by Piper and Gibbs diagrams, respectively (Srinivasamoorthy *et al.* 2014; Koffi *et al.* 2017). Urmia Lake, as the second great salt lake in the world, is one of the most important and valuable ecosystems located in the Northwest of Iran (Hassanzadeh *et al.* 2012). In the last 10 years, the surface water table in the Urmia Lake has dropped dramatically due to climate changes, continuous droughts, development of cultivated areas, land-use changes, and construction of dams (Ghale *et al.* 2018). The hypersalinity of the Urmia Lake has highlighted the critical role of groundwater in supplying the water required for drinking, agricultural, and industrial purposes in the urban areas around the lake. In the last decade, the groundwater resources in the Urmia Lake Basin have been threatened by chemical pollution due to the significant increase of agricultural activities and population around the lake. Therefore, it is vital to assess the spatio-temporal status of the quality of the multi-purpose groundwater in this area.

Considering the condition of the Urmia lake in recent years, which is threatening the groundwater quality due to the continuous drying of the salty lake, it is of great importance to investigate the groundwater quality and determine the effect of the salt water of the lake in the area. Accordingly, the present study aimed to investigate groundwater quality variation in recent years in a large area. In addition, the temporal status of groundwater for drinking purpose together with identification of sources of the major ions in the Eastern lake is well documented for the first time.


In this study, the spatio-temporal variations of groundwater quality have been evaluated using the WQI and diagram-based methods. In addition, the sources of chemical parameters were determined using the multivariate statistical techniques.

The objectives of this study were: (1) to evaluate the spatio-temporal variations of the hydrochemical parameters of groundwater, (2) to determine the status of the groundwater quality for drinking and irrigation purposes, and (3) to assess the hydrogeochemical process of groundwater in the study area. This study was carried out in Azarshahr and Ajabshir cities in 2019.

## MATERIALS AND METHODS

### Characteristics of the study area

The study area is located in East-Azerbaijan province in the North-West of Iran. The groundwater wells in the Azarshahr plane, with an area of 168 km<sup>2</sup>, and the Ajabshir plane, with an area of 144 km<sup>2</sup>, in the eastern part of Urmia Lake were selected for sampling (Figure 1). The study area has a semi-arid and cold climate. In the Azarshahr and Ajabshir planes, the average annual temperatures are 13 °C and 12 °C and the average annual precipitation is 221.2 and 300 mm, respectively (Docheshmeh Gorgij & Asghari Moghaddam 2017; Gorgij *et al.* 2017; Moghaddam *et al.* 2018). The mean water table depths of the aquifers are in the range of 0.7–29.8 m in the Azarshahr plane and 0.5–37.7 m in the Ajabshir plane (Rezaei & Gurdak 2020). In recent years, the groundwater aquifers in the study area have been faced with over-extraction due to overpopulation, industrialization, and overuse of fertilizers. Since the study area is located in the vicinity of the hypersaline region of the Urmia Lake, where there is an imbalance between precipitation and evaporation, water consumption is completely dependent on groundwater resources (Gorgij *et al.* 2019). For example, in the Azarshahr plane, 100% of the water for drinking, domestic, and industrial purposes and 80% of the water for agricultural purposes are supplied by the groundwater resources (Gorgij *et al.* 2017).



**Figure 1** | Geographic location of the study area with the sampling sites in the Azarshahr and Ajabshir planes in the vicinity of Urmia Lake.

### Sampling and analysis

Totally, 24 and 15 groundwater samples (39 samples) were collected from the Azarshahr and Ajabshir planes in October 2019, respectively. For temporal assessment of the groundwater quality, the data on hydrochemical parameters of the sampling sites from 2014 to 2018 was obtained from the Regional Water Company of East-Azerbaijan. Water of wells was pumped for 10 min to remove the impact of stagnant water (Adimalla 2019). 250 ml polyethylene bottles were washed with deionized water and used for collecting the samples. The collected samples were stored at 4 °C in a refrigerator and transferred to the laboratory to be studied for hydrochemical parameters. Accordingly, the groundwater hydrochemical parameters such as pH, electrical conductivity (EC), total dissolved solids (TDS), total hardness as CaCO<sub>3</sub> (TH), cations [potassium (K<sup>+</sup>), sodium (Na<sup>+</sup>), magnesium (Mg<sup>2+</sup>), calcium (Ca<sup>2+</sup>)], and anions [sulfate (SO<sub>4</sub><sup>2-</sup>), nitrate (NO<sub>3</sub><sup>-</sup>), chloride (Cl<sup>-</sup>), bicarbonate (HCO<sub>3</sub><sup>-</sup>), fluoride (F<sup>-</sup>)] were analyzed. After sampling, pH and EC were detected *in situ* using a portable pH/EC meter. Moreover, TDS was calculated by multiplying a factor (0.64) to EC. HCO<sub>3</sub><sup>-</sup>, SO<sub>4</sub><sup>2-</sup>, and Cl<sup>-</sup> were determined by titrimetric methods, and NO<sub>3</sub><sup>-</sup> and F<sup>-</sup> were obtained using a spectrophotometer. Na<sup>+</sup> and K<sup>+</sup> were analyzed by flame atomic absorption spectrometry, while TH, Mg<sup>2+</sup> and Ca<sup>2+</sup> were measured by EDTA titrimetric method. All the samples were tested in two replicates to ensure the accuracy of the analyses.

### Assessment of groundwater quality

#### Groundwater quality for drinking

WQI is a useful index-based method for overall assessment of groundwater quality for drinking purposes (Ramakrishnaiah *et al.* 2009; Khan & Jhariya 2017). To calculate the WQI, weights of 2–5 were assigned to the hydrochemical parameters of groundwater based on their significance in groundwater quality. In the first step, the relative weights ( $W_i$ ) were computed using (Equation (1)).

$$W_i = \frac{w_i}{\sum_{i=1}^n w_i} \quad (1)$$

The values of weights and relative weights for each parameter are presented in Table 1 (Khalid 2019).

In the second step, the quality rating scale ( $Q_i$ ) for each hydrochemical parameter was calculated by the ratio of the measured parameters' values to the relevant standard value proposed by WHO (2011)

**Table 1** | Standard values, weights, and relative weights for each hydrochemical parameter

Elements	WHO (2011) (mg/L)	Weight ( $w_i$ )	Relative weight ( $W_i$ )
pH	6.5–8.5	4	0.087
EC	1,500	4	0.087
TDS	1,000	4	0.087
SO <sub>4</sub> <sup>2-</sup>	250	5	0.109
NO <sub>3</sub> <sup>-</sup>	45	5	0.109
Cl <sup>-</sup>	250	4	0.087
HCO <sub>3</sub> <sup>-</sup>	500	3	0.065
F <sup>-</sup>	1.5	5	0.109
TH	100	3	0.065
K <sup>+</sup>	12	2	0.043
Na <sup>+</sup>	200	3	0.065
Mg <sup>2+</sup>	50	2	0.043
Ca <sup>2+</sup>	75	2	0.043
		$\sum w_i = 46$	$\sum W_i = 1$

according to (Equation (2)).

$$Q_i = \frac{C_i}{S_i} \times 100 \quad (2)$$

where,  $C_i$  and  $S_i$  are the measured value and the standard value of parameters, respectively.

In the final step, the water quality sub-index and the WQI are obtained using (Equation (3)) and (Equation (4)).

$$SI_i = Q_i \times W_i \quad (3)$$

$$WQI = \sum_{i=1}^n SI_i \quad (4)$$

The WQI was classified into five classes as presented in Table 2 (Saleem *et al.* 2016).

### Groundwater quality for irrigation

To assess the suitability of the groundwater for agricultural purposes, Wilcox diagrams were plotted using the value of sodium absorption ratio (SAR) versus EC (Wilcox 1955). SAR can be calculated by (Equation (5)).

$$SAR = \frac{Na^+}{\sqrt{\frac{Mg^{2+} + Ca^{2+}}{2}}} \quad (5)$$

**Table 2** | The WQI classification for drinking purposes

Classification	Groundwater quality
WQI < 25	Excellent groundwater
25 < WQI < 50	Good groundwater
50 < WQI < 75	Poor groundwater
75 < WQI < 100	Very poor groundwater
WQI > 100	Unsuitable groundwater

The groundwater quality for irrigation was categorized into four classes as presented in Table 3 (Fallahati *et al.* 2020).

### Controlling factors, types and facies of groundwater

Piper diagram is a useful method to determine the hydrochemical type and facies of groundwater (Patolia & Sinha 2017; Li *et al.* 2019a, 2019b; Sahoo & Khaoash 2020). According to the Piper diagram, the groundwater was classified into four dominants for major cations and anions including (a) Mg type, (b) Ca type, (c) Na & K type, (d) no dominant type, (e) SO<sub>4</sub> type, (f) HCO<sub>3</sub> type, (g) Cl type. Finally, the groundwater was classified into six faces as Face 1: Ca–Mg–HCO<sub>3</sub>, Face 2: Na–Cl, Face 3: mixed Ca–Na–HCO<sub>3</sub>, Face 4: Mixed Ca–Mg–Cl, Face 5: Ca–Cl, and Face 6: Na– HCO<sub>3</sub> (Shakerkhatibi *et al.* 2019; Sharmin *et al.* 2020).

The factors for controlling the groundwater chemistry were determined using two semilog digrams designed by Gibbs (1970). The Gibbs diagrams were plotted by TDS versus Cl/Cl + HCO<sub>3</sub> and Na/Na + Ca representing the anions and cations, respectively. According to the Gibbs diagrams, the controlling factors were grouped into three dominances including rock dominance, precipitation dominance, and evaporation dominance (He & Wu 2019; Heydarirad *et al.* 2019).

### Statistical analysis

For both Azarshahr and Ajabshir planes, multivariate statistical analyses were performed to determine the interactions among the hydrochemical parameters (Sethy *et al.* 2016; Sangsefidi *et al.* 2017). The Kolmogorov-Smirnov test was carried out to assess the normality of the value of the parameters (Haghnazar *et al.* 2021). Moreover, Pearson's correlation matrix was applied to identify the relationship among the parameters. The correlation coefficients in the range of 0.5–0.7 and greater than 0.7 were considered as moderate and significant, respectively (Mehraein *et al.* 2020). PCA was conducted to reduce the dimension of the dataset and determine the sources of hydrochemical parameters of the groundwater (Mosaferei *et al.* 2014). In the PCA calculation method, the rotation of principal components was carried out using Varmix rotation for better interpretation.

## RESULTS AND DISCUSSION

### Temporal assessment of the groundwater chemistry

Temporal assessment of the hydrochemical parameters of the groundwater in the Azarshahr and Ajabshir planes during six years (from 2014 to 2019) was studied and the values of the parameters were compared with the permissible values for drinking water proposed by WHO (2011). The mean values of the hydrochemical parameters in both planes have been presented in Table 4. The pH range of 7.2–7.5 in the groundwater of the Azarshahr and Ajabshir planes represented the alkaline groundwater in nature for six years. In addition, the dominant cations were found to be in the order of Ca<sup>2+</sup> > Na<sup>2+</sup> > Mg<sup>2+</sup> > K<sup>+</sup>, and the dominant anions had the order of HCO<sub>3</sub><sup>-</sup> > SO<sub>4</sub><sup>2-</sup> ≈ Cl<sup>-</sup> > NO<sub>3</sub><sup>-</sup> > F<sup>-</sup> for both planes from 2014 to 2019. In the Azarshahr plane, the mean values of EC, TDS, NO<sub>3</sub><sup>-</sup>, HCO<sub>3</sub><sup>-</sup>, TH, Mg<sup>2+</sup>, and Ca<sup>2+</sup> increased in the second three years (2017–2019) compared to the first three years (2014–2016).

The mean values of all the parameters, except for TH, were found to be less than the permissible levels provided by WHO (2011). The values of TH were higher than the permissible levels in the whole six years. The minimum and maximum values of TH were 1.88 and 2.07 times greater than the permissible values in 2014 and 2019, respectively. In the Ajabshir plane, the mean values of EC, TDS, TH, Na<sup>2+</sup>, and Ca<sup>2+</sup> had a similar order and an increasing trend in the second three years (2017–2019). The mean values of all the parameters, except for TH and Ca<sup>2+</sup>, exceeded the permissible levels. The maximum and minimum values of TH were

**Table 3** | The Wilcox classification for groundwater quality

Class	Water quality for irrigation
C1S1	Sweet
C1S2-C2S2-C2S1	A little salty
C1S3-C2S3-C3S1-C3S2-C3S3	Salty
C1S4-C2S4-C3S4-C4S4-C4S3-C4S2 -C4S1	Very salty

**Table 4** | Hydrochemical parameters in the groundwater of Azarshahr (Az) and Ajabshir (Aj) planes


Year	2014		2015		2016		2017		2018		2019		WHO (2011)
	Az	Aj	Az	Aj	Az	Aj	Az	Aj	Az	Aj	Az	Aj	
pH	7.5	7.4	7.4	7.2	7.5	7.5	7.4	7.4	7.2	7.2	7.2	7.2	6.5–8.5
EC	506.5	556.5	496.1	771.4	470.6	585.7	541.8	862.0	516.2	664.6	520.0	799.8	1,500
TDS	324.1	356.2	313.7	493.7	301.2	374.9	346.7	551.7	330.4	425.4	332.7	511.9	1,000
SO <sub>4</sub> <sup>2-</sup>	48.7	32.9	34.3	54.9	30.7	32.9	47.8	53.4	39.2	42.2	40.2	47.7	250
NO <sub>3</sub> <sup>-</sup>	1.7	4.9	9.1	21.6	2.7	14.2	11.5	18.4	7.7	9.6	10.4	20.9	45
Cl <sup>-</sup>	39.7	35.3	40.6	52.2	30.8	27.9	38.5	53.2	38.2	36.2	39.4	56.0	250
HCO <sub>3</sub> <sup>-</sup>	190.3	258.6	193.8	324.8	200.5	274.0	208.4	375.1	205.5	296.8	204.0	341.3	500
F <sup>-</sup>	0.4	0.6	0.3	0.4	0.3	0.3	0.4	0.4	0.3	0.3	0.4	0.4	1.5
TH	188.0	232.0	182.0	286.3	178.2	214.6	209.0	331.1	200.5	232.2	207.2	310.7	100
K <sup>+</sup>	3.4	3.2	3.5	3.9	3.1	5.5	3.3	3.2	2.8	4.8	2.8	3.1	12
Na <sup>+</sup>	29.9	26.7	29.2	48.1	25.7	35.4	27.5	45.2	26.3	44.0	23.9	41.7	200
Mg <sup>2+</sup>	11.8	16.1	12.7	24.1	12.4	14.3	14.3	25.1	13.9	16.6	14.7	19.9	50
Ca <sup>2+</sup>	54.5	58.1	51.5	74.9	50.5	62.4	59.7	90.6	57.0	65.2	58.3	91.3	75

2.14 and 3.1 times greater than the permissible values in 2016 and 2019, respectively. For Ca<sup>2+</sup>, the highest value was recorded as 1.2 times greater than the permissible value in 2016 and 2017.

The high concentration of TH, due to anthropogenic sources and geological structures, in both planes has led to deterioration of the groundwater quality in recent years. According to the mean values obtained for TH and TDS, the groundwater quality for drinking was found to be very hard (Sawyer & McCarthy 1967) and fresh (Freeze & Cherry 1979) for both planes.


### Groundwater quality assessment

The results of groundwater quality in the Azarshahr and Ajabshir planes in the six years are illustrated in Figure 2. The mean values of WQI in the six years were lower in the Azarshahr plane than in the Ajabshir plane, indicating better quality of water for drinking. The minimum and maximum mean values of WQI in the Azarshahr and Ajabshir planes were 38.8/45.65 and 47.68/65.37, respectively, showing good groundwater quality in the Azarshahr plane and good to poor groundwater quality in the Ajabshir plane.

**Figure 2** | Groundwater quality (WQI) in the Azarshahr and Ajabshir planes.

The regression lines for the data showed that the mean WQI values for both planes increased from 2014 to 2019, which indicated the decline of water quality over time.

The spatial distributions of WQI in the Azarshahr and Ajabshir planes in 2019 have been presented in Figure 3. The results indicated a very poor groundwater quality in two sampling points at the Eastern and Southern parts of



**Figure 3** | Spatial distribution of WQI: (a) Azarshahr plane, (b) Ajabshir plane.

the Azarshahr and Ajabshir planes, respectively. In the Azarshahr plane, 84, 8, and 8% of the samples had a good, poor, and very poor groundwater quality, respectively. However, in the Ajabshir plane, 20, 67, and 13% of the samples showed a good, poor, and very poor groundwater quality, respectively. These hotspots located in urban and agricultural areas of the planes revealed that anthropogenic activities were responsible for the poor and very poor quality of groundwater for drinking.

SAR and EC are two major indicators for sodium hazard and salinity of groundwater. The Wilcox diagram is an effective and simple method to assess the applicability of groundwater for irrigation by combining alkalinity and salinity (Li *et al.* 2018). The Wilcox diagrams for both planes were plotted for 2019. As shown in Figure 4(a), most of the sampling points in the Azarshahr plane fall within C2S1 and C2S2, indicating a slightly salty and almost good quality for irrigation. However, four samples categorized under C3S1 and C3S2 groups proved to be salty but usable for agriculture. According to Figure 4(b), half of the samples under C2S1 and C2S2 groups and the rest of those under C3S1 and C3S2 groups could be known as slightly salty and salty, respectively. The results showed that the level of water quality for irrigation was better in the Azarshahr plane than in the Ajabshir plane due to having fewer salty sites. The sampling sites with  $EC > 750$  in the Azarshahr plane (13, 22, 23, and 24) and Ajabshir plane (1, 2, 3, 4, 5, 6, 10, and 12), located in the eastern and southern parts of the planes, were found to be affected by leaching of soluble salts caused by anthropogenic activities (Wu *et al.* 2014; Adimalla 2019).


### Geochemical characterization of groundwater

The Piper diagrams were plotted based on the dominant dissolved cations and anions to determine the hydrochemical evolution of the groundwater parameters and overall effects of reactions among major ions (Patolia & Sinha 2017; Li *et al.* 2018). According to Figure 5, the results of the Piper trilinear diagrams for both Azarshahr and Ajabshir planes were similar in 2019. Cations fell within zones B and D, indicating calcium and no dominant groundwater types, respectively. Anions belonged to zones F and D, representing bicarbonate and no dominant groundwater types, respectively. Finally, most of the sampling sites fell into zone 1, demonstrating a Ca–Mg–HCO<sub>3</sub> groundwater face. However, two sampling sites in the Azarshahr plane and one sampling site in the Ajabshir plane fell within zone 4, indicating the mixed Ca–Mg–Cl groundwater types.

The Gibbs diagrams were employed to analyze the mechanisms of evolution in the surface water and groundwater. The Gibbs diagrams for the Azarshahr and Ajabshir planes in 2019 were plotted according to governing cations, anions, and TDS values (Figures 6 and 7, respectively). As shown in Figures 6 and 7, all the sampling sites in both planes fell within the middle part of both cation-based and anion-based diagrams, indicating rock dominance or water-rock interaction group. The results indicated that mineral weathering was a main factor in controlling the ionic composition and the chemistry of groundwater.

### Multivariate statistical analyses

Correlation analysis and PCA were conducted to determine the relationship between physicochemical variables and potential sources of minerals in the groundwater. The Pearson's correlation coefficients and PCA results,



**Figure 4** | Wilcox diagram of the sampling sites (a) Azarshahr plane, (b) Ajabshir plane.






Figure 5 | Piper diagram of the sampling sites: (a) Azarshahr plane, (b) Ajabshir plane.




Figure 6 | The Gibbs diagrams for the sampling sites of the Azarshahr plane.




Figure 7 | The Gibbs diagrams for the sampling sites of the Ajabshir plane.

with eigenvalues greater than 1 for the Azarshahr and Ajabshir planes, were obtained based on the data related to 2019 (Tables 5 and 6). In the Azarshahr plane, the results indicated four principal components describing 84.4% of the total variance. The first component (PC1) included  $\text{SO}_4$ , Cl, Ca, Mg, EC, TDS, and TH. According to the correlation matrix, Ca had significant and moderate correlations with  $\text{SO}_4$  (0.893) and Cl (0.549), respectively, indicating the dissolution of gypsum ( $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ ) and  $\text{CaCl}_2$  salt (Wu *et al.* 2009; Thilagavathi *et al.* 2017). The correlation of EC and TDS with other parameters in PC1 could be explained by the dominance of these ions caused by mineralization in the aquifers (Khalid 2019). The second component (PC2) was specified by Cl, Na, K, EC, and pH. The positive correlation between Na and K (0.702) could be related to weathering and mineralization (Bhuiyan *et al.* 2016). The third component (PC3) included K, Na, F, and  $\text{NO}_3$ . The correlation of Na and  $\text{NO}_3$  (0.521) with F and K (0.562) might be related to agricultural activities (Kawo & Karuppanan 2018). The fourth component (PC4), which was specified by TDS and  $\text{HCO}_3$ , demonstrated the geogenic/anthropogenic sources. Analysis of the Ajabshir plane showed three principal components specifying 88.4% of the total variance. According to the results shown in Table 6, the first component included  $\text{HCO}_3$ , Ca, Mg, K, EC, TDS, and TH, which illustrated that EC and TDS were controlled by other parameters in PC1. The moderate correlation between  $\text{HCO}_3$  and Mg (0.596) as well as the significant correlation of Ca and  $\text{HCO}_3$  (0.902) with  $\text{HCO}_3$  and K (0.829) indicated silicate and carbonate weathering and dissolution (Sethy *et al.* 2016; Wu *et al.* 2017). The second component (PC2) was specified by  $\text{SO}_4$ , Cl, F, and Na. The significant correlation between Na and Cl (0.932) showed the effects of the Urmia Lake on the dissolution of NaCl salt and interaction of halite rocks and groundwater (Muhammad & Husam 2011; Li *et al.* 2013). In addition, Na had a significant correlation with  $\text{SO}_4$ , indicating mirabilite ( $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$ ) due to a salt lake (Gao *et al.* 2007). Na showed a moderate correlation with F (0.589), indicating NaF salt due to the discharge of industrial wastewater from the Ajabshir Industrial City to the groundwater sources (Masindi & Abiye 2018). The third principal component (PC3) included  $\text{NO}_3$  and pH, which demonstrated anthropogenic activities such as the use of fertilizers and domestic wastewater (Gao *et al.* 2007).

### Spatial distribution of ions

The spatial distribution plots for cations and anions were extracted using the measured concentration in each sampling site for 2019 and presented in Figures S1 and S2. In the Azarshahr plane, the most concentrated areas of ions, except for Na, were located in the Eastern part. Concentration of Na in the western of the plane indicated the effects of Urmia Lake. The higher concentration of  $\text{SO}_4$ , Cl, Ca, and Mg in the western part of the plane demonstrated the types of geological layers based on the multivariate statistical results. The spatial distributions of K, F, and  $\text{NO}_3$  indicated the higher rate of agricultural activities in the eastern part of the plane. Mixed geogenic and anthropogenic activities in the eastern parts of the plane were found to be responsible for the concentrated areas of  $\text{HCO}_3$ .

In the Ajabshir plane, the concentrated areas of  $\text{HCO}_3$ , Ca, Mg, K were located in the southern and eastern parts, indicating higher amounts of weathering and natural sources in these areas. The higher concentrations of Na, Cl, and  $\text{SO}_4$  in the western part of the plane were related to Urmia Lake. The increase of Na and F concentrations in the north-west areas was due to the industrial activities in Ajabshir Industrial City. Moreover,  $\text{NO}_3$  showed a high concentration in the western and southern areas due to the anthropogenic activities in the agricultural and urban areas.

### CONCLUSION

In this study, the groundwater in the Urmia Lake Basin was subjected to spatial and temporal assessments. For this purpose, the Azarshahr and Ajabshir planes, located at the east of the lake, were selected. The results indicated that the groundwater in nature was alkaline due to the pH range of 7.2–7.5. In both planes, the value of TH was greater than the standard level within six years (2014 to 2019). The mean values of TH and TDS in the six years showed that the groundwater quality for drinking was very hard and fresh in both planes. The results of WQI revealed a good groundwater quality in the Azarshahr plane and a good to poor groundwater quality in the Ajabshir plane for drinking purposes. In all, 84% and 20% of the samples fell within the range of good groundwater quality in the Azarshahr and Ajabshir planes, respectively. Furthermore, 8 and 67% of the samples showed poor groundwater quality, and 8% and 13% of the samples proved to be in the range of very poor groundwater quality in the Azarshahr and Ajabshir planes, respectively. Using the Wilcox diagrams, the groundwater quality for irrigation purposes was found to be slightly salty and salty in both planes. The hydrochemical facies of the

**Table 5** | Correlation matrix for groundwater in Azarshahr and Ajabshir planes

<b>Azarshahr</b>	<b>HCO<sub>3</sub></b>	<b>NO<sub>3</sub></b>	<b>SO<sub>4</sub></b>	<b>Cl</b>	<b>F</b>	<b>Ca</b>	<b>Mg</b>	<b>Na</b>	<b>K</b>	<b>EC</b>	<b>PH</b>	<b>TDS</b>	<b>TH</b>
HCO <sub>3</sub>	1.000												
NO <sub>3</sub>	0.307	1.000											
SO <sub>4</sub>	0.085	0.292	1.000										
Cl	0.378	0.121	0.766	1.000									
F	0.016	0.546	0.324	0.174	1.000								
Ca	0.035	0.381	0.893	0.549	0.177	1.000							
Mg	0.048	-0.323	0.439	0.421	-0.170	0.238	1.000						
Na	0.260	0.521	0.238	0.355	0.468	0.047	-0.223	1.000					
K	-0.166	0.351	0.536	0.351	0.562	0.317	-0.121	0.702	1.000				
EC	0.135	0.298	0.859	0.658	0.252	0.811	0.185	0.335	0.515	1.000			
pH	-0.187	-0.072	-0.450	-0.580	-0.059	-0.321	0.190	-0.416	-0.455	-0.591	1.000		
TDS	0.652	0.344	0.787	0.878	0.216	0.648	0.353	0.381	0.310	0.716	-0.518	1.000	
TH	0.010	0.028	0.706	0.500	-0.031	0.669	0.749	-0.189	-0.034	0.502	-0.085	0.499	1.000
<b>Ajabshir</b>	<b>HCO<sub>3</sub></b>	<b>NO<sub>3</sub></b>	<b>SO<sub>4</sub></b>	<b>Cl</b>	<b>F</b>	<b>Ca</b>	<b>Mg</b>	<b>Na</b>	<b>K</b>	<b>EC</b>	<b>PH</b>	<b>TDS</b>	<b>TH</b>
HCO <sub>3</sub>	1.000												
NO <sub>3</sub>	0.245	1.000											
SO <sub>4</sub>	0.457	0.478	1.000										
Cl	0.562	0.481	0.871	1.000									
F	-0.259	0.303	0.413	0.450	1.000								
Ca	0.902	0.344	0.720	0.690	-0.122	1.000							
Mg	0.596	0.097	0.059	0.247	-0.160	0.310	1.000						
Na	0.634	0.498	0.789	0.932	0.589	0.715	0.161	1.000					
K	0.829	0.142	0.620	0.519	-0.326	0.882	0.287	0.594	1.000				
EC	0.930	0.400	0.732	0.816	0.047	0.942	0.506	0.829	0.830	1.000			
pH	-0.476	-0.626	-0.008	-0.072	0.259	-0.372	-0.163	-0.249	-0.314	-0.362	1.000		
TDS	0.930	0.400	0.732	0.816	0.047	0.942	0.506	0.829	0.830	1.000	-0.362	1.000	
TH	0.961	0.314	0.626	0.657	-0.168	0.952	0.586	0.650	0.846	0.962	-0.367	0.962	1.000

**Table 6** | Principal components (PCs) of groundwater in Azarshahr and Ajabshir planes

	Azarshahr plane				Ajabshir plane		
	PC1	PC2	PC3	PC4	PC1	PC2	PC3
% Total variance	43.1	20.6	11.5	9.2	60.6	17.5	10.3
HCO <sub>3</sub>	0.013	0.057	0.078	0.974	0.952	0.106	0.218
NO <sub>3</sub>	0.076	-0.024	0.866	0.280	0.050	0.449	0.836
SO <sub>4</sub>	0.859	0.411	0.275	0.020	0.448	0.805	0.042
Cl	0.610	0.557	0.030	0.396	0.460	0.873	0.088
F	0.077	0.074	0.836	-0.075	-0.404	0.852	-0.026
Ca	0.815	0.233	0.273	-0.018	0.878	0.337	0.184
Mg	0.734	-0.212	-0.345	0.085	0.596	-0.115	0.040
Na	-0.150	0.556	0.600	0.223	0.497	0.775	0.219
K	0.120	0.643	0.580	-0.285	0.894	0.152	0.052
EC	0.656	0.575	0.257	0.047	0.869	0.449	0.196
pH	-0.101	-0.914	0.032	-0.144	-0.294	0.168	-0.909
TDS	0.617	0.433	0.192	0.613	0.869	0.449	0.196
TH	0.930	-0.090	-0.105	0.025	0.944	0.238	0.163

groundwater was determined as Ca-Mg-HCO<sub>3</sub>, and weathering of minerals was specified as a controlling factor in groundwater chemistry in both planes. The results of correlation analysis, PCA, and spatial distribution demonstrated that geological structures and agricultural and anthropogenic activities were the main sources affecting the parameters in the Azarshahr plane. Moreover, weathering and dissolution, salts of Urmia Lake, fertilizers, and wastewater were found to be the main sources affecting the parameters in the Ajabshir plane.

#### DATA AVAILABILITY STATEMENT

Data cannot be made publicly available; readers should contact the corresponding author for details.

#### REFERENCES

- Abbasnia, A., Yousefi, N., Mahv, A. H., Nabizadeh, R., Radfard, M., Yousefi, M. & Alimohammadi, M. 2019 Evaluation of groundwater quality using water quality index and its suitability for assessing water for drinking and irrigation purposes: case study of Sistan and Baluchistan province (Iran). *Human and Ecological Risk Assessment: An International Journal* **25** (4), 988–1005.
- Abboud, I. A. 2018 Geochemistry and quality of groundwater of the Yarmouk basin aquifer, North Jordan. *Environmental Geochemistry and Health* **40** (4), 1405–1435.
- Abd El-Aziz, S. H. 2017 Evaluation of groundwater quality for drinking and irrigation purposes in the North-Western area of Libya (Aligeelat). *Environmental Earth Sciences* **76** (4), 1–17.
- Abtahi, M., Golchinpour, N., Yaghmaeian, K., Rafiee, M., Jahangiri-rad, M., Keyani, A. & Saeedi, R. 2015 A modified drinking water quality index (DWQI) for assessing drinking source water quality in rural communities of Khuzestan Province, Iran. *Ecological Indicators* **53**, 283–291.
- Adimalla, N. 2019 Groundwater quality for drinking and irrigation purposes and potential health risks assessment: a case study from semi-arid region of South India. *Exposure and Health* **11** (2), 109–123.
- Adimalla, N. & Qian, H. 2019 Groundwater quality evaluation using water quality index (WQI) for drinking purposes and human health risk (HHR) assessment in an agricultural region of Nanganur, south India. *Ecotoxicology and Environmental Safety* **176**, 153–161.
- Alaya, M. B., Saïdi, S., Zemni, T. & Zargouni, F. 2014 Suitability assessment of deep groundwater for drinking and irrigation use in the Djefara aquifers (Northern Gabes, South-Eastern Tunisia). *Environmental Earth Sciences* **71** (8), 3387–3421.
- Berhe, B. A. 2020 Evaluation of groundwater and surface water quality suitability for drinking and agricultural purposes in Kombolcha town area, eastern Amhara region, Ethiopia. *Applied Water Science* **10** (6), 1–17.
- Bhuiyan, M. A. H., Bodrud-Doza, M., Islam, A. T., Rakib, M. A., Rahman, M. S. & Ramanathan, A. L. 2016 Assessment of groundwater quality of Lakshimpur district of Bangladesh using water quality indices, geostatistical methods, and multivariate analysis. *Environmental Earth Sciences* **75** (12), 1–23.

- Brindha, K., Pavelic, P., Sotoukee, T., Douangsavanh, S. & Elango, L. 2017 Geochemical characteristics and groundwater quality in the Vientiane plain, Laos. *Exposure and Health* **9** (2), 89–104.
- Choramin, M., Safaei, A., Khajavi, S., Hamid, H. & Abozari, S. 2015 Analyzing and studying chemical water quality parameters and its changes on the base of Schuler, Wilcox and Piper diagrams (project: Bahamanshir River). *WALIA Journal* **31**, 22–27.
- Docheshmeh Gorgij, A. & Asghari Moghaddam, A. 2017 Assessment of saltwater pollution in Azarshahr plain groundwater, Iran: conjugating GALDIT method and geostatistics. *Journal of Water Sciences Research* **9** (1), 63–72.
- Ezugwu, C. K., Onwuka, O. S., Egbueri, J. C., Unigwe, C. O. & Ayejoto, D. A. 2019 Multi-criteria approach to water quality and health risk assessments in a rural agricultural province, Southeast Nigeria. *HydroResearch* **2**, 40–48.
- Fallahati, A., Soleimani, H., Alimohammadi, M., Dehghanifard, E., Askari, M., Eslami, F. & Karami, L. 2020 Impacts of drought phenomenon on the chemical quality of groundwater resources in the central part of Iran – application of GIS technique. *Environmental Monitoring and Assessment* **192** (1), 1–19.
- Freeze, R. A. & Cherry, J. A. 1979 *Groundwater*. Prentice-Hall, Englewood Cliffs, New Jersey, p. 604.
- Gao, X., Wang, Y., Li, Y. & Guo, Q. 2007 Enrichment of fluoride in groundwater under the impact of saline water intrusion at the salt lake area of Yuncheng basin, Northern China. *Environmental Geology* **53** (4), 795–803.
- Ghale, Y. A. G., Altunkaynak, A. & Unal, A. 2018 Investigation anthropogenic impacts and climate factors on drying up of Urmia Lake using water budget and drought analysis. *Water Resources Management* **32** (1), 325–337.
- Gibbs, R. J. 1970 Mechanisms controlling world water chemistry. *Science* **170** (3962), 1088–1090.
- Gorgij, A. D., Kisi, O. & Moghaddam, A. A. 2017 Groundwater budget forecasting, using hybrid wavelet-ANN-GP modelling: a case study of Azarshahr Plain, East Azerbaijan, Iran. *Hydrology Research* **48** (2), 455–467.
- Gorgij, A. D., Wu, J. & Moghadam, A. A. 2019 Groundwater quality ranking using the improved entropy TOPSIS method: a case study in Azarshahr plain aquifer, East Azerbaijan, Iran. *Human and Ecological Risk Assessment: An International Journal* **25** (1–2), 176–190.
- Haghnazar, H. & Saneie, M. 2019 Impacts of pit distance and location on river sand mining management. *Modeling Earth Systems and Environment* **5** (4), 1463–1472.
- Haghnazar, H., Sangsefidi, Y., Mehraein, M. & Tavakol-Davani, H. 2020 Evaluation of infilling and replenishment of river sand mining pits. *Environmental Earth Sciences* **79** (14), 1–18.
- Haghnazar, H., Pourakbar, M., Mahdavianpour, M. & Aghayani, E. 2021 Spatial distribution and risk assessment of agricultural soil pollution by hazardous elements in a transboundary river basin. *Environmental Monitoring and Assessment* **193** (4), 1–17.
- Hassanzadeh, E., Zarghami, M. & Hassanzadeh, Y. 2012 Determining the main factors in declining the Urmia Lake level by using system dynamics modeling. *Water Resources Management* **26** (1), 129–145.
- He, S. & Wu, J. 2019 Hydrogeochemical characteristics, groundwater quality, and health risks from hexavalent chromium and nitrate in groundwater of Huanhe Formation in Wuqi county, Northwest China. *Exposure and Health* **11** (2), 125–137.
- Heydarirad, L., Mosafieri, M., Pourakbar, M., Esmailzadeh, N. & Maleki, S. 2019 Groundwater salinity and quality assessment using multivariate statistical and hydrogeochemical analysis along the Urmia Lake coastal in Azarshahr plain, North West of Iran. *Environmental Earth Sciences* **78** (24), 1–16.
- Jalili, D., RadFard, M., Soleimani, H., Nabavi, S., Akbari, H., Akbari, H., Kavosi, A., Abasnia, A. & Adibzadeh, A. 2018 Data on nitrate–nitrite pollution in the groundwater resources a Sonqor plain in Iran. *Data in Brief* **20**, 394–401.
- Kawo, N. S. & Karuppanan, S. 2018 Groundwater quality assessment using water quality index and GIS technique in Modjo River Basin, Central Ethiopia. *Journal of African Earth Sciences* **147**, 300–311.
- Khalid, S. 2019 An assessment of groundwater quality for irrigation and drinking purposes around brick kilns in three districts of Balochistan province, Pakistan, through water quality index and multivariate statistical approaches. *Journal of Geochemical Exploration* **197**, 14–26.
- Khan, R. & Jhariya, D. C. 2017 Groundwater quality assessment for drinking purpose in Raipur City, Chhattisgarh using water quality index and geographic information system. *Journal of the Geological Society of India* **90** (1), 69–76.
- Khoramabadi Shams, G., Yusefzadeh, A., Godini, H., Hoseinzadeh, E. & Khoshgoftar, M. 2014 Water quality zoning based on water quality index and Wilcox index using geographic information system. *Jundishapur Journal of Health Sciences* **6** (3), e21724.
- Koffi, K. V., Obuobie, E., Banning, A. & Wohnlich, S. 2017 Hydrochemical characteristics of groundwater and surface water for domestic and irrigation purposes in Veve catchment, Northern Ghana. *Environmental Earth Sciences* **76** (4), 185.
- Li, P., Wu, J. & Qian, H. 2013 Assessment of groundwater quality for irrigation purposes and identification of hydrogeochemical evolution mechanisms in Pengyang County, China. *Environmental Earth Sciences* **69** (7), 2211–2225.
- Li, P., Wu, J. & Qian, H. 2016 Hydrochemical appraisal of groundwater quality for drinking and irrigation purposes and the major influencing factors: a case study in and around Hua County, China. *Arabian Journal of Geosciences* **9** (1), 15.
- Li, P., He, S., He, X. & Tian, R. 2018 Seasonal hydrochemical characterization and groundwater quality delineation based on matter element extension analysis in a paper wastewater irrigation area, Northwest China. *Exposure and Health* **10** (4), 241–258.
- Li, P., He, X. & Guo, W. 2019a Spatial groundwater quality and potential health risks due to nitrate ingestion through drinking water: a case study in Yan'an City on the Loess Plateau of Northwest China. *Human and Ecological Risk Assessment: an International Journal* **25** (1–2), 11–31.

- Li, P., He, X., Li, Y. & Xiang, G. 2019b Occurrence and health implication of fluoride in groundwater of loess aquifer in the Chinese Loess Plateau: a case study of Tongchuan, Northwest China. *Exposure and Health* **11** (2), 95–107.
- Masindi, K. & Abiye, T. 2018 Assessment of natural and anthropogenic influences on regional groundwater chemistry in a highly industrialized and urbanized region: a case study of the Vaal River Basin, South Africa. *Environmental Earth Sciences* **77** (20), 1–14.
- Mehraein, M., Torabi, M., Sangsefidi, Y. & MacVicar, B. 2020 Numerical simulation of free flow through side orifice in a circular open-channel using response surface method. *Flow Measurement and Instrumentation* **76**, 101825.
- Mirzabeygi, M., Abbasnia, A., Yunesian, M., Nodehi, R. N., Yousefi, N., Hadi, M. & Mahvi, A. H. 2017 Heavy metal contamination and health risk assessment in drinking water of Sistan and Baluchistan, Southeastern Iran. *Human and Ecological Risk Assessment: An International Journal* **23** (8), 1895–1905.
- Moghaddam, M. H. R., Rouhi, M. N., Sarkar, S. & Rahimpour, T. 2018 Groundwater vulnerability assessment using the DRASTIC model under GIS platform in the Ajabshir Plain, Southeast coast of Urmia Lake, Iran. *Arabian Journal of Geosciences* **11** (19), 1–15.
- Mosaferi, M., Pourakbar, M., Shakerkhatibi, M., Fatehifar, E. & Belvasi, M. 2014 Quality modeling of drinking groundwater using GIS in rural communities, Northwest of Iran. *Journal of Environmental Health Science and Engineering* **12** (1), 1–14.
- Muhammad, A. K. & Husam, A. N. 2011 Hydro-geochemical characteristics of groundwater beneath the Gaza Strip. *Journal of Water Resource and Protection* **3** (5), 341–348.
- Narsimha, A. & Sudarshan, V. 2017 Assessment of fluoride contamination in groundwater from Basara, Adilabad district, Telangana state, India. *Applied Water Science* **7** (6), 2717–2725.
- Onwuka, O. S., Ezugwu, C. K. & Ifediegwu, S. I. 2019 Assessment of the impact of onsite sanitary sewage system and agricultural wastes on groundwater quality in Ikem and its environs, south-eastern Nigeria. *Geology, Ecology, and Landscapes* **3** (1), 65–81.
- Patolia, P. & Sinha, A. 2017 Fluoride contamination in Gharbar Village of Dhanbad District, Jharkhand, India: source identification and management. *Arabian Journal of Geosciences* **10** (17), 1–10.
- Qasemi, M., Farhang, M., Biglari, H., Afsharnia, M., Ojrati, A., Khani, F., Samiee, M. & Zarei, A. 2018 Health risk assessments due to nitrate levels in drinking water in villages of Azadshahr, Northeastern Iran. *Environmental Earth Sciences* **77** (23), 1–9.
- Qasemi, M., Shams, M., Sajjadi, S. A., Farhang, M., Erfanpoor, S., Yousefi, M., Zarei, A. & Afsharnia, M. 2019 Cadmium in groundwater consumed in the rural areas of Gonabad and Bajestan, Iran: occurrence and health risk assessment. *Biological Trace Element Research* **192** (2), 106–115.
- Rahman, M. A. T., Paul, M., Bhounmik, N., Hassan, M., Alam, M. K. & Aktar, Z. 2020 Heavy metal pollution assessment in the groundwater of the Meghna Ghat industrial area, Bangladesh, by using water pollution indices approach. *Applied Water Science* **10** (8), 1–15.
- Raju, N. J., Patel, P., Gurung, D., Ram, P., Gossel, W. & Wycisk, P. 2015 Geochemical assessment of groundwater quality in the Dun valley of Central Nepal using chemometric method and geochemical modeling. *Groundwater for Sustainable Development* **1** (1–2), 135–145.
- Ramakrishnaiah, C. R., Sadashivaiah, C. & Ranganna, G. 2009 Assessment of water quality index for the groundwater in Tumkur Taluk, Karnataka State, India. *E-Journal of Chemistry* **6** (2), 523–530.
- Rao, N. S., Marghade, D., Dinakar, A., Chandana, I., Sunitha, B., Ravindra, B. & Balaji, T. 2017 Geochemical characteristics and controlling factors of chemical composition of groundwater in a part of Guntur district, Andhra Pradesh, India. *Environmental Earth Sciences* **76** (21), 1–22.
- Rawat, K. S. & Singh, S. K. 2018 Water Quality Indices and GIS-based evaluation of a decadal groundwater quality. *Geology, Ecology, and Landscapes* **2** (4), 240–255.
- Rezaei, A. & Gurdak, J. J. 2020 Large-scale climate variability controls on climate, vegetation coverage, lake and groundwater storage in the Lake Urmia watershed using SSA and wavelet analysis. *Science of the Total Environment* **724**, 138273.
- Rezaei, A. & Hassani, H. 2018 Hydrogeochemistry study and groundwater quality assessment in the North of Isfahan, Iran. *Environmental Geochemistry and Health* **40** (2), 583–608.
- Rezaei, A., Hassani, H., Hayati, M., Jabbari, N. & Barzegar, R. 2018 Risk assessment and ranking of heavy metals concentration in Iran's Rayen groundwater basin using linear assignment method. *Stochastic Environmental Research and Risk Assessment* **32** (5), 1317–1336.
- Sahoo, S. & Khaoash, S. 2020 Impact assessment of coal mining on groundwater chemistry and its quality from Brajrajnagar coal mining area using indexing models. *J Geochem Explor* **215**, 1–17.
- Sahu, P. & Sikdar, P. K. 2008 Hydrochemical framework of the aquifer in and around East Kolkata Wetlands, West Bengal, India. *Environmental Geology* **55** (4), 823–835.
- Saleem, M., Hussain, A. & Mahmood, G. 2016 Analysis of groundwater quality using water quality index: a case study of greater Noida (Region), Uttar Pradesh (UP), India. *Cogent Engineering* **3** (1), 1237927.
- Sangsefidi, Y., Mehraein, M., Ghodsian, M. & Motalebizadeh, M. R. 2017 Evaluation and analysis of flow over arced weirs using traditional and response surface methodologies. *Journal of Hydraulic Engineering* **143** (11), 04017048.
- Sawyer, C. & McCarthy, P. 1967 *Chemical and Sanitary Engineering*, 2nd edn. McGraw-Hill, New York.
- Sethy, S. N., Syed, T. H., Kumar, A. & Sinha, D. 2016 Hydrogeochemical characterization and quality assessment of groundwater in parts of Southern Gangetic Plain. *Environmental Earth Sciences* **75** (3), 232.

- Shakerkhatibi, M., Mosafari, M., Pourakbar, M., Ahmadnejad, M., Safavi, N. & Banitorab, F. 2019 [Comprehensive investigation of groundwater quality in the North-West of Iran: physicochemical and heavy metal analysis](#). *Groundwater for Sustainable Development* **8**, 156–168.
- Sharmin, S., Mia, J., Miah, M. S. & Zakir, H. M. 2020 [Hydrogeochemistry and heavy metal contamination in groundwaters of Dhaka metropolitan city, Bangladesh: assessment of human health impact](#). *HydroResearch* **3**, 106–117.
- Srinivasamoorthy, K., Gopinath, M., Chidambaram, S., Vasanthavigar, M. & Sarma, V. S. 2014 [Hydrochemical characterization and quality appraisal of groundwater from Pungar sub basin, Tamilnadu, India](#). *Journal of King Saud University-Science* **26** (1), 37–52.
- Taghipour, H., Mosafari, M., Pourakbar, M. & Armanfar, F. 2012 [Heavy metals concentrations in groundwater used for irrigation](#). *Health Promotion Perspectives* **2** (2), 205.
- Thilagavathi, R., Chidambaram, S., Thivya, C., Prasanna, M. V., Keesari, T. & Pethaperumal, S. 2017 [Assessment of groundwater chemistry in layered coastal aquifers using multivariate statistical analysis](#). *Sustainable Water Resources Management* **3** (1), 55–69.
- Ustaoglu, F., Tepe, Y. & Taş, B. 2020 [Assessment of stream quality and health risk in a subtropical Turkey river system: a combined approach using statistical analysis and water quality index](#). *Ecological Indicators* **113**, 105815.
- Wang, Y. B., Liu, C. W., Liao, P. Y. & Lee, J. J. 2014 [Spatial pattern assessment of river water quality: implications of reducing the number of monitoring stations and chemical parameters](#). *Environmental Monitoring and Assessment* **186** (3), 1781–1792.
- WHO 2011 *WHO Guidelines for Drinking-Water Quality*, 4th edn. World Health Organization, Geneva.
- Wilcox, L. 1955 *Classification and use of Irrigation Waters* (No. 969). US Department of Agriculture, Washington, DC.
- Wu, P., Tang, C., Zhu, L., Liu, C., Cha, X. & Tao, X. 2009 [Hydrogeochemical characteristics of surface water and groundwater in the Karst Basin, Southwest China](#). *Hydrological Processes: An International Journal* **23** (14), 2012–2022.
- Wu, J., Li, P., Qian, H. & Fang, Y. 2014 [Assessment of soil salinization based on a low-cost method and its influencing factors in a semi-arid agricultural area, Northwest China](#). *Environmental Earth Sciences* **71** (8), 3465–3475.
- Wu, J., Wang, L., Wang, S., Tian, R., Xue, C., Feng, W. & Li, Y. 2017 [Spatiotemporal variation of groundwater quality in an arid area experiencing long-term paper wastewater irrigation, Northwest China](#). *Environmental Earth Sciences* **76** (13), 1–14.
- Yousefi, M., Mohammadi, A. A., Yaseri, M. & Mahvi, A. H. 2017 [Epidemiology of drinking water fluoride and its contribution to fertility, infertility, and abortion: an ecological study in West Azerbaijan Province, Poldasht County, Iran](#). *Fluoride* **50** (3), 343–353.
- Zhang, Y., Wu, J. & Xu, B. 2018 [Human health risk assessment of groundwater nitrogen pollution in Jinghui canal irrigation area of the loess region, Northwest China](#). *Environmental Earth Sciences* **77** (7), 1–12.

First received 20 March 2021; accepted in revised form 31 May 2021. Available online 12 June 2021