

Enhanced biomass production of duckweeds by inoculating a plant growth-promoting bacterium, *Acinetobacter calcoaceticus* P23, in sterile medium and non-sterile environmental waters

T. Toyama, M. Kuroda, Y. Ogata, Y. Hachiya, A. Quach, K. Tokura, Y. Tanaka, K. Mori, M. Morikawa and M. Ike

ABSTRACT

Duckweed offers the promise of a co-benefit culture combining water purification with biomass production. *Acinetobacter calcoaceticus* P23 is a plant growth-promoting bacterium isolated from a duckweed, *Lemna aequinoctialis*. This study quantified its growth-promoting effect on three duckweeds (*L. aoukikusa*, *L. minor*, and *Spirodela polyrhiza*) in sterile Hoagland solution and evaluated its usefulness in duckweed culture under non-sterile conditions. P23 promoted growth of three duckweeds in sterile Hoagland solution at low to high nutrient concentrations (1.25–10 mg NO₃-N/L and 0.25–2.0 mg PO₄-P/L). It increased the biomass production of *L. aequinoctialis* 3.8–4.3-fold, of *L. minor* 2.3–3.3-fold, and of *S. polyrhiza* 1.4–1.5-fold after 7 days compared with noninoculated controls. P23 also increased the biomass production of *L. minor* 2.4-fold in pond water and 1.7-fold in secondary effluent of a sewage treatment plant under non-sterile conditions at laboratory-scale experiments. P23 rescued *L. minor* from growth inhibition caused by microorganisms indigenous to the pond water. The results demonstrate that the use of P23 in duckweed culture can improve the efficiency of duckweed biomass production, and a positive effect of P23 on duckweed-based wastewater treatment can be assumed.

Key words | *Acinetobacter calcoaceticus*, biomass production, duckweed, plant growth-promoting bacteria

T. Toyama (corresponding author)

Y. Tanaka

K. Mori

Graduate Faculty of Interdisciplinary Research,
University of Yamanashi,
4-3-11 Takeda, Kofu,
Yamanashi 400-8511,
Japan
E-mail: ttohyama@yamanashi.ac.jp

M. Kuroda

Y. Hachiya

A. Quach

K. Tokura

M. Ike

Division of Sustainable Energy and Environmental
Engineering,
Osaka University,
2-1 Yamadaoka,
Suita, Osaka 565-0871,
Japan

Y. Ogata

National Institute for Environmental Studies,
16-2 Onogawa,
Tsukuba, Ibaraki 305-8506,
Japan

M. Morikawa

Division of Biosphere Science, Graduate School of
Environmental Science,
Hokkaido University,
Kita-10 Nishi-5,
Kita-ku, Sapporo 060-0810,
Japan

INTRODUCTION

Duckweeds are the smallest and fastest-growing aquatic plants, classified in the Araceae subfamily Lemnoideae, which includes five genera: *Lemna*, *Landoltia*, *Spirodela*, *Wolffia*, and *Wolffiella* (Landolt 1986). They are useful agents for removing nitrogen (N) and phosphorus (P) from municipal (Dalu & Ndamba 2003; Ran *et al.* 2004), livestock

(Cheng *et al.* 2002; Xu & Shen 2011), and industrial (Ozengin & Elmaci 2007) wastewaters because of their high growth rate and high nutrient uptake capabilities. They are also used to clean up eutrophied water bodies (Ansari & Khan 2008, 2009). Wastewater treatment and purification of polluted waters using duckweeds offers successful, cost-effective, low-energy, and environmentally friendly options around the world.

In addition, duckweeds have recently attracted attention as a good alternative feedstock for biofuel production owing to their high growth rate and high starch accumulation

This is an Open Access article distributed under the terms of the Creative Commons Attribution Licence (CC BY 4.0), which permits copying, adaptation and redistribution, provided the original work is properly cited (<http://creativecommons.org/licenses/by/4.0/>).

doi: 10.2166/wst.2017.296

capabilities (Cui & Cheng 2014). Conversion of duckweed biomass to ethanol (Xu *et al.* 2011, 2012; Soda *et al.* 2015), butanol (Su *et al.* 2014), hydrogen gas (Xu & Deshusses 2015), bio-oil (Muradov *et al.* 2010; Duan *et al.* 2013) and liquid fuels (e.g., gasoline, diesel, and kerosene) (Baliban *et al.* 2013) has been demonstrated by using biological, pyrolysis, hydrothermal, or thermochemical processes. Duckweeds are considered to have several advantages over terrestrial energy crops: they can take up nutrients directly from water, do not need extra fertilization or irrigation, can be easily grown and harvested, and do not compete with food crop production and agricultural land use. Therefore, they have high potential as an energy crop, especially when they are grown in a co-benefit system that combines water purification with biomass production (Xu *et al.* 2011, 2012). Enhancement of duckweed growth is critical to achieving a highly efficient co-benefit system. Proposed strategies include selecting fast-growing species (Zhao *et al.* 2015; Ziegler *et al.* 2015) and optimizing growth conditions such as water depth, duckweed coverage ratio, and harvest period (Zhao *et al.* 2014a). However, different strategies to further enhance duckweed growth are highly desired.

The use of plant growth-promoting bacteria (PGPB; Bashan & Holguin 1998) may offer a rational way to further increase biomass yields. A number of PGPB have been found to improve growth and yields of terrestrial agricultural crops. Significantly increased crop productivity, with 1.1- to 2.6-fold increases in yields, has been reported in greenhouse and field trials (Lucy *et al.* 2004; Adesemoye & Kloepper 2009; Bhattacharyya & Jha 2012; Pérez-Montañón *et al.* 2014). In contrast, PGPB in aquatic plants had not been studied until we isolated *Acinetobacter calcoaceticus* P23 from a duckweed species, *Lemna aequinoctialis* (Yamaga *et al.* 2010). We initially isolated P23 as a phenol-degrading bacterium and then found that it doubled the growth rate of *L. aequinoctialis* in axenic Hoagland solution both with and without phenol. If PGPB can be used in duckweed culture, in combination with the above approaches for optimizing duckweed production, they should greatly enhance biomass productivity. However, little information is yet available on the capacity of PGPB in duckweed culture. To apply PGPB to practical duckweed culture, it is necessary to evaluate their host-plant specificity and effects on plant growth under different conditions. In particular, it is important to confirm that they can indeed promote plant growth in non-sterile environmental waters, including wastewater.

Our objectives were therefore to quantify the growth-promoting effect of P23 on different duckweeds under

various conditions in order to clarify its general usefulness in duckweed culture. We evaluated the effects of P23 on the growth of three major duckweed species, *L. aequinoctialis*, *L. minor*, and *Spirodela polyrhiza*. We quantified the growth-promoting effects in sterile Hoagland nutrient solution at various nutrient concentrations, verified the effects in real environmental water, and finally confirmed the effects in real secondary effluent from a sewage treatment plant.

MATERIALS AND METHODS

Plant materials

Bacteria-free *L. aequinoctialis*, *L. minor*, and *S. polyrhiza* were prepared by washing in 0.5% sodium hypochlorite for 3 min, then in 70% ethanol for 1 min, and finally in sterilized water three times for 1 min. Plants were aseptically cultured in flasks containing sterile Hoagland solution (36.1 mg/L KNO₃, 293 mg/L K₂SO₄, 3.87 mg/L NaH₂PO₄, 103 mg/L MgSO₄·7H₂O, 147 mg/L CaCl₂·H₂O, 3.33 mg/L FeSO₄·7H₂O, 0.95 mg/L H₃BO₃, 0.39 mg/L MnCl₂·4H₂O, 0.03 mg/L CuSO₄·5H₂O, 0.08 mg/L ZnSO₄·7H₂O, and 0.254 mg/L H₂MoO₄·4H₂O; pH 7.0) in a growth chamber (28 ± 1 °C, fluorescent lamps at a photosynthetic photon flux density of 80 μmol/m²/s, 16 L:8 D photoperiod).

Acinetobacter calcoaceticus P23 culture and preparation of cell suspensions

Strain P23 was grown in LB medium (10 g/L Bacto Peptone, 5 g/L Bacto Yeast extract, 5 g/L NaCl; pH 7.2) overnight at 28 °C, with shaking at 150 rpm. Cells were harvested by two cycles of centrifugation (10,000 ×g, room temperature, 5 min) and washing for 30 s in sterilized Hoagland solution, and then were resuspended in the same medium. The P23 cell suspension, whose optical density at 600 nm (OD₆₀₀) was first adjusted to a known value, was inoculated as described below. For determination of cell density, the cells were dried for 3 h at 90 °C and weighed. OD₆₀₀ = 1 was equivalent to 0.496 mg dry weight/mL.

Pond water and secondary effluent from a sewage treatment plant

Water collected from Inukai Pond (Suita, Osaka, Japan) was first passed through an Isopore membrane filter (pore size, 10 μm; polycarbonate; Merck Millipore) to remove

microalgae so as to avoid overgrowth during experiments. It had a pH of 7.3; undetectable $\text{NH}_4\text{-N}$, $\text{NO}_2\text{-N}$, and $\text{PO}_4\text{-P}$; and 0.01 mg/L $\text{NO}_3\text{-N}$. After incubation for 7 days at 28 °C on 0.1× LB agar plates, it was determined to have 8.0×10^5 CFU/mL of total heterotrophic bacteria. Secondary effluent was collected from a municipal sewage treatment plant in Kofu city (Yamanashi, Japan). It had a pH of 7.5, 4.27 mg/L $\text{NH}_4\text{-N}$, 0.66 mg/L $\text{NO}_2\text{-N}$, 7.72 mg/L $\text{NO}_3\text{-N}$ and 0.98 mg/L $\text{PO}_4\text{-P}$, and 8.5×10^6 CFU/mL of total heterotrophic bacteria.

Co-culture of duckweed with P23 cells in Hoagland solution under sterile conditions at various nutrient concentrations

Ten fronds of each duckweed species were placed in 300-mL flasks ($n=2$ per species per nutrient concentration, including control) containing 100 mL of sterile Hoagland solution. We then inoculated P23 cells into the flasks at a final density of 0.15 mg dry weight/mL. The Hoagland solution concentrations ranged from 0.25× to 2×; this corresponded to nutrient concentrations ranging from 1.25 mg $\text{NO}_3\text{-N/L} + 0.25$ mg $\text{PO}_4\text{-P/L}$ to 10 mg $\text{NO}_3\text{-N/L} + 2.0$ mg $\text{PO}_4\text{-P/L}$. All of the flasks were incubated without shaking in the chamber for 7 days as described above. The fronds were counted daily. After 7 days, all plants in each flask were collected, dried for 3 h at 90 °C, and weighed.

Co-culture of duckweed with P23 cells in pond water under sterile and non-sterile conditions

We examined the growth-promoting activity of P23 with *L. minor*, which is one of the most widespread duckweed species in the world (Holm et al. 1997). To examine the effects of indigenous microorganisms in pond water on duckweed growth and the ability of P23 to promote growth, we used both sterile and non-sterile pond water. Sterile pond water was prepared by filtration (pore size 0.2 μm). Both water samples were then supplemented with KNO_3 (1.25 mg $\text{NO}_3\text{-N/L}$) and NaH_2PO_4 (0.25 mg $\text{PO}_4\text{-P/L}$). Ten fronds of *L. minor* were transferred into 200-mL flasks containing 100 mL of the water samples ($n=3$), followed by inoculation with P23 cell suspension at 0.15 mg dry weight/mL. All flasks were incubated without shaking in the chamber as described above for 15 days. The number of fronds was counted daily. After 15 days, all plants in each flask were collected, dried, and weighed.

To further determine the effect of cell density on the ability of P23 to promote duckweed growth under non-sterile conditions, we inoculated P23 cell suspensions at 0, 0.025, 0.05, and 0.15 mg dry weight/mL into flasks containing 10 fronds of *L. minor* in 100 mL of non-sterile pond water ($n=3$ per cell density). All flasks were incubated without shaking in the chamber for 12 days. The fronds were counted daily. After 12 days, all plants in each flask were collected, dried, and weighed.

Co-culture of duckweed with P23 cells in secondary effluent under non-sterile conditions

To verify the ability of P23 to promote duckweed growth in real sewage effluent, we placed 10 fronds in 500-mL flasks ($n=3$) containing 200 mL of secondary effluent with or without P23 cells at a final density of 0.15 mg dry weight/mL. All flasks were incubated without shaking in the chamber as described above for 7 days. The fronds were counted daily, and the dry weights of all plants in each flask were measured after 7 days. These three co-culture experiments in Hoagland solution, pond water, or secondary effluent are summarized in Figure 1.

Statistical analysis

Each value used in the statistical analysis represents the results of two or three samples ($n=2$ or 3 replicates) per experiment. All results are expressed as mean ± standard deviation (SD). Significance ($P < 0.05$) was assessed using the *t*-test in IBM SPSS Statistics v. 22.0.

RESULTS AND DISCUSSION

Growth promotion of *L. aequinoctialis*, *L. minor*, and *S. polyrhiza* by P23 cells under sterile conditions at various nutrient concentrations

We grew each of the three duckweed species with P23 cells (0.15 mg dry weight/mL) in sterile Hoagland solution at different nutrient concentrations for 7 days. The results are summarized in Figures 2–5 and Table 1. At all nutrient concentrations, all three duckweed species grew more rapidly in the presence of P23 cells than in the absence. The increase in the number of fronds (final number minus initial 10) was 2.9–3.3-fold in *L. aequinoctialis*, 2.7–3.8-fold in *L. minor*, and 1.2–1.4-fold in *S. polyrhiza* compared with the non-inoculated controls (Table 1). Biomass production (increase in dry weight) was similarly increased 3.8–4.3-fold,

I. Co-culture of duckweed with P23 cells in Hoagland solution under sterile conditions at various nutrient concentration

- Duckweeds (3 species): *L. aequinoctialis*, *L. minor*, and *S. polyrhiza*
- Hoagland solution (5 different concentrations): 0.25 ×, 0.5 ×, 1.0 ×, 1.5 ×, and 2.0 ×
Nutrient concentration in 1.0 × Hoagland = 5 mg NO₃-N/L and 1 mg PO₄-P/L
- P23 (with/without P23): 0 and 0.15 mg dry weight/mL

II. Co-culture of duckweed with P23 cells in pond water under sterile and non-sterile conditions

- Duckweed (1 species): *L. minor*
- Pond water : sterile and non-sterile
Nutrient concentrations in pond water = 1.25 mg NO₃-N/L and 0.25 mg PO₄-P/L
- P23 (4 different cell densities): 0, 0.025, 0.05, and 0.15 mg dry weight/mL

III. Co-culture of duckweed with P23 cells in secondary effluent under non-sterile condition

- Duckweed (1 species): *L. minor*
- Secondary effluent: non-sterile
Nutrient concentrations = 4.27 mg-NH₄-N/L, 0.66 mg NO₂-N/L, 7.72 mg NO₃-N/L, and 0.98 mg PO₄-P/L
- P23 (with/without P23): 0 and 0.15 mg dry weight/mL

Figure 1 | Experimental scheme.




Figure 2 | Changes in the number of *L. aequinoctialis* fronds in culture with P23 (○, 0.15 mg dry weight/mL) and without P23 (●) in Hoagland solution (1× = 5.0 mg NO₃-N/L + 1.0 mg PO₄-P/L) at different nutrient concentrations: (a) 0.25×; (b) 0.50×; (c) 1.0×; (d) 1.5×; (e) 2.0×. Values are mean ± SD (n = 2).

2.3–3.3-fold, and 1.4–1.5-fold, respectively (Table 1). The nutrient concentration of the medium did not greatly affect the growth-promoting activity of P23 in any species.

The fact that P23 could promote the growth of all three duckweed species indicates its versatile potential to accelerate biomass production. Various duckweed species have




Figure 3 | Changes in the number of *L. minor* fronds in culture with P23 (○, 0.15 mg dry weight/mL) and without P23 (●) in Hoagland solution (1× = 5.0 mg NO₃-N/L + 1.0 mg PO₄-P/L) at different nutrient concentrations: (a) 0.25×; (b) 0.50×; (c) 1.0×; (d) 1.5×; (e) 2.0×. Values are mean ± SD (n = 2).




Figure 4 | Changes in the number of *S. polyrhiza* fronds in culture with P23 (○, 0.15 mg dry weight/mL) and without P23 (●) in Hoagland solution (1× = 5.0 mg NO₃-N/L + 1.0 mg PO₄-P/L) at different nutrient concentrations: (a) 0.25×; (b) 0.50×; (c) 1.0×; (d) 1.5×; (e) 2.0×. Values are mean ± SD (n = 2).




Figure 5 | Dry weight of duckweed biomass after 7 days in culture with P23 (□, 0.15 mg dry weight/mL) and without P23 (■) in Hoagland solution ($1\times = 5.0$ mg NO_3^-/N / $\text{L} + 1.0$ mg $\text{PO}_4\text{-P}/\text{L}$) at different nutrient concentrations: (A) 0.25 \times ; (B) 0.50 \times ; (C) 1.00 \times ; (D) 1.5 \times ; (E) 2.0 \times . Initial dry weights of 10 fronds per flask: *L. aequinoctialis*, 0.45 ± 0.07 mg; *L. minor*, 0.85 ± 0.09 mg; *S. polyrhiza*, 3.4 ± 0.14 mg. Values are mean \pm SD ($n = 2$). *Significant difference ($P < 0.05$) between values with and without P23 within a treatment.

been used in wastewater treatment, water purification, and biomass production according to wastewater, temperature, and purpose, because growth rate, nutrient removal ability, optimal growth conditions, and chemical composition differ among species (Landolt 1986; Lemon et al. 2001; Zhao et al. 2014a, 2014b; Soda et al. 2015; Ziegler et al. 2015). For example, *L. minor* and *S. polyrhiza* are suitable for culture at lower and higher temperatures, respectively, because they grow at 5 to 31 °C and 15 to above 35 °C, respectively (Landolt 1986). *S. polyrhiza* is suitable for saline wastewater treatment because of its high salt

tolerance (Sree et al. 2015). In addition, mixed cultures of duckweed species have higher nutrient removal and biomass production rates than single cultures (Zhao et al. 2014b). Therefore, it will be useful to examine a variety of species to achieve a high-efficiency culture system. In this context, the broad-spectrum growth-promoting effect of P23 is of great practical significance.

Although P23 promoted the growth of all three species, it promoted the growth of *L. aequinoctialis* and *L. minor* more than that of *S. polyrhiza*. The ability of P23 to promote the growth of *L. aequinoctialis* was the highest. The finding may result from a certain host specificity. Also, the ability of P23 to promote the growth of *L. minor* more than that of *S. polyrhiza* may relate to the evolutionarily closer relationship of *L. minor* and *L. aequinoctialis* (Les et al. 2002).

In terrestrial soils, N and P often limit plant growth. N-fixing PGPB and P-solubilizing PGPB thus play significant roles by providing nutrients that support plant growth. Their effects are strongly influenced by nutrient conditions (Adesemoye & Kloepper 2009; Bhattacharyya & Jha 2012), with much stronger effects in nutrient-deficient soils than in nutrient-rich soils (Egamberdiyeva 2007). In contrast, P23 promoted duckweed growth over a wide range of nutrient concentrations in this study, suggesting that the nature and mechanisms of duckweed growth promotion by P23 might differ from that of soil PGPB. Thus, growth-promoting mechanisms specific to aquatic plants must be explored in future studies.

Enhancement of *L. minor* growth by P23 cells in pond water and secondary effluent

In the absence of P23, the increases in frond number and biomass production of *L. minor* were significantly lower ($P < 0.05$) in non-sterile pond water than in sterile pond water (Figure 6, Table 2). This result indicates that indigenous microorganisms in the pond water inhibited the growth of *L. minor*.

In the presence of P23 in the sterile pond water, the increased frond number and biomass production of *L. minor* were increased 1.3- and 1.9-fold, respectively, relative to its absence (Figure 6, Table 2). This enhanced biomass production must result from the direct growth promotion effect of P23, as observed in sterile Hoagland solution. Although the growth of *L. minor* in the absence of P23 was less in non-sterile pond water than in sterile pond water, it is noteworthy that P23 increased the frond number and biomass production even in non-sterile pond water 3.8- and 2.4-fold, respectively (Figure 6,

Table 1 | Effects of P23 on the growth of *L. aequinoctialis*, *L. minor*, and *S. polyrhiza* in sterile Hoagland solution at indicated nutrient concentrations

| Duckweed species | Hoagland solution | Treatment | Increase in number of fronds per flask during 7 days ^a (and ratio of P23 to control) | Biomass production (increase in dry weight; mg) of plants per flask during 7 days ^a (and ratio of P23 to control) | |
|--------------------------|-------------------|-----------|---|--|--------------------|
| <i>L. aequinoctialis</i> | 0.25× | P23 | 143 ± 5.0 (3.2)* | 9.45 ± 0.71 (4.2)* | |
| | | Control | 44.5 ± 0.7 | 2.25 ± 0.28 | |
| | 0.5× | P23 | 172 ± 4.2 (3.1)* | 10.4 ± 0.07 (3.8)* | |
| | | Control | 55.0 ± 2.8 | 2.75 ± 0.28 | |
| | 1.0× | P23 | 134 ± 9.9 (2.9)* | 9.20 ± 0.35 (4.1)* | |
| | | Control | 46.0 ± 1.4 | 2.25 ± 0 | |
| | 1.5× | P23 | 130 ± 4.2 (3.3)* | 7.95 ± 0.71 (4.3)* | |
| | | Control | 39.0 ± 2.8 | 1.85 ± 0.28 | |
| | 2.0× | P23 | 121 ± 5.7 (3.3)* | 7.70 ± 0.35 (4.3)* | |
| | | Control | 36.5 ± 0.7 | 1.80 ± 0.35 | |
| | <i>L. minor</i> | 0.25× | P23 | 65.5 ± 6.3 (3.0)* | 9.95 ± 0.85 (3.1)* |
| | | | Control | 21.5 ± 0.7 | 3.25 ± 0.28 |
| 0.5× | | P23 | 85.5 ± 3.5 (3.5)* | 9.80 ± 0.35 (3.0)* | |
| | | Control | 24.5 ± 3.5 | 3.30 ± 0.07 | |
| 1.0× | | P23 | 93.5 ± 6.4 (3.8)* | 9.75 ± 0.71 (3.3)* | |
| | | Control | 24.5 ± 5.0 | 3.00 ± 0.21 | |
| 1.5× | | P23 | 86.5 ± 3.5 (3.3)* | 8.45 ± 0.28 (2.8)* | |
| | | Control | 26.5 ± 2.1 | 3.05 ± 0 | |
| 2.0× | | P23 | 71.0 ± 2.8 (2.7)* | 7.40 ± 0.21 (2.3)* | |
| | | Control | 26.5 ± 0.7 | 3.25 ± 0.28 | |
| <i>S. polyrhiza</i> | | 0.25× | P23 | 34 ± 1.4 (1.4)* | 19.0 ± 1.2 (1.5)* |
| | | | Control | 23.5 ± 3.5 | 12.4 ± 0.1 |
| | 0.5× | P23 | 43.0 ± 1.4 (1.3)* | 24.5 ± 0.4 (1.5)* | |
| | | Control | 32.5 ± 0.7 | 16.1 ± 0.5 | |
| | 1.0× | P23 | 46.5 ± 2.1 (1.4)* | 26.8 ± 0.8 (1.5)* | |
| | | Control | 32.5 ± 0.7 | 17.9 ± 0.6 | |
| | 1.5× | P23 | 53.5 ± 2.1 (1.2)* | 29.1 ± 0.8 (1.4)* | |
| | | Control | 43.5 ± 2.1 | 21.5 ± 1.0 | |
| | 2.0× | P23 | 60.5 ± 2.1 (1.4)* | 30.8 ± 0.9 (1.4)* | |
| | | Control | 44.0 ± 1.4 | 21.7 ± 0.4 | |

Values are means ± SD. Values in parentheses are the ratio of the P23 value to the control (no P23) value.

^aFinal value minus initial value.

*Significant difference ($P < 0.05$) between values with and without P23 within a treatment.

Table 2). This highly enhanced biomass production is attributable to synergism between P23's direct growth promotion effect and its indirect effect of rescue from growth inhibition by indigenous microorganisms. It is interesting that P23's growth promotion effects were greater in non-sterile pond water (3.8-fold effect on frond number increase and 2.4-fold effect on biomass production) than in sterile pond water (1.3- and 1.9-fold, respectively), indicating that P23 could be especially useful in practical culture systems. On the other hand, the size of *L. minor* grown in the sterile pond water with P23 was a little bit larger than that in the non-sterile pond water with P23. The highest biomass production of *L. minor* was reached in the sterile pond water with P23 (Figure 6, Table 2).

Terrestrial plant PGPB often fail to perform in field conditions (Lucy et al. 2004). For example, Cu-tolerant PGPB highly promoted the growth of plants in sterilized tailings, but only weakly promoted it in non-sterile conditions (Liu et al. 2014). The report suggested that terrestrial plant PGPB cannot fully express their growth-promoting effect under non-sterile conditions owing to competition with other indigenous microorganisms (Liu et al. 2014). It is notable that P23 significantly promoted the growth of *L. minor* in real pond water, which harbors complex indigenous microbial communities. This trait of the duckweed-P23 association in water confers an advantage over the terrestrial plant-PGPB association in soil.

To determine the effective cell density of P23 to promote duckweed growth under non-sterile conditions, we grew




Figure 6 | Effects of P23 (0.05 mg dry weight/mL) on the growth of *L. minor* in sterile or non-sterile pond water for 15 days. (a) Changes in number of fronds growing in sterile (■, □) or non-sterile (●, ○) pond water without (■, ●, control) or with P23 cells (□, ○). (b) Biomass production (final minus initial dry weight) of *L. minor* in sterile or non-sterile pond water with or without P23 cells. Values are mean \pm SD ($n = 3$). *Significant difference ($P < 0.05$) between dry weights.

Table 2 | Effects of P23 on the growth of *L. minor* in sterile or non-sterile pond water

| Pond water | Treatment | Increase in number of fronds per flask during 15 days ^a (and ratio of P23 to control) | Biomass production (increase in dry weight; mg) of plants per flask during 15 days ^a (and ratio of P23 to control) |
|-------------|-----------|--|---|
| Sterile | P23 | 38 \pm 6.8 (1.3)* | 22.6 \pm 2.5 (1.9)* |
| | Control | 30 \pm 4.5 | 12.0 \pm 0.2 |
| Non-sterile | P23 | 42 \pm 3.0 (3.8)* | 18.2 \pm 1.2 (2.4)* |
| | Control | 11 \pm 1.07 | 7.7 \pm 1.0 |

Values are means \pm SD. Values in parentheses are the ratio of the P23 value to the control (no P23) value.

^aFinal value minus initial value.

*Significant difference ($P < 0.05$) between values with and without P23 within a treatment.

L. minor in pond water at various densities of P23 cells. Although there was no significant effect on growth at a cell density of 0.025 mg/mL, the effect of P23 became significant ($P < 0.05$) at higher cell densities (Figure 7, Table 3). The increased frond number and biomass production were increased 2.0- and 1.4-fold, respectively, in the presence of P23 cells at 0.05 mg/mL, and 4.3- and 1.8-fold at 0.15 mg/mL. Thus, the initial inoculation density of P23 cells would be important for the significant promotion of duckweed growth in this system.

Finally, P23 at 0.15 mg/mL significantly promoted the growth of *L. minor* even in real secondary sewage effluent (Figure 8, Table 4). After 7 days' culture, the frond number and biomass production were increased about 1.7-fold by P23 inoculation relative to the control. These results demonstrate that P23 can enhance duckweed biomass production even in real-world secondary sewage effluent.

It is notable that P23 is capable of exerting synergistic effects of growth promotion and rescue from growth inhibition under non-sterile conditions, increasing biomass production of *L. minor* 2.4- and 1.7-fold in real pond water and secondary sewage effluent, respectively. This growth-promoting effect was equivalent to or higher than that of terrestrial PGPB in crops (1.1- to 2.6-fold increases in yields at greenhouse and field trials) (Lucy et al. 2004; Adesemoye & Kloepper 2009; Bhattacharyya & Jha 2012; Pérez-Montañó et al. 2014). Moreover, P23 enhanced biomass production more effectively than previous approaches of optimizing water depth (1.5-fold), duckweed coverage ratio (1.6-fold), and harvest period (1.1-fold) (Zhao et al. 2014a). Thus, aquatic PGPB technology offers hope for enhancing duckweed growth to achieve effective biomass production coupled with effective purification of eutrophic environmental water and sewage. In the next steps, it will be necessary to evaluate the sustainable effect and utilization




Figure 7 | Effects of P23 on the growth of *L. minor* in non-sterile pond water for 12 days. (a) Changes in the number of fronds growing with P23 cells at densities of 0 (□, control), 0.025 (○), 0.05 (Δ), or 0.15 (◇) mg dry weight/mL. (b) Biomass production (final minus initial dry weight) of *L. minor* in pond water with P23 cell densities of 0 (control), 0.025, 0.05, or 0.15 mg dry weight/mL during 12 days. Values are mean ± SD ($n = 3$). *Significant difference ($P < 0.05$) from control.

Table 3 | Effects of different cell densities of P23 on the growth of *L. minor* in non-sterile pond water

| Cell density of P23 (mg dry weight/L) | Increase in number of fronds per flask during 12 days ^a (and ratio of P23 to control) | Biomass production (increase in dry weight; mg) of plants per flask during 12 days ^a (and ratio of P23 to control) |
|---------------------------------------|--|---|
| 0 (control) | 22 ± 2.5 | 4.5 ± 0.4 |
| 0.025 | 24 ± 9.6 (1.1) | 5.0 ± 1.3 (1.1) |
| 0.05 | 45 ± 4.8 (2.0)* | 6.2 ± 0.4 (1.4)* |
| 0.15 | 94 ± 6.7 (4.3)* | 8.0 ± 0.1 (1.8)* |

Values are means ± SD. Values in parentheses are the ratio of the P23 value to the control (no P23) value.

^aFinal value minus initial value.

*Significant difference ($P < 0.05$) between values with and without P23 within a treatment.

of P23 in pilot-scale experiments to construct a practical PGPB-duckweed system.

CONCLUSION

Acinetobacter calcoaceticus P23, a plant growth-promoting bacterium originally isolated from *L. aequinoctialis*, significantly promoted the growth of three duckweed species (*L. aequinoctialis*, *L. minor*, and *S. polyrhiza*) and accelerated their biomass production, not only in sterilized Hoagland solution at a range of nutrient concentrations, but also in non-sterile pond water and secondary sewage




Figure 8 | Effects of P23 (0.15 mg dry weight/mL) on the growth of *L. minor* in secondary effluent for 7 days. (a) Changes in the number of fronds growing without (●) or with P23 cells (○). (b) Biomass production (final minus initial dry weight) of *L. minor* during 7 days. Values are mean ± SD ($n = 3$). *Significant difference ($P < 0.05$) between treatments.

Table 4 | Effects of P23 on the growth of *L. minor* in secondary effluent

| Treatment | Increase in number of fronds per flask during 7 days ^a (and ratio of P23 to control) | Biomass production (increase in dry weight; mg) of plants per flask during 7 days ^a (and ratio of P23 to control) |
|-----------|---|--|
| P23 | 266 ± 15 (1.7)* | 32.6 ± 2.1 (1.7)* |
| Control | 159 ± 15 | 19.0 ± 1.3 |

Values are means ± SD. Values in parentheses are the ratio of the P23 value to the control (no P23) value.

^aFinal value minus initial value.

*Significant difference ($P < 0.05$) between values with and without P23 within a treatment.

effluent. This excellent biomass productivity enhanced by P23 can be ascribed to synergistic effects of direct growth promotion and rescue from growth inhibition by indigenous microorganisms. Overall, these results strongly suggest that aquatic PGPB such as P23 offer the promise of effective biomass production in duckweed culture and seem to be beneficial for duckweed-based wastewater treatment and water purification.

ACKNOWLEDGEMENTS

This study was supported in part by the Advanced Low Carbon Technology Research and Development Program (ALCA) of the Japan Science and Technology Agency (JST), and by a Grant-in-Aid for Scientific Research (B) (No. 24310057) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

REFERENCES

- Adesemoye, A. O. & Kloepper, J. W. 2009 Plant-microbes interactions in enhanced fertilizer-use efficiency. *Applied Microbiology and Biotechnology* **85** (1), 1–12.
- Ansari, A. A. & Khan, F. A. 2008 Remediation of eutrophied water using *Lemna minor* in a controlled environment. *African Journal of Aquatic Science* **33** (3), 275–278.
- Ansari, A. A. & Khan, F. A. 2009 Remediation of eutrophied water using *Spirodela polyrhiza* L. Shleid in controlled environment. *Pan-American Journal of Aquatic Sciences* **4** (1), 52–54.
- Baliban, R. C., Elia, J. A., Floudas, C. A., Xiao, X., Zhang, Z., Li, J., Cao, H., Ma, J., Qiao, Y. & Hu, X. 2013 Thermochemical conversion of duckweed biomass to gasoline, diesel, and jet fuel: process synthesis and global optimization. *Industrial and Engineering Chemistry Research* **52** (33), 11436–11450.
- Bashan, Y. & Holguin, G. 1998 Proposal for the division of plant growth-promoting rhizobacteria into two classifications: biocontrol-PGPB (plant growth-promoting bacteria) and PGPB. *Soil Biology and Biochemistry* **30** (8), 1225–1228.
- Bhattacharyya, P. N. & Jha, D. K. 2012 Plant growth-promoting rhizobacteria (PGPR): emergence in agriculture. *World Journal of Microbiology and Biotechnology* **28** (4), 1327–1350.
- Cheng, J., Bergmann, B. A., Classen, J. J., Stomp, A. M. & Howard, J. W. 2002 Nutrient recovery from swine lagoon water by *Spirodela punctata*. *Bioresource Technology* **81**, 81–85.
- Cui, W. & Cheng, J. J. 2014 Growing duckweed for biofuel production: a review. *Plant Biology* **17** (s1), 16–23.
- Dalu, J. M. & Ndamba, J. 2003 Duckweed based wastewater stabilization ponds for wastewater treatment (a low cost technology for small urban areas in Zimbabwe). *Physics and Chemistry of the Earth* **28** (20–27), 1147–1160.
- Duan, P., Chang, Z., Xu, Y., Bai, X., Wang, F. & Zhang, L. 2013 Hydrothermal processing of duckweed: effect of reaction conditions on product distribution and composition. *Bioresource Technology* **135**, 710–719.
- Egamberdiyeva, D. 2007 The effect of plant growth promoting bacteria on growth and nutrient uptake of maize in two different soils. *Applied Soil Ecology* **36** (2–3), 184–189.
- Holm, L., Doll, J., Holm, E., Pancho, J. & Herberger, J. 1997 *World Weeds: Natural Histories and Distribution*. Wiley, New York, USA, pp. 433–441.
- Landolt, E. 1986 *Biosystematic Investigations in the Family of Duckweed (Lemnaceae) Vol. 2: a Monographic Study Vol. 1*. Veröffentlichungen des Geobotanischen Institutes der ETH, Stiftung Rübel, Zürich, Switzerland.
- Lemon, G. D., Posluszny, U. & Husband, B. C. 2001 Potential and realized rates of vegetative reproduction in *Spirodela polyrhiza*, *Lemna minor*, and *Wolffia borealis*. *Aquatic Botany* **70** (1), 79–87.
- Les, D. H., Crawford, D. J., Landolt, E., Gavel, J. D. & Kimball, R. T. 2002 Phylogeny and systematics of Lemnaceae, the duckweed family. *Systematic Botany* **27** (2), 221–240.
- Liu, W., Yang, C., Shi, S. & Shu, W. 2014 Effects of plant growth-promoting bacteria isolated from copper tailings on plants in sterilized and non-sterilized tailings. *Chemosphere* **97**, 47–53.
- Lucy, M., Reed, E. & Glick, B. R. 2004 Applications of free living plant growth-promoting rhizobacteria. *Antonie van Leeuwenhoek* **86** (2), 1–25.
- Muradov, N., Fidalgo, B., Gujar, A. C. & T-Raissi, A. 2010 Pyrolysis of fast-growing aquatic biomass – *Lemna minor* (duckweed): Characterization of pyrolysis products. *Bioresource Technology* **101**, 8424–8428.
- Ozengin, N. & Elmaci, A. 2007 Performance of duckweed (*Lemna minor* L.) on different types of wastewater treatment. *Journal of Environmental Biology* **28** (2), 307–314.
- Pérez-Montaño, F., Alías-Villegas, C., Bellogín, R. A., del Cerro, P., Espuny, M. R., Jiménez-Guerrero, I., López-Baena, F. J., Ollero, F. J. & Cubo, T. 2014 Plant growth promotion in cereal and leguminous agricultural important plants: from microorganism capacities to crop production. *Microbiological Research* **169** (5–6), 325–336.
- Ran, N., Agami, M. & Oron, G. 2004 A pilot study of constructed wetlands using duckweed (*Lemna gibba* L.) for treatment of domestic primary effluent in Israel. *Water Research* **38** (9), 2241–2248.
- Soda, S., Ohchi, T., Piradee, J., Takai, Y. & Ike, M. 2015 Duckweed biomass as a renewable biorefinery feedstock: ethanol and succinate production from *Wolffia globosa*. *Biomass and Bioenergy* **81**, 364–368.
- Sree, K. S., Adelman, K., Garcia, C., Lam, E. & Appenroth, K.-J. 2015 Natural variance in salt tolerance and induction of starch accumulation in duckweeds. *Planta* **241** (6), 1395–1404.
- Su, H., Zhao, Y., Jiang, J., Lu, Q., Li, Q., Luo, Y., Zhao, H. & Wang, M. 2014 Use of duckweed (*Landoltia punctata*) as a fermentation substrate for the production of higher alcohols as biofuels. *Energy and Fuels* **28** (5), 3206–3216.
- Xu, J. & Deshusses, M. A. 2015 Fermentation of swine wastewater-derived duckweed for biohydrogen production. *International Journal of Hydrogen Energy* **40** (22), 7028–7036.

- Xu, J. & Shen, G. 2011 Growing duckweed in swine wastewater for nutrient recovery and biomass production. *Bioresource Technology* **102** (2), 848–853.
- Xu, J., Cui, W., Cheng, J. J. & Stomp, A.-M. 2011 Production of high-starch duckweed and its conversion to bioethanol. *Biosystems Engineering* **110** (2), 67–72.
- Xu, J., Cheng, J. J. & Stomp, A.-M. 2012 Growing *Spirodela polyrhiza* in swine wastewater for the production of animal feed and fuel ethanol: a pilot study. *CLEAN-Soil, Air, Water* **40** (7), 760–765.
- Yamaga, F., Washio, K. & Morikawa, M. 2010 Sustainable biodegradation of phenol by *Acinetobacter calcoaceticus* P23 isolated from the rhizosphere of duckweed *Lemna aoukikusa*. *Environmental Science and Technology* **44** (16), 6470–6474.
- Zhao, Y., Fang, Y., Jin, Y., Huang, J., Bao, S., He, Z., Wang, F. & Zhao, H. 2014a Effects of operation parameters on nutrient removal from wastewater and high-protein biomass production in a duckweed-based (*Lemna aequinoctialis*) pilot-scale system. *Water Science and Technology* **70** (7), 1195–1204.
- Zhao, Z., Shi, H., Liu, Y., Zhao, H., Su, H., Wang, M. & Zhao, Y. 2014b The influence of duckweed species diversity on biomass productivity and nutrient removal efficiency in swine wastewater. *Bioresource Technology* **167**, 383–389.
- Zhao, Y., Fang, Y., Jin, Y., Huang, J., Bao, S., Fu, T., He, Z., Wang, F., Wang, M. & Zhao, H. 2015 Pilot-scale comparison of four duckweed strains from different genera for potential application in nutrient recovery from wastewater and valuable biomass production. *Plant Biology* **17** (s1), 82–90.
- Ziegler, P., Adelman, K., Zimmer, S., Schmidt, C. & Appenroth, K.-J. 2015 Relative *in vitro* growth rates of duckweeds (Lemnaceae)—the most rapidly growing higher plants. *Plant Biology* **17** (s1), 33–41.

First received 13 December 2016; accepted in revised form 5 May 2017. Available online 5 June 2017