

Three Biomarkers Identified from Serum Proteomic Analysis for the Detection of Early Stage Ovarian Cancer

Zhen Zhang,¹ Robert C. Bast, Jr.,² Yinhua Yu,² Jinong Li,¹ Lori J. Sokoll,¹ Alex J. Rai,¹ Jason M. Rosenzweig,¹ Bonnie Cameron,¹ Young Y. Wang,¹ Xiao-Ying Meng,³ Andrew Berchuck,⁴ Carolien van Haaften-Day,⁵ Neville F. Hacker,⁵ Henk W. A. de Bruijn,⁶ Ate G. J. van der Zee,⁶ Ian J. Jacobs,⁷ Eric T. Fung,³ and Daniel W. Chan¹

¹Department of Pathology, Biomarker Discovery Center, Johns Hopkins Medical Institutions, Baltimore, Maryland; ²M. D. Anderson Cancer Center, Houston, Texas; ³Ciphergen Biosystems, Inc., Fremont, California; ⁴Duke University Medical Center, Durham, North Carolina; ⁵The Royal Hospital for Women, Randwick, New South Wales, Australia; ⁶University Hospital Groningen, Groningen, the Netherlands; and ⁷Bart's and The London, Queen Mary's School of Medicine, London, United Kingdom

ABSTRACT

Early detection remains the most promising approach to improve long-term survival of patients with ovarian cancer. In a five-center case-control study, serum proteomic expressions were analyzed on 153 patients with invasive epithelial ovarian cancer, 42 with other ovarian cancers, 166 with benign pelvic masses, and 142 healthy women. Data from patients with early stage ovarian cancer and healthy women at two centers were analyzed independently and the results cross-validated to discover potential biomarkers. The results were validated using the samples from two of the remaining centers. After protein identification, biomarkers for which an immunoassay was available were tested on samples from the fifth center, which included 41 healthy women, 41 patients with ovarian cancer, and 20 each with breast, colon, and prostate cancers. Three biomarkers were identified as follows: (a) apolipoprotein A1 (down-regulated in cancer); (b) a truncated form of transthyretin (down-regulated); and (c) a cleavage fragment of inter- α -trypsin inhibitor heavy chain H4 (up-regulated). In independent validation to detect early stage invasive epithelial ovarian cancer from healthy controls, the sensitivity of a multivariate model combining the three biomarkers and CA125 [74% (95% CI, 52–90%)] was higher than that of CA125 alone [65% (95% CI, 43–84%)] at a matched specificity of 97% (95% CI, 89–100%). When compared at a fixed sensitivity of 83% (95% CI, 61–95%), the specificity of the model [94% (95% CI, 85–98%)] was significantly better than that of CA125 alone [52% (95% CI, 39–65%)]. These biomarkers demonstrated the potential to improve the detection of early stage ovarian cancer.

INTRODUCTION

Despite progress in cancer therapy, ovarian cancer mortality has remained virtually unchanged over the past two decades (1). Annually in the United States alone, ~23,000 women are diagnosed with the disease and almost 14,000 women die from it (1). Given our knowledge about the steep survival gradient relative to the stage at which the disease is diagnosed, it is reasonable to suggest that early detection remains the most promising approach to improve the long-term survival of ovarian cancer patients.

The relatively low prevalence (40 out of 100,000) of ovarian cancer among postmenopausal women in the general population, the lack of a clearly defined precursor lesion, and the high cost and possible complications associated with surgical confirmatory procedures have placed stringent requirements on any test intended for general population screening. Currently, none of the existing serum markers, such as CA125, CA 72–4, or macrophage colony-stimulating factor, can be

used individually for screening (2). Longitudinal studies are under way in Europe, Japan, and the United States to evaluate screening strategies using CA125 and/or transvaginal sonography (3–5) and their impact on overall cancer mortality (6). Preliminary results have shown encouraging evidence of a survival benefit among patients diagnosed through a screening regimen (3).

Reports from retrospective studies have shown that multivariate predictive models combining existing tumor markers improve cancer detection (7, 8). Recent advances in genomic and proteomic profiling technology have made it possible to apply computational methods to detect changes in protein expressions and their association to disease conditions, thereby hastening the identification of novel markers that may contribute to multimarker combinations with better diagnostic performance (9–13).

In this study, we hypothesized that comparison of protein expressions of serum specimens from patients with early stage ovarian cancer with those from healthy women could lead to the discovery of candidate biomarkers for the detection of early stage ovarian cancer. To ensure that the discovered biomarkers are truly associated with ovarian cancer rather than the result of biases in samples, profiling data of specimens from multiple institutions were used for cross-comparison and independent validation. We additionally determined the protein identities of the discovered biomarkers to allow for additional validation with independent methods and as a first step toward understanding the pathways in which they may function.

MATERIALS AND METHODS

Samples. The study involved a retrospective sample of 645 serum specimens. All were collected with institutional approval. Proteomic profiles were obtained from 503 specimens collected at four medical centers (M. D. Anderson Cancer Center, Duke University Medical Center, Groningen University Hospital, the Netherlands, and the Royal Hospital for Women, Australia). Among them, the cancer group consisted of 65 patients with stages I/II invasive epithelial ovarian cancer, 88 patients with stages III/IV invasive epithelial ovarian cancer, 28 patients with borderline tumors, and 14 patients with recurrent disease, all optimally staged by pathologists based on Fédération Internationale des Gynaeccologues et Obstetristes criteria. Among the stages I/II invasive cases, 20 were serous, 17 were mucinous, 15 were endometrioid, 8 were clear cell, 1 was carcinosarcoma, and 4 were mixed epithelial carcinoma. The samples also included 166 patients with benign pelvic masses and 142 healthy donors as controls. All of the samples were collected before the day of surgery or treatment, stored at -70°C , and thawed immediately before assay. CA125 levels had been obtained previously using a CA125II radioimmunoassay (Centocor). The clinical characteristics and age distribution of the proteomic profiling study population are summarized in Table 1.

In addition to the 503 specimens for proteomic profiling, 142 independent, archived serum specimens collected for routine clinical laboratory testing at the Johns Hopkins Medical Institutions were tested for levels of the identified biomarkers for which an immunoassay test was available. The sample included 41 healthy women, 41 patients with late-stage ovarian cancer, and groups of 20 patients each with breast, colon, and prostate cancers. All of the samples were processed promptly after collection and stored at $2-8^{\circ}\text{C}$ for a maximum of

Received 3/1/04; revised 5/17/04; accepted 5/26/04.

Grant support: National Cancer Institute Grant IP50 CA83639, UTMDACC Specialized Programs of Research Excellence in Ovarian Cancer (R. Bast, Jr., and Z. Zhang), and funding from Ciphergen Biosystems, Inc. (Z. Zhang, J. Li, A. Rai, J. Rosenzweig, B. Cameron, Y. Wang, and D. Chan).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Requests for reprints: Zhen Zhang, Center for Biomarker Discovery, Johns Hopkins Medical Institutions, 419 N. Caroline Street, Room 200, Baltimore, MD 21231. E-mail: zzhang7@jhmi.edu.

Table 1 Clinical characteristics and age distribution of 503 study samples used for proteomic profiling

Diagnostic groups and FIGO stages	n	Sample origin*				Age†		Histologic subtypes				
		A	B	C	D	Mean ± SD (median)	≥50 yr, %	Serous	Mucinous	Endometrioid	Clear cell	Others‡
Biomarker discovery sets												
Healthy control	79	30	49			39 ± 11 (38)	20.5					
Epithelial ovarian cancer	57	20	37			52 ± 16 (54)	57.9	18	24	11	4	
Stages I/II invasive cancer	42	20	22			53 ± 16 (55)	59.5	14	14	10	4	
Stage IIIA invasive cancer	2		2			54 ± 18 (54)	50.0	1		1		
Stages I/II borderline tumor	13		13			52 ± 19 (50)	53.8	3	10			
Independent validation set												
Healthy control	63				63	44 ± 10 (45)	34.9					
All epithelial ovarian cancer	138			138		57 ± 13 (57)	74.6	75	18	13	9	23
Stages I/II invasive cancer	23			23		56 ± 15 (56)	60.9	6	3	5	4	5
Stages III/IV invasive cancer	86			86		58 ± 13 (59)	80.2	50	10	7	4	15
Recurrent	14			14		52 ± 9 (52)	64.3	10		1	1	2
Stages I/II borderline tumor	12			12		55 ± 14 (57)	75.0	7	4			1
Stages III/IV borderline tumor	3			3		45 ± 14 (50)	66.7	2	1			
Benign pelvic masses	166	50	90	26		53 ± 16 (51)	55.5					

Abbreviation: FIGO, Federation Internationale des Gynaeologues et Obstetristes.

* Origin of samples: (A) Groningen University Hospital (Groningen, the Netherlands), (B) Duke University Medical Center (Durham, NC), (C) Royal Hospital for Women (Sydney, Australia), and (D) M. D. Anderson Cancer Center (Houston, TX).

† Age: Three samples with missing age data, one healthy control in biomarker discovery set and two patients with benign pelvic masses in independent validation set.

‡ Others: In the independent validation set, stage I/II invasive cancers included one carcinosarcoma and four mixed epithelial tumor; stage III/IV invasive cancers included four carcinosarcoma, six mixed epithelial tumor, and five unspecified adenocarcinoma; recurrent cancers included one mixed epithelial tumor and one unspecified adenocarcinoma; and stage I/II borderline tumors included one carcinosarcoma.

48 h before freezing at -70°C . CA125II assay was performed using a two-site immunoassay on the Tosoh AIA-600 II analyzer (Tosoh Medics).

Proteomic Expression Profiling. The ProteinChip Biomarker System (CIPHERGEN Biosystems), a platform for surface-enhanced laser desorption/ionization time-of-flight mass spectrometry, was used for protein expression profiling (14, 15). Samples from all of the centers were processed identically. To increase the total number of detectable peaks, serum samples (20 μl) were first fractionated via anion exchange chromatography by stepwise pH gradient elution (taken at pH 9/flow through, 7, 5, 4, 3, and organic solvent) using a Biomek 2000 equipped with the ProteinChip Biomarker Integration Package (CIPHERGEN Biosystems). A control sample of pooled human serum specimen (Intergen) was used for data calibration between experiments. Aliquots of each fraction were bound in triplicate with a randomized chip/spot allocation scheme to IMAC3-Cu, SAX2, H50, and WCX2 ProteinChip arrays. The energy absorbing molecule (crystallization matrix) for surface-enhanced laser desorption/ionization time-of-flight mass spectrometry, saturated sinapinic acid dissolved in 50% acetonitrile/0.5% trifluoroacetic acid, was applied promptly. The spotted arrays were read on PBS II ProteinChip readers. Instruments were monitored weekly for performance using insulin and immunoglobulin standards.

Spectra were externally calibrated, baseline subtracted, and normalized to total ion current within m/z (mass/charge) range of 1.5–150 kDa. Qualified mass peaks (signal/noise >5 ; cluster mass window at 0.3%) within the m/z range of 2–50 kDa were selected automatically. Logarithmic transformation was applied to the peak intensity before analysis for biomarker discovery. After biomarker discovery, the quality and intensity readings of the selected peaks were manually reconfirmed from raw spectra.

Multicenter Study Design. The diagram in Fig. 1 describes the design of this multicenter study and usage of samples for biomarker discovery, predictive model construction, and independent validation. To minimize the possibility of false discovery because of site-specific systematic biases from pre-analytical variables, a key feature of the design is that for biomarker discovery, the stage I/II ovarian cancer samples and healthy controls from Duke University Medical Center and Groningen University Hospital were analyzed separately according to sites. Only those potential biomarkers that were deemed statistically informative and shared the same up or down dysregulation patterns in analyses of data from both sites were additionally validated using the remaining samples at these two sites and all of the samples at the Royal Hospital for Women and M. D. Anderson Cancer Center. Samples from Johns Hopkins Medical Institutions were used only for additional independent confirmation by immunoassays. Because of insufficient specimen volume, the immunoassays were not tested on the samples from the first four institutions.

Bioinformatics Analysis. The unified maximum separability analysis algorithm implemented in the software ProPeak (3Z Informatics) was used to

analyze the peak intensity data and to select a subset of informative peaks. The unified maximum separability analysis incorporates information from the traditional multivariate statistical classification methods into the support vector machine algorithm (16) to provide a robust approach to analyzing high-dimensional expression data (12, 17–19). Bootstrap was also applied to help to identify peaks with a consistently high discriminatory power over multiple resampled subpopulations.

Protein Identification. The discovered biomarkers were purified according to individual biochemical properties using a series of protein separation procedures including anion exchange, size exclusion, and reverse-phase chromatography followed by SDS-PAGE separation. To monitor the purification process, healthy control samples were processed in parallel with the ovarian cancer samples. During each of the iterations, the new fractions were profiled on the same type of ProteinChip arrays used for discovery to monitor the presence or absence of the biomarkers of interest. The purified biomarkers were excised and digested with trypsin before being spotted onto NP20 ProteinChip arrays and read in a PBSII ProteinChip reader. The masses of the proteolytic fragments were used for database searching with the ProFound algorithm. For confirmation, the NP20 arrays containing the proteolytic fragments were analyzed by collision-induced dissociation using a Q-STAR MS/MS instrument (Applied Biosystems/MDS Sciex) equipped with a PCI 1000 ProteinChip Interface (CIPHERGEN Biosystems).

Multivariate Predictive Models. The profiling data from the two sites used for biomarker discovery were combined and then divided randomly into a training set and a test set for the derivation and testing of nonlinear multivariate predictive models using nonlinear unified maximum separability analysis. The derived models were then additionally evaluated, in the same way as the individual biomarkers, on the independent validation data that were not involved in biomarker discovery or model construction.

Statistical Analysis. The performance of the identified biomarkers in detecting early stage ovarian cancer from healthy controls was evaluated by descriptive statistics, Mann-Whitney U test or Kruskal-Wallis test, and receiver operating characteristic curves (ROCKIT; University of Chicago, Chicago, IL; Ref. 20). Performance of the nonlinear multivariate predictive models was compared with performance of the CA125 assay alone by receiver operating characteristic analysis and estimated sensitivity and specificity. All of the statistical analyses were performed using Statistica 6.1 (Statsoft).

RESULTS

Biomarker Discovery and Identification. Three potential biomarkers were discovered. Two of them were peaks from fraction pH 4 at m/z 12,828 and 28,043, both down-regulated in the cancer group, and the third was from fraction pH 9/flow through at m/z 3,272,




Fig. 1. Diagram of study design and patient flow showing the usage of samples from five academic medical centers. Numbers of samples in individual diagnostic groups from each site are indicated in parentheses. *A*, biomarker discovery and validation. *B*, derivation and validation of multivariate nonlinear predictive models.

up-regulated in the cancer group. All bound to the IMAC3-Cu (immobilized metal affinity chromatography array charged with copper ions) ProteinChip array (representative spectra in Fig. 2).

The m/z 28,043 peak was purified and identified as apolipoprotein A1 (z score = 2.38 using the ProFound algorithm; 59% coverage). Three peptides were sequenced by tandem mass spectrometry and confirmed this finding. The m/z 12,828 peak was purified and identified as a form of transthyretin (prealbumin; z -score 2.19; 91% coverage). One of the peptides was sequenced by tandem mass spectrometry and confirmed this finding. The m/z 12,828 peak copurified with a much more abundant m/z 13,900 peak, which was also identified as transthyretin. Immunoprecipitation and tandem mass spectrometry additionally determined that a truncated form of transthyretin lacking the NH₂-terminal 10 amino acids corresponded to the m/z 12,828 peak. The m/z 3,272 peak was purified from the pooled serum of ovarian cancer patients and its sequence was determined to be MNFRPGVLSRQLGLPGPPDVPDHAAYHPF, a fragment spanning amino acids 660–689 of human inter- α trypsin inhibitor, heavy chain H4 (PK-120). This result was confirmed by the analysis of pepsin digestion products of the marker.

Discriminatory Power of Individual Biomarkers. Table 2 provides the descriptive statistics (mean \pm SD, median) and results from statistical tests of these three biomarkers between healthy controls and ovarian cancer patients of different stage groups in the combined discovery sets and in the independent validation set. Within the discovery sets, the expression levels of the three biomarkers were statistically significantly different between the healthy controls and the early stage ovarian cancer patients ($P < 0.000001$ for all of the three markers). With the independent validation set, the two biomarkers at m/z 12,828 and m/z 28,043 retained their statistically discriminatory power in detecting stage I/II ovarian cancer ($P < 0.000001$

and $P < 0.000008$, respectively). However, the biomarker at m/z 3,272, with its large within-group variances, had only a marginal effect in separating healthy controls from epithelial ovarian cancer. In Table 3, the distributions of the biomarkers are compared among subgroups of the cancer samples based on stages, histological subtypes, and age.

Fig. 3, *A–D*, compares the discriminatory power of the individual biomarkers with that of CA125, using receiver operating characteristic analysis on data from patients with early stage ovarian cancer and healthy controls. CA125 and m/z 12,828 performed comparably on both the discovery and independent validation sets, whereas the other two markers had a lower area-under-curve than CA125 in one or both data sets. However, the estimated correlations among the three biomarkers and CA125 were low (data not shown), suggesting the possibility that they were complementary to each other and that a multivariate approach might outperform the single assay of CA125.

Because 27% of the samples in the healthy controls were from women age 50 or older compared with 61% of those in the early stage ovarian cancer group, we were concerned that these markers might reflect age-related changes. However, the differences between the early stage cancer samples and the healthy controls in the discovery data sets remained statistically significant in a multiple regression after adjusting for difference in age ($P = 0.000001$ for m/z 12,828 and m/z 3,272; and $P = 0.0174$ for m/z 28,043). In a stratified analysis, there were no appreciable differences in biomarker values between cancer patients of age ≥ 50 years and those below 50 years (Table 3). Previous population-based studies have shown that levels of apolipoprotein A1 actually increase slightly with age (21, 22).

Multivariate Predictive Models. The two data sets used for biomarker discovery were merged and then randomly divided into a training set and a test set. The training set had 28 ovarian cancer cases




Fig. 2. Segments of mass spectra showing the discovered biomarkers at m/z 3,272 from fraction pH 9 and at m/z 12,828 and 28,043 from fraction pH 4, all spotted on IMAC-Cu arrays. Peak m/z locations of the discovered biomarkers are marked by arrows. *Top*, two single segments showing details around the peak m/z locations. *Bottom*, stacked pseudo-gel view of segments from all of the spectra in the discovery set.

and 33 healthy controls, whereas the test set consisted of 29 cancer cases and 46 controls. The training and test data sets retained only the peak intensity values of the three discovered biomarkers and CA125 test results.

Two multivariate predictive models were constructed. The first used only the three biomarkers as its input and the second used the three biomarkers along with the CA125 level. Panels *E–H* in Fig. 3 compare the overall diagnostic performance of the two models with that of CA125 using receiver operating characteristic analysis, among which are the superimposed views of both the empirical and fitted receiver operating characteristic curves estimated using the healthy control samples against all of the stage I/II epithelial ovarian cancer samples (Fig. 3*G*) or the invasive stage I/II epithelial ovarian cancer samples (Fig. 3*H*) in the independent validation data set.

Using cutoffs that maximized the sum of sensitivity and specificity on the training data, these models were applied to the test data and the independent validation data (Table 4). For discrimination between healthy controls and stages I/II invasive ovarian cancer in the independent validation set, the multivariate model using the three biomarkers and CA125, at a sensitivity of 83% (95% CI, 61–95%), had a specificity of

94% (95% CI, 85–98%). This is statistically significantly better than CA125 at the same sensitivity of 83% (with a cutoff of 11 units/ml), which yielded a specificity of only 52% (95% CI, 39–65%). On the other hand, CA125 at the cutoff of 35 units/ml had a specificity of 97% (95% CI, 89–100%) and a corresponding sensitivity of 65% (95% CI, 43–84%). At the same fixed specificity, the multivariate model using the three biomarkers and CA125 resulted in a sensitivity of 74% (95% CI, 52–90%). The difference, however, is not statistically significant, partially because of the few stage I/II invasive cases in the independent validation set. Table 4 also lists in detail the estimated sensitivities or specificities for individual diagnostic groups in the training set, test set, and independent validation set.

It should be noted that the biomarkers, with the exception of m/z 3,272, as well as the two predictive models were moderately capable of differentiating stages I/II invasive cancer from benign cases in the independent validation set ($P = 0.002$ and 0.09 for m/z 12,828 and 28,043, respectively; and $P = 0.02$ and 0.03 for models without CA125 and with CA125, respectively).

Evaluation Using Immunoassays. The 142 archived specimens from the Johns Hopkins Medical Institutions were analyzed for apo-

Table 2 Comparison of distributions of serum levels of CA125 and the discovered biomarkers among samples from different diagnostic groups

Diagnostic groups and FIGO stages	n	Biomarker level (peak intensity)* (mean ± SD, median) and P†								
		CA125 (units/ml)	P	m/z 12,828	P	m/z 28,043	P	m/z 3,272	P	
Biomarker discovery sets										
Healthy control	79	13.0 ± 10.2 (10.2)		0.40 ± 0.09 (0.42)		0.58 ± 0.25 (0.52)		0.57 ± 1.06 (0.32)		
Epithelial ovarian cancer	57	104.1 ± 145.4 (49.7)	0.000000	0.27 ± 0.09 (0.28)	0.000000	0.36 ± 0.26 (0.29)	0.000000	2.70 ± 2.61 (1.76)	0.000000	
Stages I/II invasive cancer	42	98.2 ± 143.7 (47.4)	0.000001	0.27 ± 0.10 (0.28)	0.000001	0.34 ± 0.27 (0.26)	0.000001	2.11 ± 2.01 (1.54)	0.000007	
Stage IIIA invasive cancer	2	118.9 ± 115.5 (118.9)	n.c.	0.24 ± 0.04 (0.24)	n.c.	0.29 ± 0.16 (0.29)	n.c.	5.78 ± 7.87 (5.78)	n.c.	
Stages I/II borderline tumor	13	120.8 ± 162.9 (82.6)	0.000001	0.25 ± 0.08 (0.26)	0.000004	0.42 ± 0.22 (0.31)	0.03	4.11 ± 2.82 (3.95)	0.000000	
Independent validation set										
Healthy control	63	13.0 ± 12.2 (10.5)		0.39 ± 0.09 (0.38)		0.80 ± 0.41 (0.74)		0.64 ± 0.81 (0.28)		
All epithelial ovarian cancer	138	330.7 ± 337.2 (195.1)	0.000000	0.20 ± 0.09 (0.20)	0.000000	0.42 ± 0.30 (0.32)	0.000000	0.98 ± 1.30 (0.41)	0.09	
Stages I/II invasive cancer	23	180.5 ± 224.7 (74.5)	0.000002	0.22 ± 0.09 (0.21)	0.000000	0.42 ± 0.26 (0.32)	0.000008	1.10 ± 1.59 (0.37)	0.54	
Stages III/IV invasive cancer	86	433.2 ± 357.3 (347.4)	0.000000	0.20 ± 0.08 (0.19)	0.000000	0.39 ± 0.26 (0.31)	0.000000	0.76 ± 0.99 (0.37)	0.40	
Recurrent	14	208.2 ± 249.1 (95.4)	0.000001	0.20 ± 0.10 (0.20)	0.000000	0.43 ± 0.29 (0.36)	0.0006	1.52 ± 1.71 (0.74)	0.05	
Stages I/II borderline tumor	12	88.3 ± 152.9 (12.2)	0.11	0.22 ± 0.11 (0.23)	0.000012	0.66 ± 0.50 (0.37)	0.20	1.72 ± 1.84 (0.91)	0.01	
Stages III/IV borderline tumor	3	84.2 ± 124.2 (18.7)	0.22	0.19 ± 0.04 (0.21)	0.004	0.55 ± 0.24 (0.63)	0.27	0.64 ± 0.42 (0.60)	0.36	
Benign pelvic masses	166	36.1 ± 54.5 (15.7)	0.000000‡	0.30 ± 0.12 (0.31)	0.000000‡	0.50 ± 0.29 (0.47)	0.0009‡	2.94 ± 2.68 (2.16)	n.c.§	

Abbreviation: n.c., not computed. FIGO, Federation Internationale des Gynaecologistes et Obstetristes.

* The total ion current (area-under-spectrum curve) within the m/z range 1.5 kDa–150 kDa of all spectra were normalized to the constant 0.2 before analysis.

† P of all two-group comparisons estimated by Mann-Whitney U test; P estimated between patients with ovarian cancer and healthy controls unless specified otherwise.

‡ P estimated between patients with benign pelvic mass and patients with epithelial ovarian cancer in the independent validation set.

§ Mean level of biomarker m/z 3,272 was higher among patients with benign pelvic mass than in patients with epithelial ovarian cancer in the independent validation set.

lipoprotein A1 using a turbidimetric immunoassay performed in a microtiter plate format (Wako Chemical USA) and for transthyretin using a particle enhanced turbidimetric immunoassay performed on the Dimension RxL Instrument (Dade-Behring; Table 5). The serum levels of CA125 were up-regulated among the 41 patients with late-stage ovarian cancer compared with the 41 healthy controls (mean ± SD in units/ml, 2388 ± 4723 versus 18 ± 23, P = 0.000000), whereas levels of apolipoprotein A1 and transthyretin were down-regulated (mean ± SD in mg/dl, 122 ± 42 versus 153 ± 28, P = 0.0004, and 20 ± 7 versus 27 ± 6, P = 0.00005, respectively). The mean serum apolipoprotein A1 level among the healthy controls was not significantly different from that of patients with breast or colorectal cancer (P = 0.77 and P = 0.69, respectively) and only marginally different from that of patients with prostate cancer (P = 0.02). The mean serum transthyretin level was down-regulated among patients with colorectal cancer (P = 0.01) albeit to a lesser degree than that in patients with ovarian cancer. The differences in mean serum transthyretin levels between the healthy controls and patients with breast or prostate cancer were not significant (P = 0.51 and P = 0.22, respectively).

DISCUSSION

Differential analysis of serum protein profiles from patients with early stage ovarian cancer and healthy women revealed three biomarkers the discriminatory power of which was confirmed with samples

from multiple institutions through cross-validation and independent validation. Combined with CA125 in a multivariate predictive model, these biomarkers improve significantly on the specificity of CA125 alone although maintaining a relatively high sensitivity. Two of the markers were also evaluated using immunoassays. Although the immunoassay for transthyretin was not specific for the particular truncated form that corresponds to the m/z 12,828 peak, the results corroborated the findings from the surface-enhanced laser desorption/ionization mass spectrum data and provided preliminary analysis of tumor site specificity of these two markers.

Results from receiver operating characteristic curve analysis (Fig. 3, G and H) show that at a fixed high sensitivity, the predictive models, with or without CA125 as one of its inputs, had a much-improved specificity over that of CA125 alone. However, when the models were compared with CA125 at a fixed high specificity, the improvement in sensitivity was more moderate and tended to diminish as specificity approached 98% and above. The level of analytical variability in peak intensity measurement relative to the few early stage ovarian cancer cases in the independent validation set made it difficult to evaluate in a statistically meaningful way the diagnostic performance of the individual biomarkers or the multivariate models at extreme specificity or sensitivity values.

The differences in area-under-curve in detecting stage I/II invasive cancer from healthy controls were not statistically significant for the independent validation set. Given the fairly large absolute differences

Table 3 Comparison of distributions of serum levels of CA125 and the discovered biomarkers among different groups of epithelial ovarian cancer patients from combined discovery and independent validation sets

Groups	n	Biomarker level (peak intensity)*, (mean ± SD, median) and P†								
		CA125 (units/ml)	P	m/z 12,828	P	m/z 28,043	P	m/z 3,272	P	
By stage										
Stages I/II invasive cancer	65	127.3 ± 179.3 (55.8)		0.25 ± 0.10 (0.25)		0.37 ± 0.27 (0.28)		1.76 ± 1.92 (0.87)		
Stage I/II borderline tumor	25	105.2 ± 155.8 (34.5)	0.0001	0.24 ± 0.09 (0.25)	0.0006	0.54 ± 0.39 (0.32)	0.13	2.96 ± 2.65 (2.30)	0.0001	
Stages III/IV invasive cancer	88	426.1 ± 356.5 (321.6)		0.20 ± 0.08 (0.19)		0.38 ± 0.26 (0.31)		0.88 ± 1.50 (0.37)		
By subtype										
Serous	93	294.9 ± 328.7 (151.9)		0.21 ± 0.09 (0.20)		0.43 ± 0.32 (0.32)		1.56 ± 2.10 (0.52)		
Mucinous	42	175.4 ± 240.3 (71.7)	0.23	0.26 ± 0.09 (0.26)	0.02	0.43 ± 0.27 (0.39)	0.48	1.72 ± 2.14 (0.79)	0.88	
Endometrioid	24	225.8 ± 265.8 (93.3)		0.22 ± 0.08 (0.23)		0.36 ± 0.28 (0.27)		1.63 ± 1.88 (0.63)		
Clear cell	13	192.1 ± 264.9 (107.7)		0.22 ± 0.07 (0.22)		0.33 ± 0.21 (0.28)		0.96 ± 0.41 (1.24)		
By age										
Age ≥ 50 years	136	301.0 ± 327.0 (132.6)	0.004	0.21 ± 0.09 (0.21)	0.02	0.39 ± 0.27 (0.31)	0.52	1.46 ± 1.94 (0.54)	0.97	
Age < 50 years	59	180.2 ± 255.8 (55.8)		0.24 ± 0.10 (0.24)		0.43 ± 0.32 (0.35)		1.52 ± 1.97 (0.52)		

* The total ion current (area-under-spectrum curve) within the m/z range 1.5 kDa–150 kDa of all spectra were normalized to the constant 0.2 before analysis.

† P estimated by Mann-Whitney U test for two-group comparison and by Kruskal-Wallis test for multigroup comparison.




Fig. 3. Comparison of receiver operating characteristic curves between CA125 and three discovered biomarkers (A–D) and the two multivariate predictive models (E–H). For A–D, (1) CA125; (2) m/z 12,828; (3) m/z 28,043; and (4) m/z 3,272. For E–H, ○, CA125; □, multivariate model using the three biomarkers and CA125. G–H, the fitted receiver operating characteristic curves are superimposed by the empirical receiver operating characteristic curves. *, Part of biomarker discovery set. †, Epithelial ovarian cancer included both invasive and borderline tumors. Panels G and H: vertical and horizontal dashed lines correspond to 90% specificity and 83% sensitivity, respectively.

in area-under-curve, a likely reason would be the relatively few stage I/II invasive cases available.

Serum levels of CA125 have been used widely for distinguishing benign from malignant pelvic masses and for monitoring the clinical course of patients with ovarian cancer. However, CA125 is elevated in only about half of stage I/II ovarian cancer patients (23). Lowering the cutoff of CA125 would increase its sensitivity in detecting stage I/II cancer but result in many false positives in patients with benign conditions as well as in healthy women. The improvement in sensitivity by the multivariate models to detect early stage cancer also increased the false-positive rate among patients with benign pelvic masses, but unlike with CA125, it did not lead to appreciable concomitant loss of specificity among healthy women. At a sensitivity of 83% and specificity of 94%, the current multivariate model by itself is not suitable for general population screening. However, in a two-stage approach with a combination of a serum test followed by sonography, a high specificity might be reached at an acceptable total cost.

The identified biomarkers have been characterized generally as

acute-phase reactants. The down-regulation of transthyretin and apolipoprotein A1 in ovarian cancer patients suggests the possibility that they are byproducts of the host response to the tumor. Although recent data have supported the concept that inflammation is a critical component of tumor progression (24), such acute-phase reactants may still represent only epiphenomena because of the presence of tumor and may not be specific to a particular type of cancer (25, 26). In this study, with preliminary data from immunoassays analysis, we were able to verify that the levels of apolipoprotein A1 were not altered in the serum of breast or colon cancer patients and the levels of transthyretin were not altered in the serum of breast or prostate cancer patients.

Transthyretin and lipoprotein(a) have been separately reported to be decreased in epithelial ovarian cancer (27, 28). Transthyretin is the major carrier for serum thyroxine and tri-iodothyronine and facilitates the transport of retinol via its interaction with retinol binding protein. Transgenic mice lacking transthyretin expression have dramatically lower levels of retinol and retinol binding protein, and decreased levels of retinol binding protein as well as cellular retinol binding

Table 4 Comparison of sensitivities and specificities of two multivariate predictive models and CA125

	Specificity, %			Sensitivity, %					
	Healthy control	Benign pelvic mass	Epithelial ovarian cancer	Group by stage					
				Stages I/II invasive cancer	All stages I/II cancer*	Stage III/IV invasive cancer	Recurrent	Stages I/II borderline tumor	Stages III/IV borderline tumor
Training set									
CA125 at cutoff = 35 units/ml	97 (32/33)		68 (19/28)	64 (14/22)	67 (18/27)	100 (1/1)		80 (4/5)	
Model with three markers	88 (29/33)		89 (25/28)	86 (19/22)	89 (24/27)	100 (1/1)		100 (5/5)	
Model with three markers + CA125	97 (32/33)		100 (28/28)	100 (22/22)	100 (27/27)	100 (1/1)		100 (5/5)	
Test set									
CA125 at cutoff = 35 units/ml	98 (45/46)		66 (19/29)	70 (14/20)	64 (18/28)	100 (1/1)		50 (4/8)	
Model with three markers	91 (42/46)		83 (24/29)	75 (15/20)	82 (23/28)	100 (1/1)		100 (8/8)	
Model with three markers + CA125	98 (45/46)		86 (25/29)	80 (16/20)	86 (24/28)	100 (1/1)		100 (8/8)	
Independent validation set									
Fixed specificity for healthy controls									
CA125 at cutoff = 35 units/ml	97 (61/63)	76 (126/166)	77 (106/138)	65 (15/23)	54 (19/35)	87 (75/86)	79 (11/14)	33 (4/12)	33 (1/3)
Model with three markers	97 (61/63)	52 (86/166)	78 (107/138)	74 (17/23)	74 (26/35)	81 (70/86)	71 (10/14)	75 (9/12)	33 (1/3)
Model with three markers + CA125	97 (61/63)	45 (75/166)	78 (108/138)	74 (17/23)	66 (23/35)	85 (73/86)	79 (11/14)	50 (6/12)	33 (1/3)
Fixed sensitivity for stage I/II invasive cancer									
CA125 at cutoff = 11 units/ml	52 (33/63)†	32 (53/166)	88 (122/138)	83 (19/23)	74 (26/25)	94 (81/86)	93 (13/14)	58 (7/12)	67 (2/3)
Model with three markers	89 (56/63)†	35 (58/166)	82 (113/138)	83 (19/23)	80 (28/35)	86 (74/86)	71 (10/14)	75 (9/12)	33 (1/3)
Model with three markers + CA125	94 (59/63)†	33 (55/166)	82 (113/138)	83 (19/23)	71 (25/35)	87 (75/86)	86 (12/14)	50 (6/12)	33 (1/3)

* All stage I/II cancer included stage I/II invasive epithelial ovarian cancer and stage I/II borderline tumor.

† 95% confidence intervals for CA125 at cutoff = 11 units/ml, model with three markers, and model with three marker + CA125 were 39–65%, 78–95%, and 85–98%, respectively.

protein have been shown to be associated with an increased rate of malignant transformation of ovarian epithelium (29, 30). In addition, levels of cellular retinal binding proteins have been reported to be changed in ovarian cancer by oligonucleotide array analysis (31) and have been shown to be decreased in approximately one-third of ovarian cancers by immunohistochemistry (32).

The carboxyl portion of inter- α trypsin inhibitor, heavy chain H4, from which the m/z 3,272 biomarker is derived, has been shown to be a substrate for plasma kallikrein (33, 34). The fragment that we identified as a biomarker differs from the postulated fragment derived from cleavage by plasma kallikrein, suggesting a different protease is responsible for the generation of this biomarker. Kallikrein proteases consist of plasma kallikrein and tissue kallikreins, which have overlapping substrate specificity (35). The tissue kallikreins are products of a large multigene family that includes prostate specific antigen (hK3), a tumor marker for prostate cancer. Gene expression of seven of the tissue kallikreins has been found to be up-regulated in ovarian cancer patients, and expression of hK6, hK10, and hK11 has been found to be increased in >50% of patients with ovarian cancer (36, 37).

The identified proteins associated with the three biomarkers are all of high abundance in serum and are unlikely to be released by tumor cells. This limitation has been pointed out previously as being common among studies using mass spectrometry based proteomic profiling (25, 26, 38). However, this is not purely a limitation of the mass spectrometry technological platforms, but rather reflects the limitation of high throughput sample preprocessing technologies that would allow us to examine subsets of the serum proteome without introduc-

ing additional significant analytical variability. The use of sample prefractionation in addition to multiple array surfaces in this study is a first attempt in surface-enhanced laser desorption/ionization mass spectrometry based clinical proteomics for biomarker discovery.

Two of the markers found in this study are cleavage products of precursor proteins and present in the serum at a fraction of the concentrations of their corresponding full-length proteins. Neither of these truncation products has been reported previously, indicating that they are new candidate markers. These markers may be the product of cleavage by one or more proteases, including plasma kallikrein, tissue kallikreins, matrix metalloproteases, or prostasin, a trypsin-like serine protease that was reported recently to be increased in cases of ovarian cancer (39). The proteases that generate these markers may themselves act as markers that can be combined with those discovered here in a predictive model. More generally, these results support the increasing body of evidence indicating alterations in the balance of protease and protease inhibitor activity in the serum and tissue of patients with cancer (40, 41). Preliminary results from additional immunoprecipitation pull-down analysis suggest that the forms and relative frequencies of posttranslational modifications of inter- α trypsin inhibitor, heavy chain H4 and transthyretin may be associated with the ovarian cancer disease status (data not shown).

The use of proteomics to identify potential biomarkers has been explored previously for the detection of a number of cancers including prostate (13, 42–44), colon (45), bladder (46), breast (12), and ovarian (9, 17). For example, Petricoin *et al.* reported the use of self-organizing map coupled with genetic algorithm to search through raw mass spectrum data for informative variables and to form clusters of train-

Table 5 Comparison of distributions of serum levels of CA125 and the discovered biomarkers among 142 archived samples for independent validation using immunoassays for apolipoprotein A1 and transthyretin

Diagnostic groups	n	Age mean \pm SD	Biomarker level (concentration) (mean \pm SD, median) and P*					
			CA125 (units/ml)	P	Apolipoprotein A1 (mg/dl)	P	Transthyretin (mg/dl)	P
Healthy control	41	33 \pm 10	17.9 \pm 23.0 (10.2)		152.8 \pm 28.1 (154.3)		26.7 \pm 5.8 (25.1)	
Stage III/IV ovarian cancer	41	59 \pm 11	2387.9 \pm 4723.3 (426.6)	0.000000	121.5 \pm 41.9 (123.9)	0.0004	19.8 \pm 7.1 (18.7)	0.00005
Breast cancer	20	53 \pm 14	13.4 \pm 13.0 (7.9)	0.39	154.4 \pm 31.8 (145.9)	0.77	26.8 \pm 4.8 (26.4)	0.51
Colon cancer	20	70 \pm 15	17.0 \pm 14.2 (10.9)	0.86	144.2 \pm 39.7 (149.5)	0.69	22.4 \pm 5.1 (22.8)	0.01
Prostate cancer	20	57 \pm 7	13.1 \pm 21.2 (7.6)	0.07	138.2 \pm 21.2 (136.1)	0.02	27.8 \pm 7.8 (29.8)	0.22

* P estimated by Mann-Whitney U test.

ing samples as the basis of a predictive model. More recently, Kozak *et al.* (47) reported several panels of selected surface-enhanced laser desorption/ionization peaks for the detection of ovarian cancer. A number of these peaks and their expression patterns are consistent with those from our current or previous results (17).

Biomarker discovery using clinical proteomics involves the simultaneous analysis of expression levels of many proteins measured on a relatively few clinical samples. The difficulty in statistical analysis is complicated additionally by the possibility of nondisease-related biases and variability in data from preanalytical and analytical variables and within-group heterogeneity associated with clinical samples. A major concern has always been whether the discovered biomarkers and the derived multivariate models are truly associated with the disease process. Recent reports examined and highlighted the danger of such issues (26, 38, 48). The current study had a number of features that were designed to alleviate the impact of these factors. First, only the intensities of detectable peaks were used in analysis, which in general are less sensitive to mass shift in raw spectrum data, and the discovered biomarkers are more likely to have an identifiable biological identity. Second, biomarker discovery and construction of predictive models were done in two separate stages. During the discovery stage, only linear models were used to evaluate the informativeness of individual peaks. The bootstrap resampling was used to additionally ensure that the peak selection results are robust among multiple subpopulations. It was only after a small group of biomarkers had been selected and individually validated across the multiple data sets that the more complicated nonlinear unified maximum separability analysis predictive models were derived to combine the multiple markers. It would have been difficult to explain and validate the exact roles of individual inputs if we had started the peak selection directly with a complex nonlinear model using raw spectrum data. Third, in this study, the data sets from multiple sites were used separately to cross-validate discoveries from other data sets and to detect systematic biases in data such as those because of differences in preanalytical sample processing procedures. This differs significantly from the approaches in which data from multiple sites are pooled together and then divided through randomization into (artificially made) identically distributed training and test data sets. Considering the high cost associated with post discovery validation, however, we believe that the conservative approach that we used is more close to the real clinical environment and appropriate for biomarker discovery through expression profiling. Finally, the identification of the discovered biomarkers allowed additional confirmation through immunoassay tests.

Previous studies have illustrated the benefits of combining multiple markers and the longitudinal use of tumor markers in the detection of ovarian cancers (8, 49, 50). New biomarkers derived from proteomic analysis of clinical samples, once validated, may provide additional choices in the selection of an optimal panel of markers that, through a multivariate approach, would be capable of detecting early stage ovarian cancer in a more general population. Studies remain to be performed to evaluate these markers individually and in combination in larger populations and, most importantly, in preclinical samples from women with ovarian cancer obtained in screening trials. It should be pointed out, however, that intensity data from mass spectra measure the relative abundance of proteins. In this study, all of the surface-enhanced laser desorption/ionization data were generated under well-controlled conditions, with which we were able to demonstrate the performance of the combined three markers and their complementary value to CA125, using samples across multiple sites. The successful application of the multivariate predictive models to new clinical samples, however, will require additional work to assure the consistency of measured peak intensities between multiple runs

and across different instruments and sites. Knowledge of the identities of the biomarkers will certainly help to expedite future development of assays.

REFERENCES

- Jemal A, Thomas A, Murray T, Thun M. Cancer statistics, 2002. *CA - Cancer J Clin* 2002;52:23-47.
- Bast RC Jr, Xu FJ, Yu YH, et al. CA 125: the past and the future. *Int J Biol Markers* 1998;13:179-87.
- Jacobs IJ, Skates SJ, MacDonald N, et al. Screening for ovarian cancer: a pilot randomised controlled trial. *Lancet* 1999;353:1207-10.
- van Nagell JR Jr, DePriest PD, Reedy MB, et al. The efficacy of transvaginal sonographic screening in asymptomatic women at risk for ovarian cancer. *Gynecol Oncol* 2000;77:350-6.
- Hayashi H, Yaginuma Y, Kitamura S, et al. Bilateral oophorectomy in asymptomatic women over 50 years old selected by ovarian cancer screening. *Gynecol Obstet Investig* 1999;47:58-64.
- Menon U, Jacobs I. Screening for ovarian cancer. *Best Pract Res Clin Obstet Gynaecol* 2002;16:469-82.
- van Haften-Day C, Shen Y, Xu F, et al. OVX1, macrophage-colony stimulating factor, and CA-125-II as tumor markers for epithelial ovarian carcinoma: a critical appraisal. *Cancer (Phila)* 2001;92:2837-44.
- Zhang Z, Barnhill SD, Zhang H, et al. Combination of multiple serum markers using an artificial neural network to improve specificity in discriminating malignant from benign pelvic masses. *Gynecol Oncol* 1999;73:56-61.
- Petricoin EF, Ardekani AM, Hitt BA, et al. Use of proteomic patterns in serum to identify ovarian cancer. *Lancet* 2002;359:572-7.
- van de Vijver MJ, He YD, van't Veer LJ, et al. A gene-expression signature as a predictor of survival in breast cancer. *N Engl J Med* 2002;347:1999-2009.
- Rosenwald A, Wright G, Chan WC, et al. The use of molecular profiling to predict survival after chemotherapy for diffuse large-B-cell lymphoma. *N Engl J Med* 2002;346:1937-47.
- Li J, Zhang Z, Rosenzweig J, Wang YY, Chan DW. Proteomics and bioinformatics approaches for identification of serum biomarkers to detect breast cancer. *Clin Chem* 2002;48:1296-304.
- Qu Y, Adam BL, Yasui Y, et al. Boosted decision tree analysis of surface-enhanced laser desorption/ionization mass spectral serum profiles discriminates prostate cancer from noncancer patients. *Clin Chem* 2002;48:1835-43.
- Hutchens TW, Yip TT. New desorption strategies for the mass spectrometric analysis of micromolecules. *Rapid Commun Mass Spectrom* 1993;7:576-80.
- Fung ET, Thulasiraman V, Weinberger SR, Dalmasso EA. Protein biochips for differential profiling. *Curr Opin Biotechnol* 2001;12:65-9.
- Vapnik VN. *Statistical learning theory*. Wiley-Interscience; 1998. p. 736.
- Rai AJ, Zhang Z, Rosenzweig J, et al. Proteomic approaches to tumor marker discovery. *Arch Pathol Lab Med* 2002;126:1518-26.
- Zhang Z, Page G, Zhang H. Applying classification separability analysis to microarray data. In: Lin SM, Johnson KF, editors. *Methods of microarray data analysis: papers from CAMDA '00*. Boston: Kluwer Academic Publishers; 2001. p. 125-36.
- Zhang Z, Page G, Zhang H. Fishing expedition—a supervised approach to extract patterns from a compendium of expression profiles. In: Lin SM, Johnson KF, editors. *Microarray data analysis II: papers from CAMDA '01*. Boston: Kluwer Academic Publishers; 2002.
- Metz CE. Basic principles of ROC analysis. *Semin Nucl Med* 1978;8:283-98.
- Jungner I, Marcovina SM, Walldius G, et al. Apolipoprotein B and A-I values in 147576 Swedish males and females, standardized according to the World Health Organization-International Federation of Clinical Chemistry First International Reference Materials. *Clin Chem* 1998;44:1641-9.
- Bachorik PS, Lovejoy KL, Carroll MD, Johnson CL. Apolipoprotein B and AI distributions in the United States, 1988-1991: results of the National Health and Nutrition Examination Survey III (NHANES III). *Clin Chem* 1997;43:2364-78.
- Jacobs I, Bast RC Jr. The CA 125 tumour-associated antigen: a review of the literature. *Hum Reprod* 1989;4:1-12.
- Coussens LM, Werb Z. Inflammation and cancer. *Nature (Lond)* 2002;420:860-7.
- Diamandis EP. Proteomic patterns in serum and identification of ovarian cancer. *Lancet* 2002;360:170.
- Diamandis EP. Analysis of serum proteomic patterns for early cancer diagnosis: drawing attention to potential problems. *J Natl Cancer Inst (Bethesda)* 2004;96:353-6.
- Mahlck CG, Grankvist K. Plasma prealbumin in women with epithelial ovarian carcinoma. *Gynecol Obstet Investig* 1994;37:135-40.
- Kuesel AC, Kroft T, Prefontaine M, Smith IC. Lipoprotein(a) and CA125 levels in the plasma of patients with benign and malignant ovarian disease. *Int J Cancer* 1992;52:341-6.
- van Bennekum AM, Wei S, Gamble MV, et al. Biochemical basis for depressed serum retinol levels in transthyretin-deficient mice. *J Biol Chem* 2001;276:1107-13.
- Roberts D, Williams SJ, Cvetkovic D, et al. Decreased expression of retinol-binding proteins is associated with malignant transformation of the ovarian surface epithelium. *DNA Cell Biol* 2002;21:11-9.
- Giordano TJ, Shedden KA, Schwartz DR, et al. Organ-specific molecular classification of primary lung, colon, and ovarian adenocarcinomas using gene expression profiles. *Am J Pathol* 2001;159:1231-8.
- Cvetkovic D, Williams SJ, Hamilton TC. Loss of cellular retinol-binding protein 1 gene expression in microdissected human ovarian cancer. *Clin Cancer Res* 2003;9:1013-20.

33. Pu XP, Iwamoto A, Nishimura H, Nagasawa S. Purification and characterization of a novel substrate for plasma kallikrein (PK-120) in human plasma. *Biochim Biophys Acta* 1994;1208:338–43.
34. Nishimura H, Kakizaki I, Muta T, et al. cDNA and deduced amino acid sequence of human PK-120, a plasma kallikrein-sensitive glycoprotein. *FEBS Lett* 1995;357:207–11.
35. Diamandis EP, Yousef GM. Human tissue kallikreins: a family of new cancer biomarkers. *Clin Chem* 2002;48:1198–205.
36. Yousef GM, Polymeris ME, Yacoub GM, et al. Parallel overexpression of seven kallikrein genes in ovarian cancer. *Cancer Res* 2003;63:2223–7.
37. Yousef GM, Diamandis EP. Kallikreins, steroid hormones and ovarian cancer: is there a link? *Minerva Endocrinol* 2002;27:157–66.
38. Diamandis EP. Mass spectrometry as a diagnostic and a cancer biomarker discovery tool: opportunities and potential limitations. *Mol Cell Proteomics* 2004;3:367–78.
39. Mok SC, Chao J, Skates S, et al. Prostatein, a potential serum marker for ovarian cancer: identification through microarray technology. *J Natl Cancer Inst (Bethesda)* 2001;93:1458–64.
40. Matrisian LM, Sledge GW Jr, Mohla S. Extracellular proteolysis and cancer: meeting summary and future directions. *Cancer Res* 2003;63:6105–9.
41. Yousef GM, Diamandis EP. The new human tissue kallikrein gene family: structure, function, and association to disease. *Endocr Rev* 2001;22:184–204.
42. Petricoin EF III, Ornstein DK, Pawletz CP, et al. Serum proteomic patterns for detection of prostate cancer. *J Natl Cancer Inst* 2002;94:1576–8.
43. Banez LL, Prasanna P, Sun L, et al. Diagnostic potential of serum proteomic patterns in prostate cancer. *J Urol* 2003;170:442–6.
44. Adam BL, Qu Y, Davis JW, et al. Serum protein fingerprinting coupled with a pattern-matching algorithm distinguishes prostate cancer from benign prostate hyperplasia and healthy men. *Cancer Res* 2002;62:3609–14.
45. Lawrie LC, Curran S, McLeod HL, Fothergill JE, Murray GI. Application of laser capture microdissection and proteomics in colon cancer. *Mol Pathol* 2001;54:253–8.
46. Vlahou A, Schellhammer PF, Mendrinis S, et al. Development of a novel proteomic approach for the detection of transitional cell carcinoma of the bladder in urine. *Am J Pathol* 2001;158:1491–502.
47. Kozak KR, Amneus MW, Pusey SM, et al. Identification of biomarkers for ovarian cancer using strong anion-exchange ProteinChips: Potential use in diagnosis and prognosis. *Proc Natl Acad Sci USA* 2003;100:12343–8.
48. Baggerly KA, Morris JS, Coombes KR. Reproducibility of SELDI-TOF protein patterns in serum: comparing data sets from different experiments. *Bioinformatics* 2004;20:777–85.
49. Woolas RP, Conaway MR, Xu F, et al. Combinations of multiple serum markers are superior to individual assays for discriminating malignant from benign pelvic masses. *Gynecol Oncol* 1995;59:111–6.
50. Skates SJ, Pauler DK, Jacobs I. Screening based on the risk of cancer calculation from Bayesian hierarchical changepoint and mixture models of longitudinal markers. *J Am Stat Assoc* 2001;96:429–39.