

Physical Activity in the Prevention of Type 2 Diabetes

The Finnish Diabetes Prevention Study

David E. Laaksonen,^{1,2} Jaana Lindström,³ Timo A. Lakka,^{2,4,5} Johan G. Eriksson,³ Leo Niskanen,¹ Katja Wikström,³ Sirkka Aunola,^{6,7} Sirkka Keinänen-Kiukaanniemi,^{8,9} Mauri Laakso,⁸ Timo T. Valle,³ Pirjo Ilanne-Parikka,^{10,11} Anne Louheranta,¹² Helena Hämäläinen,⁷ Merja Rastas,^{3,7} Virpi Salminen,¹³ Zygimantas Cepaitis,³ Martti Hakumäki,^{2,12} Hannu Kaikkonen,^{8,9} Pirjo Härkönen,^{8,9} Jouko Sundvall,¹⁴ Jaakko Tuomilehto,^{3,15} and Matti Uusitupa,¹² for the Finnish Diabetes Prevention Study Group

Clinical trials have demonstrated that lifestyle changes can prevent type 2 diabetes, but the importance of leisure-time physical activity (LTPA) is still unclear. We carried out post hoc analyses on the role of LTPA in preventing type 2 diabetes in 487 men and women with impaired glucose tolerance who had completed 12-month LTPA questionnaires. The subjects were participants in the Finnish Diabetes Prevention Study, a randomized controlled trial of lifestyle changes including diet, weight loss, and LTPA. There were 107 new cases of diabetes during the 4.1-year follow-up period. Individuals who increased moderate-to-vigorous LTPA or strenuous, structured LTPA the most were 63–65% less likely to develop diabetes. Adjustment for changes in diet and body weight during the study attenuated the association somewhat (upper versus lower third: moderate-to-vigorous LTPA, relative risk 0.51, 95% CI 0.26–0.97; strenuous, structured LTPA, 0.63, 0.35–1.13). Low-intensity and lifestyle LTPA and walking also conferred benefits, consistent with the finding that the change in total LTPA (upper versus lower third: 0.34, 0.19–0.62)

was the most strongly associated with incident diabetes. Thus increasing physical activity may substantially reduce the incidence of type 2 diabetes in high-risk individuals. *Diabetes* 54:158–165, 2005

Epidemiological studies have suggested that moderate-to-vigorous leisure-time physical activity (LTPA) protects against the development of type 2 diabetes (1–6) as well as the metabolic syndrome (7), which commonly precedes diabetes and cardiovascular disease (8,9). In intervention trials, physical exercise has variably reduced body weight, visceral fat accumulation (10–13), and insulin resistance (10,11,14–17); improved glucose tolerance (12,15,17) and lipid profile (12,18); and decreased blood pressure (12,15,19). It is not yet clear to what extent concomitant weight loss and alterations in body composition influence these metabolic changes (20). Moderate-to-vigorous LTPA may be more beneficial than low-intensity LTPA with respect to metabolic and chronic disease end points (2,7,12,21).

Structured aerobic exercise has traditionally been emphasized as being effective in reducing diabetes risk, but the benefits of walking (3,5), resistance training (16,22,23), and lifestyle physical activity (24) have become increasingly evident. Based on trial and epidemiological evidence, the Centers for Disease Control and Prevention (CDC) and the American College of Sports Medicine (ACSM) have jointly published recommendations that adults should engage in at least 30 min of moderate physical activity (e.g., brisk walking) on most, and preferably all, days of the week (25).

Recently, both the Finnish Diabetes Prevention Study (DPS) (26) and the Diabetes Prevention Program (27) found that multiple relatively modest lifestyle changes, including weight loss, dietary changes more in line with current recommendations, and increased LTPA reduce the risk of type 2 diabetes by 58% in individuals with impaired glucose tolerance (IGT). Post hoc analyses of the independent role of increased physical activity in these two randomized controlled trials have not been published, however. In the Chinese Da Qing Impaired Glucose Tolerance and Diabetes Study (28), which was randomized by clinic rather than by participant, diabetes risk was reduced

From the ¹Department of Medicine, Kuopio University Hospital, Kuopio, Finland; the ²Department of Physiology, University of Kuopio, Kuopio, Finland; the ³Diabetes and Genetic Epidemiology Unit, Department of Epidemiology and Health Promotion, Helsinki, Finland; the ⁴Pennington Biomedical Research Center, Louisiana State University, Baton Rouge, Louisiana; the ⁵Kuopio Research Institute of Exercise Medicine, Kuopio, Finland; the ⁶Department of Health and Functional Capacity, National Public Health Institute, Helsinki, Finland; the ⁷Research Department, Social Insurance Institution, Turku, Finland; the ⁸Department of Public Health Science and General Practice, Unit of General Practice, Oulu University Hospital, Oulu, Finland; the ⁹Department of Sports Medicine, Oulu Deaconess Institute, Oulu, Finland; the ¹⁰Finnish Diabetes Association, Tampere, Finland; the ¹¹Tampere University Hospital Research Unit, Tampere, Finland; the ¹²Department of Clinical Nutrition and Food and Health Research Center, University of Kuopio, Kuopio, Finland; the ¹³Institute of Nursing and Health Care, Tampere, Finland; the ¹⁴Laboratory of Analytical Biochemistry, National Public Health Institute, Helsinki, Finland; and the ¹⁵Department of Public Health, University of Helsinki, Helsinki, Finland.

Address correspondence and reprint requests to Prof. Matti Uusitupa, MD, PhD, Department of Clinical Nutrition and Food and Health Research Center, University of Kuopio, P.O. Box 1627, Kuopio, Finland. E-mail: matti.uusitupa@uku.fi.

Received for publication 22 June 2004 and accepted in revised form 15 September 2004.

ACSM, American College of Sports Medicine; CDC, Centers for Disease Control and Prevention; DPS, Diabetes Prevention Study; IGT, impaired glucose tolerance; KIHJ study, Kuopio Ischemic Heart Disease Risk Factor study; LTPA, leisure-time physical activity; OGTT, oral glucose tolerance test.

© 2005 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

46% in the exercise group, 42% in the diet and exercise group, and 31% in the diet-treated group. Physical activity levels were already greater at baseline in the exercise only group, however, which may partly explain why the greatest risk reduction seemed to be in this group. Furthermore, low-intensity and moderate-to-vigorous LTPA or structured and lifestyle LTPA were not separately reported. Therefore, more evidence on the role of LTPA in the prevention of type 2 diabetes is urgently needed.

We report the results of post hoc analyses on the role of LTPA in the prevention of type 2 diabetes in the Finnish DPS. We extended the follow-up period of the original trial (26) from 3.2 to 4.1 years. We first studied the effect of the intervention on different forms of LTPA by comparing the changes in LTPA between the control and intervention groups. We then assessed the role of LTPA in preventing diabetes by examining the association of changes in LTPA during the study with the incidence of diabetes in the combined intervention and control groups. Our final goal was to examine the role of compliance with current LTPA recommendations, and more specifically the role of walking for exercise, in the prevention of diabetes.

RESEARCH DESIGN AND METHODS

The Finnish DPS is a multicenter, randomized, controlled trial designed to test the hypothesis that lifestyle changes can prevent type 2 diabetes in high-risk individuals. The design of the DPS has been described in detail elsewhere (29). Briefly, overweight or obese individuals ages 40–65 years with IGT were eligible for the study. IGT was defined as a plasma glucose concentration of 7.8–11.0 mmol/l 2 h after a 75-g oral glucose challenge in subjects whose fasting glucose concentration was <7.8 mmol/l (30). The study protocol was approved by the ethics committee of the National Public Health Institute in Helsinki, Finland, and all participants gave written informed consent.

In all, 522 individuals from five study centers were randomly assigned to the intervention ($n = 265$) or control ($n = 257$) group (26). Of these 522 subjects, 487 completed a questionnaire at baseline and at least once during follow-up quantifying the previous 12 months of LTPA and were included in the present analyses. The original trial ended after an average follow-up of 3.2 years, during which time 86 cases of incident diabetes were diagnosed (26). In this study, we extended the follow-up period to 4.1 years (range 1–6 years), during which time the original random allocation and intensive intervention was continued. In all, 107 of the 487 participants developed diabetes during the 4.1-year follow-up.

The intervention group was given detailed advice on how to achieve the goals of the intervention, which were moderate-to-vigorous exercise for at least 30 min per day; a reduction in body weight to $\geq 5\%$, total intake of fat to <30% of energy consumed, and intake of saturated fat to <10% of energy consumed; and an increase in fiber intake to at least 15 g/1,000 kcal. Endurance exercise (e.g., swimming, bicycling, jogging, aerobic ball games, skiing) was encouraged. No-cost, supervised circuit-type resistance training was offered to the intervention group 4–6 months after the randomization. Walking and lifestyle physical activity were also recommended. The control group was given general verbal and written information about exercise and diet at baseline and at subsequent annual visits, but no specific individualized programs were offered.

Assessment of leisure-time physical activity. LTPA questionnaires were filled out at baseline and during yearly follow-up visits. The percentage of those remaining in the study who completed the forms ranged from 89 to 98% throughout the study. The validated Kuopio Ischemic Heart Disease Risk Factor Study (KIHD) 12-month LTPA questionnaire was used, as previously described (31,32). The questionnaire provides detailed quantitative information on the duration, frequency, and mean intensity of the most common lifestyle and structured LTPA as recalled over the previous 12 months (33). In a sample of these overweight men and women with IGT, maximal oxygen uptake during a bicycle exercise test was $\sim 30\%$ lower than in the younger and leaner men who participated in the population-based KIHD cohort study for whom the questionnaire was designed (data not shown). We therefore decreased the METs (1 MET is defined as metabolic expenditure at rest, corresponding to an oxygen uptake of 3.5 ml O_2 /kg) assigned to different subjective intensities for given forms of physical activity by 30%. Moderate-to-vigorous LTPA was defined as ≥ 3.5 METs. LTPA was quantified in hours

per week. Men and women were assigned the same MET value for a given subjective intensity of a given physical activity.

Common moderate-to-vigorous LTPA included walking at a moderate or very strenuous intensity, bicycling, swimming, resistance training, skiing, jogging, ball games, and lifestyle activities such as chopping wood or clearing brush. Common low-intensity activities included casual or mildly exertive walking, bicycling at a leisurely intensity, gardening and yard work, and picking berries and mushrooms.

Structured LTPAs other than walking (e.g., bicycling, skiing, jogging, swimming, resistance training, ball sports, rowing) were classified as strenuous if practiced at a moderate or high subjective intensity (e.g., causing moderate or heavy sweating or breathlessness) and nonstrenuous if practiced at lower intensities. Walking for exercise was categorized separately. Lifestyle LTPA included gardening, yard work, shoveling snow, picking berries and mushrooms, hobbies and crafts, hunting, fishing, chopping wood, or clearing brush. Walking or cycling to and from work was categorized separately as commuting LTPA.

Clinical, anthropometric, and dietary assessments. At baseline and each annual visit, all study subjects underwent an oral glucose tolerance test (OGTT) and completed a 3-day food diary, as previously described (26,29,34). Changes in dietary factors and body weight have been reported elsewhere (26,34).

Biochemical assessments. Plasma glucose was measured using a glucose dehydrogenase method after precipitation of proteins by trichloroacetic acid. The result of the second OGTT was considered the baseline value. The serum insulin concentration was measured by a radioimmunoassay (Pharmacia, Uppsala, Sweden). Serum total and HDL cholesterol and triglycerides were measured by an enzymatic assay at the National Public Health Institute, Analytical Biochemistry Laboratory.

Determination of diabetes. Diabetes was defined according to the 1985 criteria of the World Health Organization (30) as either a fasting plasma glucose concentration ≥ 7.8 mmol/l or a plasma glucose concentration ≥ 11.1 mmol/l 2 h after a 75-g oral glucose challenge. Participants were asked to fast and refrain from strenuous exercise for 12 h before the OGTT. If the diagnosis of diabetes was not confirmed by a second OGTT, the subject continued in the study (26).

Statistical analysis. Two-sided t tests and χ^2 tests were used to analyze the differences between the groups at baseline and during follow-up. Intake of dietary total and saturated fat and fiber (in grams per day) was adjusted by daily energy intake with linear regression analysis before further statistical analysis (35). The change in LTPA during the trial was calculated by subtracting the baseline total LTPA and its components (in hours per week) from corresponding measures of averaged LTPA questionnaires completed during follow-up. Changes in dietary and biochemical measurements and body weight during the trial were similarly calculated. The change in LTPA for the combined intervention and control groups was categorized into thirds to assess the association of the change in LTPA with the risk of type 2 diabetes using Cox proportional hazards models. Moderate-to-vigorous LTPA was also categorized into inactive (<1 h/week) and active (≥ 2.5 h/week) groups to assess adherence to LTPA goals. Because walking for exercise is the most common form of LTPA and is emphasized in current ACSM/CDC recommendations, we also examined the association of time spent walking for exercise during follow-up with incident diabetes. The covariates for the Cox proportional hazards models were forced into the model. In analyses with fasting and 2-h insulin as covariates, the group mean was substituted for the 44 and 47 individuals with missing values in the intervention and control groups, respectively. Statistical significance was defined as $P < 0.05$. Statistical analyses were performed with SPSS 11.0 for Windows (SPSS, Chicago, IL).

RESULTS

Baseline characteristics between the intervention and control groups were similar (Table 1). Reported participation in <1 h/week of moderate-to-vigorous LTPA was 37% (91 of 249) in the intervention group and 39% (94 of 238) in the control group. Engagement in ≥ 2.5 h/week of moderate-to-vigorous LTPA was 41% in the intervention group and 40% in the control group.

Changes in LTPA during follow-up for the intervention versus control group. During the 4.1-year follow-up, moderate-to-vigorous intensity LTPA increased 0.8 h/week in the intervention group, whereas little change was reported in the control group (Table 2). Engagement in <1 h/week of moderate-to-vigorous LTPA during follow-up

TABLE 1
Baseline characteristics of participants in the intervention and control groups

	Intervention	Control
<i>n</i>	249	238
Age (years)	55.8 ± 7.2	55.0 ± 6.9
Sex (male/female)	87/162	75/163
Smokers (%)	8	11
Family history of diabetes (%)	67	61
Systolic blood pressure (mmHg)	140 ± 18	136 ± 18
Diastolic blood pressure (mmHg)	86 ± 9	86 ± 10
Antihypertensive drug treatment (%)	28	31
BMI (kg/m ²)	31.3 ± 4.5	31.2 ± 4.6
Serum total cholesterol (mmol/l)	5.6 ± 1.0	5.6 ± 0.9
Serum HDL cholesterol (mmol/l)	1.2 ± 0.3	1.2 ± 0.3
Serum triglycerides (mmol/l)	1.53 (1.15–2.00)	1.60 (1.20–2.10)
Cholesterol-lowering medication (%)	5	7
OGTT		
Fasting plasma glucose (mmol/l)	6.1 ± 0.8	6.2 ± 0.7
2-h plasma glucose (mmol/l)	8.7 (7.9–9.8)	8.8 (7.9–10.0)
Fasting serum insulin (pmol/l)	90 (69–125)	97 (69–125)
2-h serum insulin (pmol/l)	556 (340–837)	542 (403–813)
Dietary intake		
Energy (kcal)	1,695 (1,431–2,023)	1,683 (1,363–2,054)
Total fat, energy adjusted (g)	72 (63–79)	73 (66–81)
Saturated fat, energy adjusted (g)	32.0 (27.0–36.9)	33.6 (28.9–37.9)
Fiber, energy adjusted (g)	19.4 (15.7–23.8)	19.0 (15.2–23.4)
LTPA (h/week)†		
Total	5.8 (3.3–9.2)	5.5 (2.8–9.8)
Moderate and vigorous	2.6 (1.1–4.9)	2.8 (1.1–5.8)
Low intensity	2.2 (0.7–4.5)	1.9 (0.5–4.7)
Strenuous, structured	0.1 (0.0–0.8)	0.2 (0.0–0.8)
Nonstrenuous, structured	0.3 (0.0–1.1)	0.2 (0.0–0.7)
Walking for exercise	1.1 (0.3–2.7)	0.9 (0.2–2.5)
Casual or mildly exertive (%)	71	76
Moderately or very strenuous (%)	29	24
Lifestyle	1.9 (0.4–4.0)	1.8 (0.4–4.9)
Commuting on foot or by bicycle	0.0 (0.0–0.5)	0.0 (0.0–0.5)

Data are means ± SD, medians (interquartile ranges), or percent and are given for participants who provided sufficient information on LTPA. Total LTPA is the sum of moderate and vigorous plus low-intensity LTPA or the sum of strenuous, structured LTPA other than walking; nonstrenuous, structured LTPA other than walking; walking for exercise; lifestyle LTPA; and commuting LTPA. Moderate and vigorous LTPA includes walking at a moderate or very strenuous intensity, bicycling, swimming, resistance training, skiing, jogging, ball games, and lifestyle activities such as chopping wood or clearing brush. Low-intensity LTPA includes casual or mildly exertive walking, bicycling at a leisurely intensity, gardening and yard work, and picking berries and mushrooms. Strenuous, structured LTPA other than walking includes bicycling, swimming, skiing, jogging, resistance training, ball games, and rowing practiced at a moderate or high subjective intensity; nonstrenuous, structured LTPAs are those same activities practiced at lower intensities. Walking for exercise was categorized separately. Lifestyle LTPA includes gardening, yard work, shoveling snow, picking berries, hobbies and crafts, hunting, fishing, and chopping wood. Commuting LTPA was defined as walking or cycling to and from work.

was 15 vs. 25% ($P = 0.004$) in the intervention and control groups, respectively. Reported participation in ≥ 2.5 h/week of moderate-to-vigorous LTPA during follow-up in the intervention compared with the control group was 62 vs. 46% ($P < 0.001$). The clearest difference in the change in LTPA between groups, other than walking, was for strenuous, structured LTPA (Table 2).

Changes in LTPA and risk of diabetes in the combined intervention and control groups. During the follow-up period, 107 individuals developed diabetes (41 of 249 in the intervention group and 66 of 238 in the control group). In the combined groups, the change in total LTPA was more strongly associated with incident diabetes than changes in subcategories of LTPA (Table 3). Adjusting for age, sex, group, smoking status, major risk factors for diabetes at baseline (BMI, fasting and 2-h plasma glucose and insulin levels, and family history of diabetes), and baseline total LTPA, participants in the

upper third of the change in total LTPA were 80% less likely to develop diabetes than those in the lower third (relative risk [RR] 0.20, 95% confidence interval [CI] 0.10–0.41; $P < 0.001$ for linear trend).

When comparing the change in total LTPA from baseline to the first year of follow-up or from baseline to the last year of follow-up, the association of change in LTPA with diabetes was similar, but somewhat weaker, than when using averaged values of LTPA during the follow-up. Those who increased LTPA from the first to the last year of follow-up seemed to have a lower risk of diabetes than those who decreased LTPA after adjusting for baseline levels of LTPA and the change in LTPA from baseline to the first year of follow-up, but the association was not significant.

Participants reporting a change in moderate-to-vigorous LTPA in the upper third were 49–65% less likely to develop diabetes than those in the lower third in multivariate

TABLE 2
Changes in LTPA during the trial period

	Intervention	Control	<i>P</i>
<i>n</i>	249	238	
Total	0.4 ± 5.2 (−0.3 to 1.1)	−0.1 ± 5.3 (−0.7 to 0.6)	0.36
Moderate and vigorous	0.8 ± 2.7 (0.4–1.1)	0.2 ± 2.9 (−0.2 to 0.6)	0.028
Low intensity	−0.4 ± 4.6 (−1.0 to 0.2)	−0.3 ± 4.5 (−0.9 to 0.4)	0.79
Strenuous structured (other than walking)	0.6 ± 1.4 (0.5–0.8)	−0.1 ± 1.3 (−0.2 to 0.1)	<0.001
Nonstrenuous structured	−0.1 ± 1.1 (−0.3 to 0.0)	0.0 ± 0.9 (−0.1 to 0.2)	0.063
Walking for exercise	0.2 ± 2.1 (−0.1 to 0.5)	0.3 ± 2.1 (0.0–0.5)	0.91
Increase in walking intensity	121 (49)	109 (46)	0.54
Lifestyle	−0.3 ± 4.4 (−0.8 to 0.2)	−0.3 ± 4.7 (−0.9 to 0.3)	0.75
Commuting on foot or by bicycle	0.0 ± 1.0 (−0.1 to 0.1)	0.0 ± 0.9 (−0.2 to 0.1)	0.80

Data are means ± SD (95% CI) or *n* (%). LTPA data represent hours per week. *P* values were determined using Student's unpaired *t* test, except for change in walking intensity, which was determined with the χ^2 test.

models including low-intensity LTPA and its changes (Table 3). Changes in low-intensity LTPA similarly predicted a reduction in the risk of incident diabetes of 59–64%.

Increased strenuous, structured LTPA other than walking was also associated with a decreased risk of diabetes (Fig. 1A). After multivariate adjustment, including baseline BMI and its changes, the association was no longer significant. In separate analyses of the individual forms of LTPA, home exercise, resistance training, and dancing (a single category in the questionnaire) made up the greatest portion of change in strenuous LTPA. The change in this category differed between groups (median change 0.3 vs. 0.0 h/week in the intervention and control groups; *P* < 0.001). In the combined groups, the change was also inversely associated with incident diabetes (median 0.7 vs. 0.0 h/week for the upper and lower thirds; RR 0.54, 95% CI 0.32–0.91) after adjusting for age, sex, group, and respective baseline levels. Bicycling, skiing, and swimming made up smaller portions of the change in strenuous LTPA, but were also individually associated with decreased risk for diabetes (RR 0.53–0.62 for those who increased these activities versus all others). Changes in nonstrenuous, structured LTPA other than walking were not associated with diabetes risk.

An increase in walking for exercise during follow-up also decreased the risk of diabetes (Fig. 1B). Furthermore, those who increased the intensity of walking for exercise during follow-up (*n* = 230, 47%) had a 48% lower risk of diabetes than all others when adjusting for age, sex, group, baseline walking intensity, and the duration of walking for exercise and its changes (RR 0.52, 95% CI 0.33–0.83). Further adjustment for baseline dietary factors and their changes (0.62, 0.38–1.00) and BMI and its changes (0.66, 0.40–1.09) attenuated the association.

Increased lifestyle LTPA also seemed to decrease the risk of incident diabetes (Fig. 1C). Changes in commuting LTPA were not associated with incident diabetes, but only 33% of men and women engaged in commuting LTPA, and changes in commuting LTPA were small (Table 2). As an alternative approach, we categorized average engagement in commuting LTPA during follow-up into none (*n* = 253), <30 min/week (*n* = 100), and ≥30 min/week (median 1.3 h/week; *n* = 134). In models adjusting for age, sex, change in commuting LTPA from baseline, and average duration of other forms of LTPA during follow-up and their changes from baseline, at least 30 min of commuting LTPA during follow-up decreased the risk of diabetes (model 1: RR 0.54, 95% CI 0.33–0.89; model 2: 0.58, 0.34–0.97), but additional adjustment for average BMI during follow-up and its

TABLE 3
Relative risk of developing type 2 diabetes during the trial period according to tertiles of change in LTPA

	Model 1	Model 2	Model 3
Change in total LTPA			
−3.2 (−35 to −0.5)	1	1	1
0.5 (−0.5 to 1.7)	0.47 (0.28–0.79)	0.48 (0.28–0.82)	0.52 (0.31–0.89)
3.8 (1.8–19)	0.26 (0.15–0.47)	0.29 (0.16–0.53)	0.34 (0.19–0.62)
<i>P</i> for trend	<0.001	<0.001	<0.001
Change in moderate-to-vigorous LTPA (≥3.5 METs)			
−1.5 (−13.5 to −0.1)	1	1	1
0.5 (−0.1 to 1.3)	0.78 (0.46–1.33)	0.86 (0.49–1.48)	0.95 (0.54–1.65)
2.6 (1.3–14.4)	0.35 (0.18–0.65)	0.40 (0.21–0.76)	0.51 (0.26–0.97)
<i>P</i> for trend	0.001	0.004	0.037
Change in low-intensity LTPA (<3.5 METs)			
−3.2 (−34 to −1.0)	1	1	1
0.1 (−0.9 to 1.1)	0.83 (0.47–1.45)	0.85 (0.47–1.53)	0.63 (0.34–1.17)
3.1 (1.1–15.0)	0.38 (0.20–0.70)	0.41 (0.22–0.77)	0.36 (0.19–0.67)
<i>P</i> for trend	0.001	0.003	0.001

Data are relative risk (95% CI). LTPA data represent hours per week. Model 2: Adjusted variables in model 1 and baseline values and changes in dietary intake of energy, total fat, saturated fat, and fiber. Model 3: Adjusted for variables in model 2 and baseline values and changes in BMI.




FIG. 1. Association of risk of diabetes with changes in strenuous, structured LTPA other than walking (A), walking for exercise (B), and lifestyle LTPA (C) according to tertiles in the combined intervention and control groups. Medians and ranges for tertiles of changes in strenuous, structured LTPA other than walking: -0.2 h/week, -14.6 to 0.0 (reference); 0.2 h/week, 0.0 – 0.5 ; 1.1 h/week, 0.5 – 8.3 . Medians and ranges for tertiles of changes in walking for exercise: -1.1 h/week, -12.4 to -0.1 (reference); 0.2 h/week, -0.1 to 0.6 ; 1.5 h/week, 0.6 – 11.5 . Medians and ranges for tertiles of changes in lifestyle physical activity: -1.8 h/week, -37 to -0.3 (reference); 0.0 h/week, -0.3 to 0.5 ; 1.9 h/week, 0.6 – 17 . *P* values are for linear trends. Model 1: Adjusted for age, sex, group, the respective form of LTPA at baseline, and the other forms of LTPA and their changes (e.g., for strenuous structured LTPA other than walking, also adjustment for nonstrenuous structured LTPA other than walking, lifestyle LTPA, LTPA while commuting, and walking for exercise). Model 2: Adjusted variables in model 1 and baseline values and changes in dietary intake of energy, total and saturated fat, and fiber. Model 3: Adjusted for variables in model 2 and baseline values and changes in BMI.

change from baseline weakened the association (model 3: 0.72 , 0.43 – 1.22).

Adherence to leisure-time physical activity recommendations and type 2 diabetes. Men and women whose average levels of moderate-to-vigorous LTPA dur-




FIG. 2. Likelihood of developing diabetes during follow-up according to average duration of moderate and vigorous physical activity during follow-up (<1 , 1 – 2.4 , and ≥ 2.5 h/week) (A), and average duration of walking for exercise (<1 , 1 – 2.4 , and ≥ 2.5 h/week) (B). Model 1: Adjusted for age, sex, and 1) change in moderate-to-vigorous LTPA from baseline and average low-intensity LTPA during follow-up and its change from baseline, or 2) change in duration of walking for exercise from baseline and other forms of LTPA as in Fig. 1 and their changes from baseline. Model 2: Adjusted variables in model 1 and average values of dietary intake of energy, total and saturated fat, and fiber during follow-up and their changes from baseline. Model 3: Adjusted for variables in model 2 and average BMI during follow-up and its change from baseline.

ing follow-up were in compliance with current ACSM/CDC recommendations (≥ 2.5 h/week; $n = 265/487$) were 44% less likely to develop diabetes than those remaining sedentary (<1 h/week; $n = 97/487$) in models adjusting for age, sex, group, change in moderate-to-vigorous LTPA from baseline, and average low-intensity LTPA during follow-up and its change from baseline (Fig. 2A). Further adjustment for dietary variables and BMI during follow-up and changes from baseline attenuated the associations, however. Analysis of adherence to the stated goals of the trial (≥ 3.5 h/week) yielded similar results. Similarly analyzed, individuals engaging in ≥ 2.5 h/week of walking for exercise during follow-up ($n = 157/487$) were 63–69% less likely to develop diabetes than those who walked <1 h/week ($n = 192/487$) (Fig. 2B).

Influence of sex and group. At baseline, men engaged in more total LTPA (median 7.3 vs. 5.3 h/week) and moderate-to-vigorous (median 2.3 vs. 1.4 h/week) LTPA than women, but amounts of low-intensity LTPA did not differ

significantly between men and women (median 3.2 vs. 2.9 h/week). For both sexes, walking was the most common form of any LTPA, bicycling the most common form of structured LTPA other than walking, and gardening, yard work, and snow shoveling the most common lifestyle activity. The changes in the LTPA categories shown in Table 2 did not differ between sexes.

Although the trial is not powered sufficiently to study complex interactions among LTPA, group, and sex, the association of LTPA with incident diabetes was present in women in the intervention group but not in the control group, whereas the association was overall present for men in both the intervention and control groups (data not shown).

DISCUSSION

Moderate-to-vigorous and strenuous structured LTPA increased in the intervention group by ~36–48 min/week during the 4-year follow-up period. Consistently, adherence to the CDC/ACSM physical activity recommendations (≥ 2.5 h/week) during follow-up was 62% in the intervention group but only 46% in the control group. In the combined groups, an increase in moderate-to-vigorous and strenuous structured LTPA was associated with a large reduction in the risk of diabetes. Moreover, an increase in the duration and intensity of walking for exercise and even lifestyle LTPA predicted a decrease in incident diabetes.

We used the validated KIHD 12-month LTPA questionnaire (31) in this study. The questionnaire provides detailed information on the frequency, duration, and intensity of common structured and lifestyle physical activities as recalled for the preceding 12 months. The questionnaire is quite repeatable (intraclass correlation coefficient 0.58) (31). Measurement error was nonetheless large. To reduce measurement variability and better reflect the actual levels of LTPA throughout the study, we administered the questionnaire yearly and used the average of LTPA levels during follow-up. As in Finland in general (36), reported LTPA levels were quite high. Participants may nonetheless have overestimated the amount of habitual LTPA. The questionnaire was originally designed for a cohort study in men, but contains the same forms of LTPA as the validated questionnaire used in the Nurses Health Study, except for climbing stairs (37). Despite these limitations, the questionnaire seems suitable for use in lifestyle intervention trials.

The increase in strenuous, structured LTPA other than walking best represented the effect of the exercise intervention. In the combined groups, an increase in structured, strenuous LTPA seemed to decrease the risk of diabetes by 57% for the upper versus lower third after adjusting for dietary changes, but further adjustment for changes in BMI attenuated the decrease to 47%. Strenuous, structured LTPA thus likely mediated the decrease in diabetes risk in part through weight loss and weight maintenance. Weight loss is a powerful determinant of improved insulin sensitivity, as has been shown in a subgroup of the DPS (38), and plays a critical role in diabetes prevention. Our findings nonetheless indicate that LTPA may influence glucose and insulin metabolism through mechanisms other than weight loss or changes in body composition (20,39).

Both endurance LTPA and resistance training practiced at an at least moderately strenuous intensity appeared to protect against progression from IGT to diabetes. In the questionnaire, home exercise, resistance training, and dancing were part of the same category. Because this category increased during the study only in the intervention group, however, resistance training is likely to make up most of the increase. In type 2 diabetic women, adding resistance training to aerobic training enhances insulin sensitivity, decreases abdominal subcutaneous and visceral adipose tissue, and increases muscle density more than a similar quantity of aerobic training alone (23). Resistance training also improves insulin sensitivity in IGT (16). Changes in casual, structured LTPA were not associated with incident diabetes, even though low-intensity and lifestyle LTPA seemed to be protective. This may be because more time is spent in most low-intensity and lifestyle activities than in structured LTPA.

The stated goal of the LTPA intervention, moderate-to-vigorous LTPA, also increased more in the intervention group. In the combined groups, an increase in moderate-to-vigorous LTPA in the upper third was associated with a 49% lower risk of incident diabetes even after further adjustment for diet, BMI, and their changes. The risk of diabetes was consistently halved in those who complied with the CDC/ACSM recommendations during follow-up compared with risk in sedentary persons, but not after adjustment for dietary changes and weight loss.

Participants whose increase in walking for exercise was in the upper third were 59% less likely to develop diabetes than those whose change was in the lower third, independent of other factors. Moreover, a subjective increase in the pace of walking also decreased the likelihood of diabetes independently of time spent walking. A key message from a public health standpoint would be that at least 2.5 h/week of walking for exercise during follow-up seemed to decrease the risk of diabetes by 63–69%, largely independent of dietary factors and BMI. Walking for exercise is unlikely to have contributed to the overall effect of the intervention, because the median duration of walking did not change in either group. In the Nurses Health Study (3) and the Health Professionals Study (5), both the amount of walking for exercise and the pace were strongly associated with a decreased risk of diabetes, with some attenuation when adjusted for BMI. Changes in walking or bicycling to work were not associated with incident diabetes, but engagement in >30 min/week of commuting LTPA during follow-up also appeared to protect against diabetes, consistent with the findings of another Finnish study (6).

In the Da Qing Impaired Glucose Tolerance and Diabetes Study (28), incident diabetes decreased by 46% in the exercise group, 42% in the diet and exercise group, and 31% in the diet-treated group. LTPA was assessed by recall of the previous week's activities and quantified according to arbitrary units assigned to three intensities of LTPA, which also took into account duration of activity. According to this crude assessment, LTPA was already greater at baseline in the exercise group. The amount of LTPA increased during the study in within-group assessments, but not compared with the control group. Moderate-to-vigorous and low-intensity LTPA was not separately re-

ported. In the nonrandomized Malmö Feasibility Study in 260 middle-aged men with IGT, the incidence of diabetes was 50% lower in the intervention group after 5 years (40). Weight loss and increased maximal oxygen uptake independently predicted improved glucose tolerance, suggesting that increased LTPA contributed to the decreased risk of diabetes.

One interesting finding was that changes in lifestyle and low-intensity LTPA also seemed to decrease risk of diabetes independently of moderate-to-vigorous or structured LTPA, dietary factors, and weight loss. Because the amount of lifestyle or low-intensity LTPA did not change in either group, changes in low-intensity or lifestyle LTPA did not explain the decrease in incident diabetes brought about by the intervention. We (2,7,21,32) and others (rev. in 12) have generally found that more vigorous LTPA has greater benefits on metabolic and chronic disease end points. Low-intensity LTPA may nonetheless confer benefits that are independent of or complementary to those of moderate-to-vigorous LTPA. The total time or energy spent on LTPA may therefore have been more important than the intensity in these middle-aged, obese, generally unfit individuals at high risk for diabetes. Indeed, the increase in total LTPA was associated with a greater reduction in diabetes risk than increases in any of its subcategories.

The strengths of this study include its randomized controlled design (26), detailed assessment of LTPA using a validated questionnaire, and repeated measurement of LTPA. The trial was designed to assess the effect of multiple interventions on the prevention of diabetes and did not include a separate exercise-only group. Therefore, we cannot rule out residual confounding with the other interventions. Detailed assessment of adherence to the dietary and weight loss interventions nonetheless allowed statistical adjustment for the other lifestyle interventions. Another limitation of our study was the lack of serial measurements of maximal oxygen uptake or other objective measures of LTPA. The MET values assigned to different forms of LTPA and their subjective intensities are somewhat arbitrary, but the assignment of higher MET values or cutoff points $>$ or <3.5 METs has little effect on the results.

A program promoting lifestyle changes that also included structured aerobic and resistance LTPA increased participation in moderate-to-vigorous LTPA and especially in strenuous, structured LTPA. Increased moderate-to-vigorous and strenuous, structured LTPA strongly decreased the risk of type 2 diabetes, as did walking for exercise, low-intensity LTPA, and lifestyle LTPA. Our findings extend those from prospective cohort studies indicating that LTPA decreases the risk of diabetes through mechanisms beyond weight loss alone. Compliance with current physical activity recommendations may substantially reduce the incidence of type 2 diabetes and should be widely encouraged, especially in high-risk individuals. Low-intensity physical activity may also be beneficial.

ACKNOWLEDGMENTS

This study was supported by the Academy of Finland (Grants 8473/2298, 40758/5767, 38387/54175, 46558, and

40758), the Ministry of Education, the Novo Nordisk Foundation, the Yrjö Jahnsson Foundation, the Finnish Diabetes Research Foundation, the EVO fund of the Kuopio University Hospital (Grant 5106 to M.U.), and the Juho Vainio Foundation. The funding organizations did not participate in the design or conduct of the study; in the collection, analysis, or interpretation of the data; or in the preparation, review, or approval of the manuscript.

We are indebted to Kirsi Frantsi, Olli Heinonen, Katri Hemio, Pirjo Härkönen, Pia Högström, Anja Ilmanen, Kaija Kettunen, Paivi Kleemola, Anna Korhonen, Arja Kylliäinen, Marjukka Lauhkonen, Pirjo Lehto, Liisa Mikkola, Paula Nyholm, and Arja Putila for their skillful assistance in performing the study; to Profs. Marja-Riitta Taskinen and Antti Aro for their participation on the end points committee; and to Dr. William C. Knowler for his important contribution to the planning of our study.

REFERENCES

- Helmrich SP, Ragland DR, Leung RW, Paffenbarger RS Jr: Physical activity and reduced occurrence of non-insulin-dependent diabetes mellitus. *N Engl J Med* 325:147–152, 1991
- Lynch J, Helmrich SP, Lakka TA, Kaplan GA, Cohen RD, Salonen R, Salonen JT: Moderately intense physical activities and high levels of cardiorespiratory fitness reduce the risk of non-insulin-dependent diabetes mellitus in middle-aged men. *Arch Intern Med* 156:1307–1314, 1996
- Hu FB, Sigal RJ, Rich-Edwards JW, Colditz GA, Solomon CG, Willett WC, Speizer FE, Manson JE: Walking compared with vigorous physical activity and risk of type 2 diabetes in women: a prospective study. *JAMA* 282:1433–1439, 1999
- Wei M, Gibbons LW, Mitchell TL, Kampert JB, Lee CD, Blair SN: The association between cardiorespiratory fitness and impaired fasting glucose and type 2 diabetes mellitus in men. *Ann Intern Med* 130:89–96, 1999
- Hu FB, Leitzmann MF, Stampfer MJ, Colditz GA, Willett WC, Rimm EB: Physical activity and television watching in relation to risk for type 2 diabetes mellitus in men. *Arch Intern Med* 161:1542–1548, 2001
- Hu G, Qiao Q, Silventoinen K, Eriksson JG, Jousilahti P, Lindstrom J, Valle TT, Nissinen A, Tuomilehto J: Occupational, commuting, and leisure-time physical activity in relation to risk for type 2 diabetes in middle-aged Finnish men and women. *Diabetologia* 46:322–329, 2003
- Laaksonen DE, Lakka HM, Salonen JT, Niskanen LK, Rauramaa R, Lakka TA: Low levels of leisure-time physical activity and cardiorespiratory fitness predict development of the metabolic syndrome. *Diabetes Care* 25:1612–1618, 2002
- Laaksonen DE, Lakka HM, Niskanen LK, Kaplan GA, Salonen JT, Lakka TA: Metabolic syndrome and development of diabetes mellitus: application and validation of recently suggested definitions of the metabolic syndrome in a prospective cohort study. *Am J Epidemiol* 156:1070–1077, 2002
- Lakka HM, Laaksonen DE, Lakka TA, Niskanen LK, Kumpusalo E, Tuomilehto J, Salonen JT: The metabolic syndrome and total and cardiovascular disease mortality in middle-aged men. *JAMA* 288:2709–2716, 2002
- Rice B, Janssen I, Hudson R, Ross R: Effects of aerobic or resistance exercise and/or diet on glucose tolerance and plasma insulin levels in obese men. *Diabetes Care* 22:684–691, 1999
- Ross R, Dagnone D, Jones PJ, Smith H, Paddags A, Hudson R, Janssen I: Reduction in obesity and related comorbid conditions after diet-induced weight loss or exercise-induced weight loss in men: a randomized, controlled trial. *Ann Intern Med* 133:92–103, 2000
- Department of Health and Human Services: *Physical Activity and Health: A Report of the Surgeon General*. Atlanta, GA, Dept. of Health and Human Services, 1996
- Irwin ML, Yasui Y, Ulrich CM, Bowen D, Rudolph RE, Schwartz RS, Yukawa M, Aiello E, Potter JD, McTiernan A: Effect of exercise on total and intra-abdominal body fat in postmenopausal women: a randomized controlled trial. *JAMA* 289:323–330, 2003
- Hellenius ML, Brismar KE, Berglund BH, de Faire UH: Effects on glucose tolerance, insulin secretion, insulin-like growth factor I and its binding protein, IGFBP-1, in a randomized controlled diet and exercise study in healthy, middle-aged men. *J Intern Med* 238:121–130, 1995
- Dengel DR, Galecki AT, Hagberg JM, Pratley RE: The independent and combined effects of weight loss and aerobic exercise on blood pressure

- and oral glucose tolerance in older men. *Am J Hypertens* 11:1405–1412, 1998
16. Eriksson J, Tuominen J, Valle T, Sundberg S, Sovijarvi A, Lindholm H, Tuomilehto J, Koivisto V: Aerobic endurance exercise or circuit-type resistance training for individuals with impaired glucose tolerance? *Horm Metab Res* 30:37–41, 1998
 17. Kelley DE, Goodpaster BH: Effects of physical activity on insulin action and glucose tolerance in obesity. *Med Sci Sports Exerc* 31:S619–S623, 1999
 18. Kraus WE, Houmard JA, Duscha BD, Knetzger KJ, Wharton MB, McCartney JS, Bales CW, Henes S, Samsa GP, Otvos JD, Kulkarni KR, Slentz CA: Effects of the amount and intensity of exercise on plasma lipoproteins. *N Engl J Med* 347:1483–1492, 2002
 19. Whelton SP, Chin A, Xin X, He J: Effect of aerobic exercise on blood pressure: a meta-analysis of randomized, controlled trials. *Ann Intern Med* 136:493–503, 2002
 20. Ross R: Does exercise without weight loss improve insulin sensitivity? *Diabetes Care* 26:944–945, 2003
 21. Lakka TA, Laaksonen DE, Lakka H-M, Mannikko N, Niskanen LK, Rauramaa R, Salonen JT: Sedentary lifestyle, poor cardiorespiratory fitness and the metabolic syndrome. *Med Sci Sports Exerc* 35:1279–1286, 2003
 22. Castaneda C, Layne JE, Munoz-Orians L, Gordon PL, Walsmith J, Foldvari M, Roubenoff R, Tucker KL, Nelson ME: A randomized controlled trial of resistance exercise training to improve glycemic control in older adults with type 2 diabetes. *Diabetes Care* 25:2335–2341, 2002
 23. Cuff DJ, Meneilly GS, Martin A, Ignaszewski A, Tildesley HD, Frohlich JJ: Effective exercise modality to reduce insulin resistance in women with type 2 diabetes. *Diabetes Care* 26:2977–2982, 2003
 24. Dunn AL, Marcus BH, Kampert JB, Garcia ME, Kohl HW 3rd, Blair SN: Comparison of lifestyle and structured interventions to increase physical activity and cardiorespiratory fitness: a randomized trial. *JAMA* 281:327–334, 1999
 25. Centers for Disease Control and Prevention: Surgeon General's report on physical activity and health. *JAMA* 276:522, 1996
 26. Tuomilehto J, Lindström J, Eriksson JG, Valle TT, Hämäläinen H, Ilanne-Parikka P, Keinänen-Kiukaanniemi S, Laakso M, Louheranta A, Rastas M, Salminen V, Uusitupa M: Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med* 344:1343–1350, 2001
 27. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM: Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 346:393–403, 2002
 28. Pan XR, Li GW, Hu YH, Wang JX, Yang WY, An ZX, Hu ZX, Lin J, Xiao JZ, Cao HB, Liu PA, Jiang XG, Jiang YY, Wang JP, Zheng H, Zhang H, Bennett PH, Howard BV: Effects of diet and exercise in preventing NIDDM in people with impaired glucose tolerance: the Da Qing IGT and Diabetes Study. *Diabetes Care* 20:537–544, 1997
 29. Eriksson J, Lindström J, Valle T, Aunola S, Hamalainen H, Ilanne-Parikka P, Keinänen-Kiukaanniemi S, Laakso M, Lauhkonen M, Lehto P, Lehtonen A, Louheranta A, Mannelin M, Martikkala V, Rastas M, Sundvall J, Turpeinen A, Viljanen T, Uusitupa M, Tuomilehto J: Prevention of type II diabetes in subjects with impaired glucose tolerance: the Diabetes Prevention Study (DPS) in Finland: study design and 1-year interim report on the feasibility of the lifestyle intervention programme. *Diabetologia* 42:793–801, 1999
 30. World Health Organization: *Diabetes Mellitus: Report of a WHO Study Group*. Geneva, World Health Org., 1985 (Tech. Rep. Ser., no. 727)
 31. Lakka TA, Salonen JT: Intra-person variability of various physical activity assessments in the Kuopio Ischaemic Heart Disease Risk Factor Study. *Int J Epidemiol* 21:467–472, 1992
 32. Lakka TA, Venäläinen JM, Rauramaa R, Salonen R, Tuomilehto J, Salonen JT: Relation of leisure-time physical activity and cardiorespiratory fitness to the risk of acute myocardial infarction. *N Engl J Med* 330:1549–1554, 1994
 33. Pereira MA, FitzerGerald SJ, Gregg EW, Joswiak ML, Ryan WJ, Suminski RR, Utter AC, Zmuda JM: A collection of physical activity questionnaires for health-related research. *Med Sci Sports Exerc* 29:S1–S205, 1997
 34. Lindström J, Louheranta A, Mannelin M, Rastas M, Salminen V, Eriksson J, Uusitupa M, Tuomilehto J: The Finnish Diabetes Prevention Study (DPS): lifestyle intervention and 3-year results on diet and physical activity. *Diabetes Care* 26:3226–3229, 2003
 35. Willett WC: Nutritional epidemiology. In *Monographs in Epidemiology and Biostatistics*. 2nd ed. Kelsey JL, Marmot MC, Stolley PD, Vessey MP, Eds. Oxford, Oxford University Press, 1998, p. 1–514
 36. Barengo NC, Nissinen A, Tuomilehto J, Pekkarinen H: Twenty-five-year trends in physical activity of 30- to 59-year-old populations in eastern Finland. *Med Sci Sports Exerc* 34:1302–1307, 2002
 37. Wolf AM, Hunter DJ, Colditz GA, Manson JE, Stampfer MJ, Corsano KA, Rosner B, Kriska A, Willett WC: Reproducibility and validity of a self-administered physical activity questionnaire. *Int J Epidemiol* 23:991–999, 1994
 38. Uusitupa M, Lindi V, Louheranta A, Salopuro T, Lindström J, Tuomilehto J: Long-term improvement in insulin sensitivity by changing lifestyles of persons with impaired glucose tolerance. *Diabetes* 52:2532–2538, 2003
 39. Henriksen EJ: Effects of acute exercise and exercise training on insulin resistance (Invited Review). *J Appl Physiol* 93:788–796, 2002
 40. Eriksson KF, Lindgarde F: Prevention of type 2 (non-insulin-dependent) diabetes mellitus by diet and physical exercise: the 6-year Malmo feasibility study. *Diabetologia* 34:891–898, 1991