

Rosiglitazone Treatment in Zucker Diabetic Fatty Rats Is Associated With Ameliorated Cardiac Insulin Resistance and Protection From Ischemia/Reperfusion-Induced Myocardial Injury

Tian-Li Yue, Weike Bao, Juan-Li Gu, Jianqi Cui, Ling Tao, Xin-Liang Ma, Eliot H. Ohlstein, and Beat M. Jucker

The mechanism responsible for the enhanced myocardial susceptibility to ischemic insult in patients with type 2 diabetes is not clear. The present study examines the effect of rosiglitazone treatment on cardiac insulin sensitization and its association with cardioprotection from ischemia/reperfusion injury in an animal model of diabetes. Male Zucker diabetic fatty (ZDF) rats were treated with rosiglitazone ($3 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$ orally) or vehicle for 8 days before undergoing 30 min of coronary artery ligation, followed by reperfusion for 4 h (apoptosis) or 24 h (infarction). Rosiglitazone reduced the blood levels of glucose, triglycerides, and free fatty acids; enhanced cardiac glucose oxidation; and increased Akt phosphorylation (Akt-pS473) 2.1-fold and Akt kinase activity 1.8-fold in the ischemic myocardium. The phosphorylation of two downstream targets of Akt, glycogen synthase kinase- β and FKHR (forkhead transcription factor), was also enhanced by 2- and 2.9-fold, respectively. In rosiglitazone-treated rats, the number of apoptotic cardiomyocytes and the myocardial infarct size were decreased by 58 and 46%, respectively, and the myocardial contractile dysfunction was improved. Blockade of the insulin-Akt signaling pathway by wortmannin in the 8-day rosiglitazone-treated ZDF rats resulted in a markedly diminished cardioprotective effect of rosiglitazone. In addition, 8-day rosiglitazone treatment in Zucker lean rats or 2-day rosiglitazone treatment in ZDF rats, both of which showed no change in whole-body insulin sensitivity, resulted in a significant reduction in cardiac infarct size, but to a lesser degree when compared with that observed in 8-day rosiglitazone-treated ZDF rats. These results suggest that chronic treatment with rosiglitazone protects the heart

against ischemia/reperfusion injury in ZDF rats, and that the enhanced cardiac protection observed after rosiglitazone treatment might be attributable in part to an improvement in cardiac insulin sensitivity. *Diabetes* 54:554–562, 2005

Hear disease is the leading cause of death among patients with diabetes (rev. in 1). Relative to nondiabetic individuals, patients with diabetes demonstrate impaired recruitment of contractile reserve in noninfarct segment (2), greater reduction in global left ventricular (LV) function (3), and increased incidence of congestive heart failure (4) and death (5) after an index myocardial infarction. A significant increase in the number of necrotic cardiomyocytes in ventricular myocardial biopsies obtained from diabetic patients was also reported, reflecting a primary impairment in myocardial ischemic tolerance (6,7). Therefore, type 2 diabetes has been recognized as an independent risk factor for ischemic cardiomyopathy (8).

Insulin resistance is known as a central pathophysiological feature of type 2 diabetes. Insulin resistance in the heart has been demonstrated in obese rats, associated with decreased GLUT4 protein content and impaired GLUT4 translocation to the sarcolemma (9,10). Cardiac insulin resistance was also demonstrated in diabetic patients with cardiac diseases (11–13). A recent study used direct techniques to measure the response of the arterial-coronary sinus glucose balance to an increase in local intracoronary insulin concentration, and it demonstrated that cardiac muscles of type 2 diabetic patients are insulin resistant during fasting relative to those of nondiabetic control subjects (14). These data suggest that a potential underlying mechanism related to the increased cardiac susceptibility to ischemic insult in type 2 diabetic patients may be caused in part by cardiac insulin resistance. However, the impact of type 2 diabetes on the insulin response system of the myocardium in vivo and the increased susceptibility to ischemic injury have not been fully examined.

In contrast to the overwhelming clinical data indicating that the diabetic heart is more sensitive to ischemic injury, experimental evidence from a number of studies using animal models of diabetes showed no change (15) or an

From the Department of Investigative and Cardiac Biology, GlaxoSmithKline Pharmaceuticals, King of Prussia, Pennsylvania.

Address correspondence and reprint requests to Tian-Li Yue, PhD, Department of Investigative and Cardiac Biology, GlaxoSmithKline, 709 Swedeland Rd., P.O. Box 1539, UW2510, King of Prussia, PA 19406. E-mail: tian-li_yue@gsk.com.

Received for publication 10 February 2004 and accepted in revised form 27 September 2004.

L.T. and X.-L. M. are currently affiliated with the Department of Emergency Medicine, Thomas Jefferson University, Philadelphia, Pennsylvania.

DAPI, 4',6-diamidino-2-phenylindole; GSK, glycogen synthase kinase; LV, left ventricular; NMR, nuclear magnetic resonance; PI, phosphatidylinositol; PPAR, peroxisome proliferator-activated receptor; TUNEL, terminal deoxynucleotidyl transferase-mediated dUTP nick-end labeling.

© 2005 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

increase (16) or a decrease (17,18) in susceptibility to ischemia. The reasons for such diverse effects may be due to the difference in the length or severity of the diabetic state, the severity of the ischemic insult, or the levels of exogenous substrates (16). Additionally, a number of *in vivo* studies in nondiabetic animals (19–22) and *ex vivo* studies in diabetic animals (23,24) have shown the cardioprotective effect of thiazolidinediones. Although there is a clear beneficial effect of thiazolidinediones in protecting insulin-sensitive hearts from ischemia-reperfusion injury, the mechanism for this protection is less clear. It has been postulated that thiazolidinediones may inhibit the inflammatory response after such injury in the heart (20–22,24). However, the degree of *in vivo* cardioprotection from ischemia/reperfusion insult and the role of insulin sensitization in the heart after thiazolidinedione treatment in diabetic animals is not known.

The aim of this study was to determine whether rosiglitazone, a peroxisome proliferator-activated receptor (PPAR)- γ agonist, protects the heart against ischemia/reperfusion injury in the Zucker diabetic fatty (ZDF) rat, and, if so, whether this cardioprotection is associated with an improvement in cardiac insulin sensitivity.

RESEARCH DESIGN AND METHODS

This study was conducted in accordance with the guidelines for care and use of laboratory animals of the U.S. National Institutes of Health. Male ZDF rats (ZDF/Gmi-*fa/fa*) and lean Zucker (*fa/?*) rats (Genetic Models) at age 12–14 weeks were used ($n = 245$). Unless otherwise indicated, rosiglitazone maleate (GlaxoSmithKline) was administered at $3 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$ via oral gavage for 7 days, and the eighth dose was given 1 h before ischemia. The animals were premedicated with atropine sulfate (0.04 mg/kg intramuscularly) and then anesthetized with Nembutal (60 mg/kg i.p.; Abbott Laboratories). Trachea were intubated and ventilated with a Harvard small animal respirator. The electrocardiogram and the body temperature of the rats were monitored throughout the experiment. Coronary artery (left anterior descending artery) occlusion (30 min) and reperfusion (4 h for apoptosis and 24 h for infarct and functional assessment) were induced by inflating and then deflating a non-traumatic balloon occluder that was fixed on the left anterior descending artery. The successful performance of coronary occlusion and reperfusion was verified by visual inspection of color in the apex and typical electrocardiogram changes.

Assessment of cardiac glucose versus fat oxidation. In a separate group of rosiglitazone- and vehicle-treated ZDF rats, indwelling carotid artery and jugular vein catheters were surgically implanted. Upon recovery from surgery, rats were fasted overnight before baseline or normoglycemic-hyperinsulinemic cardiac glucose oxidation measurements. A normoglycemic-hyperinsulinemic ($10 \text{ mU} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$, Humulin Regular; Eli Lilly, Indianapolis, IN) clamp was initiated in awake rats by administering a variable rate [$1\text{-}^{13}\text{C}$]glucose (99% enriched, 20% wt/vol; Cambridge Isotope Laboratories, Cambridge, MA) infusion at 2.5 min after the commencement of a primed ($150 \text{ mU} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ for 0–45 s, $75 \text{ mU} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ for 45–90 s) continuous insulin infusion. Plasma glucose concentrations were clamped at normoglycemic levels for 75 min. Blood samples were drawn at baseline, 7.5 min, 15 min, and every 15 min thereafter for immediate assessment of plasma glucose and lactate concentrations (2300 STAT Plus analyzer; YSI, Yellow Springs, OH). Baseline measurements were performed in awake rats by maintaining a [$1\text{-}^{13}\text{C}$]glucose ($10 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$)/somatostatin ($1.5 \mu\text{g}/\text{min}$) infusion for 120 min. At the end of the clamp experiments, rats were anesthetized with Nembutal at 50 mg/kg i.v., and the heart was rapidly removed and frozen in liquid nitrogen.

Cardiac tissue extracts were prepared for high-field nuclear magnetic resonance (NMR) analysis by homogenizing $\sim 0.3 \text{ g}$ of the heart in a similar fashion as previously described (25). NMR analysis was performed at 9.4 Tesla (WB-400 Avance NMR spectrometer; Bruker Medical, Billerica, MA). Proton-observed, carbon-enhanced spectroscopy was performed on tissue extract samples for measuring [$3\text{-}^{13}\text{C}$]lactate, [$3\text{-}^{13}\text{C}$]alanine, and [$4\text{-}^{13}\text{C}$]glutamate fractional enrichments (25). The relative cardiac glucose and fat/ketone oxidation rates in terms of acetyl-CoA units were assessed from the metabolite pool enrichments. The degree to which the [$4\text{-}^{13}\text{C}$]glutamate pool was diluted from the [$3\text{-}^{13}\text{C}$]pyruvate (equivalent to the [$3\text{-}^{13}\text{C}$]lactate and

[$3\text{-}^{13}\text{C}$]alanine pool enrichment) pool depends on the extent of label dilution from unlabeled fat oxidation present. Therefore, the relative glucose oxidation (%) is defined as $\{[4\text{-}^{13}\text{C}]\text{glutamate}/([3\text{-}^{13}\text{C}]\text{lactate} + [3\text{-}^{13}\text{C}]\text{alanine})/2\} \times 100$, and relative fat/ketone oxidation (%) is calculated as $(1 - \text{relative glucose oxidation}) \times 100$.

Western blot analysis for determination of Akt, Akt-pS473, glycogen synthase kinase-3 β -pS9, and FKHR-pS256. Myocardial Akt, phospho-Akt (Akt-pS473), phospho-FKHR (forkhead transcription factor; FKHR-pS256), and phospho-glycogen synthase kinase (GSK)-3 β (GSK-3 β -pS9) in the ischemic myocardium after 4-h reperfusion were determined by Western blot analysis as reported previously (26). Briefly, protein extracts ($50\text{--}100 \mu\text{g}$) from ischemic myocardium collected at 4 h after reperfusion were fractionated on 10% polyacrylamide gel under reducing conditions, transferred onto nitrocellulose membranes, and probed with the following primary antibodies against Akt: Akt-pS473, FKHR-pS256, and GSK-3 β -pS9 (all antibodies from Cell Signaling). Blots were incubated with horseradish peroxidase-conjugated second antibody, and the signal was detected with LumiGlo chemiluminescent reagent (Cell Signaling). β -Actin was used for normalization.

Akt kinase activity assay. Ischemic myocardial tissues collected 4 h after reperfusion were lysed and immunoprecipitated with anti-Akt antibody and then used to measure Akt kinase activity using an Akt kinase assay kit (Cell Signaling) with GSK-3 α/β as an exogenous substrate, according to the manufacturer's instructions.

Determination of myocardial apoptosis. The apoptotic myocytes in ischemic myocardium after 4-h reperfusion were determined by terminal deoxynucleotidyl transferase-mediated dUTP nick-end labeling (TUNEL) assay with a cell death detection kit (Roche), as described previously (26). Briefly, the tissue sections after permeability treatment were incubated with TUNEL reaction mixture containing terminal deoxynucleotidyl transferase and fluorescein-dUTP (deoxyuridine triphosphate). For double labeling, the tissue sections after incubation with TUNEL reaction mixture and blockade of endogenous peroxidase, and nonspecific binding sites were incubated with the monoclonal antibody against α -sarcomeric actin (Sigma). After washing, the slides were incubated with biotinylated secondary antibody, and the cardiomyocytes were finally visualized with Texas red avidin. For detection of total nuclei, the slides were covered with the mounting medium containing 4',6-diamidino-2-phenylindole (DAPI). At least six slides per block were evaluated. For each slide, 15 fields were randomly chosen, and a total of $2.5\text{--}3 \times 10^3$ myocytes per slide were counted. We determined the index of apoptosis, i.e., the number of TUNEL-positive myocytes divided by the total myocytes that were stained with anti- α -sarcomeric actin (see Fig. 4C) per field.

DNA ladder. DNA ladder experiments were performed as described previously (27). In brief, ischemic myocardial tissues collected at 4 h after reperfusion were minced in lysis buffer on ice, and proteinase K ($100 \mu\text{g}/\text{ml}$) was added. After incubation at 55°C with shaking for 18 h, DNA was extracted with phenol/chloroform three times, precipitated in ethanol, treated with DNA-free RNase, re-extracted, and precipitated again. DNA concentration was determined, and $10 \mu\text{g}$ of DNA was used for electrophoreses on a 1.8% agarose gel.

Determination of myocardial infarction. The ischemic area (area at risk) was distinguished from the area not at risk by Evans blue dye staining, and the infarcted portion of the myocardium (necrotic area) was determined by the triphenyl tetrazolium chloride method as described in detail previously (20). The three portions (i.e., area not at risk, area at risk, and necrotic area) of the left ventricle were quantified by use of Image-Pro software.

Assessment of myocardial contractile function. LV pressure and the arterial blood pressure were measured via two 1.4 F Millar Mikrotip catheter transducers that were inserted into the LV cavity through the right carotid artery and the right femoral artery, respectively. LV pressure and arterial blood pressure were digitally processed via a hemodynamic analyzing system (Gould 3P 6600). Mean arterial blood pressure, LV systolic pressure, LV end diastolic pressure, positive and negative maximal values of the first derivative of LV pressure ($+dp/dt$ and $-dp/dt$), and heart rate were derived by computer algorithms (20).

Statistical analysis. Data are expressed as the means \pm SE and analyzed by one-way ANOVA with subsequent post hoc paired comparisons or by unpaired Student's *t* test. Differences with a value of $P < 0.05$ were considered statistically significant.

RESULTS

Rosiglitazone improves whole-body and cardiac insulin sensitivity in ZDF rats. As shown in Table 1, treatment with rosiglitazone for 8 days significantly reduced the circulating levels of glucose, triglycerides, and free

TABLE 1
Effect of rosiglitazone on blood levels of glucose, triglycerides, and free fatty acids

Group	Glucose (mg/dl)	Triglycerides (mg/dl)	Free fatty acids (mmol/l)
Zucker lean	137 ± 4	44 ± 2	0.9 ± 0.05
ZDF	235 ± 12*	504 ± 41*	1.7 ± 0.12*
ZDF + rosiglitazone	179 ± 13†‡	303 ± 39*‡	1.1 ± 0.06‡

Data are means ± SD. Blood samples were collected from the animals that were fasted overnight (15 h). The glucose, triglycerides, and free fatty acid measurements were performed on an Olympus AU640 chemistry analyzer. * $P < 0.01$ vs. Zucker lean rats; † $P < 0.05$ vs. Zucker lean rats; ‡ $P < 0.01$ vs. ZDF rats ($n = 14-15$).

fatty acids in ZDF rats. During a normoglycemic-hyperinsulinemic clamp, the 8-day rosiglitazone-treated animals exhibited an increased insulin-stimulated glucose infusion rate from 13.2 ± 1.4 to 19.1 ± 2.4 $\text{mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ for vehicle and rosiglitazone, respectively ($P < 0.03$, $n = 8$). NMR experiments revealed that glucose oxidation under baseline conditions represented a small fraction of the entire substrate oxidation in the heart, and 8-day treatment with rosiglitazone had no effect on this measure (Fig. 1). After insulin stimulation, both 8-day rosiglitazone and vehicle groups exhibited an increase in relative cardiac glucose oxidation. However, the rosiglitazone-treated animals exhibited a markedly increased insulin-sensitized glucose oxidation rate versus vehicle-treated animals (Fig. 1). Conversely, fat/ketone oxidation in the heart was lower in rosiglitazone-treated animals versus the vehicle group (41.2 ± 4.8 vs. $54.7 \pm 0.9\%$ acetyl-CoA source, respectively; $P < 0.02$; $n = 8$).

Rosiglitazone activates Akt signal pathway in ischemic myocardium. As shown in Fig. 2, treatment with rosiglitazone for 8 days resulted in a significant increase in Akt phosphorylation (Akt-pS473) in the ischemic myocardium (Fig. 2A). Quantitative measurement by Western blot analysis showed a 2.1-fold increase in Akt-pS473 in the rosiglitazone-treated group versus the vehicle group ($P <$




FIG. 1. Relative cardiac glucose oxidation at baseline or during a normoglycemic-hyperinsulinemic clamp. The relative cardiac glucose oxidation was expressed as the percent of acetyl-CoA units oxidized by glucose versus fat/ketone metabolism. The infusion experiments were performed in awake ZDF rats and the measurements were performed in whole heart in vehicle- and rosiglitazone (ROSI)-treated animals ($n = 8$).

0.01, $n = 8$) (Fig. 2B). The result of the Akt kinase assay (Fig. 2C) was consistent with the Western blot analysis, showing a 1.8-fold increase in cardiac Akt activity in rosiglitazone-treated animals compared with the vehicle group ($P < 0.01$, $n = 8$). There was no difference in myocardial levels of total Akt between the rosiglitazone-treated and the vehicle groups (Fig. 2A).

GSK-3 and FKHR are two downstream targets of Akt. As shown in Fig. 3B and C, treatment with rosiglitazone enhanced the phosphorylation of GSK-3 (GSK-3 β -pS9) by 2-fold ($P < 0.05$, $n = 8$) and FKHR (FKHR-pS256) by 2.9-fold ($P < 0.01$, $n = 8$) in the ischemic myocardium compared with the vehicle group, respectively. Figure 3A is a representative Western blot from one of two experiments showing similar results.

Rosiglitazone attenuates ischemia/reperfusion-induced myocardial apoptosis. Myocardial tissue from sham-operated ZDF rats exhibited no DNA ladder formation (Fig. 4A, lane 2), but clear DNA ladder formation was observed in myocardial tissue from the hearts subjected to ischemia/reperfusion and receiving vehicle (Fig. 4A, lanes 3–8). In contrast, DNA ladder formation was less clear or almost not detectable in the ischemic myocardium from six ZDF rats treated with rosiglitazone (Fig. 4A, lane 9–14).

In addition, the TUNEL assay demonstrated that TUNEL-positive-staining myocytes (apoptotic myocytes) in ischemic myocardium reached $21.3 \pm 3.5\%$ in the vehicle-treated ZDF rats (Fig. 4B). The TUNEL-positive myocytes were mainly localized in salvaged areas surrounding the infarcted tissues (data not shown). The number of TUNEL-positive myocytes in the ischemic regions from rosiglitazone-treated ZDF rats was reduced to $8.9 \pm 2.5\%$ ($n = 6$, $P < 0.01$ vs. vehicle) (Fig. 4B). Under the same conditions (i.e., 30-min ischemia followed by 4-h reperfusion), the number of TUNEL-positive myocytes in the ischemic myocardium from vehicle-treated lean Zucker rats was significantly lower ($14.3 \pm 2.1\%$) compared with the vehicle-treated ZDF rats ($P < 0.01$, $n = 6$), and this number was reduced to $8.1 \pm 1.2\%$ in rosiglitazone-treated lean rats ($P < 0.05$ vs. lean + vehicle) (Fig. 4B). Figure 4C is representative photomicrographs of in situ detection of apoptotic myocytes (TUNEL positive) in the heart tissue from ZDF rats subjected to 30 min of ischemia and 4 h of reperfusion and treated with vehicle (column V, rows 1–3) or rosiglitazone (column R, rows 1–3) for 8 days, or sham operated (column S, rows 1–3).

Rosiglitazone reduces ischemia/reperfusion-induced myocardial infarction. There was no difference in ischemic area (area at risk) expressed as percent of left ventricle (Fig. 5A, left side), indicating a comparable degree of ischemic insult between vehicle- and rosiglitazone-treated groups after left anterior descending artery occlusion in ZDF rats. Treatment with rosiglitazone for 8 days reduced ischemia/reperfusion-induced cardiac infarct size (% ischemic area) by 45.5%, from $60.8 \pm 2.5\%$ (vehicle) to $33.1 \pm 3.7\%$ (rosiglitazone) ($n = 10$, $P < 0.01$) (Fig. 5A, right side).

Ischemia/reperfusion-induced myocardial infarct size in the lean Zucker rats was significantly smaller than the infarct size in the ZDF rats ($P < 0.01$) (Fig. 5A, right side), although the ischemic area (% left ventricle) was similar




FIG. 2. Effect of rosiglitazone (ROSI) on Akt phosphorylation (Akt-pS473) in ZDF rat hearts subjected to ischemia (30 min)/reperfusion (4 h). **A:** Representative Western blots showing that treatment with rosiglitazone enhanced the expression of Akt-pS473 in the ischemic myocardium. Equal amounts of proteins extracted from ischemic myocardium were loaded into each lane. The sham group expressed low levels of Akt-pS473 in heart; therefore, data are not shown. The same blots after detection of Akt-pS473 were stripped and reblotted with anti-Akt antibody, showing no difference in the total Akt levels between the two groups. **B:** Quantitative results of measurement of Akt-pS473 using β -actin for normalization ($n = 8$). **C:** Phospho-Akt activity in the ischemic myocardium determined by Akt kinase assay kit (Cell Signaling) ($n = 8$). I/R, ischemia/reperfusion; Std, standards.

between the two groups (Fig. 5, left side). In rosiglitazone-treated lean rats, the ischemia/reperfusion-induced cardiac infarction was reduced by 26.5% compared with the vehicle group ($P < 0.05$, $n = 8$).

Rosiglitazone improves ischemia/reperfusion-induced myocardial dysfunction. Treatment with rosiglitazone for 8 days significantly improved cardiac functional contraction in ZDF rats at 24 h after reperfusion, as shown in Fig. 5B. The values of LV systolic pressure, $+dP/dt$, and $-dP/dt$ were significantly enhanced in the rosiglitazone-treated group compared with the vehicle group (all $P < 0.05$, $n = 9$). Additionally, the mean arterial blood pressure was also improved in the rosiglitazone-treated group versus the vehicle-treated group, but no difference in the heart rate between the two groups was observed (data not shown).

Wortmannin attenuates the cardioprotective effect of rosiglitazone. As shown in Fig. 6, pretreatment of the ZDF rats with wortmannin, a phosphatidylinositol (PI) 3-kinase inhibitor, 15 min before reperfusion (26) markedly attenuated the cardioprotective effect of 8-day rosiglitazone treatment ($>90\%$ reduction). However, wortmannin alone had no effect on ischemia/reperfusion-induced myocardial infarction.

Rosiglitazone cardioprotection is independent of insulin sensitization. To further differentiate the non-insulin-related cardioprotective effect of rosiglitazone, a separate group of ZDF rats were treated with rosiglitazone at $3 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$ for 2 days. There was no difference in glucose infusion rates between the rosiglitazone-treated and vehicle-treated groups (15.5 ± 1.2 and $15.4 \pm 0.6 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$, respectively) during a normoglycemic-hyper-

insulinemic clamp ($n = 10$) (Fig. 7A), indicating that 2-day rosiglitazone treatment did not change insulin resistance in the rats. The ischemia/reperfusion-induced cardiac infarct size (% ischemic area) was reduced by 22.1%, from $58.6 \pm 1.8\%$ (vehicle) to $46.2 \pm 1.7\%$ (rosiglitazone; $n = 9$, $P < 0.05$) (Fig. 7B). This reduction was significantly less when compared with the 8-day rosiglitazone-treated group ($P < 0.05$). Additionally, there was no activation in myocardial level of Akt-pS473 in rosiglitazone-treated animals, as shown in Fig. 7C. The apoptotic myocytes in the ischemic myocardium of ZDF rats was $16.7 \pm 5.6\%$ ($n = 4$, reduced by 21.6% compared with the vehicle group).

DISCUSSION

Treatment with rosiglitazone for 8 days in ZDF rats resulted in a decrease in the in vivo ischemia/reperfusion-induced cardiac infarct size by 46% and apoptosis by 58%. This cardiac protection was associated with improved LV function as well as increased cardiac insulin sensitivity. There was a significant increase in insulin-sensitized cardiac glucose oxidation and a reduction in fat/ketone oxidation in rosiglitazone-treated ZDF rats versus the vehicle group. Concomitantly, a significantly enhanced level of Akt signaling in myocardium was observed in rosiglitazone-treated animals.

In ZDF rats, GLUT4 protein expression is decreased and insulin-stimulated cardiac glucose uptake is impaired (10,23). Previous studies have argued for a potential role of glucose in cardioprotection from ischemic injury (28–30). It was shown that cardiac myocytes in the presence of glucose and glycolytic inhibitors such as 2-deoxyglucose




FIG. 3. Effect of rosiglitazone (ROSI) on phosphorylation of GSK-3 β and FKHR (forkhead transcription factor) in the ischemic myocardium. **A:** Representative Western blots showing an enhancement in phosphorylation of GSK-3 (GSK-3 β -pS9) and FKHR (FKHR-pS256) in ischemic myocardium after 4-h reperfusion from rosiglitazone-treated ZDF rats compared with the vehicle-treated animals. **B** and **C:** Quantitative densitometry data are presented as the means \pm SE ($n = 8$). I/R, ischemia/reperfusion.

or iodoacetate reversed the protective effect of glucose on hypoxic injury (28). In isolated rat hearts, cardiac output after ischemia/reperfusion was increased when glucose was added to lactate in the absence of insulin (29). Conversely, if glucose is absent in the presence of hypoxia for 30 min followed by 60 min of reperfusion, LV function is impaired to a greater extent (30). A recent *in vitro* study has demonstrated that rosiglitazone treatment results in the normalization of insulin-stimulated glucose uptake in isolated perfused obese Zucker rat hearts (23). As a result, the increased insulin-sensitized cardiac glucose oxidation

observed in the present study may in part contribute to the cardioprotection by rosiglitazone in the ZDF rats. However, it is also known that ischemia/reperfusion results in an increase in free fatty acid levels, which aggravates the severity of myocardial infarction (31). Therefore, the cardioprotection observed after the 8-day rosiglitazone treatment in the present study might be indirectly mediated by lowering systemic free fatty acid levels.

Akt signaling has been widely recognized as an important downstream modulator of the insulin receptor, and it plays a critical role in maintaining insulin sensitivity in humans (32). Akt2-deficient mice show resistance to insulin's effects on glucose metabolism and develop frank diabetes (33,34). A recent study reported that a mutation in the gene encoding the Akt2 in a family resulted in severe insulin resistance and diabetes, providing an example of a monogenic inherited defect in postreceptor insulin signaling that leads to human insulin resistance and diabetes (32). An increasing body of evidence also suggests that Akt signaling plays an important cytoprotective role against ischemic injury in the heart. In cultured rat neonatal cardiomyocytes, activation of Akt significantly reduced H₂O₂-induced apoptosis, and blockade of Akt signaling by PI 3-kinase inhibitor or overexpression of a dominant-negative mutant of PI 3-kinase abolished the protective effect of insulin (35). In isolated perfused hearts or *in vivo* rat studies, insulin administration reduced ischemia/reperfusion-induced infarct size, and this protective effect was abolished if the upstream or downstream signaling in the Akt pathway was inhibited by pharmacological inhibitor, suggesting that the cardioprotective effect of insulin is mainly mediated via Akt signaling (26). It has been reported that the cytoprotective effect of Akt signaling is attributable to the inhibition of GSK-3 and FKHR, two downstream targets of Akt known to be involved in cell death. GSK-3 is inactivated by Akt via phosphorylation at serine 9 or 21 (36). Both the *in vitro* and *in vivo* studies have demonstrated that inactivation of GSK-3 enhanced cell survival (37,38), improved postischemic cardiac function, and reduced infarct size (39). Phosphorylation of FKHR by Akt keeps this transcriptional factor retention in the cytoplasm and away from the nucleus, where it triggers activation of genes critical for inducing cell death (40). The present study demonstrated a significant increase in the levels of GSK-3 β -pS9 and FKHR-pS256 in the ischemic myocardium from rosiglitazone-treated ZDF rats, further suggesting a role of the Akt signaling pathway in PPAR- γ -mediated cardioprotection. It is interesting to note in the current study that treatment with rosiglitazone did not change the total Akt level in the heart, but only the active form of Akt (Fig. 2A), suggesting that PPAR- γ -mediated activation of Akt may serve as a protective mechanism when the heart is under an insulting stress.

Previous studies have shown that PPAR- γ agonists protected the heart against ischemia/reperfusion injury in isolated perfused hearts (24) as well as *in vivo* in the nondiabetic pig (19) and rat (20–22). The enhanced insulin sensitivity by chronic treatment with troglitazone was found in nondiabetic pigs, and it was assumed to contribute to the improved recovery of LV function after cardiac ischemia (19). However, other actions of the PPAR- γ agonist that do not appear to be linked to insulin sensi-




FIG. 4. Protection by rosiglitazone (ROSI) of myocardial apoptosis after ischemia and reperfusion in ZDF rats. **A:** Electrophoretic analysis of internucleosomal DNA extracted from ischemic myocardium from ZDF rats exposed to 30 min of ischemia and 4 h of reperfusion in the presence of vehicle (lanes 3–8) or rosiglitazone (lanes 9–14). The first lane is a DNA size marker, and the second lane is from sham-operated controls. **B:** Quantitative measurement of the percentage of apoptotic cardiomyocytes (nuclei staining positive for TUNEL) in ischemic tissue sections from ZDF rats or lean rats subjected to ischemia (30 min)/reperfusion (4 h) in the absence or presence of rosiglitazone, or the sham-operated control ($n = 6$). **C:** Representative photomicrographs of in situ detection of DNA fragments (TUNEL assay) in heart tissue from ZDF rats that were treated with vehicle (column V, rows 1–3) or rosiglitazone (column R, rows 1–3) for 8 days and subjected to 30 min of ischemia and 4 h of reperfusion. S, sham-operated (column S, rows 1–3). Column V, row 1; column R, row 1; and column S, row 1, depict the total nuclei stained with DAPI. Column V, row 2; column R, row 2; and column S, row 2, are the same sections of column V, row 1; column R, row 1; and column S, row 1, respectively, depicting the apoptotic nuclei stained with fluorescein isothiocyanate (TUNEL-positive nuclei). Column V, row 3; column R, row 3; and column S, row 3, are the same sections of column V, row 1; column R, row 1; and column S, row 1, respectively. They were double-labeled with anti- α -sarcomeric actin and visualized with Texas red avidin (see RESEARCH DESIGN AND METHODS). I/R, ischemia/reperfusion.

zation, such as anti-inflammatory activities (20–22) and inhibition of c-Jun NH₂-terminal kinase/activating protein 1 pathway (24), have been suggested as potential mechanisms responsible for PPAR- γ -mediated cardioprotection.

Nevertheless, the present study provides several lines of evidence suggesting a potential association between cardiac insulin resistance and enhanced susceptibility to ischemic insult. First, the ischemia/reperfusion-induced




FIG. 5. A: Protection by rosiglitazone (ROSI) against ischemia/reperfusion-induced myocardial infarction in ZDF and lean Zucker rats. Rosiglitazone was administered at $3 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$ via oral gavage for 8 days, and the last dose was given 1 h before ischemia. Animals were subjected to ischemia for 30 min and killed 24 h after reperfusion. The left side of the figure shows the ischemic area, expressed as percent left ventricle. The right side is the infarct size, expressed as percent ischemic area ($n = 8-10$). **B:** Rosiglitazone improves ischemia/reperfusion-induced cardiac dysfunction. The ZDF rats were treated with rosiglitazone or vehicle, as described in A, and the myocardial contractile function was determined at 24 h after reperfusion ($n = 9$). LVSP, LV systolic pressure.

myocardial injury (apoptosis and infarction) was greater in the ZDF rats than in the lean controls, suggesting a role for insulin resistance in the enhanced myocardial susceptibility to ischemia. Second, a greater degree of cardioprotection by rosiglitazone was observed in ZDF versus lean controls, suggesting that the improved cardiac insulin sensitivity provided the additional degree of cardioprotection. This finding is consistent with our previous study in nondiabetic Lewis rats (20). The cardioprotection by rosiglitazone against ischemia/reperfusion injury in Lewis rats was significant (24% reduction versus vehicle) but much less compared with that observed in ZDF rats in the present study (46%) while the animals were treated with the same dose regimen ($3 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$ orally for 8 days) (20).

To further differentiate the insulin-sensitizing from the non-insulin-sensitizing mechanism in the cardioprotective role of rosiglitazone, a 2-day rosiglitazone dosing regimen was also used in the present study. The lack of insulin sensitization in the ZDF rats under this 2-day dosing regimen was associated with significantly less cardiopro-




FIG. 6. Treatment with wortmannin (WN) attenuates the cardioprotection by rosiglitazone (ROSI). The ZDF rats received rosiglitazone for 8 days, as described in the legend for Fig. 5. Wortmannin ($15 \mu\text{g}/\text{kg}$) or vehicle was administered intravenously 15 min before reperfusion. The animals were killed 24 h later, and the ischemic area and infarct size were determined ($n = 8-10$).




FIG. 7. Effect of 2-day rosiglitazone (ROSI) treatment on whole-body insulin sensitivity (A), ischemia/reperfusion-induced myocardial infarction size (B), and cardiac Akt phosphorylation (C). ZDF rats were treated with vehicle or rosiglitazone ($3 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$, orally) for 2 days, and whole-body glucose disposal was measured during a normoglycemic-hyperinsulinemic clamp ($n = 10$) (A), or rats were subjected to myocardial ischemia (30 min) followed by reperfusion for 24 h to determine infarct size ($n = 9$, $P < 0.05$ vs. vehicle) (B) or reperfusion for 4 h to measure cardiac Akt phosphorylation ($n = 4$, lanes 3–6) (C). Lanes 1–2 are samples from the ZDF rats treated with rosiglitazone for 8 days. * $P < 0.05$ vs. vehicle. GIR, glucose infusion rate; P-Akt, Akt phosphorylation (Akt-pS473).

tection than that observed after an 8-day rosiglitazone treatment. Nevertheless, approximately half of the protection seen after an 8-day rosiglitazone treatment was observed after only a 2-day rosiglitazone treatment. Finally, pretreatment of ZDF rats with wortmannin at a dose that completely blocked Akt phosphorylation and activation by insulin, as demonstrated previously (26), remarkably diminished but did not abolish the cardioprotective effect of rosiglitazone. These results suggest that although non-insulin-related mechanisms appear to be involved in the cardioprotection by rosiglitazone, the improvement in insulin sensitization appears to be associated with enhanced cardioprotection against ischemia/reperfusion-induced injury in rosiglitazone-treated ZDF rats. It is not

clear at present to what extent insulin- and non-insulin-related mechanisms play a role in the cardioprotection observed in the diabetic and nondiabetic condition. Further studies are necessary to clearly define and differentiate the direct role for both insulin- and non-insulin-related cardioprotective mechanisms.

In conclusion, chronic treatment with rosiglitazone protects the heart from ischemia/reperfusion-induced cardiac injury and improves cardiac contractile dysfunction in ZDF rats. The improvement in cardiac insulin sensitivity after an 8-day rosiglitazone treatment, as reflected by increased Akt signaling and glucose oxidation, was associated with this cardioprotection.

REFERENCES

1. Taegtmeyer H, McNulty P, Young ME: Adaptation and maladaptation of the heart in diabetes. Part I. General concepts. *Circulation* 105:1727–1733, 2002
2. Takahashi N, Iwasaka T, Sugiura T, Hasegawa T, Tarumi N, Kimura Y, Kurihara S, Onoyama H, Inada M: Left ventricular regional function after acute anterior myocardial infarction in diabetic patients. *Diabetes Care* 12:630–635, 1989
3. Iwasaka T, Takahashi N, Nakamura S, Sugiura T, Tarumi N, Kimura Y, Okubo N, Taniguchi H, Matsui Y, Inada M: Residual left ventricular pump acute myocardial infarction in NIDDM patients. *Diabetic Care* 15:1522–1526, 1992
4. O'Connor CM, Hathaway WR, Bates ER, Leimberger JD, Sigmon KN, Kereiakes DJ, George BS, Samaha JK, Abbottsmith CW, Candela RJ, Topol EJ, Califf RM: Clinical characteristics and long-term outcome in patients in whom congestive heart failure develops after thrombolytic therapy for acute myocardial infarction: development of a predictive model. *Am Heart J* 133:663–673, 1997
5. Miettinen H, Lehto S, Salomaa V, Mahonen M, Niemela M, Haffner SM, Pyorala K, Tuomilehto J: Impact of diabetes on mortality after the first myocardial infarction: the FINMONICA Myocardial Infarction Register Study Group. *Diabetes Care* 21:69–75, 1998
6. Frustaci A, Kajstura J, Chimenti C, Jakoniuk I, Leri A, Maseri A, Nadal-Ginard B, Anversa P: Myocardial cell death in human diabetes. *Circ Res* 87:1123–1132, 2000
7. Woodfield SL, Lundergan CF, Reiner JS, Greenhouse SW, Thompson MA, Rohrbeck SC, Deychak Y, Simoons ML, Califf RM, Topol EJ, Ross AM: Angiographic findings and outcome in diabetic patients treated with thrombolytic therapy for acute myocardial infarction: the GUSTO-1 experience. *J Am Coll Cardiol* 28:1661–1669, 1996
8. Domanski M, Krause-Steinrauf H, Deedwania P, Follmann D, Ghali JK, Gilbert E, Haffner S, Katz R, Lindenfeld J, Lowes BD, Martin W, McGrew F, Bristow MR: The effect of diabetes on outcomes of patients with advanced heart failure in the BEST trial. *J Am Coll Cardiol* 42:914–922, 2003
9. Utriainen T, Takala T, Luotolahti M: Insulin resistance characterizes glucose uptake in skeletal muscle but not the heart in NIDDM. *Diabetologia* 41:555–559, 1998
10. Zaninetti D, Greco-Perotto R, Assimacopoulos-Jeannot F, Jeanrenaud B: Dysregulation of glucose transport and transporters in perfused hearts of genetically obese (fa/fa) rats. *Diabetologia* 32:56–60, 1989
11. Volpio-Pulkki L-M, Nuutila P, Knuuti MJ, Ruotsalainen U, Haaparanta M, Teras M, Wegelius U, Koivisto VA: Heart and skeletal muscle glucose disposal in type 2 diabetic patients as determined by positron emission tomography. *J Nucl Med* 34:2064–2067, 1993
12. Paternostro G, Camici PG, Lammerstma AA, Marinho N, Baliga RR, Kooner JS, Kadda GK, Ferrannini E: Cardiac and skeletal muscle insulin resistance in patients with coronary heart diseases: a study with positron emission tomography. *J Clin Invest* 98:2094–2099, 1996
13. Ohtake T, Yokoyama I, Watanabe T, Momose T, Serezawa T, Nishikawa J, Sasaki Y: Myocardial glucose metabolism in non-insulin-dependent diabetes mellitus patients evaluated by FDG-PET. *J Nucl Med* 36:456–463, 1995
14. Jagasia D, Whiting JM, Concato J, Pfau S, McNulty PH: Effect of non-insulin-dependent diabetes mellitus on myocardial insulin responsiveness in patients with ischemic heart disease. *Circulation* 103:1734–1739, 2001
15. Lopaschuk GD, Saddik M, Barr R, Huang L, Barder DD, Muzyke RA: Effects of high levels of fatty acids on functional recovery of ischemic hearts from diabetic rats. *Am J Physiol* 263:E1046–E1053, 1992
16. Paulson DJ: The diabetic heart is more sensitive to ischemic injury. *Cardiovasc Res* 34:104–112, 1997

17. Feuvray D, Lopaschuk GD: Controversies on the sensitivity of the diabetic heart to ischemic injury: the sensitivity of the diabetic heart to ischemic injury is decreased. *Cardiovasc Res* 34:113–120, 1997
18. Wang P, Chatham JC: Onset of diabetes in Zucker diabetic fatty (ZDF) rats leads to improved recovery of function after ischemia in the isolated perfused heart. *Am J Physiol Endocrinol Metab* 286:E725–E736, 2004
19. Zhu P, Lu L, Xu Y, Schwartz GG: Troglitazone improves recovery of left ventricular function after regional ischemia in pigs. *Circulation* 101:1165–1171, 2000
20. Yue TL, Chen J, Bao W, Narayanan PK, Bril A, Jiang W, Lysko PG, Gu JL, Boyce R, Zimmerman DM, Hart TK, Buckingham RE, Ohlstein EH: In vivo myocardial protection from ischemia/reperfusion injury by the peroxisome proliferator-activated receptor- γ agonist rosiglitazone. *Circulation* 104:2588–2594, 2001
21. Wayman NS, Hattori Y, McDonald MC, Mota-Filipe H, Cuzzocrea S, Pisano B, Chatterjee PK, Thiemermann C: Ligands of the peroxisome proliferator-activated receptors (PPAR- γ and PPAR- α) reduce myocardial infarct size. *FASEB J* 16:1027–1040, 2002
22. Ito H, Nakano A, Kinoshita M, Matsumori A: Pioglitazone, a peroxisome proliferator-activated receptor- γ agonist, attenuates myocardial ischemia/reperfusion injury in a rat model. *Lab Invest* 83:1715–1721, 2003
23. Sidell RJ, Cole MA, Draper NJ, Desrois M, Buckingham RE, Clarke K: Thiazolidinedione treatment normalizes insulin resistance and ischemic injury in the Zucker fatty rat heart. *Diabetes* 51:1110–1117, 2002
24. Khandoudi N, Delerive P, Berrebi-Bertrand I, Buckingham RE, Staels B, Bril A: Rosiglitazone, a peroxisome proliferator-activated receptor- γ , inhibits the Jun NH₂-terminal kinase/activating protein 1 pathway and protects the heart from ischemia/reperfusion injury. *Diabetes* 51:1507–1514, 2002
25. Jucker BM, Rennings JM, Cline GW, Shulman GI: ¹³C and ³¹P NMR studies on the effects of increased plasma free fatty acids on intramuscular glucose metabolism in the awake rat. *J Biol Chem* 272:10464–10473, 1997
26. Gao F, Gao E, Yue TL, Ohlstein EH, Lopez BL, Christopher TA, Ma XL: Nitric oxide mediates the antiapoptotic effect of insulin in myocardial ischemia-reperfusion. *Circulation* 105:1497–1502, 2000
27. Yue TL, Ma XL, Wang XK, Romanic AM, Liu GL, Loudon C, Gu JL, Kumar S, Poste George, Ruffolo RR, Feuerstein GZ: Possible involvement of stress-activated protein kinase signaling pathway and Fas receptor expression in prevention of ischemia/reperfusion-induced cardiomyocyte apoptosis by carvedilol. *Circ Res* 82:166–174, 1998
28. Malhotra R, Brosius FC: Glucose uptake and glycolysis reduce hypoxia-induced apoptosis in cultured neonatal cardiac myocytes. *J Biol Chem* 274:12567–12575, 1999
29. Temsah RM, Netticadan T, Kawabata K-I, Dhalla NS: Lack of oxygen and glucose contributes to I/R-induced changes in cardiac SR function. *Am J Physiol (Cell Physiol)* 283:C1306–C1312, 2002
30. Van Rooyen J, McCarthy J, Opie LH: Increased glycolysis during ischemia mediates the protective effect of glucose and insulin in the isolated rat heart despite the presence of cardiodepressant exogenous substrates. *Cardiovasc J S Afr* 13:103–109, 2002
31. Van Der Vusse GJ, Glatz JFC, Stam CG, Reneman RS: Fatty acid homeostasis in the normoxic and ischemic heart. *Physiol Rev* 72:881–940, 1992
32. George S, Rochford JJ, Wolfrum C, Gray SL, Schinner S, Wilson JC, Soos MA, Murgatroyd PR, Williams RM, Acerini CL, Dunger DB, Barford D, Umpleby AM, Wareham NJ, Cavies HA, Schafer AJ, Stoffel M, O'Rahilly S, Barroso I: A family with severe insulin resistance and diabetes due to a mutation in AKT2. *Science* 304:1325–1328, 2004
33. Cho H, Mu J, Kim JK, Thorvaldsen JL, Chu Q, Crenshaw EB, Kaestner KH, Bartolomei MS, Shulman GI, Birnbaum MJ: Insulin resistance and a diabetes mellitus-like syndrome in mice lacking the protein kinase Akt2 (PKB β). *Science* 292:1728–1731, 2001
34. Garofalo RS, Orena SJ, Pafidi K, Torchia AJ, Stock JL, Hildebrandt AL, Coskran T, Black SC, Brees DJ, Wicks JR, McNeish JD, Coleman KG: Severe diabetes, age-dependent loss of adipose tissue, and mild growth deficiency in mice lacking Akt2/PKB β . *J Clin Invest* 112:197–208, 2003
35. Aikawa R, Nawano M, Gu Y, Katagiri H, Asano T, Zhu W, Nagai R, Komuro I: Insulin prevents cardiomyocytes from oxidative stress-induced apoptosis through activation of P13 kinase/Akt. *Circulation* 102:2873–2879, 2000
36. Cohen P, Frame S: The renaissance of GSK3. *Nat Rev Mol Cell Biol* 2:769–776, 2001
37. Pap M, Cooper GM: Role of glycogen synthase kinase-3 in the phosphatidylinositol-3-kinase/Akt cell survival pathway. *J Biol Chem* 273:19929–19932, 1998
38. Hoeflich KP, Luo J, Rubie EA, Tsao MS, Jin O, Woodgett JR: Requirement for glycogen synthase kinase-3 β in cell survival and NF- κ B activation. *Nature* 406:86–90, 2000
39. Tong H, Imahashi K, Steenbergen C, Murphy E: Phosphorylation of glycogen synthase kinase-3 β during preconditioning through a phosphatidylinositol-3-kinase-dependent pathway is cardioprotection. *Circ Res* 90:377–379, 2002
40. Brunet A, Bonni A, Zigmond MJ, Lin MZ, Juo P, Hu LS, Anderson MJ, Arden KC, Blenis J, Greenberg ME: Akt promotes cell survival by phosphorylating and inhibiting a Forkhead transcription factor. *Cell* 96:857–868, 1999