

SLC12A3 (Solute Carrier Family 12 Member [Sodium/Chloride] 3) Polymorphisms Are Associated With End-Stage Renal Disease in Diabetic Nephropathy

Jae Hyeon Kim,^{1,2} Hyoung Doo Shin,³ Byung Lae Park,³ Min Kyong Moon,^{1,2} Young Min Cho,^{3,4} Young Hwan Hwang,⁴ Kook Whan Oh,⁴ Seong Yeon Kim,⁴ Hong Kyu Lee,⁴ Curie Ahn,⁴ and Kyong Soo Park^{3,4}

Diabetic nephropathy is the most common cause of end-stage renal disease (ESRD). Genetic susceptibility plays an important role in the development and progression of diabetic nephropathy. Previous studies have revealed that polymorphisms in the *SLC12A3* (solute carrier family 12 member [sodium/chloride] 3) gene, which encodes solute carrier family 12 member 3, might contribute to genetic susceptibility to diabetic nephropathy and essential hypertension. In this study, we examined whether the *SLC12A3* gene locus is associated with ESRD resulting from diabetic nephropathy. We genotyped 11 common single nucleotide polymorphisms (SNPs) in the *SLC12A3* gene in 177 patients with ESRD due to type 2 diabetes and 184 patients with diabetic retinopathy but with no signs of renal involvement. Three SNPs (g.34372G>A [Arg913Gln], g.39143G>A, and g.41727C>T) were found to be associated with ESRD due to diabetic nephropathy. These three SNPs were in complete linkage disequilibrium. Haplotype 4 in block 2 (18806C, 21822C, 34372A, 39143A, 39240T, 39375C, and 41727T) showed a significant association with ESRD due to type 2 diabetes ($P = 0.0028$). These results suggest that the *SLC12A3* gene locus is associated with ESRD due to diabetic nephropathy. *Diabetes* 55:843–848, 2006

Diabetic nephropathy is the most common cause of end-stage renal disease (ESRD) (1,2). Strong evidence has been provided by epidemiological (3) and familial (4,5) studies suggesting that genetic susceptibility plays an important role in the pathogenesis of diabetic nephropathy; however, the causative genes remain elusive. The *SLC12A3* (solute carrier family

12 member [sodium/chloride] 3) gene, at chromosome 16q13, which encodes a thiazide-sensitive sodium chloride cotransporter (MIM600968), has been shown to be a new candidate gene for diabetic nephropathy (6,7) and hypertension (8–10). It was shown that loss-of-function mutations in the *SLC12A3* gene cause Gitelman's syndrome, which is inherited as an autosomal-recessive trait and is characterized by low blood pressure due to renal sodium wasting, hypokalemia, metabolic alkalosis, hypocalciuria, and volume depletion (11). On the other hand, substitution of Arg913 to Gln in the *SLC12A3* gene (Arg904Gln in the previous report), has been reported to predispose essential hypertension in the Swedish and Japanese (8,9). In addition, blood pressure reduction by thiazides was found to be dependent on genetic variations in the *SLC12A3* gene (10). Recently, a genome-wide association study using single nucleotide polymorphisms (SNPs) showed that the gene encoding *SLC12A3* (especially Arg913Gln) may contribute to genetic susceptibility to diabetic nephropathy in the Japanese (6). In this study, we investigated the associations between *SLC12A3* polymorphisms and ESRD due to type 2 diabetic nephropathy in the Korean population.

RESULTS

We studied 361 patients with type 2 diabetes, comprising 177 patients with ESRD resulting from diabetic nephropathy (the ESRD group) and 184 patients with diabetic retinopathy and duration of diabetes of >15 years but without evidence of renal disease (the control group). The clinical characteristics of these groups are summarized in Table 1. Among the identified 52 polymorphisms in the *SLC12A3* gene, as determined by sequencing of 48 Japanese DNA samples (12), we selected four SNPs (g.34372 G>A [Arg913Gln], g.39143 G>A, g.39240 C>T, and g.41727 C>T) that had been previously reported to be associated with diabetic nephropathy (6). We also selected an additional seven common SNPs based on frequency (frequency >0.05 in regulatory region, frequency >0.1 in introns) and linkage disequilibrium (LD) status (12). Two SNPs were in the promoter region, three in exons, and six in introns (Fig. 1A). Haplotype blocks were constructed by using LD patterns among the 11 SNPs genotyped. There was one break in the LD patterns; accordingly, two blocks were found in this study (Fig. 1C). We selected common haplotypes (frequency >0.05) from each block, which accounted for >95% of observed haplotypes (Fig. 1B). The

From the ¹Department of Internal Medicine, Seoul National University Boramae Hospital, Seoul, Korea; the ²Genome Research Center for Diabetes and Endocrine Disease, Clinical Research Institute, Seoul National University Hospital, Seoul, Korea; the ³Department of Genetic Epidemiology, SNP genetics, Seoul, Korea; and the ⁴Department of Internal Medicine, Seoul National University College of Medicine, Seoul, Korea.

Address correspondence and reprint requests to Kyong Soo Park, MD, PhD, or Curie Ahn, MD, PhD, Department of Internal Medicine, Seoul National University College of Medicine, 28 Yongon-Dong Chongno-Gu, Seoul, 110-744, Korea. E-mail: kspark@snu.ac.kr or curie55@naver.com.

Received for publication 8 August 2005 and accepted in revised form 28 November 2005.

C.A. and K.S.P. jointly supervised this project.

ESRD, end-stage renal disease; LD, linkage disequilibrium; SNP, single nucleotide polymorphism.

© 2006 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

TABLE 1
Clinical characteristics of the study subjects


	Control	ESRD	P
n	184	177	
Sex (M/F)	70/114	105/72	<0.001
Age (years)	64 ± 7	61 ± 9	0.002
Duration of diabetes (years)	20 ± 5	18 ± 8	0.006
BMI (kg/m ²)	23.9 ± 2.6	23.2 ± 2.8	0.015
Systolic blood pressure (mmHg)	131 ± 16	150 ± 21	<0.001
Diastolic blood pressure (mmHg)	77 ± 9	83 ± 12	<0.001
Fasting plasma glucose (mmol/l)	8.1 ± 2.4	8.2 ± 3.2	NS
HbA _{1c} (%)	7.7 ± 1.2	7.2 ± 1.5	<0.001
Serum creatinine (μmol/l)	75 ± 12	622 ± 212	<0.001

Data are means ± SD.

genotype distributions of the 11 SNPs were in Hardy-Weinberg equilibrium.

Three individual polymorphisms (g.34372G>A [Arg913Gln], g.39143G>A, and g.41727C>T) of the 11 genotyped SNPs were found to be associated with ESRD caused by diabetic nephropathy (Table 2). Allele frequencies of g.34372G>A (Arg913Gln) and g.41727C>T were found to be significantly different between ESRD patients and control subjects after correction for multiple comparisons (corrected P = 0.033, odds ratio 2.30 [95% CI 1.32–4.00]; corrected P = 0.044, 2.20 [1.27–3.80], respectively; Table 2). These three SNPs were in complete LD (|D'| = 1) and were determinants of haplotype 4 in block 2 (Table 3). An analysis of the frequencies of common haplotypes from each block revealed a significant association between haplotype 4 (18806C, 21822C, 34372A, 39143A, 39240T, 39375C, and 41727T) in block 2 with ESRD due to diabetic nephropathy, even after Bonferroni correction (corrected P = 0.031; Table 3).

A Map of *SLC12A3* (solute carrier family 12 member 3) on chromosome 16q13 (48 kb)



B Haplotype of *SLC12A3*

Block1					Block2									
Hap.	-1991C>A	-950A>G	+1784C>T	+1918A>G	Freq.	Hap.	+18806C>T	+21822C>T	+34372G>A	+39143G>A	+39240C>T	+39375C>T	+41727C>T	Freq.
BL1-ht1	C	G	C	A	0.422	BL2-ht1	T	C	G	G	C	C	C	0.428
BL1-ht2	C	G	C	G	0.226	BL2-ht2	C	T	G	G	C	C	C	0.223
BL1-ht3	C	A	T	A	0.110	BL2-ht3	C	C	G	G	C	T	C	0.180
BL1-ht4	C	A	C	A	0.088	BL2-ht4	C	C	A	A	T	C	T	0.087
BL1-ht5	A	G	C	A	0.087	BL2-ht5	C	C	G	G	T	T	C	0.051
BL1-ht6	C	A	C	G	0.068	BL2-ht6	C	C	G	G	C	C	C	0.017

C LDs among SNPs in *SLC12A3* visualized by Haploview




FIG. 1. Gene maps (A), haplotypes (B), and LD patterns (C) of the *SLC12A3* gene. A: Polymorphisms genotyped in the *SLC12A3* gene. Coding exons are marked by shaded blocks. *SNPs that have been previously associated with diabetic nephropathy (6). The first nucleotide of the transcription start site is denoted as nucleotide plus one (+1; reference sequence of *SLC12A3*: NM_000339). B: Common haplotypes of *SLC12A3* gene. Haplotypes and their frequencies were analyzed by the algorithm developed by Schaid et al. (haplo.score) (21). C: LD among SNPs in *SLC12A3* visualized by Haploview. There was one break in the LD patterns; accordingly, two blocks were found in this study.

TABLE 2
Associations between *SLC12A3* SNPs and ESRD due to diabetic nephropathy

g.-1991C>A	CC	CA	AA	<i>P</i>	C	A	<i>P</i>	OR (95% CI)
ESRD	154 (87.0)	22 (12.4)	1 (0.5)	0.182	0.93	0.07	0.149	0.63 (0.37–1.08)
Control	147 (79.9)	36 (19.6)	1 (0.5)		0.90	0.10		
g.-950A>G	AA	AG	GG	<i>P</i>	A	G	<i>P</i>	OR (95% CI)
ESRD	85 (48.6)	74 (42.3)	16 (9.1)	0.077	0.70	0.30	0.023*	1.47 (1.05–2.06)
Control	108 (60.0)	62 (34.3)	10 (5.6)		0.77	0.23		
g.1784C>T	CC	CT	TT	<i>P</i>	C	T	<i>P</i>	OR (95% CI)
ESRD	139 (78.5)	34 (19.2)	4 (2.3)	0.825	0.88	0.12	0.527	1.16 (0.73–1.85)
Control	148 (80.9)	32 (17.5)	3 (1.6)		0.90	0.10		
g.1918A>G	AA	AG	GG	<i>P</i>	A	G	<i>P</i>	OR (95% CI)
ESRD	91 (51.7)	67 (38.1)	18 (10.2)	0.197	0.71	0.29	0.916	0.98 (0.71–1.35)
Control	87 (47.3)	85 (46.2)	12 (6.5)		0.70	0.30		
g.18806C>T	CC	CT	TT	<i>P</i>	C	T	<i>P</i>	OR (95% CI)
ESRD	69 (39.2)	76 (43.2)	31 (17.6)	0.129	0.61	0.39	0.042*	0.74 (0.55–0.99)
Control	54 (29.5)	87 (47.5)	42 (23.0)		0.53	0.47		
g.21822C>T	CC	CT	TT	<i>P</i>	C	T	<i>P</i>	OR (95% CI)
ESRD	98 (56.0)	71 (40.6)	6 (3.4)	0.464	0.76	0.24	0.369	1.18 (0.83–1.67)
Control	114 (62.0)	63 (34.2)	7 (3.8)		0.79	0.21		
g.34372G>A	GG	GA	AA	<i>P</i>	G	A	<i>P</i>	OR (95% CI)
ESRD	136 (77.7)	37 (21.1)	2 (1.1)	0.009*	0.88	0.12	0.003	2.30 (1.32–4.00)
Control	164 (89.6)	18 (9.8)	1 (0.5)		0.95	0.05		
g.39143G>A	GG	GA	AA	<i>P</i>	G	A	<i>P</i>	OR (95% CI)
ESRD	137 (77.4)	34 (19.2)	6 (3.4)	0.017*	0.87	0.13	0.005*	2.05 (1.23–3.41)
Control	163 (88.6)	17 (9.2)	4 (2.2)		0.93	0.07		
g.39240C>T	CC	CT	TT	<i>P</i>	C	T	<i>P</i>	OR (95% CI)
ESRD	124 (71.7)	47 (27.2)	2 (1.2)	0.350	0.85	0.15	0.462	1.17 (0.77–1.80)
Control	140 (76.5)	39 (21.3)	4 (2.2)		0.87	0.13		
g.39375C>T	CC	CT	TT	<i>P</i>	C	T	<i>P</i>	OR (95% CI)
ESRD	98 (56.3)	70 (40.2)	6 (3.4)	0.793	0.76	0.24	0.716	0.94 (0.67–1.32)
Control	102 (55.4)	73 (39.7)	9 (4.9)		0.75	0.25		
g.41727C>T	CC	CT	TT	<i>P</i>	C	T	<i>P</i>	OR (95% CI)
ESRD	137 (77.4)	38 (21.5)	2 (1.1)	0.013*	0.88	0.12	0.004	2.20 (1.27–3.80)
Control	162 (89.0)	19 (10.4)	1 (0.5)		0.94	0.06		

Genotype distributions are shown as *n* (%). *P* values, odds ratios (ORs), and 95% CIs were calculated using the χ^2 test. **P* values were not significant after correction for multiple comparisons (significant *P* < 0.0045).

DISCUSSION

In this study, we found that genetic variations of the *SLC12A3* gene are associated with ESRD resulting from diabetic nephropathy in the Korean population.

We found that three SNPs (g.34372G>A [Arg913Gln], g.39143G>A, and g.41727C>T) and haplotype 4 (18806C, 21822C, 34372A, 39143A, 39240T, 39375C, and 41727T) in block 2 were significantly associated with ESRD caused by diabetic nephropathy. The variants of three SNPs (34372A, 39143A, and 41727T) were in complete LD and were determinants of haplotype 4 in block 2 (Table 3). Association between haplotype 4 in block 2 and ESRD is more significant than each of three SNPs (g.34372G>A [Arg913Gln], g.39143G>A, and g.41727C>T).

Our results are consistent with the previous observation that the substitution of Arg913 to Gln in the *SLC12A3* gene was reported to increase the risk of developing essential hypertension (8,9), considering elevated blood pressure is one of the major risk factors for the development and progression of diabetic nephropathy (13).

However, our results are quite opposite to Japanese data showing that the substitution of Arg913 to Gln is

associated with reduced risk of development of diabetic nephropathy (6,7). Although overall frequencies of four SNPs (g.34372 G>A [Arg913Gln], g.39143 G>A, g.39240 C>T, and g.41727 C>T), which had been previously reported to be associated with diabetic nephropathy, in our control group (5, 7, 13, and 6%, respectively) were similar to those of Japanese (8, 9, 12, and 8%, respectively), there were significant differences in the nephropathy group between two studies (12 vs. 3%, 13 vs. 4%, 15 vs. 7%, and 12 vs. 3%, respectively) (6). In addition, the frequency of haplotype 4 in block 2 in the control group (5.4%) was also similar to that of Japanese (6.0%, haplotype 7 + haplotype 9), whereas it was different in the nephropathy group between two studies (11.9 vs 0.9%) (6). At this time, the explanation for the conflicting results between the two studies is unclear, although we selected patients with ESRD due to diabetic nephropathy for nephropathy cases, whereas patients with overt proteinuria and patients undergoing renal replacement therapy were included in the previous report (6).

The use of prevalent ESRD patients as cases may lead to strong survival bias. However, after dividing the patients with

TABLE 3
Association between SLC12A3 haplotype blocks and end-stage renal disease due to diabetic nephropathy

Block 1 Haplotype	Locus						Number (% Frequency [95% CI])	P	OR (95% CI)
	-1991	-950	1784	1918	ESRD	Control			
BL1-ht1	C	G	C	A	156 (44.1 [38.9-49.3])	165 (44.8 [39.7-49.9])	0.835	0.97 (0.73-1.30)	
BL1-ht2	C	G	C	G	66 (18.6 [14.5-22.7])	81 (22.0 [17.8-26.2])	0.352	0.84 (0.59-1.21)	
BL1-ht3	C	A	T	A	42 (11.9 [8.5-15.3])	38 (10.3 [7.2-13.4])	0.51	1.17 (0.73-1.86)	
BL1-ht4	C	A	C	G	36 (10.2 [7.0-13.4])	28 (7.6 [4.9-10.3])	0.226	1.38 (0.82-2.31)	
BL1-ht5	A	G	C	A	24 (6.8 [4.2-9.4])	38 (10.3 [7.2-13.4])	0.089	0.63 (0.37-1.08)	
BL1-ht6	C	A	C	A	28 (7.9 [5.1-10.7])	18 (4.9 [2.7-7.1])	0.097	1.67 (0.91-3.08)	

Block 2 Haplotype	Locus						Number (% Frequency [95% CI])	P	OR (95% CI)	
	18806	21822	34372	39143	39240	39375				41727
BL2-ht1	T	C	G	G	C	C	139 (39.3 [34.2-44.4])	170 (46.2 [41.0-51.4])	0.06	0.75 (0.56-1.01)
BL2-ht2	C	T	G	G	C	C	86 (24.3 [19.8-28.8])	76 (20.7 [16.5-24.9])	0.241	1.23 (0.87-1.75)
BL2-ht3	C	C	G	G	C	T	67 (18.9 [14.8-23.0])	62 (16.8 [12.9-20.7])	0.466	1.15 (0.79-1.69)
BL2-ht4	C	C	A	A	T	C	42 (11.9 [8.5-15.3])	20 (5.4 [3.0-7.8])	0.0028	2.34 (1.35-4.08)
BL2-ht5	C	C	G	G	T	T	12 (3.4 [1.5-5.3])	25 (6.8 [4.2-9.4])	0.05*	0.48 (0.24-0.97)

The haplotype frequencies are estimates made according to unphased data and are not based on actual haplotype counts. Each haplotype with a frequency >0.05 is shown. P values of haplotype associations were analyzed by the algorithm developed by Schaid et al. (haplo.score) (21). *P values were not significant after correction for multiple comparisons (significant P < 0.0045). Shaded alleles revealed rare alleles in their loci.

ESRD into quartiles by duration of ESRD, we could not find any association of *SLC12A3* with the duration of ESRD (data not shown). Therefore, it is less likely that there is a spurious association due to survival bias between *SLC12A3* and ESRD attributed to diabetic nephropathy.

At the moment, functional significance of substitution of Arg913 to Gln is unclear, although there was a suggestion that the Arg913Gln in the *SLC12A3* gene might be a gain-of-function polymorphism and increase the risk of developing essential hypertension (9). Nevertheless, in this study, we found no association between Arg913Gln and blood pressure in either control subjects or ESRD cases (data not shown). Hence, further biological and/or functional evidence would be needed to confirm the suggestive association of *SLC12A3* polymorphisms with ESRD resulting from diabetic nephropathy that has been reported in this study.

In conclusion, we found that SNPs and haplotypes of the *SLC12A3* gene, especially Arg913Gln, are significantly associated with ESRD caused by diabetic nephropathy in the Korean population.

RESEARCH DESIGN AND METHODS

We studied 361 unrelated patients with type 2 diabetes comprising two groups according to the following criteria: 1) the control group ($n = 184$): patients with diabetic retinopathy and a duration of diabetes >15 years but with no sign of renal involvement, i.e., a urinary albumin-to-creatinine ratio <30 mg/g and with a creatinine clearance (using the Cockcroft equation [14]) of >60 ml/min per m^2 ; or 2) the ESRD group ($n = 177$): patients with diabetic retinopathy and ESRD due to type 2 diabetes, as indicated by a creatinine clearance rate of <15 ml/min per m^2 or being under renal replacement therapy. We excluded subjects with ESRD with any one of the following criteria: 1) no retinopathy, 2) a consistent hematuria history before renal function deterioration, 3) no history of proteinuria before renal function deterioration, 4) reduced kidney size, and 5) evidence of other systemic or primary glomerular diseases. The patients with ESRD were recruited from four university hospitals and three dialysis clinics, and the control subjects were recruited from two hospital clinics from June 2003 to July 2004. All subjects enrolled in this study were ethnic Koreans. Type 2 diabetes was diagnosed using World Health Organization criteria (15). Subjects with positive GAD antibodies and an age at diagnosis of type 2 diabetes of <30 years, as well as those who started insulin treatment within 1 year of diagnosis and were ketosis prone, were excluded. The institutional review board of the Clinical Research Institute at Seoul National University Hospital approved the study protocol, and informed consent for genetic analysis was obtained from each subject.

Genotyping for SNPs in the *SLC12A3* gene. Since there is a close genetic relationship between Koreans and Japanese (16,17), we used Japanese sequencing data to select common polymorphisms in the *SLC12A3* gene (12). We selected four SNPs (g.34372 G>A [Arg913Gln], g.39143 G>A, g.39240 C>T, and g.41727 C>T) that had been previously reported to be associated with diabetic nephropathy (6), and we also selected seven additional common SNPs based on frequency (frequency >0.05 in regulatory region, frequency >0.1 in introns) and LD status ($r^2 > 0.5$) among the identified polymorphisms in the *SLC12A3* gene, as determined by sequencing of 48 Japanese DNA samples (12) (Fig. 1A). The 11 polymorphisms were genotyped by fluorescence polarization detection or single-base extension (18). For genotyping polymorphic sites, amplifying primers and probes were designed for TaqMan (19). Primer sets for the amplification and sequencing analysis of *SLC12A3* gene were designed based on GenBank sequences (ref. genome seq. NC_000016, released on 29 December 2004). Information regarding the primers used is available on our website (<http://www.snp-genetics.com/reference/SLC12A3.doc>). The TaqMan method and single-base extension both achieved a >98% success rate for genotype.

Statistics. Differences in genotype frequencies were compared using the χ^2 test. We used Bonferroni correction for the adjustment of multiple comparisons (11 independent tests in genotype and 11 independent tests in haplotype). For association analyses between SNPs in *SLC12A3* and blood pressure, we adjusted for age, sex, and BMI using a general linear regression procedure. Haplotype structures were visualized by Haploview software (20). Haplotype frequencies and *P* values of haplotype associa-

tions were analyzed by the algorithm developed by Schaid et al. (haplo.score) (21). A *P* value <0.05 was considered statistically significant.

ACKNOWLEDGMENTS

This work was supported by a grant from the Korean Health 21 R & D Project, Korean Ministry of Health & Welfare (00-PJ3-PG6-GN07-001).

We thank Drs. Sung Kyun Kim, Woo Kyung Jung, Ji-Eun Oh, Eung Taek Kang, Joong Geon Lee, and Jin Cheol Kim for help in recruiting study subjects.

REFERENCES

1. US Renal Data System: USRDS 2004 Annual Data Report. *Am J Kidney Dis* 45 (Suppl. 1):S8–S280, 2005
2. Kim SY, Jin DC, Bang BK: Current status of dialytic therapy in Korea. *Nephrology (Carlton)* 8 (Suppl.):S2–S9, 2003
3. Krolewski AS, Warram JH, Christlieb AR, Busick EJ, Kahn CR: The changing natural history of nephropathy in type 1 diabetes. *Am J Med* 87:785–794, 1985
4. Seaquist ER, Goetz FC, Rich S, Barbosa J: Familial clustering of diabetic kidney disease: evidence for genetic susceptibility to diabetic nephropathy. *N Engl J Med* 320:1161–1165, 1989
5. Borch-Johnsen K, Norgaard K, Hommel E, Mathiesen ER, Jensen JS, Deckert T, Parving H-H: Is diabetic nephropathy an inherited complication? *Kidney Int* 41:719–722, 1992
6. Tanaka N, Babazono T, Saito S, Sekine A, Tsunoda T, Haneda M, Tanaka Y, Fujioka T, Kaku K, Kawamori R, Kikkawa R, Iwamoto Y, Nakamura Y, Maeda S: Association of solute carrier family 12 (sodium/chloride) member 3 with diabetic nephropathy, identified by genome-wide analyses of single nucleotide polymorphisms. *Diabetes* 52:2848–2853, 2003
7. Nishiyama K, Tanaka Y, Nakajima K, Mokubo A, Atsumi Y, Matsuoka K, Watada H, Hirose T, Nomiya T, Maeda S, Kawamori R: Polymorphism of the solute carrier family 12 (sodium/chloride transporters) member 3, *SLC12A3*, gene at exon 23 (+78G/A: Arg913Gln) is associated with elevation of urinary albumin excretion in Japanese patients with type 2 diabetes: a 10-year longitudinal study. *Diabetologia* 48:1335–1338, 2005
8. Melander O, Orho-Melander M, Bengtsson K, Lindblad U, Rastam L, Groop L, Hulthen UL: Genetic variants of thiazide-sensitive NaCl-cotransporter in Gitelman's syndrome and primary hypertension. *Hypertension* 36:389–394, 2000
9. Matsuo A, Katsuya T, Ishikawa K, Sugimoto K, Iwashima Y, Yamamoto K, Ohishi M, Rakugi H, Ogihara T: G2736A polymorphism of thiazide-sensitive Na-Cl cotransporter gene predisposes to hypertension in young women. *J Hypertens* 22:2123–2127, 2004
10. Matayoshi T, Kamide K, Takiuchi S, Yoshii M, Miwa Y, Takami Y, Tanaka C, Banno M, Horio T, Nakamura S, Nakahama H, Yoshihara F, Inenaga T, Miyata T, Kawano Y: The thiazide-sensitive Na(+)-Cl(-) cotransporter gene, C1784T, and adrenergic receptor-beta3 gene, T727C, may be gene polymorphisms susceptible to the antihypertensive effect of thiazide diuretics. *Hypertens Res* 27:821–833, 2004
11. Simon DB, Nelson-Williams C, Bia MJ, Ellison D, Karet FE, Molina AM, Vaara I, Iwata F, Cushner HM, Koolen M, Gainza FJ, Gitelman HJ, Lifton RP: Gitelman's variant of Bartter's syndrome, inherited hypokalaemic alkalosis, is caused by mutations in the thiazide-sensitive Na-Cl cotransporter. *Nat Genet* 12:24–30, 1996
12. Kokubo Y, Kamide K, Inamoto N, Tanaka C, Banno M, Takiuchi S, Kawano Y, Tomoike H, Miyata T: Identification of 108 SNPs in TSC, WNK1, and WNK4 and their association with hypertension in a Japanese general population. *J Hum Genet* 49:507–515, 2004
13. Mogensen CE: Combined high blood pressure and glucose in type 2 diabetes: double jeopardy: British trial shows clear effects of treatment, especially blood pressure reduction. *BMJ* 317:693–694, 1998
14. Durakovic Z: Creatinine clearance in the elderly: a comparison of direct measurement and calculation from serum creatinine. *Nephron* 44:66–69, 1986
15. Alberti KG, Zimmet PZ: Definition, diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabet Med* 15:539–553, 1998
16. Rolf B, Horst B, Eigel A, Sagansermisri T, Brinkmann B, Horst J: Microsatellite profiles reveal an unexpected genetic relationship between Asian populations. *Hum Genet* 102:647–652, 1998
17. Miller RV, Phillips MS, Jo I, Donaldson MA, Studebaker JF, Addleman N, Alfisi SV, Ankener WM, Bhatti HA, Callahan CE, Carey BJ, Conley CL, Cyr JM, Derohannessian V, Donaldson RA, Elosua C, Ford SE, Forman AM, Gelfand CA, Grecco NM, Gutendorf SM, Hock CR, Hozza MJ, Hur S, In SM,

- Jackson DL, Jo SA, Jung SC, Kim S, Kimm K, Kloss EF, Koboldt DC, Kuebler JM, Kuo FS, Lathrop JA, Lee JK, Leis KL, Livingston SA, Lovins EG, Lundy ML, Maggan S, Minton M, Mockler MA, Morris DW, Nachtman EP, Oh B, Park C, Park CW, Pavelka N, Perkins AB, Restine SL, Sachidanandam R, Reinhart AJ, Scott KE, Shah GJ, Tate JM, Varde SA, Walters A, White JR, Yoo YK, Lee JE, Boyce-Jacino MT, Kwok PY, the SNP Consortium Allele Frequency Project: High-density single-nucleotide polymorphism maps of the human genome. *Genomics* 86:117–126, 2005
18. Vreeland WN, Meagher RJ, Barron AE: Multiplexed, high-throughput genotyping by single-base extension and end-labeled free-solution electrophoresis. *Anal Chem* 74:4328–4333, 2002
19. Livak KJ: Allelic discrimination using fluorogenic probes and the 5' nuclease assay. *Genet Anal* 14:143–149, 1999
20. Barrett JC, Fry B, Maller J, Daly MJ: Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics* 21:263–265, 2005
21. Schaid DJ, Rowland CM, Tines DE, Jacobson RM, Poland GA: Score tests for association of traits with haplotypes when linkage phase is ambiguous. *Am J Hum Genet* 70:425–434, 2002