

Paradoxical Changes in Muscle Gene Expression in Insulin-Resistant Subjects After Sustained Reduction in Plasma Free Fatty Acid Concentration

Mandeep Bajaj,^{1,2} Rafael Medina-Navarro,³ Swangjit Suraamornkul,² Christian Meyer,^{2,4} Ralph A. DeFronzo,² and Lawrence J. Mandarino³

Lipid oversupply plays a role in developing insulin resistance in skeletal muscle, decreasing expression of nuclear-encoded mitochondrial genes, and increasing extracellular matrix remodeling. To determine if a decrease in plasma lipid content reverses these abnormalities, insulin-resistant subjects with a family history of type 2 diabetes had euglycemic clamps and muscle biopsies before and after acipimox treatment to suppress free fatty acids. Free fatty acids fell from 0.584 ± 0.041 to 0.252 ± 0.053 mmol/l ($P < 0.001$) and glucose disposal increased from 5.28 ± 0.46 to 6.31 ± 0.55 mg · kg⁻¹ · min⁻¹ ($P < 0.05$) after acipimox; intramuscular fatty acyl CoA decreased from 10.3 ± 1.9 to 4.54 ± 0.82 pmol/mg muscle ($P < 0.01$). Paradoxically, expression of PGC-1 and nuclear-encoded mitochondrial genes decreased after acipimox, and expression of collagens I and III α -subunits (82- and 21-fold increase, respectively, $P < 0.05$), connective tissue growth factor (2.5-fold increase, $P < 0.001$), and transforming growth factor- β 1 increased (2.95-fold increase, $P < 0.05$). Therefore, a reduction in lipid supply does not completely reverse the molecular changes associated with lipid oversupply in muscle. Changes in expression of nuclear-encoded mitochondrial genes do not always correlate with changes in insulin sensitivity. *Diabetes* 56:743–752, 2007

Lipid oversupply leading to tissue lipid accumulation has received significant attention as a potential etiological factor in the development of insulin resistance in skeletal muscle. This excess of lipids either can derive from the diet (1), a decreased ability of muscle to use lipid (2,3), or both. Lipid oversupply leads to an increase in intramyocellular lipid in the form of triglycerides, diacylglycerol, fatty acyl CoA, and ceramides (4–12). The latter three can interfere directly or

indirectly with various aspects of insulin signaling (9–19), thus producing insulin resistance. Diacylglycerol and fatty acyl CoAs can activate protein kinase C isoforms that produce insulin resistance by means of serine phosphorylation of insulin receptor substrate 1 (9,14). Ceramides activate phosphatases that inactivate Akt and also may activate protein kinase C (15–18).

Mitochondrial dysfunction, which is thought to be involved in the pathogenesis of insulin resistance (20,21), is characterized by a coordinated pattern of decreased nuclear-encoded mitochondrial gene expression, potentially driven by reduced expression of the transcriptional coactivator PGC-1 (22,23). We observed an inverse correlation between plasma free fatty acids (FFAs) and PGC-1 mRNA (L.J.M., Mary-Elizabeth Patti, unpublished data), suggesting that lipid oversupply itself might result in suppression of nuclear-encoded mitochondrial gene expression and mitochondrial dysfunction (23). Experimental lipid oversupply produced by an infusion of a triglyceride emulsion is well known to produce insulin resistance (24–26) accompanied by decreased insulin signaling and glycogen formation. Therefore, we further showed that lipid oversupply brought about by a lipid infusion also suppresses PGC-1 mRNA and produces a decrease in nuclear-encoded mitochondrial gene expression (27). In light of these data, the first purpose of the present study was to determine whether an experimental decrease in plasma FFAs and tissue lipid content in insulin-resistant subjects would improve PGC-1 and nuclear-encoded mitochondrial gene expression in skeletal muscle.

In our previous study, which was designed to determine whether experimental lipid oversupply alters nuclear-encoded mitochondrial gene expression, we showed that lipid infusion produced profound increases in the mRNA and protein expression of a variety of extracellular matrix proteins (27). Subsequently, we went on to show that collagen expression was increased in skeletal muscle from insulin-resistant obese nondiabetic and type 2 diabetic individuals (28). These changes in gene expression appeared to be driven by an increase in the expression of connective tissue growth factor (CTGF) (CTGF/CCN2), a member of the CCN gene family that mediates fibrotic responses (29). Therefore, the second purpose of this study was to determine whether a decrease in plasma FFAs also induces changes in expression of extracellular matrix genes.

Acipimox, a nicotinic acid analog that reduces plasma FFAs by suppressing lipolysis (30), improves insulin sensitivity (31,32). In the present study, we used acipimox

From the ¹University of Texas Medical Branch, Galveston, Texas; the ²University of Texas Health Science Center, San Antonio, Texas; the ³Center for Metabolic Biology, Arizona State University, Tempe, Arizona; and the ⁴Carl T. Hayden VA Medical Center, Phoenix, Arizona.

Address correspondence and reprint requests to Lawrence J. Mandarino, PhD, Director, Center for Metabolic Biology, Professor and Chair, Department of Kinesiology, Professor, School of Life Sciences, Arizona State University, P.O. Box 874501, Tempe, AZ 85287-4501. E-mail: lawrence.mandarino@asu.edu.

Received for publication 20 June 2006 and accepted in revised form 18 December 2006.

CTGF, connective tissue growth factor; FFA, free fatty acid; OGTT, oral glucose tolerance test; TGF, transforming growth factor.

DOI: 10.2337/db06-0840

© 2007 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

treatment to produce a sustained decrease in plasma FFAs in insulin-resistant individuals who are genetically predisposed to develop type 2 diabetes. Such individuals, who have two or more primary relatives with type 2 diabetes, are characterized by insulin resistance, increased plasma FFAs, and decreased insulin signaling in skeletal muscle (33,34). We hypothesized that in these insulin-resistant subjects, acipimox would decrease plasma FFAs, concomitantly decrease myocellular fatty acyl CoAs while improving insulin sensitivity, and increase expression of PGC-1 α and nuclear-encoded mitochondrial genes. We further hypothesized that reduced FFA levels would result in decreased expression of extracellular matrix genes.

RESEARCH DESIGN AND METHODS

Ten subjects with at least two first-degree relatives with type 2 diabetes took part in the study. Data for in vivo glucose metabolism and plasma FFA and adiponectin concentrations for 8 of the 10 subjects were included in a previous publication (35). None of the subjects had any significant medical problems, and their weight was stable for at least 3 months before the study. No subject was taking any medications known to affect glucose metabolism. None of the subjects participated in any heavy exercise, and they were instructed not to engage in vigorous exercise for at least 3 days before the study. The purpose, nature, and potential risks of the study were explained to all subjects, and written consent was obtained before their participation. The protocol was approved by the Institutional Review Board of the University of Texas Health Science Center at San Antonio.

Study design. Three weeks before study, all subjects met with a dietitian and were instructed to consume a weight-maintaining diet containing 50% carbohydrate, 30% fat (10% saturated fat, 10% polyunsaturated fat, and 10% monounsaturated fat), and 20% protein. During the week before the start of acipimox treatment, all subjects received 1) a 75-g oral glucose tolerance test (OGTT) and 2) a euglycemic insulin clamp study (36) in combination with 3-[³H]glucose to examine hepatic and peripheral tissue sensitivity to insulin and with vastus lateralis muscle biopsy for assessment of gene expression changes and fatty acyl CoA concentrations (see below). The euglycemic insulin clamp study was started at 0600 h and the OGTT at 0800 h after a 10- to 12-h overnight fast.

After completion of these studies, subjects were started on acipimox, 250 mg orally every 6 h (0600, 1200, 1800, and 2400 h) for 7 days. The last dose of acipimox was administered at 0600 h on day 8. At 0600 h on day 5, all subjects were admitted to the General Clinical Research Center, where they remained until completion of the study on day 8. While in the General Clinical Research Center, subjects received a weight-maintaining diet of the same composition and were encouraged to ambulate freely. There was no change in body weight in any subject between the pre- and post-acipimox studies. The OGTT was repeated on day 7 and the euglycemic-hyperinsulinemic clamp/vastus lateralis muscle biopsy study was repeated on day 8 after a 10- to 12-h overnight fast. Acipimox (Olbetam, Pharmacia Upjohn, Milan, Italy) was used under investigational new drug no. 60,687 (to R.A.D.).

OGTT. After obtaining baseline blood samples for determination of plasma glucose, adiponectin, and FFAs, subjects underwent a 75-g OGTT.

Hyperinsulinemic-euglycemic clamp. Insulin sensitivity was assessed with a euglycemic insulin clamp, as previously described (36), using an insulin infusion rate of 80 mU \cdot min⁻¹ \cdot m⁻² body surface area, with infusion of 3-[³H]glucose for measuring glucose disposal and endogenous glucose production. A needle biopsy of the vastus lateralis was obtained under local anesthesia in the fasted condition, immediately before the hyperinsulinemic-euglycemic clamp. Thirty minutes after the start of insulin infusion, a second percutaneous muscle biopsy was obtained from the contralateral vastus lateralis muscle. After acipimox treatment, biopsies were performed in contralateral legs at sites several centimeters distant to the pretreatment biopsy sites to avoid any effects of the biopsy procedure itself on the tissue. Continuous indirect calorimetry, using a ventilated hood system (Deltatrac II; Sensor Medics, Yorba Linda, CA), was performed during the last 40 min of the basal period and during the last 30 min of the insulin clamp.

Analytical determinations. Plasma glucose was measured by the glucose oxidase method (Beckman Instruments, Fullerton, CA). Plasma insulin concentration was measured by radioimmunoassay (Diagnostic Products, Los Angeles, CA). Tritiated glucose specific activity was determined on deproteinized plasma samples as previously described (22). Plasma FFA concentration was determined by an enzymatic calorimetric quantification method (Wako Chemicals, Nuess, Germany). Plasma adiponectin concentration was measured by radioimmunoassay (Linco Research, St. Charles, MO).

Fatty acyl CoA analysis. Eight of the 10 volunteers had sufficient biopsy material for assay of long-chain fatty acyl CoA concentrations in addition to mRNA assay. Long-chain acyl CoA standards were purchased from Avanti (Polar lipids; Alabaster, AL). Acetonitrile was from Burdick and Jackson (Muskegon, MI). Other chemicals and solvents were from Sigma (St. Louis, MO).

Sample extraction. Frozen muscle specimens (30–100 mg wet weight) were successively homogenized together with a C12 internal standard in a glass-glass homogenizer using 2 ml of 100 mmol/l KH₂PO₄, pH 4.6, 2 ml of 2-propanol, and 4 ml acetonitrile. Saturated aqueous ammonium sulfate (200 μ l) was added and the sample was re-homogenized. The resulting suspension was vortexed for 5 min and then centrifuged for 5 min at 1,900g. The supernatant was kept on ice, while the precipitate was re-homogenized using one-half volumes of the above solvents. The two resulting supernatants were combined, and 5% trichloroacetic acid was added to 40 ml. After periodic re-homogenization of the cold extracts for 5 min, the samples were centrifuged for 1 h at 3,000g at 4°C. Elution solvent (300 μ l; 70% buffer A:30% acetonitrile, see below) was added, and the supernatant was transferred to small tubes. The pH of the resulting extract was between 5 and 6. Samples were injected on the high-performance liquid chromatography (see below) directly. In the present work, aliquots of 100 μ l were injected immediately after addition of 2 μ l of 1.5 mmol/l EDTA, which reduced and stabilized the baseline.

High-performance liquid chromatography. All procedures were carried out on a Thermo Finnigan Surveyor high-performance liquid chromatography System coupled with a Photodiode Array Detector and controlled by Xcalibur Software 1.3 (Bellefonte, PA). The separation was carried out using a Waters SunFire C18 3.5 μ m, 2.1 \times 100 mm column (Waters, Milford, MA) and a C18 10-mm precolumn (Waters). Chromatography was performed using a gradient system including two mobile phases: 1) 10 mmol/l ammonium acetate/10 mmol/l potassium phosphate, pH 5.8, and 2) 100% acetonitrile. The starting conditions were 70% A:30% B, with a gradient to 60% A:40% B over 5 min; 54% A:46% B at 14 min; 38% A:62% B at 19 min; and finally back down to 70% A:30% B after 30 min. The flow rate was variable, starting at 0.3 ml/min and reaching a flow of 0.5 ml/min at 14 min and 0.6 ml/min at 19 min, where it was sustained until 30 min. Thereafter, it was reduced to 0.3 ml/min. Separation occurred with a column temperature of 40°C. The program was a modification of the procedure original of Corkey et al. (37).

Gene expression analysis. RNA was isolated from muscle biopsies using a commercial kit (RNA STAT-60 Total RNA/mRNA Isolation Reagent; Tel Test, Friendswood, TX) and purified before use (RNeasy Mini kit; Qiagen, Valencia, CA). Quantitative real-time PCR was used to quantify mRNA. The specific genes analyzed and the primers used for this analysis are given in Table 1. A Bio-Rad ICycler (Bio-Rad Laboratories, Hercules, CA) was used to quantify mRNA levels relative to the average value for β -actin, GAPDH, and cyclophilin as controls. Three genes were chosen as controls to minimize variability. Relative changes in mRNA levels were analyzed by the $\Delta\Delta$ Ct method (38).

Calculations. Under steady-state postabsorptive conditions, the rate of endogenous glucose appearance is calculated as the [3-³H]glucose infusion rate (dpm/min) divided by the steady-state plasma [3-³H]glucose specific activity (dpm/mg). During the euglycemic insulin clamp, the rate of total-body glucose appearance was calculated using Steele's equation (39), using a distribution volume of 250 ml/kg. Endogenous glucose production was calculated by subtracting the exogenous glucose infusion rate from glucose appearance. The rate of insulin-mediated total-body glucose disposal was determined by adding the rate of residual endogenous glucose production to the exogenous glucose infusion rate.

Statistical analysis. The effect of acipimox on muscle fatty acyl CoA concentrations was assessed by paired *t* test. The effect of acipimox on mRNA expression was expressed as fold change compared with pretreatment values, and statistical significance compared with no change (value of 1.0) was assessed by the *z* test. Correlation analysis was performed by the Pearson product-moment method.

RESULTS

Subject characteristics and OGTT. Subject characteristics are given in Table 2. All subjects had normal glucose tolerance and had at least two primary relatives with type 2 diabetes. Subjects were only mildly overweight, with an average BMI of 26.5 \pm 0.8 kg/m². After acipimox treatment, the baseline fasting plasma FFA concentration was reduced from 0.584 \pm 0.041 to 0.252 \pm 0.053 mmol/l (*P* < 0.01). Fasting plasma insulin, glucose, and adiponectin concentrations did not significantly change after acipimox

TABLE 1
Primers used in quantitative real-time PCR

CTGF	Forward: TAGCCTCAATTTCTGAACAC Reverse: TTGACGGACTGTCATTCTAT
CPT1 β (64)	Forward: GAGGCCTCAATGACCAGAATG Reverse: GTGGACTCGCTGGTACAGGAA
Citrate synthase (64)	Forward: TTGGCTGACCTGATACCTAAGG Reverse: CACCATACATCATGTCCACAGTG
FN (65)	Forward: CCTCAATTGTTGTTGCTGGA Reverse: GCTACCTTCTACTGATGGCGAATAG
COL1 α 1 (65)	Forward: CCTCAAGGGCTCCAACGAG Reverse: TCAATCACTGTCTTGCCCCA
COL3 α 1	Forward: TCTCCTGGAAGATTGGTTTA Reverse: CACCATAACATGCGTCTTTA
TGF β 1 (65)	Forward: GCGTGCTAATGGTGGAACC Reverse: CGGAGCTCTGATGTGTTGAAGA
PORIN	Forward: CAGTGCCAAATCAAAGCTGA Reverse: CCTGATGTCCAAGCAAGGTT
COX6	Forward: AAGGACGTTGGTGTGAGGT Reverse: TTTCTTTGATCAGCCACAG
COX7	Forward: GTTTGTACTTTGGCTATGCATT Reverse: TGGCATATGAGTTCTAGTTTGA
NRF1	Forward: TGACATTGGAACAGTGACAT Reverse: AATGCAGTTTCTTCCACCAAT
NRF2	Forward: AAATTGAGATTGATGGAACAGAGAA Reverse: TATGGCCTGGCTTACACATCA
MtTFA	Forward: AGATTCCAAGAAGCTAAGGGTATT Reverse: TTTCAGAGTCAGACAGATTTTCCA
PGC1 α	Forward: TTTCTTTTGGCCATGGAATC Reverse: GAAAGAACCGCTGAACAAGC
PGC1 β	Forward: CTGCATTGATAACAGCCTTA Reverse: AAAGGACAGCAGTTTCAAGT
GAPDH	Forward: AACCTGCCAAATATGATGAC Reverse: TCATACCAGGAAATGAGCTT
β -ACTIN	Forward: AAAGTGGAAACGGTGAAGGTG Reverse: AGAGAAGTGGGGTGGCTTTT
Cyclophilin	Forward: GCCATGGAGCGCTTTGG Reverse: CCACAGTCAGCAATGGTGATC
NADH Fe-S	Forward: TTTACTCCAAGCCTGTTGTT Reverse: CCACATAAATCACCATGACA
NADH α 1	Forward: AATTGGCGACTTCGATATTA Reverse: CAGCACTGAAGTTGTTCAAA
NADH flavo	Forward: TCATGAATAAGCCCTCAGAT Reverse: CACAAACACGTCAAATCAT
NQO1 (64)	Forward: GGAGTAAGAAGGCAGTGCCTTTT Reverse: GGCCAGAGAATGACATTCATGT
SOD2 (66)	Forward: GCTTGTCCAAATCAGGATCCA Reverse: GCGTGCTCCACACATCA
OGG1 (66)	Forward: ATTCCAAGGTGTGCGACTGCT Reverse: GCGGGCGATGTTGTTGTT
GPX1	Forward: AGTTTGGGCATCAGGAGAACG Reverse: CAGGAAGGCGAAGAGAGGGT

CTGF, connective tissue growth factor; CPT1 β , carnitine palmitoyl-transferase 1 β ; CS, citrate synthase; FN, fibronectin; COL1 α 1, collagen I α 1-subunit; COL3 α 1, collagen III α 1; PORIN, porin; COX6, cytochrome C oxidase subunit 6; COX7, cytochrome C oxidase subunit 7; NRF1, nuclear respiratory factor 1; NRF2, nuclear respiratory factor 2; MtTFA, mitochondrial transcription factor A; PGC1 α , peroxisome proliferative-activated receptor- γ coactivator 1 α ; PGC1 β , peroxisome proliferative-activated receptor- γ coactivator 1 β ; GAPDH, glyceraldehyde phosphate dehydrogenase; β -ACTIN, β -actin; NADH Fe-S, NADH dehydrogenase iron-sulfur protein; NADH α 1, NADH dehydrogenase α 1 subunit; NADH flavo, NADH dehydrogenase flavoprotein subunit 1; NQO1, NAD(P)H quinone oxidoreductase; SOD2, superoxide dismutase 2; OGG1, 8-oxoguanine DNA glycosylase; GPX1, glutathione peroxidase 1. Where noted, primers were used as reported in the literature. Otherwise, primers were custom designed or, in the case of GPX1, courtesy of Dr. A. Civitarese.

TABLE 2
Subject characteristics

	Acipimox treatment	
	Before	After
Age (years)	41 \pm 4	—
Sex (F/M)	4/6	—
BMI (kg/m ²)	26.5 \pm 0.8	26.4 \pm 0.8
Fasting insulin (pmol/l)	38 \pm 4	40 \pm 6
Fasting glucose (mmol/l)	5.2 \pm 0.1	5.2 \pm 0.1
Fasting FFAs (mmol/l)	0.584 \pm 0.041	0.252 \pm 0.053§
Clamp FFAs (mmol/l)	0.156 \pm 0.016†	0.125 \pm 0.014†§
Fasting adiponectin (μ g/ml)	7.4 \pm 0.8	7.0 \pm 1.0

Data are means \pm SE. † P < 0.01 basal vs. clamp; § P < 0.01 after vs. before acipimox.

treatment (Table 2). During the OGTT, the mean plasma glucose (7.6 \pm 0.3 vs. 7.0 \pm 0.5, P < 0.01) and FFA (0.288 \pm 0.031 vs. 0.154 \pm 0.02, P < 0.01) concentrations were significantly reduced after acipimox treatment. The mean plasma insulin concentration during the OGTT did not change significantly after acipimox treatment (Table 2).

Effects of acipimox on FFA and insulin action in vivo. To determine how acipimox altered insulin action in vivo, euglycemic clamps (80 mU/m² per min) with indirect calorimetry were performed before and after 7 days of treatment. These data are given in Table 3. The steady-state plasma glucose concentrations during the 120-min euglycemic insulin clamp were similar before and after acipimox treatment (5.1 \pm 0.1 vs. 5.1 \pm 0.1 mmol/l). The steady-state plasma insulin concentrations during the insulin clamp studies also were similar before and after acipimox (870 \pm 53 vs. 838 \pm 65 pmol/l). After acipimox treatment, fasting plasma FFA concentrations (determined before the start of the euglycemic clamp) decreased from 0.584 \pm 0.041 to 0.252 \pm 0.053 mmol/l (P < 0.001 vs. before acipimox). Before acipimox, the plasma FFAs during the euglycemic clamp declined from 0.584 \pm 0.041 to 0.156 \pm 0.016 mmol/l (P < 0.001). After acipimox, FFAs decreased from 0.252 \pm 0.053 to 0.125 \pm 0.014 mmol/l (P < 0.01) during the clamp.

In keeping with the lack of effect of acipimox on fasting plasma glucose, fasting endogenous glucose production was unaffected by acipimox (Table 3). However, whereas before acipimox treatment insulin infusion during the clamp decreased endogenous glucose production to 0.190 \pm 0.073 mg \cdot kg⁻¹ \cdot min⁻¹, after acipimox insulin suppressed endogenous glucose production to 0.013 \pm 0.010 mg \cdot kg⁻¹ \cdot min⁻¹ (P < 0.05 vs. before acipimox). In a similar manner, acipimox increased insulin stimulation of glucose disposal, from 5.28 \pm 0.46 mg \cdot kg⁻¹ \cdot min⁻¹ before to 6.31 \pm 0.55 mg \cdot kg⁻¹ \cdot min⁻¹ (P < 0.01) after acipimox. Acipimox increased the rate of systemic glucose oxidation under basal conditions from 1.33 \pm 0.08 to 1.52 \pm 0.09 mg \cdot kg⁻¹ \cdot min⁻¹ (P < 0.05) and decreased lipid oxidation from 0.77 \pm 0.07 to 0.66 \pm 0.04 mg \cdot kg⁻¹ \cdot min⁻¹ (P < 0.05). Finally, acipimox treatment had no effect on insulin-stimulated glucose oxidation (2.90 \pm 0.18 mg \cdot kg⁻¹ \cdot min⁻¹ before or 2.87 \pm 0.22 mg \cdot kg⁻¹ \cdot min⁻¹ after acipimox) but increased insulin-stimulated nonoxidative glucose metabolism from 2.51 \pm 0.35 to 3.12 \pm 0.43 mg \cdot kg⁻¹ \cdot min⁻¹ (P < 0.01).

Fatty acyl CoAs. Under basal conditions, before acipimox treatment, the total fatty acyl CoA concentration in muscle was 10.3 \pm 1.9 pmol/mg muscle (wet weight). The

TABLE 3
Effect of acipimox and insulin on glucose metabolism

	Endogenous glucose production	Glucose disposal	Glucose oxidation	Nonoxidative glucose metabolism	Fat oxidation
Pre-acipimox					
Basal	1.94 ± 0.09		1.33 ± 0.08	0.59 ± 0.10	0.77 ± 0.07
Insulin	0.1 ± 0.07	5.28 ± 0.46	2.90 ± 0.18†	2.51 ± 0.35†	0.18 ± 0.05†
Post-acipimox					
Basal	1.97 ± 0.10		1.52 ± 0.09§	0.46 ± 0.09	0.66 ± 0.04§
Insulin	0.01 ± 0.01§	6.31 ± 0.55‡‡	2.87 ± 0.22†	3.12‡ ± 0.43†‡	0.18 ± 0.04†

Data are means ± SE. †*P* < 0.01 vs. basal, ‡*P* < 0.05 vs. pre-acipimox, §*P* < 0.01 vs. pre-acipimox.

species of fatty acyl CoAs that could be identified based on coelution with standards included palmitoleoyl, palmitoyl, oleoyl, and stearoyl. Linoleoyl and arachidonoyl CoA were coeluted with these high-performance liquid chromatography conditions and were quantified together. Before acipimox treatment, insulin infusion decreased muscle total fatty acyl CoA concentrations from 10.3 ± 1.9 to 4.06 ± 0.78 pmol/mg (*P* < 0.01). Acipimox treatment itself decreased basal total fatty acyl CoA levels to 4.54 ± 0.82 pmol/mg (*P* < 0.01 vs. pre-acipimox levels). After acipimox, insulin infusion decreased total fatty acyl CoA concentration to 2.76 ± 0.57 pmol/mg (*P* < 0.05 vs. basal post-acipimox). The concentration of the individual fatty acyl CoA species is given in Table 4. Before acipimox, insulin decreased the concentration of all fatty acyl CoA species that were identified; the decreases in palmitoyl, oleoyl, and linoleoyl plus arachidonoyl CoA were significant. Under basal conditions, acipimox significantly decreased palmitoleoyl, palmitoyl, oleoyl, and stearoyl CoA. Under basal conditions, either before or after acipimox treatment, there was no correlation between the plasma FFA concentration and muscle total fatty acyl CoA concentration. However, at the end of 2 h of insulin infusion, plasma FFA and muscle fatty acyl CoA concentrations were positively correlated, with this relationship reaching statistical significance after acipimox treatment (*R* = 0.70, *P* < 0.05, Fig. 1).

Effect of acipimox and insulin on mRNA expression of PGC-1 and nuclear-encoded mitochondrial genes. The mRNA levels for PGC-1α and -β isoforms as well as an array of nuclear-encoded mitochondrial genes were determined using quantitative real-time PCR. The effects of acipimox on basal (no insulin) mRNA levels are shown in Fig. 2. Surprisingly, PGC-1α and NRF-1 mRNA were significantly decreased after acipimox (*P* < 0.05). PGC-1β and mRNA decreased, but this decrease was not statistically significant. Other mRNAs that decreased under basal

conditions after acipimox treatment included NRF-2, porin, cytochrome c oxidase subunit 6, mitochondrial transcription factor A (mtTFA), carnitine palmitoyl transferase 1B, NADH dehydrogenase flavoprotein 1, and citrate synthase.

The effects of insulin on expression of nuclear-encoded mitochondrial genes are shown in Fig. 3. Before acipimox treatment, insulin had little effect on mRNA expression for nuclear-encoded mitochondrial genes, except for PGC-1α mRNA, which increased 1.64 ± 0.35 (*P* < 0.05 vs. basal). After acipimox, however, in keeping with improved systemic insulin sensitivity, insulin increased the expression of several genes, including PGC-1α, NRF-1, NRF-2, and porin (all *P* < 0.05, Fig. 3).

Effect of acipimox and insulin on mRNA expression of extracellular matrix genes and genes involved in free radical scavenging. Acipimox treatment had profound effects on the suppression of extracellular matrix genes in skeletal muscle under basal conditions (Fig. 4). Under basal conditions after acipimox, both collagens I and III α-subunits, as well as fibronectin mRNA, increased many fold, with some individuals having increases of over 100-fold in the mRNA for the collagen genes. For example, the range in increase for the collagen 3α1 subunit was 2.21- to 104-fold, and the range for collagen 1α1 was 2.42- to 405-fold. Interestingly, after acipimox treatment, insulin suppressed collagen subunit mRNA (Fig. 4).

Because an increase in mRNA for matrix proteins often reflects the action of transforming growth factor (TGF)-β and CTGF, we also quantified the mRNA for these two genes. CTGF mRNA was increased 2.53 ± 0.47 by acipimox (*P* < 0.001, Fig. 5). Such an extracellular matrix gene, CTGF response often is in response to increase TGF-β signaling. To gain evidence regarding this potential mechanism, TGF-β1 mRNA was quantified. Under basal conditions, TGF-β1 mRNA was also increased 2.95 ± 1.11-fold (*P* < 0.05, Fig. 5) after acipimox treatment. After acipimox,

TABLE 4
Myocellular fatty acyl CoA concentrations

	Palmitoleoyl	Linoleoyl and arachidonoyl	Palmitoyl	Oleoyl	Stearoyl	Total
Pre-acipimox						
Basal	0.19 ± 0.10	3.50 ± 1.33	1.89 ± 0.57	3.28 ± 1.18	1.49 ± 0.34	10.3 ± 1.9
Insulin	0.007 ± 0.005*	2.16 ± 0.65	0.57 ± 0.11*	0.69 ± 0.26*	0.64 ± 0.15†	4.06 ± 0.78†
Post-acipimox						
Basal	0.05 ± 0.05	1.50 ± 0.59‡	0.61 ± 0.19‡†	1.33 ± 0.31‡	1.05 ± 0.31	4.54 ± 0.82§
Insulin	0.07 ± 0.07	1.08 ± 0.45	0.44 ± 0.17	0.53 ± 0.17	0.63 ± 0.15	2.76 ± 0.57*

Data are means ± SE in units of picomoles per milligram of muscle wet weight. **P* < 0.05, †*P* < 0.01 vs. basal, ‡*P* < 0.05 vs. pre-acipimox, §*P* < 0.01 vs. pre-acipimox.




FIG. 1. Relationships between plasma FFA concentrations and muscle total fatty acyl CoA concentrations. Data for both basal and clamp (insulin) data are shown before (*A* and *C*) and after (*B* and *D*) acipimox treatment. Of the 10 subjects studied, 8 had sufficient biopsy material to complete fatty acyl CoA determinations.




FIG. 2. Effect of acipimox treatment on basal mRNA expression of nuclear-encoded mitochondrial genes. The dashed line indicates no change in gene expression. * $P < 0.05$, ** $P < 0.01$ vs. pre-acipimox. For abbreviations, see the Table 1 footnote.




FIG. 3. Effect of insulin on mRNA expression of nuclear-encoded mitochondrial genes before (A) and after (B) acipimox treatment. The dashed line indicates no change in gene expression. * $P < 0.05$, ** $P < 0.01$ insulin vs. basal. For abbreviations, see the Table 1 footnote.

insulin significantly decreased the mRNA for CTGF (0.43 ± 0.15 -fold insulin vs. basal, $P < 0.001$). Because of the increases in these indicators of an inflammatory response, we also assayed the mRNA expression for superoxide dismutase 2 (SOD2), 8-oxoguanine DNA glycosylase (OGG1), glutathione peroxidase 1 (GPX1), and NAD(P)H ubiquinone oxidoreductase (NQO1, cytoplasmic free radical scavenger). These results are shown in Fig. 6. Of these genes, only NQO1 increased significantly after acipimox. Interestingly, before acipimox, of these genes, insulin increased only NQO1 mRNA ($P < 0.05$), whereas after acipimox, insulin significantly increased not only NQO1, but also the mRNA for SOD2 and GPX1 (both $P < 0.05$).

DISCUSSION

Insulin resistance in skeletal muscle is associated with increased muscle lipid content and can be reproduced by experimental increases in plasma lipids. On a cellular level, insulin resistance is accompanied by increased myocellular fatty acyl CoA levels (31,40) and mitochondrial abnormalities (20,21), while on a molecular level it is

characterized by decreased expression of nuclear-encoded mitochondrial genes (22,23) and increased expression of extracellular matrix genes and CTGF (27). The present study was undertaken to determine whether a decrease in plasma FFA concentration and improvement in insulin sensitivity brought about by treating genetically predisposed normal glucose tolerant insulin-resistant subjects with acipimox could reverse the biochemical and molecular abnormalities associated with lipid-induced insulin resistance.

In the present study, acipimox treatment for 7 days improved insulin-stimulated glucose disposal by ~20% in normal glucose tolerant insulin-resistant subjects with a strong family history of type 2 diabetes. This result is similar to that observed in previous studies using acipimox in insulin-resistant subjects (32,35). This improvement in insulin sensitivity was accompanied by a 60% decrease in plasma FFAs under postabsorptive conditions, without changing the fasting plasma insulin concentration. Thus, this pharmacological maneuver successfully brought




FIG. 4. Effect of acipimox treatment under basal conditions (A) or insulin before and after acipimox (B) on mRNA expression of extracellular matrix genes. The dashed line indicates no change in gene expression. * $P < 0.05$, ** $P < 0.01$ vs. pre-acipimox or basal vs. insulin. For abbreviations, see the Table 1 footnote.




FIG. 5. Effect of acipimox treatment under basal conditions (A) or insulin before and after acipimox (B) on mRNA expression of CTGF and TGF- β 1. The dashed line indicates no change in gene expression. * $P < 0.05$, ** $P < 0.01$ vs. pre-acipimox or basal versus insulin. For abbreviations, see the Table 1 footnote.

about the changes required to test the hypotheses proposed in the present study.

One purpose of this study was to determine whether a decrease in plasma FFAs resulted in a decrease in myocellular fatty acyl CoAs. The results of the present study answer this question in the affirmative; after acipimox, total fatty acyl CoA concentrations in skeletal muscle decreased by ~50–60%. The decline in muscle fatty acyl CoAs is similar to what was observed after acipimox treatment of insulin-resistant patients with type 2 diabetes (31), who also had improved insulin sensitivity. This decline also is consistent with the hypothesis that increased levels of fatty acyl CoAs are linked with or causally related to insulin resistance. The decrease in fatty acyl CoA was not limited to any particular fatty acid chain length, since similar decreases were seen with all the species that could be detected using this method.

The results of this study also provide insight into the potential mechanism regulating myocellular fatty acyl CoA

levels. Insulin infusion, before acipimox treatment, suppressed plasma FFA concentrations by ~75% and muscle fatty acyl CoA levels by 60%. Acipimox treatment by itself also decreased the postabsorptive levels of both plasma FFAs and fatty acyl CoA to a similar extent. These data suggest that fatty acyl CoA levels in muscle may closely track plasma FFA availability. Supporting this was the decreased effect of insulin, after acipimox, on already suppressed plasma FFA and muscle fatty acyl CoA concentrations. Despite this, there was no significant correlation between plasma FFA and muscle fatty acyl CoA content under basal conditions before or after acipimox treatment. However, under conditions where insulin further suppressed lipolysis after treatment with acipimox (in other words, during conditions of minimum plasma FFA availability), fatty acyl CoA concentrations in muscle were significantly correlated with plasma FFAs. Therefore, although the present data do not address abnormalities in fatty acid oxidation in insulin resistance, they emphasize




FIG. 6. Effect of acipimox treatment under basal conditions (A) or insulin before and after acipimox (B) on mRNA expression of genes involved in free radical scavenging or DNA repair. The dashed line indicates no change in gene expression. * $P < 0.05$ vs. pre-acipimox or basal vs. insulin. For abbreviations, see the Table 1 footnote.

that myocellular fatty acyl CoA concentrations track plasma FFA availability when plasma FFAs are maximally suppressed. Therefore, if increased myocellular fatty acyl CoA plays a causal role in the pathogenesis of insulin resistance, it follows that simply decreasing the plasma FFA concentration could be a viable means of improving insulin sensitivity.

In undertaking a previous experiment (23), we found that plasma FFA concentrations were inversely correlated with the mRNA expression of PGC-1 α (L.J.M., Mary-Elizabeth Patti, unpublished data), and the decrease in PGC-1 α mRNA was accompanied by decreased mRNA expression of a number of nuclear-encoded mitochondrial genes involved in electron transport and oxidative phosphorylation (22,23). Subsequently, we showed that increased plasma FFAs and induced experimental insulin resistance drove down the expression of PGC-1 α and nuclear-encoded mitochondrial genes (27). Thus, we expected that in the current experiment, in which we improved insulin sensitivity by decreasing plasma FFAs, the mRNA expression of PGC-1 and nuclear-encoded mitochondrial genes would increase. In general, we found the opposite. The effect of acipimox treatment on PGC-1 and nuclear-encoded mitochondrial gene expression was clear; 9 of 13 mRNAs that were quantified decreased significantly in muscle obtained under postabsorptive conditions after acipimox. The decrease in citrate synthase provides at least indirect evidence against the notion that an increase in mitochondrial number was associated with an improvement in insulin sensitivity. It is possible that this pattern of decrease in mRNA expression is not reflected in protein abundance or activity, but unfortunately, there was not sufficient muscle biopsy material remaining after quantification of fatty acyl CoA concentration and the mRNA expression of multiple genes to perform determinations of protein concentration. Further studies integrating measurements of protein abundance and respiratory chain complex activities will be required to completely understand these events.

The present results with respect to the decrease in PGC-1 expression also differ from those in an earlier study by another group (41). In that study, lipolysis was decreased acutely in healthy subjects using nicotinic acid, and muscle PGC-1 α mRNA increased acutely (within hours). Differences in study design make it difficult to compare those results with findings from the current study. The present study was undertaken to improve insulin sensitivity in insulin-resistant subjects by suppressing plasma FFA concentrations to test whether such a change was accompanied by increased PGC-1 α expression. To do this required a longer-term suppression of FFAs than was used in the acute study of Watt et al. (41). Also, healthy subjects may respond differently than insulin-resistant subjects with a family history of type 2 diabetes. Regardless, it is conceivable that suppression of plasma FFAs in the present study may have led to an acute increase in PGC-1 expression that in fact led to increased PGC-1 protein and perhaps improved mitochondrial function, but a subsequent decrease in expression of PGC-1 and nuclear-encoded mitochondrial genes may have occurred as a compensatory homeostatic effort. Only detailed time course experiments can dissect these possibilities.

In an earlier experiment in which lipid was infused to create insulin resistance, we observed marked increases in mRNA and protein expression of extracellular matrix

genes (27). Based on those results, we predicted that decreasing plasma FFAs would decrease the expression of extracellular matrix genes. Again, this prediction was not upheld by experimental evidence. Rather than decreasing the expression of these genes, acipimox treatment, which reduced plasma FFA and myocellular fatty acyl CoA concentrations, dramatically increased extracellular matrix gene expression. Also increased were TGF- β 1 and CTGF mRNA, and these two factors often initiate a remodeling of the extracellular matrix. Taken together with our earlier results, the present data suggest that changes in insulin sensitivity in either direction induced by changes in plasma FFAs are accompanied by a remodeling of the extracellular matrix. Matrix turnover, with release or sequestering of growth factors such as TGF- β (42,43), as well as changes in cell-matrix interactions and reorganization of the cytoskeleton (44–46), can have profound effects on the behavior of cells. These findings suggest that insulin action at the cellular or molecular level may be connected to the composition and turnover of the extracellular matrix. It must be noted that many factors, in addition to TGF- β , can influence CTGF expression and its subsequent effects on extracellular matrix gene expression. CTGF expression is increased by high glucose (47,48), as well as being elevated in fatty liver disease (49,50), and atherosclerotic plaques (51). A possible link between fatty acid metabolism and CTGF expression is indicated by the observation that sphingosine 1-phosphate, signaling through sphingosine 1-phosphate receptors, directly increases CTGF mRNA in mesangial cells (52), smooth muscle (53), fibroblasts (54), and endothelial cells (55). Another possibility is that extracellular fatty acids are signaling directly through cell surface receptors, such as the toll-like receptors or others, which are present in skeletal muscle (56,57). Toll-like receptors, part of the innate immune system, engage saturated and polyunsaturated fatty acid ligands (58,59), and there is evidence that saturated and polyunsaturated fatty acids signal through toll-like receptor 4 (TLR4) to produce opposing effects (60). It should be recognized that, in the present study and the previous study infusing lipid (27), the composition of individual FFA species in the plasma have changed as dramatically as the total concentration. It is possible that engagement of different combinations of toll-like receptors by a different array of fatty acid ligands might produce similar effects on the extracellular matrix even though total FFA concentrations diverge. Intriguingly, in one study (61), lipopolysaccharide and polyunsaturated fatty acids, signaling through TLR4, activated nuclear factor κ B and increased collagen production.

Despite the similarities between the results of our two studies, there were a number of differences in experimental design that lead to the opposite changes in insulin sensitivity. In the study by Richardson et al. (27), lipid was infused into healthy subjects for 48 h, increasing not only plasma FFAs, but also plasma triglyceride levels. In the present study, 7 days of acipimox treatment only altered plasma FFA concentrations, and the study was performed in insulin-resistant subjects with a strong family history of type 2 diabetes. In addition, with the design of the current study, a direct effect of acipimox cannot be ruled out. Regardless of these differences, the present study shows that improved insulin sensitivity can be dissociated, at least temporally, from an increase in mRNA expression of PGC-1 and other nuclear-encoded mitochondrial genes. Moreover, both increased and

decreased insulin sensitivity were associated with profound increases in expression of extracellular matrix genes and CTGF, indicating changes in the immediate environment of the cell, that is, the extracellular matrix, with which the cell constantly interacts.

Irrespective of their origin, since these changes in gene expression may have represented an inflammatory response, we also determined the mRNA expression for four genes encoding proteins involved in free radical scavenging or DNA repair (SOD2, OGG1, GPX1, and NQO1). Of these, mRNA expression for the cytoplasmic free radical scavenger NQO1 [NAD(P)H ubiquinone oxidoreductase 1] was increased by 50% after acipimox. If the protein and activity for this gene are also increased, this would be consistent with the inflammatory-like response in the muscle observed after 7 days of acipimox treatment.

Although our observations link changes in expression of genes encoding mitochondrial and extracellular matrix proteins, the present data cannot directly address the mechanisms connecting these phenomena. However, the results of some other recent studies may provide insight into this. First, a connection between the composition of the extracellular matrix and mitochondrial function was provided by data from a collagen 6-null mouse that is a model of Bethlem muscular dystrophy (62). The muscular dystrophy in the mouse is due to a severe disruption of mitochondrial function that can be restored in Col 6-null myoblasts by culturing the cells on a matrix containing collagen 6 (62). Second, a more recent study from van Waveren et al. (63) provided evidence, using cultured cells in which oxidative phosphorylation was inhibited chemically or by genetic manipulation, that dysfunction in oxidative phosphorylation increases CTGF expression and extracellular matrix turnover on a transcriptional level. This opens the possibility that changes in oxidative phosphorylation can affect extracellular matrix composition directly.

In summary, the present results confirm that a reduction in myocellular fatty acyl CoA concentrations is elicited by a decrease in lipolysis and plasma FFAs and that this is accompanied by improved insulin sensitivity. However, the changes in expression of nuclear-encoded mitochondrial and extracellular matrix genes are paradoxical when viewed in the context that of earlier findings. These seemingly paradoxical results will require further study to define the role of changes in nuclear mitochondrial gene expression in the setting of changing insulin sensitivity. The profound changes in expression of extracellular matrix gene expression associated with changes, either up or down, in insulin sensitivity provide new evidence in favor of a role for remodeling of the extracellular matrix in insulin sensitivity in skeletal muscle, although the mechanism responsible for such a role is not known.

REFERENCES

- Stettler R, Ith M, Acheson KJ, Decombaz J, Boesch C, Tappy L, Binnert C: Interaction between dietary lipids and physical inactivity on insulin sensitivity and on intramyocellular lipids in healthy men. *Diabetes Care* 28:1404–1409, 2005
- Colberg S, Simoneau J-A, Thaete FL, Kelley DE: Impaired FFA utilization by skeletal muscle in women with visceral obesity. *J Clin Invest* 95:1846–1853, 1995
- Kelley DE, Simoneau J-A: Impaired FFA utilization by skeletal muscle in NIDDM. *J Clin Invest* 94:2349–2356, 1994
- Adams JM 2nd, Pratipanawatr T, Berria R, Wang E, DeFronzo RA, Sullards MC, Mandarino LJ: Ceramide content is increased in skeletal muscle from obese insulin-resistant humans. *Diabetes* 53:25–31, 2004
- Bachmann O, Dahl D, Brechtel K, Machann J, Haap M, Maier T, Loviscach M, Stumvoll M, Claussen C, Schick F, Haring H, Jacob S: Effects of intravenous and dietary lipid challenge on intramyocellular lipid content and the relation with insulin sensitivity in humans. *Diabetes* 50:2579–2584, 2001
- Goodpaster BH, Theriault R, Watkins S, Kelley DE: Intramuscular lipid content is increased in obesity and decreased by weight loss. *Metabolism* 49:467–472, 2000
- Itani SI, Ruderman NB, Schmieder F, Boden G: Lipid-induced insulin resistance in human muscle is associated with changes in diacylglycerol, protein kinase C, and IkappaB-alpha. *Diabetes* 51:2005–2011, 2002
- Krassak M, Falk Petersen K, Dresner A, DiPietro L, Vogel SM, Rothman DL, Roden M, Shulman GI: Intramyocellular lipid concentrations are correlated with insulin sensitivity in humans: a ¹H NMR spectroscopy study. *Diabetologia* 42:113–116, 1999
- Laybutt DR, Schmitz-Peiffer S, Ruderman NB, Chisholm D, Biden T, Kraegen EW: Activation of protein kinase C ϵ may contribute to muscle insulin resistance induced by lipid accumulation during chronic glucose infusion in rats (Abstract). *Diabetes* 46 (Suppl. 1):241A, 1997
- Aguirre V, Werner ED, Giraud J, Lee YH, Shoelson SE, White MF: Phosphorylation of Ser307 in insulin receptor substrate-1 blocks interactions with the insulin receptor and inhibits insulin action. *J Biol Chem* 277:1531–1537, 2002
- Gao Z, Zhang X, Zuberi A, Hwang D, Quon MJ, Lefevre M, Ye J: Inhibition of insulin sensitivity by free fatty acids requires activation of multiple serine kinases in 3T3-L1 adipocytes. *Mol Endocrinol* 18:2024–2034, 2004
- Le Marchand-Brustel Y, Gual P, Gremeaux T, Gonzalez T, Barres R, Tanti JF: Fatty acid-induced insulin resistance: role of insulin receptor substrate 1 serine phosphorylation in the retroregulation of insulin signalling. *Biochem Soc Trans* 31:1152–1156, 2003
- Dresner A, Laurent D, Marcucci M, Griffon ME, Dufour S, Cline GW, Slezak LA, Andersen DK, Hundal RS, Rothman DL, Petersen KF, Shulman GI: Effects of free fatty acids on glucose transport and IRS-1-associated phosphatidylinositol 3-kinase activity. *J Clin Invest* 103:253–259, 1999
- Powell DJ, Turban S, Gray A, Hajdouch E, Hundal HS: Intracellular ceramide synthesis and protein kinase C ζ activation play an essential role in palmitate-induced insulin resistance in rat L6 skeletal muscle cells. *Biochem J* 382:619–629, 2004
- Schmitz-Peiffer C, Craig DL, Biden TJ: Ceramide generation is sufficient to account for the inhibition of the insulin-stimulated PKB pathway in C2C12 skeletal muscle cells pretreated with palmitate. *J Biol Chem* 274:24202–24210, 1999
- Schmitz-Peiffer C, Oakes ND, Browne CL, Kraegen EW, Biden TJ: Alterations in the expression and cellular localization of protein kinase C isozymes epsilon and theta are associated with insulin resistance in skeletal muscle of the high-fat-fed rat. *Diabetes* 46:169–178, 1997
- Schubert K, Scheid M, Duronio V: Ceramide inhibits protein kinase B/Akt by promoting dephosphorylation of serine 473. *J Biol Chem* 275:13330–13335, 2000
- Teruel T, Hernandez R, Lorenzo M: Ceramide mediates insulin resistance by tumor necrosis factor- α in brown adipocytes by maintaining Akt in an inactive dephosphorylated state. *Diabetes* 50:2563–2571, 2001
- Kashyap SR, Belfort R, Berria R, Suraamornkul S, Pratipanawatr T, Finlayson J, Barrentine A, Bajaj M, Mandarino L, DeFronzo R, Cusi K: Discordant effects of a chronic physiological increase in plasma FFA on insulin signaling in healthy subjects with or without a family history of type 2 diabetes. *Am J Physiol Endocrinol Metab* 287:E537–E546, 2004
- Kelley DE, He J, Menshikova EV, Ritov VB: Dysfunction of mitochondria in human skeletal muscle in type 2 diabetes. *Diabetes* 51:2944–2950, 2002
- Petersen KF, Dufour S, Befroy D, Garcia R, Shulman GI: Impaired mitochondrial activity in the insulin-resistant offspring of patients with type 2 diabetes. *N Engl J Med* 350:664–671, 2004
- Mootha VK, Lindgren CM, Eriksson KF, Subramanian A, Sihag S, Lehar J, Puigserver P, Carlsson E, Ridderstrale M, Laurila E, Houstis N, Daly MJ, Patterson N, Mesirov JP, Golub TR, Tamayo P, Spiegelman B, Lander ES, Hirschhorn JN, Altshuler D, Groop LC: PGC-1 α -responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nat Genet* 34:267–273, 2003
- Patti ME, Butte AJ, Crunkhorn S, Cusi K, Berria R, Kashyap S, Miyazaki Y, Kohane I, Costello M, Saccone R, Landaker EJ, Goldfine AB, Mun E, DeFronzo R, Finlayson J, Kahn CR, Mandarino LJ: Coordinated reduction of genes of oxidative metabolism in humans with insulin resistance and diabetes: potential role of PGC1 and NRF1. *Proc Natl Acad Sci U S A* 100:8466–8471, 2003
- Boden G, Jadali F, White J, Liang Y, Mozzoli M, Chen X, Coleman E, Smith C: Effect of fat on insulin-stimulated carbohydrate metabolism in normal men. *J Clin Invest* 88:960–966, 1991

25. Kelley DE, Mookan M, Simoneau J-A, Mandarino LJ: Interaction between glucose and free fatty acid metabolism in human skeletal muscle. *J Clin Invest* 92:91–98, 1993
26. Thiebaut D, DeFronzo RA, Jacot E, Golay A, Acheson K, Maeder E, Jequier E, Felber JP: Effect of long chain triglyceride infusion on glucose metabolism in man. *Metabolism* 31:1128–1136, 1982
27. Richardson DK, Kashyap S, Bajaj M, Cusi K, Mandarino SJ, Finlayson J, DeFronzo RA, Jenkinson CP, Mandarino LJ: Lipid infusion decreases the expression of nuclear encoded mitochondrial genes and increases the expression of extracellular matrix genes in human skeletal muscle. *J Biol Chem* 280:10290–10297, 2005
28. Berria R, Wang L, Richardson DK, Finlayson J, Belfort R, Pratipanawatr T, De Filippis EA, Kashyap S, Mandarino LJ: Increased collagen content in insulin-resistant skeletal muscle. *Am J Physiol Endocrinol Metab* 290: E560–E565, 2006
29. Perbal B: CCN proteins: multifunctional signalling regulators. *Lancet* 363:62–64, 2004
30. Saloranta C, Franssila-Kallunki A, Ekstrand A, Taskinen MR, Groop L: Modulation of hepatic glucose production by non-esterified fatty acids in type 2 (non-insulin-dependent) diabetes mellitus. *Diabetologia* 34:409–415, 1991
31. Bajaj M, Suraamornkul S, Romanelli A, Cline GW, Mandarino LJ, Shulman GI, DeFronzo RA: Effect of a sustained reduction in plasma free fatty acid concentration on intramuscular long-chain fatty Acyl-CoAs and insulin action in type 2 diabetic patients. *Diabetes* 54:3148–3153, 2005
32. Vaag A, Skott P, Damsbo P, Gall MA, Richter EA, Beck-Nielsen H: Effect of the antilipolytic nicotinic acid analogue acipimox on whole-body and skeletal muscle glucose metabolism in patients with non-insulin-dependent diabetes mellitus. *J Clin Invest* 88:1282–1290, 1991
33. Gulli G, Ferrannini E, Stern M, Haffner S, DeFronzo RA: The metabolic profile of NIDDM is fully established in glucose-tolerant offspring of two Mexican-American NIDDM parents. *Diabetes* 41:1575–1586, 1992
34. Vaag A, Henriksen JE, Beck-Nielsen H: Decreased insulin activation of glycogen synthase in skeletal muscles in young nonobese Caucasian first-degree relatives of patients with non-insulin-dependent diabetes mellitus. *J Clin Invest* 89:782–788, 1992
35. Bajaj M, Suraamornkul S, Kashyap S, Cusi K, Mandarino L, DeFronzo RA: Sustained reduction in plasma free fatty acid concentration improves insulin action without altering plasma adipocytokine levels in subjects with strong family history of type 2 diabetes. *J Clin Endocrinol Metab* 89:4649–4655, 2004
36. DeFronzo RA, Tobin JD, Andres R: Glucose clamp technique: a method for quantifying insulin secretion and resistance. *Am J Physiol* 237:E214–E223, 1979
37. Corkey BE, Brandt M, Williams RJ, Williamson JR: Assay of short-chain acyl coenzyme A intermediates in tissue extracts by high-pressure liquid chromatography. *Anal Biochem* 118:30–41, 1981
38. Livak KJ, Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25:402–408, 2001
39. DeBodo RC, Steele R, Altszuler N, Dunn A, Bishop JS: On the hormonal regulation of carbohydrate metabolism; studies with C14 glucose. *Recent Prog Horm Res* 19:445–488, 1963
40. Houmard J, Tanner CJ, Yu C, Cunningham PG, Pories W, MacDonald K, Sulman G: Effect of weight loss on insulin sensitivity and intramuscular long-chain fatty acyl-CoAs in morbidly obese subjects. *Diabetes* 51:2959–2963, 2002
41. Watt MJ, Southgate RJ, Holmes AG, Febbraio MA: Suppression of plasma free fatty acids upregulates peroxisome proliferator-activated receptor (PPAR) alpha and delta and PPAR coactivator 1alpha in human skeletal muscle, but not lipid regulatory genes. *J Mol Endocrinol* 33:533–544, 2004
42. Fowlkes JL, Winkler MK: Exploring the interface between metallo-proteinase activity and growth factor and cytokine bioavailability. *Cytokine Growth Factor Rev* 13:277–287, 2002
43. Koli K, Saharinen J, Hyytiainen M, Penttinen C, Keski-Oja J: Latency, activation, and binding proteins of TGF-beta. *Microsc Res Tech* 52:354–362, 2001
44. Davis GE, Senger DR: Endothelial extracellular matrix: biosynthesis, remodeling, and functions during vascular morphogenesis and neovessel stabilization. *Circ Res* 97:1093–1107, 2005
45. Hai CM, Gu Z: Caldesmon phosphorylation in actin cytoskeletal remodeling. *Eur J Cell Biol* 85:305–309, 2006
46. Varon C, Tatin F, Moreau V, Van Obberghen-Schilling E, Fernandez-Sauze S, Reuzeau E, Kramer I, Genot E: Transforming growth factor beta induces rosettes of podosomes in primary aortic endothelial cells. *Mol Cell Biol* 26:3582–3594, 2006
47. McLennan SV, Wang XY, Moreno V, Yue DK, Twigg SM: Connective tissue growth factor mediates high glucose effects on matrix degradation through tissue inhibitor of matrix metalloproteinase type 1: implications for diabetic nephropathy. *Endocrinology* 145:5646–5655, 2004
48. Sakamoto N, Sugimura K, Kawashima H, Tsuchida K, Takemoto Y, Naganuma T, Tatsumi S, Nakatani T: Influence of glucose and inflammatory cytokines on TGF-beta1 and CTGF mRNA expressions in human peritoneal mesothelial cells. *Int J Mol Med* 15:907–911, 2005
49. Paradis V, Perlemuter G, Bonvoust F, Dargere D, Parfait B, Vidaud M, Conti M, Huet S, Ba N, Buffet C, Bedossa P: High glucose and hyperinsulinemia stimulate connective tissue growth factor expression: a potential mechanism involved in progression to fibrosis in nonalcoholic steatohepatitis. *Hepatology* 34:738–744, 2001
50. Sato W, Horie Y, Kataoka E, Ohshima S, Dohmen T, Iizuka M, Sasaki J, Sasaki T, Hamada K, Kishimoto H, Suzuki A, Watanabe S: Hepatic gene expression in hepatocyte-specific Pten deficient mice showing steatohepatitis without ethanol challenge. *Hepatology* 34:256–265, 2006
51. Cicha I, Yilmaz A, Klein M, Raitheil D, Brigstock DR, Daniel WG, Goppelt-Strube M, Garlich CD: Connective tissue growth factor is overexpressed in complicated atherosclerotic plaques and induces mononuclear cell chemotaxis in vitro. *Arterioscler Thromb Biol* 25:1008–1013, 2005
52. Katsuma S, Ruike Y, Yano T, Kimura M, Hirasawa A, Tsujimoto G: Transcriptional regulation of connective tissue growth factor by sphingosine 1-phosphate in rat cultured mesangial cells. *FEBS Lett* 579:2576–2582, 2005
53. Chowdhury I, Chaour B: Regulation of connective tissue growth factor (CTGF/CCN2) gene transcription and mRNA stability in smooth muscle cells: involvement of RhoA GTPase and p38 MAP kinase and sensitivity to actin dynamics. *Eur J Biochem* 271:4436–4450, 2004
54. Zhu MJ, Kim CD, Kwon YB, Kye KC, Chen YY, Lee WH, Lee S, Lim JS, Seo YJ, Suhr KB, Park JK, Lee JH: Induction of connective tissue growth factor expression by sphingosylphosphorylcholine in cultured human skin fibroblasts. *Exp Dermatol* 14:509–514, 2005
55. Muehlich S, Schneider N, Hinkmann F, Garlich CD, Goppelt-Strube M: Induction of connective tissue growth factor (CTGF) in human endothelial cells by lysophosphatidic acid, sphingosine-1-phosphate, and platelets. *Atherosclerosis* 175:261–268, 2004
56. Boyd JH, Divangahi M, Yahiaoui L, Gvozdic D, Qureshi S, Petrof BJ: Toll-like receptors differentially regulate CC and CXC chemokines in skeletal muscle via NF-kB and calcineurin. *Infect Immun* 74:6829–6838, 2006
57. Schreiner B, Voss J, Wischhusen J, Dombrowski Y, Steinle A, Lochmuller H, Dalakas M, Melms A, Wiendl H: Expression of toll-like receptors in inflammatory and HIV myopathies, mediates IL-8 release and up-regulation of NKG2D-ligands. *FASEB J* 20:118–120, 2006
58. Lee JY, Sohn KH, Rhee SH, Hwang D: Saturated fatty acids, but not unsaturated fatty acids, induce the expression of cyclooxygenase-2 mediated through Toll-like receptor 4. *J Biol Chem* 276:16683–16689, 2001
59. Lee JY, Zhao L, Youn HS, Weatherill AR, Tapping R, Feng L, Lee WH, Fitzgerald KA, Hwang DH: Saturated fatty acid activates but polyunsaturated fatty acid inhibits Toll-like receptor 2 dimerized with Toll-like receptor 6 or 1. *J Biol Chem* 279:16971–16979, 2004
60. Weatherill AR, Lee JY, Zhao L, Lemay DG, Youn HS, Hwang DH: Saturated and polyunsaturated fatty acids reciprocally modulate dendritic cell functions mediated through TLR4. *J Immunol* 174:5390–5397, 2005
61. Jia Y, Turek JJ: Altered NF-kappaB gene expression and collagen formation induced by polyunsaturated fatty acids. *J Nutr Biochem* 16:500–506, 2005
62. Irwin WA, Bergamin N, Sabatelli P, Reggiani C, Megighian A, Merlini L, Braghetta P, Columbaro M, Volpin D, Bressan GM, Bernardi P, Bonaldo P: Mitochondrial dysfunction and apoptosis in myopathic mice with collagen VI deficiency. *Nat Genet* 35:367–371, 2003
63. van Waveren C, Sun Y, Cheung HS, Moraes CT: Oxidative phosphorylation dysfunction modulates expression of extracellular matrix-remodeling genes and invasion. *Carcinogenesis* 27:409–418, 2006
64. Sparks LM, Xie H, Koza RA, Mynatt R, Hulver MW, Bray GA, Smith SR: A high-fat diet coordinately downregulates genes required for mitochondrial oxidative phosphorylation in skeletal muscle. *Diabetes* 54:1926–1933, 2005
65. Eikmans M, Sijpkens YW, Baelde HJ, de Heer E, Paul LC, Bruijn JA: High transforming growth factor-beta and extracellular matrix mRNA response in renal allografts during early acute rejection is associated with absence of chronic rejection. *Transplantation* 73:573–579, 2002
66. Civitarese AE, Ukropcova B, Carling S, Hulver M, DeFronzo RA, Mandarino L, Ravussin E, Smith SR: Role of adiponectin in human skeletal muscle bioenergetics. *Cell Metab* 4:75–87, 2006