

# Full-Length Adiponectin Attenuates Insulin Signaling and Inhibits Insulin-Stimulated Amino Acid Transport in Human Primary Trophoblast Cells

Helen N. Jones,<sup>1,2</sup> Thomas Jansson,<sup>1,3</sup> and Theresa L. Powell<sup>1,3</sup>

**OBJECTIVE**—Maternal adiponectin levels are reduced and placental nutrient transporters are upregulated in obesity and gestational diabetes mellitus; however, the effects of adiponectin on placental function are unknown. We hypothesized that adiponectin regulates placental amino acid transport.

**RESEARCH DESIGN AND METHODS**—Human primary trophoblast cells were cultured and incubated with globular adiponectin (gAd) or full-length adiponectin (fAd) alone or in combination with insulin. System A and L amino acid transport and SNAT1, SNAT2, and SNAT4 isoform expression was measured. The activity of the AMP-activated protein kinase (AMPK), phosphatidylinositol 3 kinase–AKT, and peroxisome proliferator-activated receptor- $\alpha$  (PPAR $\alpha$ ) signaling pathways was determined.

**RESULTS**—In the absence of insulin, gAd stimulated AMPK Thr172 phosphorylation, SNAT2 protein expression, and system A activity. This effect appeared to be mediated by interleukin-6 release and signal transducer and activator of transcription 3 (STAT3) signaling because gAd failed to stimulate system A in cells in which STAT3 had been silenced using small interfering RNA. fAd alone had no effect on system A activity or SNAT expression. Insulin increased AKT and insulin receptor substrate 1 (IRS-1) phosphorylation, system A activity, and SNAT2 expression. When combined with insulin, gAd did not affect system A activity or SNAT expression. In contrast, fAd abolished insulin-stimulated AKT Thr308 and IRS-1 Tyr612 phosphorylation, system A activity, and SNAT2 expression. Furthermore, fAd increased PPAR $\alpha$  expression and PPAR $\alpha$  (Ser21) phosphorylation.

**CONCLUSIONS**—In contrast to the insulin-sensitizing actions of adiponectin in liver and muscle reported in the literature, fAd attenuates insulin signaling in primary human trophoblast cells. As a result, fAd inhibits insulin-stimulated amino acid transport, which may have important implications for placental nutrient transport and fetal growth in pregnancy complications associated with altered maternal adiponectin levels. *Diabetes* 59: 1161–1170, 2010

From the <sup>1</sup>Department of Obstetrics and Gynecology, University of Cincinnati, Cincinnati, Ohio; the <sup>2</sup>Center for Molecular Fetal Therapy, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio; and the <sup>3</sup>Center for Pregnancy and Newborn Research, Department of Obstetrics and Gynecology, University of Texas Health Science Center, San Antonio, Texas.

Corresponding author: Helen N. Jones, helen.jones@cchmc.org.  
Received 1 June 2009 and accepted 27 January 2010. Published ahead of print at <http://diabetes.diabetesjournals.org> on 11 February 2010. DOI: 10.2337/db09-0824.

© 2010 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

More than half of American women enter pregnancy overweight or obese and 5–10% of all pregnant women develop gestational diabetes mellitus (1). These women are more likely to give birth to babies with increased birth weight (1), traumatic birth injuries (2), and increased risk of developing metabolic syndrome (3). The cause of fetal overgrowth remains to be established; however, the primary determinant of fetal growth is nutrient supply, which is dependent on placental nutrient transport. Placental amino acid (4) and GLUT activity (5,6) as well as the expression of fatty acid-binding proteins in the placenta (7) are increased in diabetes, suggesting that an upregulation of placental nutrient transport capacity may contribute to fetal overgrowth (8). However, the underlying mechanisms in these pregnancy complications are poorly understood. Obesity in pregnancy (9,10) and gestational diabetes mellitus (11) are associated with low maternal circulating levels of adiponectin, but the impact of adiponectin on placental nutrient transport functions is unknown.

Adiponectin is a protein hormone produced in adipose tissue. The full-length form (fAd), which consists of an NH<sub>2</sub>-terminal sequence, a complement C1q-like domain, and a COOH-terminal globular domain (12), can multimerize to form multimeric structures. A truncated form containing only the COOH-terminal portion (globular adiponectin [gAd]) is produced by proteolytic cleavage and is biologically active, although the relative serum levels of the two forms in normal physiology and pathology are unknown (13). The two receptors, AdipoR1 and AdipoR2, have distinct expression patterns and bind the two forms of adiponectin with different affinity (14). AdipoR1 is ubiquitously expressed, with the highest levels in skeletal muscle, and binds primarily gAd (14). AdipoR2 is abundantly expressed in the liver and binds fAd with higher affinity than gAd (15). Adiponectin exerts its cellular effects by activating AMP-activated protein kinase (AMPK), p38 mitogen-activated protein kinase, and peroxisome proliferator-activated receptor- $\alpha$  (PPAR $\alpha$ ); however, details of the signaling transduction pathway remain to be determined (15). In skeletal muscle, there is significant cross-talk between AdipoR1 and AdipoR2 signaling because both receptors activate PPAR $\alpha$  and AMPK to increase glucose uptake and fatty acid oxidation (15,16). In the liver, however, AdipoR1 and AdipoR2 have distinct signaling pathways: AdipoR1 signaling is believed to be mediated via AMPK-reducing gluconeogenesis, whereas AdipoR2 activates PPAR $\alpha$  to increase fatty acid oxidation (16).

Adiponectin has an insulin-sensitizing action and circulating adiponectin levels are reduced in insulin-resistant

states such as obesity and type 2 diabetes (17,18). The mechanisms underlying the increase in insulin sensitivity in response to adiponectin remains to be fully established and may include a multitude of pathways (19–21). For example, in skeletal muscle, adiponectin stimulates glucose transport by increased GLUT4 translocation (22), activates insulin signaling, and upregulates molecules involved in fatty acid transport, fatty acid oxidation, and energy dissipation, resulting in decreased triglyceride content (20). In the liver, adiponectin increases  $\beta$ -oxidation and decreases gluconeogenesis (12).

Although earlier reports suggested that adiponectin is produced and secreted by the human placenta (23,24), more recent studies show that adiponectin mRNA is not expressed in the placenta (25,26). However, the placenta is a likely target for circulating adiponectin because AdipoR2 mRNA is expressed in the human placenta (23,27), and AdipoR2 protein has been localized to the syncytiotrophoblast (23). Both AdipoR1 and AdipoR2 proteins were expressed in cytotrophoblast cells freshly isolated from human placenta as well as in human primary trophoblast cells in culture (28). Apart from reports that adiponectin stimulates placental cytokine production (29), reduces the gene expression of GLUT isoform 3 and lipoprotein lipase in rat placenta (23), and inhibits endocrine function (28), effects of adiponectin on placental function remain unknown.

There are many observations in the literature implicating changes in placental amino acid transport in the regulation of fetal growth (30). For example, placental system A amino acid transport is downregulated prior to the onset of impaired fetal growth in a rat model of intrauterine growth restriction (31). We recently reported markedly increased placental system A amino acid transport capacity associated with fetal overgrowth in mice fed a high-fat diet (32). Amino acid transport system A is ubiquitously expressed and mediates the cellular uptake of small, neutral amino acids by cotransporting sodium. There are three isoforms of system A: SNAT1, SNAT2, and SNAT4 (30). In placental cells, the system A amino acid transporter is regulated by several hormones including insulin, cortisol, and leptin (30). Another key amino acid transporter is system L, which transports large neutral amino acids in a sodium-independent manner (33,34). Regulation of system L is dependent on which light chain is present, and studies show that regulation of large neutral amino acid transporter 1 involves protein kinase C or intracellular calcium concentrations (34). To the best of our knowledge, the regulation of amino acid transporter function by adiponectin has not been studied in any tissue. Interestingly, administration of gAd in pregnant rats was recently reported to decrease placental gene expression of GLUT isoform 3 and lipoprotein lipase (23), compatible with a role of maternal adiponectin in the regulation of placental nutrient transport.

The aim of this study was to determine the effects of gAd and fAd, with or without insulin, on intracellular signaling and amino acid transport in cultured primary human trophoblast cells. We tested the hypothesis that adiponectin downregulates placental amino acid transport mediated by inhibition of insulin signaling.

## RESEARCH DESIGN AND METHODS

**Cytotrophoblast isolation.** Isolation of cytotrophoblasts from normal-term placentas was based on the protocol published by Kliman et al. (35) and performed as we reported previously (36,37). Cells were plated on 35-mm

culture dishes ( $5 \times 10^6$  cells/dish) and six-well plates ( $2 \times 10^6$  cells/well) in 5% CO<sub>2</sub>, 95% atmosphere air at 37°C. Media were changed daily for 90 h. At 66 h after plating, cells were exposed to control media or control media plus gAd or fAd for 24 h. A subset of cells was pretreated with 1 nmol/l insulin for 4 h, gAd or fAd was subsequently added, and cells were incubated for an additional 20 h. At 90 h after plating, cells were used in amino acid uptake assays or expression studies, and culture media were collected for cytokine analysis. The concentration of insulin used corresponds to normal postprandial insulin levels in pregnant women (38).

**Small interfering RNA.** Dharmafect2 transfection reagent (Thermo Scientific, Rockford, IL) and small interfering RNA (siRNA), targeted against glyceraldehyde-3-phosphate dehydrogenase, signal transducer and activator of transcription 3 (STAT3), or a scrambled sequence, were added to cells after 20 h in culture, incubated for 24 h, and removed, and fresh media were added to wells (36). After a total of 66 h in culture, cells were treated with gAd or control media for 24 h and used in the amino acid uptake assay.

**Amino acid uptake assay.** The activity of the system A and L amino acid transporters was measured in cultured trophoblast cells as previously described (36,37). Cells in duplicate were incubated in buffers with and without Na<sup>+</sup> (iso-osmotic choline replacement) and in the presence and absence of 2-amino-2-norbornane-carboxylic acid (BCH; 64  $\mu$ mol/l). Radioactivity in cell lysates was counted in a liquid scintillation counter and mediated uptake calculated as pmol/(min  $\times$  mg protein) using standards with known amounts of isotope.

**Western blot.** Protein expression of the system A transporter isoforms SNAT1, SNAT2, SNAT4; phospho-AKT Ser473 and phospho-AKT Thr308; phospho-AMPK Thr172; AMPK; STAT3; phospho-STAT3 Tyr705; suppressor of cytokine signaling 3 (SOCS3); PPAR $\alpha$ ; phospho-PPAR $\alpha$  Ser21; insulin receptor substrate (IRS-1; Tyr612); and  $\beta$ -actin was analyzed using Western blotting. A polyclonal SNAT2 antibody was generated in rabbit by Dr. P.D. Prasad (Medical College of Georgia), and affinity-purified polyclonal anti-SNAT1 and -SNAT4 antibodies were generated in rabbits by Eurogentec (Seraing, Belgium). Antibodies targeted against phospho-AKT Ser473, phospho-AKT Thr308, phospho-AMPK, AMPK, SOCS3, and STAT3 were purchased from Cell Signaling, and anti- $\beta$ -actin antibodies were obtained from Sigma. Protein concentrations were determined by Bradford assay, and Western blotting was performed as previously described (36). Analysis of the blots was performed by densitometry using Alpha Imager (Alpha Innotech, San Leandro, CA).


**Real-time PCR.** After exposure of syncytiotrophoblast cells to insulin (1 nmol/l), globular adiponectin (5  $\mu$ g/ml), full-length adiponectin (5  $\mu$ g/ml), or control media for 24 h, total RNA was isolated using TRIzol reagent (Invitrogen, Carlsbad CA) and reverse transcribed using a Quantitect Reverse Transcription kit (Qiagen, Valencia, CA). Proprietary Quantitect Primer Assays for SNAT1, SNAT2, and SNAT4 and succinate dehydrogenase were obtained from Qiagen. Quantitative PCR was performed in triplicate in 96-well plates following the standard Quantitect SYBR PCR protocol in a 7300 (Applied Biosystems, Foster City, CA). For analysis, SNAT1, SNAT2, and SNAT4 were standardized against succinate dehydrogenase expression. As negative controls, preparations lacking RNA were used. RNAs were assayed from six independent biological replicates. The RNA levels are expressed as a ratio, using the comparative cycle threshold method for comparing relative expression results between treatments in real-time PCR.

**Cytokine enzyme-linked immunosorbent assays.** Syncytiotrophoblast cell culture media were collected after 24-h incubation with 5  $\mu$ g/ml gAd or fAd. Interleukin-6 (IL-6) and tumor necrosis factor- $\alpha$  concentrations in media were determined using colorimetric ELISA (Thermo Scientific), following instructions provided by the manufacturer.

**Data presentation and statistics.** Data are presented as means  $\pm$  SEM. Statistical significance of differences between control and treated cells was assessed using repeated-measures ANOVA or Student *t* test. A *P* value  $< 0.05$  was considered significant. *n* = number of individual placentas that cells were isolated from in each treatment group. Experiments were run on duplicate sets of cells from each placenta.

## RESULTS

The effect of adiponectin on amino acid uptake in cultured human primary trophoblast cells was highly dependent on the presence or absence of insulin. In the absence of insulin, gAd significantly (*P*  $< 0.05$  repeated-measures ANOVA [RMANOVA], *n* = 6) increased system A amino acid transporter activity, as measured by sodium-dependent methylaminoisobutyric acid (MeAIB) uptake (Fig. 1A). However, system L amino acid transport activity was not altered by gAd (Fig. 1C). Both MeAIB and leucine



**FIG. 1.** Sodium-dependent  $^{14}\text{C}$ -MeAIB (A and B) and BCH-inhibitable  $^3\text{H}$ -leucine (C and D) uptake after incubation of cultured trophoblast cells in control media, gAd (A and C), or fAd (B and D) for 24 h. Data are mean  $\pm$  SEM for cells isolated from six different placentas. gAd significantly ( $P < 0.05$ ) stimulated MeAIB uptake in a dose-dependent manner (RMANOVA with post hoc tests,  $*P < 0.05$ ,  $**P < 0.01$ ).

uptake in fAd-treated cells remained at control levels (Fig. 1B and D). Insulin in concentrations corresponding to postprandial levels in pregnant women (38) markedly increased ( $P < 0.05$ , RMANOVA,  $n = 6$ ) system A transport activity (Fig. 2A and B), in agreement with previous studies (39,40), with no effect on system L (data not shown). gAd did not modify the insulin-stimulated system A amino acid transport activity (Fig. 2A). However, fAd abolished the insulin-stimulated system A activity ( $P < 0.05$ , RMANOVA,  $n = 6$ ) (Fig. 2B).


Because altered mRNA and protein expression of specific SNAT isoforms, in particular SNAT2, constitutes a key mechanism by which placental system A activity is regulated (32,36), we determined the gene and protein expression of SNAT isoforms in response to insulin, gAd, and fAd. *SNAT1*, *SNAT2*, and *SNAT4* RNA expression levels were not altered by gAd (Fig. 2C). In contrast, insulin significantly increased mRNA expression of all SNAT isoforms (Fig. 2C). Furthermore, in the presence of insulin, fAd significantly decreased the expression levels of *SNAT2* and *SNAT4* mRNA, but not *SNAT1* RNA, compared with both fAd and insulin treatment alone (Fig. 2C). Indeed, *SNAT1*, *SNAT2*, and *SNAT4* mRNA levels after fAd treatment in the presence of insulin were not significantly different from control cells (Fig. 2C). A significant increase in SNAT2 protein expression was observed after insulin, gAd, and insulin/gAd incubations (Fig. 2D and E). Whereas fAd alone had no effect on SNAT2 protein expression, fAd completely reversed the insulin-stimulated increase in SNAT2 expression (Fig. 2E).

AMPK is one of the key mediators of adiponectin signaling in muscle and liver (15,16). We determined the

effect of gAd and fAd on AMPK activation, which is dependent on the phosphorylation of the Thr172 residue (41). Incubation with gAd alone for 24 h increased the level of phospho-AMPK, but this was not observed in the presence of insulin (Fig. 3). On the other hand, fAd did not alter the phospho-AMPK expression, in the presence or absence of insulin. fAd alone significantly increased the total expression of AMPK, but this effect was lost in the presence of insulin (Fig. 3).

It is well established that adiponectin may affect the synthesis/release of proinflammatory cytokines in other tissues (42–44). Furthermore, Lappas et al. (29) showed an increase in cytokine secretion after exposure of placental villous fragments to gAd. Because we recently reported that IL-6 and TNF- $\alpha$  stimulate system A transporter activity in cultured primary human trophoblast cells (36), it is possible that the effects of gAd on system A activity may be mediated by the release of proinflammatory cytokines. We demonstrated that gAd significantly increased the levels of IL-6 and TNF- $\alpha$  produced by the trophoblast cells (Fig. 4A and B). In contrast, fAd dramatically reduced the levels of IL-6 produced, whereas TNF- $\alpha$  levels were increased (Fig. 4C and D).

STAT3 is a key component in the IL-6 signaling pathway (45) and the stimulating effect of IL-6 on system A activity in cultured human primary trophoblast cells is mediated by STAT3 activation (36). To investigate whether the effects of gAd (without insulin) on system A transport were mediated via STAT3, we transfected cultured primary human trophoblast cells with siRNA targeting STAT3, resulting in a 70% reduction in STAT3 protein expression, (36). The significant stimulation of MeAIB




**FIG. 2.** Sodium-dependent  $^{14}\text{C}$ -MeAIB uptake after incubation of cultured trophoblast cells in control media, insulin (1 nmol/l), gAd (5  $\mu\text{g}/\text{ml}$ ) (A), or fAd (5  $\mu\text{g}/\text{ml}$ ) (B) for 24 h. Subsets of cells were pretreated with insulin (1 nmol/l) for 4 h and then exposed to 5  $\mu\text{g}/\text{ml}$  gAd (IgAd) or fAd (IfAd) for an additional 20 h. Data are mean  $\pm$  SEM for cells isolated from six different placentas. Insulin alone, gAd alone, and insulin + gAd ( $P < 0.01$ ) significantly stimulated MeAIB uptake; however, insulin + fAd significantly ( $P < 0.001$ ) reduced MeAIB uptake compared with insulin alone (RMANOVA with post hoc tests,  $*P < 0.05$ ,  $**P < 0.01$ ). C: Summary data of real-time PCR of system A amino acid transporter isoforms *SNAT1*, *SNAT2*, and *SNAT4* after incubation of cultured trophoblast cells as indicated. Data are mean  $\pm$  SEM for six placentas. Insulin significantly increased *SNAT* gene expression. In the presence of fAd, the effect of insulin on the mRNA expression of *SNAT2* and *SNAT4* was significantly reduced compared with both fAd and insulin treatment alone (RMANOVA,  $P < 0.01$ ; Tukey-Kramer multiple comparisons post tests,  $*P < 0.05$ ,  $**P < 0.01$ ). D: Representative Western blot of *SNAT1*, *SNAT2*, and *SNAT4* expression after incubation of cultured trophoblast cells as indicated. E: Summary of *SNAT2* protein expression levels.  $n = 6$  for each treatment, RMANOVA,  $P < 0.01$ ; Tukey-Kramer multiple comparisons post tests,  $*P < 0.05$ ,  $**P < 0.01$ .

uptake observed after gAd treatment alone was abolished in cells in which STAT3 was silenced (Fig. 5). These results clearly demonstrate that STAT3 is critical in mediating the effect of gAd on system A amino acid transport activity.

One key mechanism involved in the insulin-sensitizing effect of adiponectin in liver and muscle is the ability to enhance insulin-stimulated AKT phosphorylation (46). To investigate possible interactions between adiponectin and insulin signaling in trophoblast cells, we measured protein expression of phospho-AKT Ser473 and phospho-AKT Thr308 after incubation in insulin with or without adiponectin. Incubation of cultured primary trophoblast cells with insulin significantly increased the levels of phospho-AKT Ser473 (Fig. 6A and B); however, incubation with gAd or fAd with or without insulin did not alter the expression levels. As expected, phospho-AKT (Thr308) levels were increased by addition of insulin. In contrast, gAd or fAd


alone did not alter phospho-AKT Thr308 expression. Furthermore, addition of gAd to insulin-stimulated cells did not affect AKT phosphorylation at Thr308. In contrast, fAd completely reversed insulin-stimulated Thr308 phosphorylation of AKT (Fig. 6A and B).

We investigated intermediaries that may be involved in modulating the insulin-signaling pathway in response to fAd. Insulin resistance is commonly associated with increased expression of the p85 $\alpha$  subunit of phosphatidylinositol 3-kinase (47), phosphatase and tensin homolog (48), SOCS3 (49), and phospho-STAT3 (50), which all attenuate insulin signaling. However, in our study, protein expression of phosphatase and tensin homolog, SOCS3, phospho-STAT3, and p85 was not different after exposure to fAd plus insulin compared with insulin alone (data not shown). Adiponectin has been shown to activate IRS-1 (46), and we explored the possibility that fAd alters the activation status of IRS-1 by determining the expression of




**FIG. 3.** Summary of phospho-AMPK and AMPK protein expression after incubation of cultured trophoblast cells with control media, insulin (1 nmol/l), gAd (5  $\mu$ g/ml), or fAd (5  $\mu$ g/ml) for 24 h. Subsets of cells were pretreated with insulin (1 nmol/l) for 4 h and then exposed to 5  $\mu$ g/ml gAd (IgAd) or fAd (IfAd) for an additional 20 h.  $n = 5$  placentas for each treatment. Phospho-AMPK expression was significantly increased by gAd (RMANOVA,  $P < 0.01$ ). AMPK expression was significantly increased by fAd (RMANOVA,  $P < 0.002$ ). Tukey-Kramer multiple comparisons post tests, \* $P < 0.05$ .


IRS-1 phosphorylated at Tyr612. In cells treated with insulin plus fAd, phosphorylation of IRS-1 at Tyr612 was significantly reduced compared with insulin-treated cells (Fig. 7A and B). Because IRS-1 is activated when phos-



**FIG. 5.** A: Representative Western blot showing reduction in protein expression of STAT3 compared with control (C) using three unique siRNAs (1, 2, 3) targeted against STAT3. B: System A activity in control, siRNA STAT3 knockdown, and gAd-treated cells. Data are mean  $\pm$  SEM for six placentas. Knocking down STAT3 significantly (RMANOVA,  $P = 0.002$ ) reduced gAd stimulation of system A activity. Tukey-Kramer multiple comparisons post tests, \*\* $P < 0.01$ .



**FIG. 4.** A: IL-6 secretion was significantly (Student paired  $t$  test,  $P < 0.01$ ) increased after incubation of cultured trophoblast cells with gAd (1  $\mu$ g/ml). Data are mean  $\pm$  SEM for six placentas. B: TNF- $\alpha$  secretion was significantly (Student paired  $t$  test,  $P < 0.01$ ) increased after incubation of cultured trophoblast cells with gAd (1  $\mu$ g/ml). Data are mean  $\pm$  SEM for six placentas. C: IL-6 secretion was significantly (RMANOVA,  $P < 0.01$ ) reduced in a dose-dependent manner after incubation of cultured trophoblast cells with fAd (1  $\mu$ g/ml). Data are mean  $\pm$  SEM for six placentas. D: TNF- $\alpha$  secretion was significantly (RMANOVA,  $P < 0.01$ ) increased in a dose-dependent manner after incubation of cultured trophoblast cells with fAd. Data are mean  $\pm$  SEM for six placentas. \*\* $P < 0.01$ .



**FIG. 6.** Representative Western blots (A) and summary data (B) of phospho-AKT Ser473 and phospho-AKT Thr308 protein expression after incubation of cultured trophoblast cells with control media, insulin (1 nmol/l), gAd (5  $\mu$ g/ml), or fAd (5  $\mu$ g/ml) for 24 h. Subsets of cells were pretreated with insulin (1 nmol/l) for 4 h and then exposed to 5  $\mu$ g/ml gAd (IgAd) or fAd (IfAd) for an additional 20 h.  $n = 5$  placentas for each treatment. Insulin significantly increased expression of both phospho-AKT Ser473 ( $P < 0.01$ ) and Thr308 ( $P < 0.01$ ). Addition of fAd to insulin-stimulated cells significantly ( $P < 0.01$ ) reduced expression of phospho-AKT Thr308 compared with insulin treatment alone. Tukey-Kramer multiple comparisons post tests, \* $P < 0.05$ , \*\* $P < 0.01$ .

phorylated at this site, these data suggest that fAd inhibits IRS-1 activity in trophoblast cells.

PPAR $\alpha$  plays a critical role in the signal transduction pathway of adiponectin receptors, in particular AdipoR2 (16). Recent data show that tissue-specific overexpression of PPAR $\alpha$  in skeletal (51) and cardiac muscle (52) decreases insulin sensitivity, mediated by inhibition of insulin signaling (52). This raises the possibility that the inhibition of insulin signaling by fAd in trophoblast cells may be mediated by activation of PPAR $\alpha$ . In addition to ligand binding, PPAR $\alpha$  can be activated by at least two mechanisms: increased total expression and phosphorylation at Ser6, Ser12, and Ser21, which enhances PPAR $\alpha$  stability and decreases degradation (53,54). We observed a significant increase in the protein expression of PPAR $\alpha$  and phospho-PPAR $\alpha$  Ser21 after incubation of trophoblast cells in insulin plus fAd compared with control or insulin-treated cells (Fig. 7C and D).

## DISCUSSION


We demonstrate that, in contrast to the insulin-sensitizing actions of adiponectin in liver and muscle, fAd attenuates insulin signaling in primary human trophoblast cells. As a result, fAd inhibits insulin-stimulated amino acid transport. To the best of our knowledge, this is the first report of regulation of amino acid transport by adiponectin, in any tissue.

It is well established that insulin stimulates placental system A amino acid transport (39,40); however, the underlying mechanism has not previously been explored. We found that physiological levels of insulin increased the mRNA expression of all system A isoforms. In contrast, only SNAT2 was upregulated at the protein level, indicating that changes in SNAT2 protein expression are important in mediating the regulation of system A activity by insulin. These findings are in line with observations that SNAT2 appears to be a highly regulated SNAT isoform both in the placenta (31,32,55,56) and in other tissues and cells such as the mammary gland (56) and 3T3-L1 adipocytes (57).

Incubation of cultured human primary trophoblast cells in gAd in the absence of insulin increased AMPK phosphorylation, in agreement with previous studies in other tissues (15,16). Silencing of STAT3 completely abolished the stimulation of system A amino acid transporter activity by gAd. These data demonstrate that STAT3 activation is critical in mediating the stimulating effect of gAd on system A activity in the absence of insulin, findings that are in agreement with studies showing that STAT3 mediates the cellular effects of gAd in mouse cardiac fibroblast cells (58). STAT3 constitutes a key component in IL-6 signaling (45), and we show that gAd treatment enhanced trophoblast IL-6 secretion, observations that reflect a report by Lappas et al. demonstrating increased IL-6 production in placental villous fragments treated with gAd (29). Furthermore, we recently reported that IL-6 increases system A amino acid transport activity in cultured human primary trophoblast cells (36). Collectively, these observations suggest that gAd stimulates system A activity in cultured trophoblast cells by releasing IL-6, resulting in STAT3 activation. However, it cannot be excluded that mediators other than IL-6 could have contributed to the STAT3-mediated stimulation of system A in response to gAd. The molecular mechanisms linking AMPK activation, IL-6 release, and STAT3 phosphorylation in trophoblast cells remains to be established but may involve mitogen-activated protein kinase and nuclear factor- $\kappa$ B signaling as shown in other tissues (42–44,59).

The effect of gAd on trophoblast signaling and amino acid transport was distinctly different in the presence of physiological concentrations of insulin compared with incubations with gAd alone. Indeed, when insulin was present, gAd did not increase AMPK phosphorylation. Furthermore, gAd failed to enhance insulin-stimulated AKT phosphorylation. This is in contrast to effects in C2C12 myotubes in which gAd further enhanced insulin-stimulated AKT phosphorylation (46), but is consistent with effects in HeLa cells (46) and L6 muscle cells (60). Thus, the effects of gAd on the insulin-signaling pathway are tissue specific, and our data suggest that gAd is not an insulin sensitizer in human primary trophoblast cells.

Unlike gAd, fAd treatment caused a reduction in IL-6 secretion, consistent with an anti-inflammatory role for fAd in the placenta similar to that seen in monocytic cells (42,44). In contrast to the lack of effect of gAd treatment on insulin-stimulated cells, fAd treatment abolished the stimulation of system A transport activity and SNAT2 expression elicited by insulin. Importantly, our results demonstrate that this effect is mediated via cross-talk between the full-length adiponectin signaling pathway and the insulin-signaling pathway. In trophoblast cells stimulated by insulin, fAd significantly reduced the phosphorylation of both IRS-1 and AKT



**FIG. 7.** A: Representative Western blot of phospho-IRS-1 Tyr612 protein and  $\beta$ -actin expression after incubation of cultured trophoblast cells with control media (C), insulin (I; 1 nmol/l) for 24 or 4 h pretreatment with insulin (1 nmol/l) followed by fAd (5  $\mu$ g/ml) for 20 h. B: Summary of phospho-IRS-1 Tyr612 protein expression. RMANOVA,  $P < 0.01$ ,  $n = 5$  placentas for each treatment; Tukey-Kramer multiple comparisons post tests,  $**P < 0.01$ . C: Representative Western blots of PPAR $\alpha$ , phospho-PPAR $\alpha$  Ser21, and  $\beta$ -actin protein expression after incubation of cultured trophoblast cells with control media (C), insulin (I; 1 nmol/l) for 24 or 4 h pretreatment with insulin (1 nmol/l) followed by fAd (5  $\mu$ g/ml) for 20 h. D: Summary of PPAR $\alpha$  and phospho-PPAR $\alpha$  Ser21 protein expression. RMANOVA,  $P < 0.01$ ,  $n = 5$  placentas for each treatment; Tukey-Kramer multiple comparisons post tests,  $*P < 0.05$ .

compared with cells treated with insulin alone. The importance of AKT phosphorylation in mediating the effects of insulin on system A amino acid transport in L6 muscle cells was demonstrated by a recent report by Green et al. (61). Thus, our data show that fAd inhibits insulin signaling in cultured human primary trophoblast cells, which is in contrast to the insulin-sensitizing effect of fAd in skeletal muscle (15).

Activation of PPAR $\alpha$  may link fAd to inhibition of insulin signaling and system A amino acid transport. It has been demonstrated that PPAR $\alpha$  activation inhibits insulin/IGF-1-stimulated IRS-1 and AKT phosphorylation in several cell types (62,63). We demonstrated that treatment of cultured trophoblast cells with fAd in the presence of insulin led to an increase in both PPAR $\alpha$  protein expression and phosphorylated PPAR $\alpha$  Ser21 compared with either control or insulin-treated cells. Further studies are needed to establish an unequivocal cause-and-effect relationship between PPAR $\alpha$  activation and inhibition of insulin signaling in response to fAd and to identify the molecular mechanisms linking PPAR $\alpha$  to the insulin-signaling pathway.

One of the strengths of our study is that functional data (amino acid transport activity) have been obtained in primary human trophoblast cells in culture using concentrations of insulin and adiponectin within the physiological range, which contributes to the physiological relevance of our results. Because fAd was the only form of adiponectin that altered trophoblast function in the presence of physiological concentrations of insulin, we propose that the attenuation of placental insulin signaling and trophoblast

amino acid transport by fAd constitutes the important biological effect in vivo.

Our data show that fAd attenuated insulin signaling in primary human trophoblast cells and reversed insulin-stimulated system A activity. Interestingly, McDonald and Wolfe (28) recently reported that adiponectin inhibits endocrine functions of cultured human primary trophoblast cells, as evidenced by a decreased synthesis of human chorion gonadotropin and human placental lactogen. These authors did not identify the mechanisms involved; however, it is possible that inhibition of insulin signaling, as demonstrated in the current study, may mediate the inhibitory effect of adiponectin on trophoblast endocrine function because it is well established that insulin stimulates trophoblast production of hormones, including human placental lactogen (64) and human chorion gonadotropin (65).

Insulin had a marked effect on the response of cultured human trophoblast cells to gAd and fAd. There may be a multitude of mechanisms underlying the regulation of adiponectin responsiveness by insulin. Human trophoblast cells have been shown to express functional AdipoR1 and AdipoR2 (28). Because AdipoR1 binds primarily gAd and AdipoR2 binds fAd with higher affinity than gAd, it is possible that insulin affects the responsiveness to adiponectin by regulating the relative abundance of the two adiponectin receptors in trophoblast cells. In support of this hypothesis, insulin increased AdipoR2 expression threefold in muscle cells while reducing AdipoR1 expression by 50% (60), which was associated with an increased sensitivity to fAd and resistance to the effects of gAd

(60,66). Considering that both AdipoR1 and AdipoR2 are believed to mediate cellular effects by promoting increased insulin sensitivity, it may be expected that deletion of the receptors would result in insulin resistance. However, deletion of AdipoR2 in transgenic mice caused enhanced insulin sensitivity, rather than insulin resistance (67–69). One explanation for the unexpected phenotype of the AdipoR2 knockout mouse could be activation of AdipoR2 inhibits insulin signaling in some tissues, possibly including the placenta. Emerging evidence shows that downstream signaling of the adiponectin receptors involves APPL1 (adaptor protein containing pleckstrin homology domain) (70). Indeed, APPL1 associates with the adiponectin receptors and mediates adiponectin signaling and its effects on metabolism (71). APPL1 also functions in the insulin-signaling pathway and is an important mediator of adiponectin-dependent insulin sensitization in skeletal muscle (70). Chronic insulin treatment results in translocation of this signaling molecule to the nucleus, making it unavailable for downstream propagation of the adiponectin signal (70). Therefore, it is possible that the inhibitory effect of insulin on gAd AMPK activation in primary trophoblast cells is mediated through APPL1 translocation to the nucleus. Recently, APPL2, an isoform of APPL1, was identified (72) and shown to suppress adiponectin and insulin signaling by sequestering APPL1. We speculate that differences between placenta and muscle in the expression and/or function of APPL1 and APPL2 result in distinct responses to adiponectin.

In conclusion, fAd inhibits insulin-stimulated trophoblast system A amino acid transport. Our data indicate that the underlying mechanism involves activation of PPAR $\alpha$ , which inhibits IRS-1 and AKT phosphorylation, resulting in reversal of insulin-stimulated SNAT2 expression and system A activity. Fetal growth is highly dependent on the capacity of the placenta to transport amino acids (8). Because insulin and IGF-I are well-established stimulators of placental nutrient transporters and fetal growth, the finding that fAd attenuates placental insulin signaling and amino acid transport may have important implications for placental nutrient transport and fetal growth in pregnancies associated with altered maternal adiponectin levels.

#### ACKNOWLEDGMENTS

T.L.P. was supported by a grant from the National Institute of Child Health and Human Development (NICHD; R03HD058030). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NICHD or the National Institutes of Health.

No potential conflicts of interest relevant to this article were reported.

#### REFERENCES

- Catalano PM. Increasing maternal obesity and weight gain during pregnancy: the obstetric problems of plentitude. *Obstet Gynecol* 2007;110:743–744
- Lam MH, Wong GY, Lao TT. Reappraisal of neonatal clavicular fracture: relationship between infant size and risk factors. *J Reprod Med* 2002;47:903–908
- Boney CM, Verma A, Tucker R, Vohr BR. Metabolic syndrome in childhood: association with birth weight, maternal obesity, and gestational diabetes mellitus. *Pediatrics* 2005;115:e290–e296
- Jansson T, Ekstrand Y, Björn C, Wennergren M, Powell TL. Alterations in the activity of placental amino acid transporters in pregnancies complicated by diabetes. *Diabetes* 2002;51:2214–2219
- Jansson T, Wennergren M, Powell TL. Placental glucose transport and GLUT 1 expression in insulin-dependent diabetes. *Am J Obstet Gynecol* 1999;180:163–168
- Gaither K, Quraishi AN, Illsley NP. Diabetes alters the expression and activity of the human placental GLUT1 glucose transporter. *J Clin Endocrinol Metab* 1999;84:695–701
- Magnusson AL, Waterman LJ, Wennergren M, Jansson T, Powell TL. Triglyceride hydrolase activities and expression of fatty acid binding proteins in the human placenta in pregnancies complicated by intrauterine growth restriction and diabetes. *J Clin Endocrinol Metab* 2004;89:4607–4614
- Jansson T, Powell TL. Human placental transport in altered fetal growth: does the placenta function as a nutrient sensor? A review. *Placenta* 2006;27(Suppl.):91–97
- Jansson N, Nilsselt M, Gellerstedt M, Wennergren M, Rossander-Hulthén L, Powell TL, Jansson T. Maternal hormones linking maternal body mass index and dietary intake to birth weight. *Am J Clin Nutr* 2008;87:1743–1749
- Ramsay JE, Ferrell WR, Crawford L, Wallace AM, Greer IA, Sattar N. Maternal obesity is associated with dysregulation of metabolic, vascular, and inflammatory pathways. *J Clin Endocrinol Metab* 2002;87:4231–4237
- Ranheim T, Haugen F, Staff AC, Braekke K, Harsem NK, Drevon CA. Adiponectin is reduced in gestational diabetes mellitus in normal weight women. *Acta Obstet Gynecol Scand* 2004;83:341–347
- Kadowaki T, Yamauchi T. Adiponectin and adiponectin receptors. *Endocr Rev* 2005;26:439–451
- Waki H, Yamauchi T, Kamon J, Kita S, Ito Y, Hada Y, Uchida S, Tsuchida A, Takekawa S, Kadowaki T. Generation of globular fragment of adiponectin by leukocyte elastase secreted by monocytic cell line THP-1. *Endocrinology* 2005;146:790–796
- Yamauchi T, Kamon J, Ito Y, Tsuchida A, Yokomizo T, Kita S, Sugiyama T, Miyagishi M, Hara K, Tsunoda M, Murakami K, Ohteki T, Uchida S, Takekawa S, Waki H, Tsuno NH, Shibata Y, Terauchi Y, Froguel P, Tobe K, Koyasu S, Taira K, Kitamura T, Shimizu T, Nagai R, Kadowaki T. Cloning of adiponectin receptors that mediate antidiabetic metabolic effects. *Nature* 2003;423:762–769
- Yamauchi T, Kadowaki T. Physiological and pathophysiological roles of adiponectin and adiponectin receptors in the integrated regulation of metabolic and cardiovascular diseases. *Int J Obes* 2008;32:S13–S18
- Yamauchi T, Kamon J, Minokoshi Y, Ito Y, Waki H, Uchida S, Yamashita S, Noda M, Kita S, Ueki K, Eto K, Akanuma Y, Froguel P, Foufelle F, Ferre P, Carling D, Kimura S, Nagai R, Kahn BB, Kadowaki T. Adiponectin stimulates glucose utilization and fatty-acid oxidation by activating AMP-activated protein kinase. *Nat Med* 2002;8:1288–1295
- Weyer C, Funahashi T, Tanaka S, Hotta K, Matsuzawa Y, Pratley RE, Tataranni PA. Hypoadiponectinemia in obesity and type 2 diabetes: close association with insulin resistance and hyperinsulinemia. *J Clin Endocrinol Metab* 2001;86:1930–1935
- Tschritter O, Fritsche A, Thamer C, Haap M, Shirkavand F, Rahe S, Staiger H, Maerker E, Häring H, Stumvoll M. Plasma adiponectin concentrations predict insulin sensitivity of both glucose and lipid metabolism. *Diabetes* 2003;52:239–243
- Fruebis J, Tsao TS, Javorschi S, Ebbets-Reed D, Erickson MR, Yen FT, Bihain BE, Lodish HF. Proteolytic cleavage product of 30-kDa adipocyte complement-related protein increases fatty acid oxidation in muscle and causes weight loss in mice. *Proc Natl Acad Sci U S A* 2001;98:2005–2010
- Yamauchi T, Kamon J, Waki H, Terauchi Y, Kubota N, Hara K, Mori Y, Ide T, Murakami K, Tsuboyama-Kasaoka N, Ezaki O, Akanuma Y, Gavrilova O, Vinson C, Reitman ML, Kagechika H, Shudo K, Yoda M, Nakano Y, Tobe K, Nagai R, Kimura S, Tomita M, Froguel P, Kadowaki T. The fat-derived hormone adiponectin reverses insulin resistance associated with both lipodystrophy and obesity. *Nat Med* 2001;7:941–946
- Berg AH, Combs TP, Du X, Brownlee M, Scherer PE. The adipocyte-secreted protein Acrp30 enhances hepatic insulin action. *Nat Med* 2001;7:947–953
- Ceddia RB, Somwar R, Maida A, Fang X, Bikopoulos G, Sweeney G. Globular adiponectin increases GLUT4 translocation and glucose uptake but reduces glycogen synthesis in rat skeletal muscle cells. *Diabetologia* 2005;48:132–139
- Caminos JE, Nogueiras R, Gallego R, Bravo S, Tovar S, García-Caballero T, Casanueva FF, Diéguez C. Expression and regulation of adiponectin and receptor in human and rat placenta. *J Clin Endocrinol Metab* 2005;90:4276–4286
- Chen J, Tan B, Kareris E, Zervou S, Digby J, Hillhouse EW, Vatish M, Randeve HS. Secretion of adiponectin by human placenta: differential modulation of adiponectin and its receptors by cytokines. *Diabetologia* 2006;49:1292–1302
- Corbetta S, Bulfamante G, Cortelazzi D, Barresi V, Cetin I, Mantovani G, Bondioni S, Beck-Peccoz P, Spada A. Adiponectin expression in human



- fetal tissues during mid- and late gestation. *J Clin Endocrinol Metab* 2005;90:2397–2402
26. Pinar H, Basu S, Hotmire K, Laffineuse L, Presley L, Carpenter M, Catalano PM, Hauguel-de Mouzon S. High molecular mass multimer complexes and vascular expression contribute to high adiponectin in the fetus. *J Clin Endocrinol Metab* 2008;93:2885–2890
  27. Meller M, Qiu C, Vadachkoria S, Abetew DF, Luthy DA, Williams MA. Changes in placental adipocytokine gene expression associated with gestational diabetes mellitus. *Phys Res* 2006;55:501–512
  28. McDonald EA, Wolfe MW. Adiponectin attenuation of endocrine function within human term trophoblast cells. *Endocrinology* 2009;150:4358–4365
  29. Lappas M, Permezel M, Rice GE. Leptin and adiponectin stimulate the release of proinflammatory cytokines and prostaglandins from human placenta and maternal adipose tissue via nuclear factor-kappaB, peroxisomal proliferator-activated receptor-gamma and extracellularly regulated kinase 1/2. *Endocrinology* 2005;146:3334–3342
  30. Jones HN, Powell TL, Jansson T. Regulation of placental nutrient transport: a review. *Placenta* 2007;8–9:763–774
  31. Jansson N, Pettersson J, Haafiz A, Ericsson A, Palmberg I, Tranberg M, Ganapathy V, Powell TL, Jansson T. Down-regulation of placental transport of amino acids precedes the development of intrauterine growth restriction in rats fed a low protein diet. *J Physiol* 2006;576:935–946
  32. Jones HN, Woollett LA, Barbour N, Prasad PD, Powell TL, Jansson T. High-fat diet before and during pregnancy causes marked up-regulation of placental nutrient transport and fetal overgrowth in C57/BL6 mice. *FASEB J* 2009;23:271–278
  33. Pineda M, Fernández E, Torrents D, Estévez R, López C, Camps M, Lloberas J, Zorzano A, Palacín M. Identification of a membrane protein, LAT-2, that co-expresses with 4F2 heavy chain, an L-type amino acid transport activity with broad specificity for small and large zwitterionic amino acids. *J Biol Chem* 1999;274:19738–19744
  34. Okamoto Y, Sakata M, Ogura K, Yamamoto T, Yamaguchi M, Tasaka K, Kurachi H, Tsurudome M, Murata Y. Expression and regulation of 4F2hc and hLAT1 in human trophoblasts. *Am J Physiol* 2002;282:C196–C204
  35. Kliman HJ, Nestler JE, Sermasi E, Sanger JM, Strauss JF III. Purification, characterization, and in vitro differentiation of cytotrophoblasts from human term placentae. *Endocrinology* 1986;118:1567–1582
  36. Jones HN, Jansson T, Powell TL. IL-6 stimulates system A amino acid transporter activity in trophoblast cells through STAT3 and increased expression of SNAT2. *Am J Physiol Cell* 2009;297:C1228–C1235
  37. Roos S, Kanai Y, Prasad PD, Powell TL, Jansson T. Regulation of placental amino acid transporter activity by mammalian target of rapamycin. *Am J Physiol* 2009;296:C142–C150
  38. Phelps RL, Metzger BE, Freinkel N. Carbohydrate metabolism in pregnancy: XVII, diurnal profiles of plasma glucose, insulin, free fatty acids, triglycerides, cholesterol, and individual amino acids in late normal pregnancy. *Am J Obstet Gynecol* 1981;140:730–736
  39. Karl PI, Alpy KL, Fisher SE. Amino acid transport by the cultured human placental trophoblast: effect of insulin on AIB transport. *Am J Physiol* 1992;262:C834–C839
  40. Jansson N, Greenwood SL, Johansson BR, Powell TL, Jansson T. Leptin stimulates the activity of the system A amino acid transporter in human placental villous fragments. *J Clin Endocrinol Metab* 2003;88:1205–1211
  41. Hawley SA, Davison M, Woods A, Davies SP, Beri RK, Carling D, Hardie DG. Characterization of the AMP-activated protein kinase from rat liver and identification of threonine 172 as the major site at which it phosphorylates AMP-activated protein kinase. *J Biol Chem* 1996;271:27879–27887
  42. Haugen F, Drevon CA. Activation of nuclear factor-kappaB by high molecular weight and globular adiponectin. *Endocrinology* 2007;148:5478–5486
  43. Tang CH, Chiu YC, Tan TW, Yang RS, Fu WM. Adiponectin enhances IL-6 production in human synovial fibroblast via an AdipoR1 receptor, AMPK, p38, and NF-kappa B pathway. *J Immunol* 2007;179:5483–5492
  44. Abke S, Neumeier M, Weigert J, Wehrwein G, Eggenhofer E, Schäffler A, Maier K, Aslanidis C, Schölmerich J, Buechler C. Adiponectin-induced secretion of interleukin-6 (IL-6), monocyte chemoattractant protein-1 (MCP-1, CCL2) and interleukin-8 (IL-8, CXCL8) is impaired in monocytes from patients with type I diabetes. *Cardiovasc Diabetol* 2006;5:17–24
  45. Pedersen BK. IL-6 signalling in exercise and disease. *Biochem Soc Trans* 2007;35:1295–1297
  46. Wang C, Mao X, Wang L, Liu M, Wetzel MD, Guan KL, Dong LQ, Liu F. Adiponectin sensitizes insulin signaling by reducing p70 S6 kinase-mediated serine phosphorylation of IRS-1. *J Biol Chem* 2007;282:7991–7996
  47. Luo J, Field SJ, Lee JY, Engelman JA, Cantley LC. The p85 regulatory subunit of phosphoinositide 3-kinase down-regulates IRS-1 signaling via the formation of a sequestration complex. *J Cell Biol* 2005;170:455–464
  48. Leslie NR, Downes CP. PTEN function: how normal cells control it and tumour cells lose it. *Biochem J* 2004;382:1–11
  49. Howard JK, Flier JS. Attenuation of leptin and insulin signaling by SOCS proteins. *Trends Endocrinol Metab* 2006;17:365–371
  50. Kim K, Kim HY, Son EJ, Heo J, Cheong J. Oleic acid inhibits hepatic insulin signaling through deregulation of STAT3 activation and C/EBPalpha expression. *Cell Signal* 2009;21:1269–1276
  51. Finck BN, Bernal-Mizrachi C, Han DH, Coleman T, Sambandam N, LaRiviere LL, Holloszy JO, Semenkovich CF, Kelly DP. A potential link between muscle peroxisome proliferator-activated receptor-alpha signaling and obesity-related diabetes. *Cell Metab* 2005;1:133–144
  52. Park SY, Cho YR, Finck BN, Kim HJ, Higashimori T, Hong EG, Lee MK, Danton C, Deshmukh S, Cline GW, Wu JJ, Bennett AM, Rothmel B, Kalinowski A, Russell KS, Kim YB, Kelly DP, Kim JK. Cardiac-specific overexpression of peroxisome proliferator-activated receptor-alpha causes insulin resistance in heart and liver. *Diabetes* 2005;54:2514–2524
  53. Yoon MJ, Lee GY, Chung JJ, Ahn YH, Hong SH, Kim JB. Adiponectin increases fatty acid oxidation in skeletal muscle cells by sequential activation of AMP-activated protein kinase, p38 mitogen-activated protein kinase, and peroxisome proliferator-activated receptor alpha. *Diabetes* 2006;55:2562–2570
  54. Cho MC, Lee K, Paik SG, Yoon DY. Peroxisome proliferators-activated receptor (PPAR) modulators and metabolic disorders. *PPAR Res* 2008;2008:1–14
  55. Nelson DM, Smith SD, Furesz TC, Sadvovsky Y, Ganapathy V, Parvin CA, Smith CH. Hypoxia reduces expression and function of system A amino acid transporters in cultured term human trophoblasts. *Am J Physiol Cell Physiol* 2003;284:C310–C315
  56. López A, Torres N, Ortiz V, Alemán G, Hernández-Pando R, Tovar AR. Characterization and regulation of the gene expression of amino acid transport system A (SNAT2) in rat mammary gland. *Am J Physiol Endocrinol Metab* 2006;291:E1059–E1066
  57. Hatanaka T, Hatanaka Y, Tsuchida J, Ganapathy V, Setou M. Amino acid transporter ATA2 is stored at the trans-Golgi network and released by insulin stimulus in adipocytes. *J Biol Chem* 2006;281:39273–39284
  58. Liao W, Yu C, Wen J, Jia W, Li G, Ke Y, Zhao S, Campell W. Adiponectin induces interleukin-6 production and activates STAT3 in adult mouse cardiac fibroblasts. *Biol Cell* 2009;101:263–272
  59. Craig R, Larkin A, Mingo AM, Thuerlauf DJ, Andrews C, McDonough PM, Glembotski CC. p38 MAPK and NF-kappa B collaborate to induce interleukin-6 gene expression and release: evidence for a cytoprotective autocrine signaling pathway in a cardiac myocyte model system. *J Biol Chem* 2000;275:23814–23824
  60. Fang X, Palanivel R, Zhou X, Liu Y, Xu A, Wang Y, Sweeney G. Hyperglycemia- and hyperinsulinemia-induced alteration of adiponectin receptor expression and adiponectin effects in L6 myoblasts. *J Mol Endocrinol* 2005;35:465–476
  61. Green CJ, Göransson O, Kular GS, Leslie NR, Gray A, Alessi DR, Sakamoto K, Hundal HS. Use of Akt inhibitor and a drug-resistant mutant validates a critical role for protein kinase B/Akt in the insulin-dependent regulation of glucose and system A amino acid uptake. *J Biol Chem* 2008;283:27653–27667
  62. Martínez de Ubago M, García-Oya I, Pérez-Pérez A, Canfrán-Duque A, Quintana-Portillo R, Rodríguez de Fonseca F, González-Yanes C, Sánchez-Margalet V. Oleylethanolamide, a natural ligand for PPAR-alpha, inhibits insulin receptor signalling in HTC rat hepatoma cells. *Biochim Biophys Acta* 2009;1791:740–745
  63. Grabacka M, Plonka PM, Urbanska K, Reiss K. Peroxisome proliferator-activated receptor alpha activation decreases metastatic potential of melanoma cells in vitro via down-regulation of Akt. *Clin Cancer Res* 2006;12:3028–3036
  64. Fuglsang J, Møller N, Ledet T, Ovesen P. Effects of glucose, glycerol, 3-hydroxybutyrate, insulin, and leptin on placental growth hormone secretion in placental explants. *Horm Metab Res* 2008;40:189–193
  65. Ren SG, Braunstein GD. Insulin stimulates synthesis and release of human chorionic gonadotropin by choriocarcinoma cell lines. *Endocrinology* 1991;128:1623–1629
  66. Berg AH, Combs TP, Scherer PE. ACRP30/adiponectin: an adipokine regulating glucose and lipid metabolism. *Trends Endocrinol Metab* 2002;13:84–89
  67. Bjursell M, Ahnmark A, Bohlöly-Y M, William-Olsson L, Rhedin M, Peng XR, Ploj K, Gerdin AK, Arnerup G, Elmgren A, Berg AL, Oscarsson J, Lindén D. Opposing effects of adiponectin receptors 1 and 2 on energy metabolism. *Diabetes* 2007;56:583–593
  68. Yamauchi T, Nio Y, Maki T, Kobayashi M, Takazawa T, Iwabu M, Okada-Iwabu M, Kawamoto S, Kubota N, Kubota T, Ito Y, Kamon J,

- Tsuchida A, Kumagai K, Kozono H, Hada Y, Ogata H, Tokuyama K, Tsunoda M, Ide T, Murakami K, Awazawa M, Takamoto I, Froguel P, Hara K, Tobe K, Nagai R, Ueki K, Kadowaki T. Targeted disruption of AdipoR1 and AdipoR2 causes abrogation of adiponectin binding and metabolic actions. *Nat Med* 2007;13:332–339
69. Liu Y, Michael MD, Kash S, Bensch WR, Monia BP, Murray SF, Otto KA, Syed SK, Bhanot S, Sloop KW, Sullivan JM, Reifel-Miller A. Deficiency of adiponectin receptor 2 reduces diet-induced insulin resistance but promotes type 2 diabetes. *Endocrinology* 2007;148:683–692
70. Deepa SS, Dong LQ. APPL1: role in adiponectin signaling and beyond. *Am J Physiol Endocrinol Metab* 2009;296:E22–E36
71. Mao X, Kikani CK, Riojas RA, Langlais P, Wang L, Ramos FJ, Fang Q, Christ-Roberts CY, Hong JY, Kim RY, Liu F, Dong LQ. APPL1 binds to adiponectin receptors and mediates adiponectin signalling and function. *Nat Cell Biol* 2006;8:516–523
72. Wang C, Xin X, Xiang R, Ramos FJ, Liu M, Lee HJ, Chen H, Mao X, Kikani CK, Liu F, Dong LQ. Yin-Yang regulation of adiponectin signaling by APPL isoforms in muscle cells. *J Biol Chem* 2009;284:31608–31615