

Aerobic Exercise Overcomes the Age-Related Insulin Resistance of Muscle Protein Metabolism by Improving Endothelial Function and Akt/Mammalian Target of Rapamycin Signaling

Satoshi Fujita,^{1,2} Blake B. Rasmussen,^{3,4} Jerson G. Cadenas,^{1,2} Micah J. Drummond,³ Erin L. Glynn,³ Fred R. Sattler,² and Elena Volpi^{1,2}

Muscle protein metabolism is resistant to insulin's anabolic effect in healthy older subjects. This is associated with reduced insulin vasodilation. We hypothesized that aerobic exercise restores muscle protein anabolism in response to insulin by improving vasodilation in older subjects. We measured blood flow, endothelin-1, Akt/mammalian target of rapamycin (mTOR) signaling, and muscle protein kinetics in response to physiological local hyperinsulinemia in two groups of older subjects following a bout of aerobic exercise (EX group: aged 70 ± 2 years; 45-min treadmill walk, 70% heart rate max) or rest (CTRL group: aged 68 ± 1 years). Baseline endothelin-1 was lower and blood flow tended to be higher in the EX group, but protein kinetics was not different between groups. Insulin decreased endothelin-1 ($P < 0.05$) in both groups, but endothelin-1 remained higher in the CTRL group ($P < 0.05$) and blood flow increased only in the EX group (EX group: 3.8 ± 0.7 to 5.3 ± 0.8 ; CTRL group: 2.5 ± 0.2 to 2.6 ± 0.2 ml \cdot min⁻¹ \cdot 100 ml leg⁻¹). Insulin improved Akt phosphorylation in the EX group and increased mTOR/S6 kinase 1 phosphorylation and muscle protein synthesis (EX group: 49 ± 11 to 89 ± 23 ; CTRL group: 58 ± 8 to 57 ± 12 nmol \cdot min⁻¹ \cdot 100 ml leg⁻¹) in the EX group only ($P < 0.05$). Because breakdown did not change, net muscle protein balance became positive only in the EX group ($P < 0.05$). In conclusion, a bout of aerobic exercise restores the anabolic response of muscle proteins to insulin by improving endothelial function and Akt/mTOR signaling in older subjects. *Diabetes* 56:1615–1622, 2007

From the ¹Department of Internal Medicine, University of Texas Medical Branch, Galveston, Texas; the ²Department Medicine, University of Southern California, Los Angeles, California; the ³Department of Physical Therapy, University of Texas Medical Branch, Galveston, Texas; and the ⁴Department of Kinesiology and Biological Sciences, University of Southern California, Los Angeles, California.

Address correspondence and reprint requests to Elena Volpi, MD, PhD, Division of Geriatric Medicine and Sealy Center on Aging, University of Texas Medical Branch, 301 University Blvd., Galveston, TX 77555-0460. E-mail: evolpi@utmb.edu.

Received for publication 9 November 2006 and accepted in revised form 1 March 2007.

Published ahead of print at <http://diabetes.diabetesjournals.org> on 9 March 2007. DOI: 10.2337/db06-1566.

4EBP1, 4E-binding protein 1; FSR, fractional synthetic rate; ICG, indocyanine green; mTOR, mammalian target of rapamycin; S6K1, S6 kinase 1.

© 2007 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Sarcopenia is an age-dependent loss of skeletal muscle mass, strength, and quality, which may lead to weakness of the lower extremities, slowing of gait speed, and increased risk of falls (1–3). Sarcopenia is a multifactorial disorder, and evidence is accumulating that a reduced response of skeletal muscle to anabolic stimuli is an important contributing factor. For example, we and others (4,5) have previously shown that healthy aging is associated with a blunted muscle protein anabolic response to hyperaminoacidemia with hyperinsulinemia. Nonetheless, muscle protein synthesis and breakdown in the fasting state may not necessarily change with age (6) and may respond normally to a simple amino acid stimulus (7–10).

Insulin is a potent anabolic stimulus for skeletal muscle. We have recently shown that physiological hyperinsulinemia increases skeletal muscle protein synthesis and anabolism in young healthy subjects, as long as blood flow and amino acid delivery to the muscle are stimulated by insulin (11). Thus, it appears that the insulin-induced modulation of muscle perfusion and nutrient availability is necessary for the anabolic response of muscle protein synthesis to insulin. Furthermore, insulin and amino acids both can stimulate the mammalian target of rapamycin (mTOR) signaling pathway (12). Specifically, insulin promotes the phosphorylation of Akt, an upstream regulator of mTOR, and enhances mTOR signaling to its downstream effectors 4E-binding protein 1 (4EBP1) and ribosomal S6 kinase 1 (S6K1), which are key regulators of translation initiation and protein synthesis (13). On the other hand, amino acids (in particular leucine) can directly activate mTOR kinase activity, resulting in an even more potent stimulation of translation initiation, elongation, and protein synthesis (13).

Healthy aging is also associated with a decreased insulin-induced vasodilation due to dysfunction of the endothelial-dependent vasodilation (14–16), which is normally initiated by insulin via activation of the endothelial NO synthase (17,18). In a recent study of healthy older subjects, we showed that such a vasomotor dysfunction is associated with increased endothelin-1 concentrations and a selective resistance of skeletal muscle protein synthesis to the anabolic action of insulin (16). This novel finding raises the question as to whether interventions that are known to improve endothelial-dependent vasodilation

TABLE 1
Characteristics of the study subjects

Subject characteristics	Control	Exercise	<i>P</i>
<i>n</i>	7	6	
Sex (men/women)	5/2	5/1	0.61
AGE	70 ± 2	68 ± 1	0.52
Body weight (kg)	80 ± 3	85 ± 3	0.41
Height (cm)	172 ± 5	180 ± 2	0.37
BMI (kg/m ²)	27.7 ± 1.5	26.7 ± 1.1	0.73
Leg volume (ml)	9,547 ± 460	9,697 ± 414	0.87

Data are means ± SE.

and/or insulin sensitivity may also improve muscle protein synthesis in response to insulin in older subjects.

Aerobic exercise improves endothelial function in older subjects (15) and is also important in the prevention and treatment of type 2 diabetes as it increases insulin sensitivity and glucose tolerance (19–21). These positive effects may be demonstrated in both patients with type 2 diabetes and normal subjects even after a single bout of aerobic exercise and persist for at least 48 h after exercise (21–27). On the other hand, as opposed to progressive resistance exercise, moderate or even strenuous aerobic exercise has only minimal direct effects on muscle protein metabolism, because it stimulates muscle protein synthesis only during the exercise bout, and these effects dissipate in the immediate postexercise period (28).

We hypothesized that a single bout of aerobic exercise performed before an insulin challenge will significantly improve endothelial function and insulin-induced vasodilation and, as a result, will improve the anabolic response of skeletal muscle protein synthesis to insulin in healthy older subjects. To test this hypothesis, we measured leg blood flow, endothelin-1 concentrations, phosphorylation of the Akt/mTOR signaling pathway, and muscle protein kinetics in healthy, nondiabetic, untrained older subjects at baseline and during an insulin challenge that was preceded by either a bout of moderate aerobic exercise (~20 h before the beginning of the measurements) or rest.

RESEARCH DESIGN AND METHODS

We studied 13 older subjects (10 men and 3 women) from the Los Angeles metropolitan area and the greater Houston/Galveston area. All subjects provided informed written consent before participating in the study, which was approved by the institutional review boards of the University of Southern California (Los Angeles, CA) and the University of Texas Medical Branch (Galveston, TX).

All subjects were healthy, nondiabetic, and physically active (no impairments in the activities of daily living or instrumental activities of daily living) but were not engaged in an exercise training program. Screening was performed with clinical history, physical examination, and laboratory tests including blood count, liver and kidney function tests, coagulation profile, fasting blood glucose, oral glucose tolerance tests, hepatitis B/C and HIV screening, thyroid-stimulating hormone, lipid profile, urinalysis, drug screening, and resting electrocardiogram. Subjects with normal results were randomly assigned to either the aerobic exercise group (EX group: 45-min treadmill walk at 70% of maximum heart rate [HR_{max}]) performed the afternoon before the experiment) or a control group (CTRL group) not exercising on the day before the experiment. Subjects randomized to the EX group underwent exercise stress testing at least 1 week before the experiment to exclude individuals with silent cardiac ischemia and measure HR_{max} and oxygen consumption ($V_{O_{2max}}$). Characteristics of subjects meeting eligibility criteria are summarized in Table 1.

We measured blood flow, muscle protein, amino acid, and glucose kinetics in the postabsorptive basal state (0–300 min) and during insulin infusion (300–480 min) in the EX and CTRL groups. The afternoon before the study, each subject was admitted to the University of Southern California or the University of Texas Medical Branch General Clinical Research Center. At




FIG. 1. Study design. Blood and muscle sampling are indicated by arrows. A detailed description of the study design is provided in the text.

1600 h, subjects assigned to the EX group performed a 45-min treadmill walk at 70% HR_{max} . Briefly, the heart rate was continuously monitored throughout the exercise bout, while treadmill speed and inclination were adjusted to reach and maintain the heart rate at 70% HR_{max} . At 1830 h, all subjects were fed a standard dinner (one-third of their estimated daily caloric requirements) and given a snack at 2200 h, after which only water ad libitum was allowed until the end of the experiment. The next morning, polyethylene catheters were inserted into a forearm vein for tracer and dextrose infusion, in a contralateral hand or wrist vein for arterialized blood sampling, and in the femoral artery and vein of one leg for blood sampling. The arterial catheter was also used for infusion of insulin (Humulin R; Eli Lilly, Indianapolis, IN) and indocyanine green (ICG) (Akorn, Buffalo Grove, IL).

At 0730 h, after drawing a blood sample to measure background phenylalanine enrichment and ICG concentration, a primed ($2 \mu\text{mol/kg}$) continuous infusion of L-[ring-¹³C₆]phenylalanine ($0.05 \mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$; Cambridge Isotope Laboratories, Andover, MA) was started and maintained at a constant rate until the end of the experiment (Fig. 1). After 2 h, the first muscle biopsy was taken from the lateral portion of the vastus lateralis of the leg with the femoral catheters, using a 5-mm Bergström biopsy needle, aseptic procedure, and local anesthesia with 1% lidocaine injected subcutaneously and on the muscle fascia. The muscle sample (150–400 mg) was quickly rinsed with ice-cold saline, blotted, and immediately frozen in liquid nitrogen and stored at -80°C until analysis. At 230 min, continuous ICG infusion was started in the femoral artery (0.5 mg/min) and maintained until 270 min. During ICG infusion, blood samples were taken four times, at 10-min intervals, from femoral and hand veins to measure ICG concentration. Subsequently, between 270 and 300 min, four blood samples were taken from the femoral artery and vein to measure glucose concentrations and phenylalanine concentrations and enrichments; four other samples were drawn from hand and femoral veins to measure systemic and femoral insulin concentration, respectively. At 300 min, a second muscle biopsy was taken as previously described.

Immediately after the second biopsy, an insulin infusion was initiated directly into the femoral artery ($0.15 \text{ mU} \cdot \text{min}^{-1} \cdot 100 \text{ ml leg}^{-1}$) to expose the leg muscle to postprandial insulin concentrations. This technique allowed us to avoid a major systemic hyperinsulinemia and the consequent dramatic reduction in blood amino acid concentrations (11,29). During insulin infusion, blood samples (0.5 ml) were taken every 5–10 min to monitor plasma glucose concentration, and 20% dextrose was infused at a variable rate as necessary to maintain plasma glucose at the basal value (euglycemic clamp).

Between 410 and 450 min, ICG was again infused to measure leg blood flow, and blood samples were taken between 420 and 450 min and 450 and 480 min to measure ICG concentrations, phenylalanine and glucose enrichments and concentrations, and insulin concentrations, as described for the basal period. At 480 min, before stopping the tracer and insulin infusion, a third muscle biopsy was taken as described above.

Analyses. Plasma insulin and endothelin-1 concentrations were determined by enzyme-linked immunosorbent assay (Linco Research, St. Charles, MO). Plasma glucose concentration was measured using an automated glucose analyzer (YSI, Yellow Springs, OH). Serum ICG concentration was measured spectrophotometrically (Beckman Coulter, Fullerton, CA) at $\lambda = 805 \text{ nm}$ (30,31).

Blood phenylalanine concentrations and enrichments were determined by gas chromatography–mass spectrometry (Agilent Technologies, Palo Alto,

CA) as previously described (32). We measured only phenylalanine concentration because it is a good predictor of insulin-induced concentration changes in all essential amino acids (33).

Muscle tissue free amino acids and proteins were extracted as previously described (32). Intracellular free phenylalanine concentrations and enrichments of were determined by gas chromatography–mass spectrometry as previously described (32). Mixed muscle protein–bound phenylalanine enrichment was analyzed by gas chromatography–mass spectrometry after protein hydrolysis and amino acid extraction (32) using the external standard curve approach (34).

Total and phosphorylated Akt, mTOR, 4EBP1, and S6K1 in skeletal muscle samples were determined by SDS-PAGE and immunoblotting (BioRad, Hercules, CA) as previously described (35). The phospho and total primary antibodies used are listed in Fig. 3. Total content for each measured protein was detected using an antibody dilution of 1:1,000. Phosphorylation and total protein were normalized to a rodent standard (Precision Plus protein standard; BioRad). Final data were expressed as normalized protein phosphorylation relative to normalized total protein.

Calculations. Muscle phenylalanine kinetic parameters were calculated using both the two- and the three-pool arteriovenous balance models because each model provides unique information regarding leg plasma (two-pool) and intracellular (three-pool) amino acid kinetics (11). The models' assumptions and validation are extensively reviewed (36). The two- and three-pool model parameters were calculated as follows.

$$\text{Phenylalanine delivery to the leg} = F_{\text{in}} = \text{BF} \times C_A \quad (1)$$

$$\text{Phenylalanine output from the leg} = F_{\text{out}} = \text{BF} \times C_V \quad (2)$$

$$\text{Net balance} = \text{NB} = \text{BF} \times (C_A - C_V) \quad (3)$$

$$\text{Total leg rate of appearance} = \text{total } R_a = \text{BF} \times C_A \times E_A/E_V \quad (4)$$

$$\text{Release in blood from proteolysis} = \text{leg } R_a = \text{total } R_a - F_{\text{in}} = \text{BF} \times C_A \times [(E_A/E_V) - 1] \quad (5)$$

$$\text{Rate of disappearance from blood} = \text{leg } R_d = \text{leg } R_a + \text{NB} = \text{BF} \times [(C_A \times E_A/E_V) - C_V] \quad (6)$$

$$\text{Muscle inward transport} = F_{M,A} = [(C_V \times \{E_M - E_V\}/\{E_A - E_M\}) + C_V] \times \text{BF} \quad (7)$$

$$\text{Muscle outward transport} = F_{V,M} = [(C_V \times \{E_M - E_V\}/\{E_A - E_M\}) + C_A] \times \text{BF} \quad (8)$$

$$\text{Arteriovenous shunting} = F_{V,A} = F_{\text{in}} - F_{M,A} \quad (9)$$

$$\text{Release from muscle protein breakdown} = F_{M,0} = F_{M,A} \times [(E_A/E_M) - 1] \quad (10)$$

$$\text{Utilization for muscle protein synthesis} = F_{0,M} = F_{M,0} + \text{NB} \quad (11)$$

$$\text{Intracellular amino acid availability} = F_{M,A} + F_{M,0} \quad (12)$$

where C_A , C_V is plasma phenylalanine concentrations in the femoral artery and vein, respectively. E_A , E_V , and E_M are free phenylalanine enrichments (tracer-to-tracee ratio) in femoral artery and vein and in muscle, respectively. BF is leg blood flow calculated from steady-state ICG concentrations (30,31). Data were expressed per 100 ml of leg volume.

We also measured the fractional synthetic rate (FSR) of mixed muscle proteins by measuring the incorporation rate of the phenylalanine tracer into the proteins ($\Delta E_p/t$) and using the precursor-product model to calculate the synthesis rate as follows (37): $\text{FSR} = (\Delta E_p/t)/[(E_{M(1)} + E_{M(2)})/2] \times 60 \times 100$. ΔE_p is the increment in protein-bound phenylalanine enrichment between two sequential biopsies, t is the time between the two sequential biopsies, $E_{M(1)}$ and $E_{M(2)}$ are the phenylalanine enrichments in the free intracellular pool in the two sequential biopsies. Data are expressed in the percentage per hour.

Leg glucose utilization was calculated as net glucose uptake across the leg: $\text{leg glucose uptake} = \text{BF} \times (C_A - C_V)$.

To determine the degree of muscle tissue exposure to insulin, we calculated the insulin delivery rate to the leg to account for recycling from the systemic circulation (16). Further, changes in blood flow can affect insulin concentration during constant infusion. Because insulin was infused in the artery, arterial insulin concentration was not measurable, and insulin delivery was estimated using the femoral insulin concentration (Ins_{FV}) as follows: $\text{insulin delivery} = \text{Ins}_{FV} \times \text{BF}$.

Since some insulin is taken up by the muscle cells after binding to the insulin receptor, this method may slightly underestimate the total insulin delivery rate, but for the above reasons we found it preferable to simply relying on the calculated dose.

Statistical analysis. Sex differences between groups were analyzed using the χ^2 test. Subjects' characteristics and baseline values for all measured variables were analyzed using one-way ANOVA. The effects of the prior bout of exercise on the response variables of the basal period and during insulin infusion were compared using ANOVA for repeated measures, the main effects being group (the EX and CTRL groups) and time (basal and insulin infusion). Post hoc testing was performed using the Tukey-Kramer test. If baseline values were significantly different or tended to be different ($P < 0.10$), we also performed an ANCOVA using baseline values as covariates. Differences were considered significant at $P < 0.05$. P for trend was set at $P < 0.10$. Data are the means \pm SE.

RESULTS

Insulin and glucose. Blood glucose and insulin concentrations and kinetics across the leg are shown in Table 2. Systemic and femoral vein insulin concentrations were not different at baseline and significantly increased in both groups during insulin infusion, but the change was larger in the CTRL group ($P < 0.05$). However, because of the

TABLE 2
Endothelial function parameters, insulin and glucose concentrations, and kinetics at baseline and during insulin infusion

	CTRL group		EX group		P		
	Basal	Insulin	Basal	Insulin	Group	Time	Interaction
Endothelial function							
Endothelin-1 (pg/ml)	3.53 \pm 0.72	2.46 \pm 0.39	1.85 \pm 0.08	1.60 \pm 0.10	<0.01	0.02	0.14
Blood flow (ml \cdot min ⁻¹ \cdot 100 ml leg ⁻¹)	2.6 \pm 0.2	2.7 \pm 0.2	3.8 \pm 0.7	5.3 \pm 0.8	0.02	<0.01	<0.01
Insulin							
Systemic concentration (μ U/ml)	8.5 \pm 1.0	24.0 \pm 2.4	6.3 \pm 0.8	15.0 \pm 1.2	<0.01	<0.0001	0.06
Femoral vein concentration (μ U/ml)	8.5 \pm 1.0	83.4 \pm 5.2	6.3 \pm 0.8	46.7 \pm 5.1	<0.001	<0.0001	0.001
Delivery to the leg (μ U \cdot min ⁻¹ \cdot 100 ml leg ⁻¹)	22 \pm 3	218 \pm 11	23 \pm 4	255 \pm 57	0.48	<0.0001	0.45
Glucose							
Femoral artery concentration (mmol/l)	4.7 \pm 0.2	4.5 \pm 0.2	4.9 \pm 0.1	5.0 \pm 0.2	0.19	0.44	0.36
Infusion rate (μ mol \cdot kg ⁻¹ \cdot min ⁻¹)*	—	10.4 \pm 0.8	—	13.6 \pm 0.9	0.02	—	—
Leg uptake (μ mol \cdot min ⁻¹ \cdot 100 ml leg ⁻¹)*	—	1.8 \pm 0.4	—	3.6 \pm 0.6	0.03	—	—

Data are means \pm SE. Statistical analysis performed using ANOVA with repeated measures for all parameters except those marked with *, which were analyzed using one-way ANOVA. Boldface data indicate statistical significance.




FIG. 2. Plasma endothelin-1 concentrations, blood flow, and muscle protein FSR in two groups of older subjects at baseline and during insulin infusion performed ~20 h following rest (Control: $n = 7$) or aerobic exercise (Exercise: $n = 6$). Values are means \pm SE. * $P < 0.05$ vs. basal, # $P < 0.05$ vs. control. □, basal; ■, insulin.

differences in leg blood flow (see below), insulin delivery to the leg was not different between groups at baseline and significantly increased during insulin infusion in both groups, with no differences between groups. Plasma arterial glucose concentrations were not different between groups at baseline and did not change during insulin infusion with euglycemic clamp. However, the exogenous glucose infusion rate and leg glucose uptake during insulin infusion were significantly higher in the EX group ($P < 0.05$).

Endothelin-1 concentrations and leg blood flow. Plasma endothelin-1 concentrations in the femoral vein and leg blood flow are shown in Table 2 and Fig. 2. Endothelin-1 concentrations were significantly lower ($P <$

0.05) in the EX group at baseline and significantly decreased during insulin infusion in both groups ($P = 0.02$) with no time-by-group interaction. Leg blood flow at baseline tended to be higher in the EX than CTRL group ($P = 0.08$). In response to insulin, leg blood flow significantly increased only in the EX group ($P < 0.05$), while no change was observed in the CTRL group. ANCOVA using baseline values as covariates confirmed the results for both blood flow and endothelin-1.

Phenylalanine concentrations and enrichments. Average phenylalanine concentrations in the femoral artery and vein and in the muscle are reported in Table 3. Phenylalanine concentrations in the artery and in the vein were not different at baseline and slightly, but significantly, decreased in both groups with no differences between groups. Phenylalanine concentrations in muscle were significantly lower in the EX group at baseline ($P < 0.01$) and decreased during insulin infusion in both groups with no time-by-group interaction. ANCOVA using the baseline values as covariates confirmed these results. Phenylalanine enrichments in the femoral artery and vein and in the muscle tissue were not different between groups in the basal period and were at steady state during both sampling periods (data not shown). Phenylalanine enrichment in the femoral artery and vein increased slightly but significantly ($P < 0.01$) during insulin infusion in both groups, with no group differences. Phenylalanine enrichment in the muscle tissue tended to increase in both groups ($P < 0.06$) with no significant group differences either at baseline or during insulin infusion.

Amino acid kinetics. Leg and muscle phenylalanine kinetics are shown in Table 4. All kinetic parameters were not significantly different between groups in the basal period. With insulin infusion, phenylalanine delivery to the leg increased in the EX group only ($P < 0.01$). Phenylalanine output from the leg increased in the EX group and decreased in the CTRL group, resulting in a significant treatment-by-group interaction ($P < 0.01$). Because of a trend for basal phenylalanine delivery and output to be higher in the EX group ($P = 0.07$), an ANCOVA was carried out for both parameters using the basal values as covariates and confirmed the ANOVA results. Phenylalanine leg R_a significantly decreased in the CTRL group ($P = 0.04$), whereas no change was observed in the EX group. In contrast, phenylalanine leg R_a increased only in the EX group during insulin infusion ($P < 0.01$), whereas it did not change in the CTRL group.

Muscle phenylalanine inward and outward transport decreased in the CTRL group and increased in the EX group, resulting in significant treatment-by-group interactions ($P < 0.05$). ANCOVA performed on these two parameters, because of a trend for the baseline values to be higher in the EX group ($P = 0.07$), confirmed the ANOVA results. Phenylalanine intracellular availability increased with insulin in the EX group, whereas it decreased in the

TABLE 3
Phenylalanine concentrations at baseline and during insulin infusion

	CTRL group		EX group		<i>P</i>		
	Basal	Insulin	Basal	Insulin	Group	Time	Interaction
Femoral artery ($\mu\text{mol/l}$)	61 \pm 2	56 \pm 2	62 \pm 2	59 \pm 2	0.31	<0.001	0.11
Femoral vein ($\mu\text{mol/l}$)	67 \pm 2	56 \pm 1	68 \pm 1	58 \pm 2	0.59	<0.0001	0.65
Muscle tissue ($\mu\text{mol/l}$)	127 \pm 11	95 \pm 11	85 \pm 5	81 \pm 8	0.02	0.05	0.12

Data are means \pm SE. Boldface data indicate statistical significance.

TABLE 4
Phenylalanine kinetics at baseline and during insulin infusion

	CTRL group		EX group		Group	Time	Interaction
	Basal	Insulin (nmol · min ⁻¹ · 100 ml leg ⁻¹)	Basal	Insulin (nmol · min ⁻¹ · 100 ml leg ⁻¹)			
Delivery to the leg	156 ± 12	145 ± 8	234 ± 43	309 ± 47	0.01	<0.01	<0.01
Output from the leg	171 ± 13	149 ± 10	255 ± 46	300 ± 45	0.02	0.23	<0.01
Release in the blood from proteolysis (R _a)	63 ± 10	44 ± 9	66 ± 14	70 ± 19	0.32	0.21	0.04
Disappearance from the blood (R _d)	48 ± 9	40 ± 7	45 ± 11	79 ± 19	0.22	0.02	<0.01
Muscle inward transport	113 ± 20	75 ± 11	193 ± 46	250 ± 47	0.01	0.47	0.02
Muscle outward transport	128 ± 21	79 ± 11	214 ± 48	241 ± 45	0.76	0.62	0.05
Arteriovenous shunting	43 ± 11	70 ± 12	41 ± 13	59 ± 28	0.03	0.06	0.77
Intracellular availability	186 ± 27	136 ± 21	263 ± 58	330 ± 60	0.03	0.49	<0.01
Release from muscle protein breakdown	73 ± 9	61 ± 14	70 ± 14	80 ± 23	0.55	0.95	0.18
Utilization for muscle protein synthesis	58 ± 8	57 ± 12	49 ± 11	89 ± 23	0.44	0.02	0.03
Net balance	-15 ± 2	-2 ± 2	-21 ± 4	9 ± 4	0.43	<0.0001	<0.01

Data are means ± SE. Boldface data indicate statistical significance.

CTRL group, resulting in a significant time-by-group interaction ($P < 0.01$). Phenylalanine arteriovenous shunting increased significantly in both groups with no group differences. Phenylalanine release from proteolysis did not change significantly with insulin infusion in either group. Phenylalanine utilization for muscle protein synthesis increased significantly during insulin infusion in the EX group ($P = 0.03$), but it did not change in the CTRL group. As a result, phenylalanine net balance across the leg improved significantly in both groups in response to insulin ($P < 0.0001$), but the increase was significantly larger in the EX group such that net balance became positive only in the EX group ($P < 0.01$), indicating net muscle protein anabolism.

Mixed muscle FSR. The mixed muscle protein FSR (%/h) (Fig. 2) was not different between groups in the basal state (CTRL group: 0.063 ± 0.007 ; EX group: 0.062 ± 0.005). Insulin infusion significantly increased FSR in the EX group ($P < 0.05$) but not in the CTRL group (CTRL group: 0.062 ± 0.005 ; EX group: 0.099 ± 0.005).

Insulin and mTOR signaling proteins. Insulin and mTOR signaling is reported in Fig. 3. Akt/PKB^{Ser473} phosphorylation increased significantly with insulin (time effect $P < 0.05$), but the response was significantly greater in the EX group ($P < 0.05$). Insulin infusion significantly increased mTOR^{Ser2448} phosphorylation only in the EX group ($P < 0.05$). 4EBP1^{Thr37/46} phosphorylation increased in both groups during insulin infusion (time effect $P < 0.05$), with no difference between groups. S6K1^{Thr389} phosphorylation increased significantly with insulin (time effect $P < 0.05$), but the increase was larger in the EX group ($P < 0.05$).

DISCUSSION

Our study indicates for the first time that a single bout of moderate aerobic exercise overcomes the muscle protein insulin resistance and restores the physiological anabolic response of muscle protein synthesis to insulin in older individuals. These exciting results indicate that a moderate-intensity aerobic exercise that can be performed by most ambulatory older individuals may be an important tool to combat sarcopenia and frailty of aging. The effect of aerobic exercise on muscle protein anabolism in these older individuals appears to be due, at least in part, to improvements in endothelial function, insulin-induced vasodilation, and nutrient delivery. Specifically, we found that muscle protein synthesis and net balance significantly increased during insulin infusion only if the infusion was preceded by a bout of aerobic exercise. Such an effect was directly associated with an increase in blood flow, which, in turn, was accompanied by a significant increase in amino acid delivery and transport into the muscle tissue, and by enhanced mTOR signaling. In contrast, insulin did not exert any effect on either blood flow or muscle amino acid and protein turnover in the subjects resting before the insulin challenge.

These novel data also indicate for the first time that insulin-induced vasodilation is an important contributor to muscle protein anabolism in older subjects. If we interpret these results in light of our recently published study (11) in young adults demonstrating that physiological hyperinsulinemia stimulates muscle protein synthesis as long as it increases blood flow and amino acid delivery to the muscle tissue, we conclude that vasodilation is a major mechanism by which insulin stimulates muscle protein




FIG. 3. Phosphorylation of Akt/PKB, mTOR, 4E-BP1, and S6K1 in two groups of older subjects at baseline and during insulin infusion performed ~20 h following rest (Control; $n = 5$ for Akt/PKB and mTOR; $n = 4$ for 4E-BP1 and S6K1) or aerobic exercise (Exercise; $n = 6$). We used the following phospho and total primary antibodies (Cell Signaling, Beverly, MA): phospho-mTOR (Ser²⁴⁴⁸, cat. no. 2971, lot no. 9; 1:1,000), phospho-p70 S6K1 (Thr³⁸⁹, cat. no. 9234, lot no. 2; 1:500), phospho-Akt (Ser⁴⁷³, cat. no. 4058, lot no. 6; 1:500), and phospho-4EBP1 (Thr^{37/46}, cat. no. 2971; 1:1,000). Anti-rabbit IgG horseradish peroxidase-conjugated secondary antibody was purchased from Amersham Bioscience (1:2,000). Values are means \pm SE. * $P < 0.05$ vs. basal, # $P < 0.05$ vs. control. □, basal; ■, insulin. Bas, basal; Ins, insulin; MW, molecular weight.

synthesis throughout the adult life. Insulin-induced vasodilation appears to be more rate limiting for muscle protein anabolism than for glucose uptake. Studies on the regulation of vascular tone and glucose metabolism by insulin have uncovered a fascinating picture whereby insulin appears to rapidly recruit nutritive capillaries, which precedes the insulin's effects on intracellular signaling and glucose uptake in muscle (38). However, the overall effect of insulin-induced vasodilation on muscle glucose metabolism is more evident when glucose uptake is high (i.e., when the artery-to-vein glucose concentration gradient is high) (39,40). This occurs because an increased blood flow reduces the arteriovenous glucose gradient and allows for a more uniform exposure of all muscle tissue to adequate blood glucose concentrations necessary to sustain the higher rate of uptake (41). The metabolic effects of insulin-induced vasodilation appear magnified when examining muscle amino acid uptake and utilization and appear to drive the overall muscle protein anabolic response (present study and 11,16,29).

The vascular tone is determined by the fine balance of a complex network of vasodilators and vasoconstrictors, many of which are directly produced by the endothelium (42). Insulin increases muscle perfusion and capillary recruitment by stimulating NO production via activation of endothelial NO synthase (17,18), and it is also well recognized that aging reduces endothelial-derived vasodilation (14–16). The negative effect of aging appears to be also

associated with an increased plasma endothelin-1 concentration (16), which is another likely cause of the age-related endothelial dysfunction. Endothelin-1 is produced by vascular endothelial cells (43) and is one of the most powerful endothelial-derived vasoconstrictors (42). Interestingly, recent studies have shown that endothelin-1 receptor blockade enhances the vasodilatory action of insulin in humans (44), and aerobic exercise training for 3 months significantly reduces plasma endothelin-1 concentration in both younger and older subjects (45). Our current study is the first to report that a single bout of aerobic exercise reduces plasma endothelin-1 concentration to youthful levels and normalizes the vasodilatory and muscle protein anabolic response to insulin in older individuals. Considering all these data, it appears that endothelin-1 is potentially a suitable marker of age-related endothelial dysfunction and muscle protein insulin resistance. Further studies are needed to determine the sensitivity and specificity of endothelin-1 as a marker of muscle protein catabolism in older individuals.

Although our data strongly suggest that aerobic exercise normalized the anabolic response of muscle to insulin by improving endothelium-dependent vasodilation, we also found an improvement in insulin signaling. This was associated with a higher regional and whole-body insulin-stimulated glucose uptake in the exercise group, which is consistent with previous reports indicating that a single bout of aerobic exercise enhances insulin sensitivity for up

to 48 h in both healthy subjects and in patients with type 2 diabetes (21–27). However, if we analyze the Akt/mTOR signaling results in light of the amino acid and protein kinetics data, interpretation becomes more complicated. In the control subjects, local hyperinsulinemia stimulated muscle Akt^{Ser473}, 4EBP1^{Thr37/46}, and S6K1^{Thr389} phosphorylation but had no effect on amino acid availability, mTOR^{Ser2448} phosphorylation, or muscle protein synthesis. This is consistent with previously published data (5,16) and suggests that increases in insulin signaling in the absence of increased amino acid availability may be insufficient to fully stimulate the mTOR signaling pathway (13). Aerobic exercise enhanced Akt^{Ser473} and S6K1^{Thr389} phosphorylation, amino acid availability (consequent to improved vasodilation), and mTOR^{Ser2448} phosphorylation in response to insulin, resulting in the stimulation of muscle protein synthesis. Thus, by merely considering the present data one can definitely conclude that aerobic exercise restores the normal muscle protein anabolic response to hyperinsulinemia. However, the question remains as to whether this effect of exercise on mTOR signaling and protein synthesis is due to its effect on vasodilation (and amino acid availability), insulin signaling, or both.

Some indirect answers can be derived from the analysis of recent data published by our group, which allow a better understanding of the complex mechanisms underlying the anabolic effect of insulin on muscle proteins. Insulin sensitivity of glucose metabolism is not correlated with the insulin sensitivity of protein metabolism, as the insulin-induced protein anabolic response in muscle is preserved in uncontrolled type 2 diabetes (46). Additionally, we have shown in healthy, insulin-sensitive, younger subjects that increasing insulin doses, which increase glucose uptake in a dose-dependent manner, are unable to stimulate muscle protein anabolism if muscle perfusion and amino acid availability are not simultaneously enhanced (11). Thus, improved insulin signaling following exercise is unlikely the only major mechanism allowing for the restoration of the physiological muscle protein anabolic response to insulin, since improvements in vasodilation and the consequent increases in amino acid availability appear to play an essential role as well.

However, it is also important to underscore how the current data make it apparent that increases in tissue perfusion and amino acid availability are necessary, but not sufficient, to induce an anabolic response of muscle proteins. Indeed, the EX group tended to have a higher baseline blood flow and amino acid delivery, but this did not result in a higher baseline mTOR signaling or muscle protein synthesis. Therefore, it is likely that for insulin to acutely increase human skeletal muscle protein synthesis and anabolism it has to increase not only muscle perfusion and amino acid delivery but also provide adequate signals for an increase in muscle protein translation. Additionally, the higher baseline blood flow with no differences in muscle protein turnover or signaling also indicates that the effect of exercise on the vascular tone is direct and not mediated by increases in insulin sensitivity or muscle metabolism.

Finally, our finding of enhanced Akt signaling following exercise is consistent with some (47,48), but not all (49,50), previous reports. However, in the negative studies (49,50) insulin signaling had been measured immediately after exercise (<6 h), while we measured it 20 h postexercise (48). These data suggest that insulin signaling may

be enhanced by aerobic exercise in a time-dependent manner. Future studies on the time course of exercise-induced enhancement of insulin signaling are warranted.

In summary, a single bout of aerobic exercise overcomes the age-related insulin resistance of muscle protein synthesis by reducing endothelial dysfunction, improving muscle perfusion and amino acid availability for the muscle tissue, and enhancing activation of the Akt/mTOR signaling pathway and muscle protein synthesis. Clinical studies are needed to investigate whether aerobic exercise may prevent the loss of muscle mass with aging by improving the muscle anabolic response during feeding.

ACKNOWLEDGMENTS

This study was supported by the National Institute on Aging (grants R01 AG018311 and P30 AG024832); the Robert E. and May R. Wright Foundation; the National Institute for Arthritis and Musculoskeletal and Skin Diseases (grant R01 AR049877); the Shared Instrumentation Grant Program, National Center for Research Resources, NIH (grant S10 RR16650); and General Clinical Research Center Grants from the National Center for Research Resources, NIH (grants M01 RR00043 [University of Southern California] and M01 RR00073 [University of Texas Medical Branch]).

We thank the study subjects for their participation, Jeanine Cordero for technical assistance, and the nurses and personnel of the General Clinical Research Center of the University of Southern California and the University of Texas Medical Branch for their assistance with the clinical conduct of this study. Preliminary data from this study were presented at the Experimental Biology Meeting 2005.

REFERENCES

- Volpi E, Nazemi R, Fujita S: Muscle tissue changes with ageing. *Curr Opin Clin Nutr Metab Care* 7:405–410, 2004
- Tinetti ME, Williams CS: Falls, injuries due to falls, and the risk of admission to a nursing home. *N Engl J Med* 337:1279–1284, 1997
- Wolfson L, Judge J, Whipple R, King M: Strength is a major factor in balance, gait, and the occurrence of falls. *J Gerontol A Biol Sci Med Sci* 50 Spec No:64–67, 1995
- Volpi E, Mittendorfer B, Rasmussen BB, Wolfe RR: The response of muscle protein anabolism to combined hyperaminoacidemia and glucose-induced hyperinsulinemia is impaired in the elderly. *J Clin Endocrinol Metab* 85:4481–4490, 2000
- Guillet C, Prod'homme M, Balage M, Gachon P, Giraudet C, Morin L, Grizard J, Boirie Y: Impaired anabolic response of muscle protein synthesis is associated with S6K1 dysregulation in elderly humans. *FASEB J* 18:1586–1587, 2004
- Volpi E, Sheffield-Moore M, Rasmussen BB, Wolfe RR: Basal muscle amino acid kinetics and protein synthesis in healthy young and older men. *JAMA* 286:1206–1212, 2001
- Volpi E, Ferrando AA, Yeckel CW, Tipton KD, Wolfe RR: Exogenous amino acids stimulate net muscle protein synthesis in the elderly. *J Clin Invest* 101:2000–2007, 1998
- Volpi E, Mittendorfer B, Wolf SE, Wolfe RR: Oral amino acids stimulate muscle protein anabolism in the elderly despite higher first pass splanchnic extraction. *Am J Physiol Endocrinol Metab* 277:E513–E520, 1999
- Volpi E, Kobayashi H, Mittendorfer B, Sheffield-Moore M, Wolfe RR: Essential amino acids are primarily responsible for the amino acid stimulation of muscle protein anabolism in healthy elderly adults. *Am J Clin Nutr* 78:250–258, 2003
- Paddon-Jones D, Sheffield-Moore M, Zhang XJ, Volpi E, Wolf SE, Aarsland A, Ferrando AA, Wolfe RR: Amino acid ingestion improves muscle protein synthesis in the young and elderly. *Am J Physiol Endocrinol Metab* 286:E321–E328, 2004
- Fujita S, Rasmussen BB, Cadenas JG, Grady JJ, Volpi E: The effect of insulin on human skeletal muscle protein synthesis is modulated by insulin-induced changes in muscle blood flow and amino acid availability. *Am J Physiol Endocrinol Metab* 291:E745–E754, 2006

12. Proud CG: Regulation of protein synthesis by insulin. *Biochem Soc Trans* 34:2–6, 2006
13. Wang X, Proud CG: The mTOR pathway in the control of protein synthesis. *Physiology* 21:362–369, 2006
14. Meneilly GS, Elliot T, Bryer-Ash M, Floras JS: Insulin-mediated increase in blood flow is impaired in the elderly. *J Clin Endocrinol Metab* 80:1899–1903, 1995
15. DeSouza CA, Shapiro LF, Clevenger CM, Dinunno FA, Monahan KD, Tanaka H, Seals DR: Regular aerobic exercise prevents and restores age-related declines in endothelium-dependent vasodilation in healthy men. *Circulation* 102:1351–1357, 2000
16. Rasmussen BB, Fujita S, Wolfe RR, Mittendorfer B, Roy M, Rowe VL, Volpi E: Insulin resistance of muscle protein metabolism in aging. *FASEB J* 20:768–769, 2006
17. Steinberg HO, Brechtel G, Johnson A, Fineberg N, Baron AD: Insulin-mediated skeletal muscle vasodilation is nitric oxide dependent: a novel action of insulin to increase nitric oxide release. *J Clin Invest* 94:1172–1179, 1994
18. Scherrer U, Randin D, Vollenweider P, Vollenweider L, Nicod P: Nitric oxide release accounts for insulin's vascular effects in humans. *J Clin Invest* 94:2511–2515, 1994
19. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM, the Diabetes Prevention Program Research Group: Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 346:393–403, 2002
20. Tuomilehto J, Lindstrom J, Eriksson JG, Valle TT, Hamalainen H, Ilanne-Parikka P, Keinanen-Kiukkaanniemi S, Laakso M, Louheranta A, Rastas M, Salminen V, Uusitupa M, the Finnish Diabetes Prevention Study Group: Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med* 344:1343–1350, 2001
21. Henriksen EJ: Invited review: effects of acute exercise and exercise training on insulin resistance. *J Appl Physiol* 93:788–796, 2002
22. Perseghin G, Price TB, Petersen KF, Roden M, Cline GW, Gerow K, Rothman DL, Shulman GI: Increased glucose transport-phosphorylation and muscle glycogen synthesis after exercise training in insulin-resistant subjects. *N Engl J Med* 335:1357–1362, 1996
23. Borghouts LB, Keizer HA: Exercise and insulin sensitivity: a review. *International Journal of Sports Medicine* 21:1–12, 2000
24. Goodyear LJ, Kahn BB: Exercise, glucose transport, and insulin sensitivity. *Annu Rev Med* 49:235–261, 1998
25. Mikines KJ, Sonne B, Tronier B, Galbo H: Effects of acute exercise and detraining on insulin action in trained men. *Journal of Applied Physiology* 66:704–711, 1989
26. Wojtaszewski JF, Hansen BF, Kiens B, Richter EA: Insulin signaling in human skeletal muscle: time course and effect of exercise. *Diabetes* 46:1775–1781, 1997
27. Wojtaszewski JF, Hansen BF, Gade Kiens B, Markuns JF, Goodyear LJ, Richter EA: Insulin signaling and insulin sensitivity after exercise in human skeletal muscle. *Diabetes* 49:325–331, 2000
28. Sheffield-Moore M, Yeckel CW, Volpi E, Wolf SE, Morio B, Chinkes DL, Paddon-Jones D, Wolfe RR: Postexercise protein metabolism in older and younger men following moderate-intensity aerobic exercise. *Am J Physiol Endocrinol Metab* 287:E513–E522, 2004
29. Bell JA, Fujita S, Volpi E, Cadenas JG, Rasmussen BB: Short-term insulin and nutritional energy provision does not stimulate muscle protein synthesis if blood amino acid availability decreases. *Am J Physiol Endocrinol Metab* 289:999–1006, 2005
30. Jorfeldt L, Juhlin-Dannfelt A: The influence of ethanol on splanchnic and skeletal muscle metabolism in man. *Metabolism* 27:97–106, 1978
31. Jorfeldt L, Wahren J: Leg blood flow during exercise in man. *Clin Sci* 41:459–473, 1971
32. Wolfe RR: *Radioactive and Stable Isotope Tracers in Biomedicine: Principles and Practice of Kinetic Analysis*. New York, Wiley-Liss, 1992
33. Denne SC, Liechty EA, Liu YM, Brechtel G, Baron AD: Proteolysis in skeletal muscle and whole body in response to euglycemic hyperinsulinemia in normal adults. *Am J Physiol* 261:E809–E814, 1991
34. Calder AG, Anderson SE, Grant I, McNurlan MA, Garlick PJ: The determination of low d5-phenylalanine enrichment (0.002–0.09 atom percent excess), after conversion to phenylethylamine, in relation to protein turnover studies by gas chromatography/electron ionization mass spectrometry. *Rapid Commun Mass Sp* 6:421–424, 1992
35. Dreyer HC, Fujita S, Cadenas JG, Chinkes DL, Volpi E, Rasmussen BB: Resistance exercise increases AMPK activity and reduces 4E-BP1 phosphorylation and protein synthesis in human skeletal muscle. *J Physiol (Lond)* 576:2–24, 2006
36. Wolfe RR, Chinkes DL: *Isotope Tracers in Metabolic Research: Principles and Practice of Kinetic Analysis*. Hoboken, NJ, John Wiley & Sons, 2004
37. Toffolo G, Foster DM, Cobelli C: Estimation of protein fractional synthetic rate from tracer data. *Am J Physiol* 264:E128–E135, 1993
38. Vincent MA, Dawson D, Clark AD, Lindner JR, Rattigan S, Clark MG, Barrett EJ: Skeletal muscle microvascular recruitment by physiological hyperinsulinemia precedes increases in total blood flow. *Diabetes* 51:42–48, 2002
39. Baron AD, Tarshoby M, Hook G, Lazaridis EN, Cronin J, Johnson A, Steinberg HO: Interaction between insulin sensitivity and muscle perfusion on glucose uptake in human skeletal muscle: evidence for capillary recruitment. *Diabetes* 49:768–774, 2000
40. Vincent MA, Clerk LH, Lindner JR, Klibanov AL, Clark MG, Rattigan S, Barrett EJ: Microvascular recruitment is an early insulin effect that regulates skeletal muscle glucose uptake in vivo. *Diabetes* 53:1418–1423, 2004
41. Clerk LH, Vincent MA, Lindner JR, Clark MG, Rattigan S, Barrett EJ: The vasodilatory actions of insulin on resistance and terminal arterioles and their impact on muscle glucose uptake. *Diabetes Metab Res Rev* 20:3–12, 2004
42. Cardillo C, Kilcoyne CM, Cannon RO III, Panza JA: Interactions between nitric oxide and endothelin in the regulation of vascular tone of human resistance vessels in vivo. *Hypertension* 35:1237–1241, 2000
43. Boulanger C, Luscher TF: Release of endothelin from the porcine aorta. Inhibition by endothelium-derived nitric oxide. *J Clin Invest* 85:587–590, 1990
44. Mather KJ, Mirzamohammadi B, Lteif A, Steinberg HO, Baron AD: Endothelin contributes to basal vascular tone and endothelial dysfunction in human obesity and type 2 diabetes. *Diabetes* 51:3517–3523, 2002
45. Maeda S, Tanabe T, Miyauchi T, Otsuki T, Sugawara J, Iemitsu M, Kuno S, Ajsaka R, Yamaguchi I, Matsuda M: Aerobic exercise training reduces plasma endothelin-1 concentration in older women. *J Appl Physiol* 95:336–341, 2003
46. Bell JA, Volpi E, Fujita S, Cadenas JG, Sheffield-Moore M, Rasmussen BB: Skeletal muscle protein anabolic response to increased energy and insulin is preserved in poorly controlled type 2 diabetes. *J Nutr* 136:1249–1255, 2006
47. Christ-Roberts CY, Pratipanawatr T, Pratipanawatr W, Berria R, Belfort R, Mandarino LJ: Increased insulin receptor signaling and glycogen synthase activity contribute to the synergistic effect of exercise on insulin action. *J Appl Physiol* 95:2519–2529, 2003
48. Wadley GD, Konstantopoulos N, Macaulay L, Howlett K, Garnham A, Hargreaves M, Cameron-Smith D: Increased insulin-stimulated Akt pSer473 and cytosolic SHP2 protein abundance in human skeletal muscle following acute exercise and short-term training. *J Appl Physiol* 102:1624–1631, 2007
49. Wojtaszewski JF, Hansen BF, Gade Kiens B, Markuns JF, Goodyear LJ, Richter EA: Insulin signaling and insulin sensitivity after exercise in human skeletal muscle. *Diabetes* 49:325–331, 2000
50. Howlett KF, Sakamoto K, Yu H, Goodyear LJ, Hargreaves M: Insulin-stimulated insulin receptor substrate-2-associated phosphatidylinositol 3-kinase activity is enhanced in human skeletal muscle after exercise. *Metabolism* 55:1046–1052, 2006