

Inhibition of Lipolysis Stimulates Peripheral Glucose Uptake but Has No Effect on Endogenous Glucose Production in HIV Lipodystrophy

Birgitte Lindegaard,^{1,2} Christian Frøsig,³ Anne Marie W. Petersen,^{1,2} Peter Plomgaard,^{1,2} Susanne Ditlevsen,⁴ Bettina Mittendorfer,⁵ Gerrit Van Hall,^{1,2} Jørgen F.P. Wojtaszewski,³ Bente K. Pedersen^{1,2}

HIV-infected patients with lipodystrophy (HIV lipodystrophy) are insulin resistant and have elevated plasma free fatty acid (FFA) concentrations. We aimed to explore the mechanisms underlying FFA-induced insulin resistance in patients with HIV lipodystrophy. Using a randomized, placebo-controlled, cross-over design, we studied the effects of an overnight acipimox-induced suppression of FFAs on glucose and FFA metabolism by using stable isotope-labeled tracer techniques during basal conditions and a two-stage euglycemic-hyperinsulinemic clamp (20 and 50 mU insulin/m² per min, respectively) in nine patients with nondiabetic HIV lipodystrophy. All patients received anti-retroviral therapy. Biopsies from the vastus lateralis muscle were obtained during each stage of the clamp. Acipimox treatment reduced basal FFA rate of appearance by 68.9% (95% CI 52.6–79.5) and decreased plasma FFA concentration by 51.6% (42.0–58.9) (both, $P < 0.0001$). Endogenous glucose production was not influenced by acipimox. During the clamp, the increase in glucose uptake was significantly greater after acipimox treatment compared with placebo (acipimox: 26.85 $\mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ [18.09–39.86] vs. placebo: 20.30 $\mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ [13.67–30.13]; $P < 0.01$). Insulin increased phosphorylation of Akt Thr³⁰⁸ and glycogen synthase kinase-3 β Ser⁹, decreased phosphorylation of glycogen synthase (GS) site 3a + b, and increased GS activity (percent I-form) in skeletal muscle ($P < 0.01$). Acipimox decreased phosphorylation of GS (site 3a + b) ($P < 0.02$) and increased GS activity ($P < 0.01$) in muscle. The present study provides direct evidence that suppression of lipolysis in patients with HIV lipodystrophy improves insulin-stimulated peripheral glucose uptake. The

increased glucose uptake may in part be explained by increased dephosphorylation of GS (site 3a + b), resulting in increased GS activity. *Diabetes* 56:2070–2077, 2007

HIV-associated lipodystrophy (HIV lipodystrophy) is characterized by various degrees of subcutaneous fat loss, visceral fat accumulation, and metabolic disturbances, including peripheral and hepatic insulin resistance (1–3) and elevated plasma free fatty acids (FFAs) (4). Increased plasma FFA concentration is due to increased lipolysis (5,6), with increased net release of FFAs from adipose tissue (5,7). In other insulin-resistant states (e.g., obesity and type 2 diabetes), increased rates of lipolysis and plasma FFA concentration are related to the degree of insulin resistance (8–10). Both acute and chronic suppression of lipolysis can improve insulin sensitivity in HIV lipodystrophy (11,12). However, the nature of improved insulin resistance (i.e., whether due to changes in hepatic or peripheral insulin sensitivity) has not been determined, and the molecular mechanism behind FFA-induced insulin resistance is still incompletely understood. FFAs have been proposed to induce skeletal muscle insulin resistance by inhibiting insulin-stimulated glucose transport/phosphorylation (13–17) and glycogen synthesis (13–15,17–20). Recently, it has been demonstrated that FFAs interfere with the proximal steps of the insulin signaling cascade by impairing the insulin-stimulated tyrosine phosphorylation of insulin receptor substrate (IRS)-1, IRS-1-associated phosphatidylinositol-3-kinase activity (13,16,21–24), and phosphorylation of Akt (Ser⁴⁷³ and Thr³⁰⁸) (21,24), resulting in decreased GLUT4 translocation to the plasma membrane. FFAs also influence the distal part of the insulin pathway. Previous studies (13–15,17–19) consistently find that FFAs can reduce glycogen synthesis via inhibition of insulin-stimulated glycogen synthase (GS) activity, but the mechanism behind reduced GS activity is unknown (24).

In HIV lipodystrophy, both oxidative and nonoxidative glucose metabolism are impaired (25–27). A recent study (27) shows that patients with HIV lipodystrophy have a decreased insulin-stimulated GS activity in skeletal muscle. This defect in insulin action is associated with reduced dephosphorylation on sites 2 + 2a, 3a, and 3a + b, most likely via impaired signaling at the level of glycogen synthase kinase (GSK)-3 α Ser²¹ and GSK-3 β Ser⁹ and Akt

From the ¹Centre of Inflammation and Metabolism, Department of Infectious Diseases, Copenhagen, Denmark; ²The Copenhagen Muscle Research Centre, Rigshospitalet Copenhagen, Copenhagen, Denmark; ³The Copenhagen Muscle Research Centre, Section of Human Physiology, Department of Exercise and Sport Sciences, University of Copenhagen, Copenhagen, Denmark; the ⁴Department of Biostatistics, University of Copenhagen, Copenhagen, Denmark; and the ⁵School of Medicine, Washington University, St. Louis, Missouri.

Address correspondence and reprint requests to Birgitte Lindegaard, Centre of Inflammation and Metabolism at the Department of Infectious Diseases M7641, Rigshospitalet, Blegdamsvej 9, DK-2100 Copenhagen Ø, Denmark. E-mail: blm@rh.dk.

Received for publication 1 February 2007 and accepted in revised form 9 May 2007.

Published ahead of print at <http://diabetes.diabetesjournals.org> on 29 June 2007. DOI: 10.2337/db07-0144.

B.L. and C.F. contributed equally to this work.

ARV, antiretroviral; FFA, free fatty acid; G6P, glucose-6-phosphate; GS, glycogen synthase; GSK, glycogen synthase kinase; IRS, insulin receptor substrate.

© 2007 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

but not further upstream at the level of IRS-1-associated phosphatidylinositol-3-kinase activity (27).

The present study aimed to explore the mechanisms underlying the increase in insulin sensitivity after acute suppression of lipolysis by acipimox in patients with HIV lipodystrophy and fasting hyperinsulinemia. We hypothesized that acipimox treatment would enhance insulin sensitivity by reducing endogenous glucose production and increasing peripheral glucose uptake. Furthermore, we hypothesized that increased insulin-stimulated peripheral glucose uptake would involve improved distal insulin signaling at the level of Akt, GSK-3, and GS in muscle.

RESEARCH DESIGN AND METHODS

Twenty-one HIV-positive men were recruited from the outpatient clinic of the Department of Infectious Disease (Rigshospitalet, Copenhagen, Denmark). Subjects underwent a medical examination, and standard blood tests were performed to determine eligibility. The following were the inclusion criteria: 1) >18 years of age; 2) stable antiretroviral (ARV) therapy, with no changes in ARV therapy during the preceding 8 weeks; and 3) fasting plasma insulin ≥ 104 pmol/l. If the fasting plasma insulin concentration was <104 pmol/l, a standard 75-g oral glucose tolerance test was made and the subject was included if insulin levels were >521 pmol/l after 120 min (11,28); 4) lipodystrophy (at least one moderate sign of lipodystrophy [fat loss] in face, arms, buttocks, or legs based on a physical examination by a single investigator [B.L.]). The degree of lipodystrophy was rated as previously described (29).

The exclusion criteria were 1) severe cardiovascular diseases; 2) hospitalization within the last 8 weeks; 3) fasting glucose ≥ 7 mmol/l; and 4) concurrent therapy with insulin, antidiabetes agents, testosterone, growth hormone, glucocorticoids, anabolic steroids, or anticoagulation. Of 21 patients who were recruited, 11 fulfilled the inclusion criteria and 2 declined to participate. Consequently, nine patients were included in the study. The following were reasons for exclusion: plasma insulin <104 pmol/l ($n = 6$), severe cardiovascular diseases ($n = 2$), and no lipodystrophy by score ($n = 2$). Two of six patients with plasma insulin <104 pmol/l had lipodystrophy by the lipodystrophy score system. All patients received a combination of ARV therapies. All, except two, received protease inhibitors. Written informed consent was obtained according to the requirements from the local ethical committee.

Body composition. Fat and fat-free tissue masses of the whole body, trunk, and extremities were measured using a dual-energy X-ray absorptiometry scanner (Norland XR 36, software version 3.94; Norland, Fort Atkinson, WI) (1).

Stable isotope infusion study. The study was a randomized, double-blind, cross-over study. Each subject was studied twice (at least 2 weeks apart), once while receiving acipimox and once while receiving placebo (Fig. 1). The ARV medication was taken 3.5–4.5 h before commencing the clamp, and the ARV combination was not changed throughout the study. Acipimox was administered in doses of 250 mg at 7:00 P.M. and 12:00 A.M. the night before and at 7:00 A.M. and 9:00 A.M. on the study day (30,31). At 7:00 A.M. on the study day after an overnight fast, an intravenous catheter was inserted into an antecubital vein of one arm for infusion of stable isotope-labeled tracers, insulin, and dextrose (20%). A second intravenous catheter was inserted into a hand vein of the contralateral arm, which was wrapped in a heating blanket to obtain arterialized blood samples. After baseline blood samples were obtained at 8:00 A.M., a primed 16- $\mu\text{mol/kg}$ constant infusion ($0.22 \mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) of [6,6- $^2\text{H}_2$]-glucose (Cambridge Isotopes Laboratories, Cambridge, MA) was started and maintained for 8.5 h to determine glucose kinetics. At 9:30 A.M., infusion of [2,2- $^2\text{H}_2$]-palmitate (Cambridge Isotopes Laboratories) at a constant rate of $0.04 \mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ was started to determine fatty acid kinetics, as previously described (3,32,33). At 10:30 A.M., 2.5 h after the start of the isotope infusion (basal period), a two-stage euglycemic-hyperinsulinemic clamp was initiated and continued for 6 h, modified after the protocol (34). During stage 1 of the clamp (from 10:30 A.M. to 12:30 P.M.), insulin (Actrapid 100 IU/ml; Novo Nordisk) was infused at a rate of 20 mU/m^2 per min (initiated with a two-step priming dose of 80 mU/m^2 per min for 5 min followed by 40 mU/m^2 per min for 5 min). During stage 2 of the clamp (from 12:30 to 4:30 P.M.), insulin was infused at a rate of 50 mU/m^2 per min (initiated with a two-step priming dose of 200 mU/m^2 per min for 5 min followed by 100 mU/m^2 per min for 5 min). Euglycemia was achieved by a variable rate of infusion of 20% glucose enriched to $\sim 2.5\%$ with [6,6- $^2\text{H}_2$]-glucose to maintain a blood glucose concentration at 5.5 mmol/l . The infusion of [6,6- $^2\text{H}_2$]-glucose was decreased by 50% of the initial infusion rate during stage 1 and 75% of basal during stage 2 to steadily maintain the plasma glucose enrichment by




FIG. 1. Schematic representation of the study. **A:** Administration of acipimox/placebo and time schedule. Acipimox and placebo were administered overnight. **B:** The experimental study day. Whole-body glucose metabolism was measured by combined euglycemic-hyperinsulinemic clamp and stable isotope tracer method. Insulin was infused for 120 min at a rate of 20 mU/m^2 per min (stage 1) and for 240 min at a rate of 50 mU/m^2 per min (stage 2). Steady-state measurements of glucose and palmitate concentrations and kinetics were carried out during the last 30 min of the basal period and during each stage of the insulin clamp every 10 min. Arrows show time points for blood and skeletal muscle biopsy samples.

accounting for the expected decline in hepatic glucose production. The infusion of [2,2- $^2\text{H}_2$]-palmitate was decreased by 50% of the initial infusion rate during stages 1 and 2 because of the expected decline in whole-body lipolytic rate. One subject could not complete the clamp study for technical reasons. Results during the clamp therefore represent data from eight subjects only.

Muscle biopsy. Muscle biopsies from m. vastus lateralis were obtained with a modified Bergström needle (including suction) under local anesthesia with 2% lidocaine within the first 30 min of the basal stage and at the end of stages 1 and 2 of the clamp. Muscle tissue was immediately frozen in liquid nitrogen and stored at -80°C .

Blood samples. Arterialized blood samples were taken before beginning the isotope infusion to determine background glucose and palmitate enrichments. To determine glucose and palmitate concentrations and kinetics, arterialized blood samples were taken every 10 min during the last 30 min of the basal period and each stage of the insulin clamp (i.e., basal: 120, 130, 140, and 150 min; stage 1: 240, 250, 260, and 270 min; and stage 2: 480, 490, 500, and 510 min after the start of the tracer infusion) (3,32) (Fig. 1). Venous blood samples were taken at 0, 30, 90, 120, 150, 180, 210, 240, 270, 300, 330, 390, 450, and 510 min. All blood samples were drawn into tubes containing EDTA and centrifuged. Plasma was stored at -80°C . Plasma concentration of FFAs and glycerol were determined using an automatic analyzer (Cobas fara; Roche, Basel, Switzerland; FFA: NEFA C; Wako Chemicals, Neuss, Germany). Insulin was measured by enzyme-linked immunosorbent assay (detection limit, 3.0 pg/ml ; DakoCytomation, Cambridgeshire, U.K.). All determinations were run as duplicates, and mean values were calculated. To maintain euglycemia, arterialized blood glucose was measured every 5 min in the 1st hour of each clamp stage and every 10 min throughout the rest of the clamp and was analyzed immediately (EML 105 radiometer; Radiometer Medical A/S, Copenhagen, Denmark).

Total, HDL, and LDL cholesterol (mmol/l); triglycerides (mmol/l); and glucose (mmol/l) measurements on the day of inclusion were determined using standard laboratory procedures. CD4^+ cell counts were calculated by flow cytometry and HIV RNA copies (lower limit of detection: 20 copies/ml) were measured by the Amplicor HIV Monitor (Roche Molecular Systems, Branchburg, NJ).

Tracer analyses

Glucose. Enrichment of glucose was measured as previously described in detail (35). In short, $20 \mu\text{l}$ plasma was vigorously shaken with $100 \mu\text{l}$ ethyl acetate to remove plasma lipids. After centrifugation ($15,000 \text{ rpm}$ at 5 min) the ethyl acetate was disposed and $20 \mu\text{l}$ 8 mol NaOH , $1 \text{ mol K}_2\text{HPO}_4$, and $10 \mu\text{l}$ benzoylchlorid were added, and the solution was vortexed for 5 min. After the

vortex, the solution was naturalized by adding 10 µl 1.4 H₃PO₄, and the glucose derivate, hexo-benzoyl glucose, was extracted by a short vigorous shake with 500 µl ethylacetate. The ethylacetate phase was separated by centrifugation and transferred to a new eppendorf tube, and the ethylacetate was evaporated under a stream of nitrogen. The glucose derivate was dissolved in the high-performance liquid chromatography buffer. The solution was filtered before being analyzed on the liquid chromatography-mass spectrometry system (aQa; Finnegan, Manchester, U.K.).

Palmitate. Concentration and enrichment was determined by gas chromatography-mass spectrometry analysis (Automass II; Finnegan, Paris, France) as described by Patterson et al. (36).

The fatty acid concentrations were determined by using an internal standard (1,2,3,4-¹³C₄-palmitic acid; Sigma Aldrich, MO), which was related to a standard curve. To determine plasma palmitate enrichment, the sample was separated on a 30-m capillary fused-silica column (Rtx-2330; Restex, Forney, TX) with an injector temperature at 50°C increased to 200°C at a rate of 20°C per min. The oven was set at the following program: initial 50°C increasing to 150°C at 10°C per min and then slowly increased to 163°C with a rate of 2°C per min. To clean the column, the temperature was increased to 260°C at a rate of 20°C per min and maintained at 260°C for 7 min. The mass spectrometer was scanned in a profile mode in a range from 287 to 311 amu.

Calculations. Tracer-to-tracee ratios for both glucose and palmitate were calculated as the ratio between the master peak (M) and the enriched peak (M + 2) after subtracting the background enrichment. The calculations are based on a steady-state assumption. For glucose the total rate of appearance (*R_a*) was calculated as the following:

$$R_{a(\text{endogenous})} = \frac{F_{\text{total}}}{E_{\text{glucose}}} - GIR \quad (1)$$

and *R_d* as the following:

$$R_d = \frac{F_{\text{total}}}{E_{\text{glucose}}} \quad (2)$$

where *R_a* is rate of appearance (µmol · kg body wt⁻¹ · min⁻¹). *F_{total}* is the total infusion rate of glucose tracer (µmol · kg⁻¹ · min⁻¹). *E_{glucose}* is the enrichment of glucose in plasma (tracer-to-tracee ratio). *GIR* is the exogenous glucose administered during the clamp. *R_d* is rate of disappearance (µmol · kg body wt⁻¹ · min⁻¹).

For palmitate, *R_a* was calculated as the following:

$$R_a = R_d = \frac{F}{E_{\text{palmitate}}}$$

where *F* is the infusion rate of ²H₂-palmitate (µmol · kg body wt⁻¹ · min⁻¹) and *E_{palmitate}* is the tracer-to-tracee ratio of plasma palmitate. The exact amount of tracer in the infusate was measured, as described above, for plasma palmitate concentration and enrichment, and this value was used in the calculation. *R_a* is equal to *R_d* during isotopic and physiologic steady state.

SDS-PAGE and Western blotting. Muscle homogenate proteins were separated using 7.5% Bis-tris gels (Invitrogen, Taastrup, Denmark) and transferred to polyvinylidene fluoride membranes (immobilon transfer membrane; Millipore, Glostrup, Denmark). After blocking (Tris-buffered saline with Tween with 2% skim milk), the membranes were incubated with primary antibodies (Tris-buffered saline with Tween with 2% skim milk) followed by incubation in horseradish peroxidase-conjugated secondary antibody (Tris-buffered saline with Tween with 2% skim milk) (Amersham Pharmacia Biotech, Buckinghamshire, U.K.). Following detection and quantification using a charge-coupled device image sensor and 1D software (Kodak Image Station, 2,000 mm; Kodak, Ballerup, Denmark), the protein content was finally expressed in arbitrary units relative to a human skeletal muscle standard. The antibodies used for detection of phosphorylation of AKT and GSK-3 were as follows: AKT Thr³⁰⁸ (Upstate Biotechnology, Lake Placid, MA) and GSK-3β Ser⁹ (~35-kDa band using the anti-GSK-3α/β Ser²¹/Ser⁹ pan antibody; New England Biolabs, Beverly, MA). For detection of phosphorylation of glycogen synthase at site 3a + 3b (Ser⁶⁴⁰) and Ser⁶⁴⁴, an antibody raised in sheep, as described (37), was used.

GS activity. GS activity was measured in muscle homogenates using a Unifilter 350 microtiter plate assay (Whatman, Frisenette, Ebeltoft, Denmark). The assay ran in triplicate based on the original protocol described by Thomas et al. (38). GS activity was determined in the presence of 8 or 0.02 mmol/l glucose-6-phosphate (G6P), representing the maximal GS activity and the G6P-independent activity, respectively. The percent I-form of GS was calculated as 100 times the activity in the presence of 0.02 mmol/l G6P divided by the activity at 8.0 mmol/l G6P.

Statistics. Values of FFAs, glycerol, *R_a*, *R_d*, and insulin were naturally log transformed to achieve homoscedacity and an approximate normal distribution. GS activity (I-form) and GSK-3β were analyzed with and without log

TABLE 1
Clinical characteristics of the study patients

Characteristics	
<i>n</i>	9
Age (years)	52.3 ± 2.9
HIV infection duration (years)	16.3
LogHIV RNA (copies/ml)	1.90 ± 0.55
CD4 + cell counts (cells/µl)	732 ± 118
HAART combination	
PI, NRTI	3
PI, NRTI, NNRTI	4
NRTI, NNRTI	1
NRTI	1
Body weight (kg)	81.9 ± 3.2
BMI (kg/m ²)	25.0 ± 0.8
Body fat (%)	19.9 ± 2.2
Fat mass (kg)	16.8 ± 2.4
Lean body mass (kg)	62.5 ± 12.7
Truncal fat mass (kg)	10.4 ± 1.6
Limb fat mass (kg)	5.5 ± 0.95
Trunk-to-limb ratio	2.00 ± 0.36
Fasting glucose (mmol/l)	5.5 ± 0.23
Fasting insulin (pmol/l)	86 ± 18
Fasting triglycerides (mmol/l)	3.07 ± 0.74
Fasting total cholesterol (mmol/l)	6.1 ± 0.48
Fasting HDL cholesterol (mmol/l)	1.4 ± 0.35
Fasting LDL cholesterol (mmol/l)	3.73 ± 0.74
Fasting FFAs (µmol/l)	412 ± 61.3

Data are means ± SE. HAART, highly active antiretroviral therapy; PI, protease inhibitor; NRTI, nucleoside reverse transcriptase inhibitor; NNRTI, nonnucleoside reverse transcriptase inhibitor.

transformation. Statistical analyses of substrate concentration and kinetics were carried out using a general linear mixed model. A random subject-specific component was introduced that allowed adjusting for the interindividual variations. The effect of treatment was estimated using a binary variable (acipimox or placebo) and the effect of insulin infusion using a categorical variable (none, low [20 units/l], or high [50 units/l] infusion level), and the model allowed for an interaction term between the two where needed. Time in minutes was entered as a continuous variable in the analysis of glucose, insulin, FFAs, and glycerol. Likelihood ratio tests were applied to assess statistical significance. The fit of the general linear model was evaluated by testing the residuals for normality and by inspection of the residual plots. Linear regression was used to quantify the proportion of the effect of increased dephosphorylation of GS site 3a + b on *R_d* that could be explained by acipimox by the decrease in the effect estimate when treatment entered the model. The model was adjusted for insulin infusion. For the analysis, the procedure lme in R (version 2.1) was used (39). Results are presented as estimate (95% CI), unless otherwise stated.

RESULTS

Baseline characteristics. Clinical characteristics of subjects are shown in Table 1. All subjects had lipodystrophy, as indicated by their trunk-to-limb fat ratio, elevated triglycerides (plasma triglycerides >1.7 mmol/l) (40), and plasma FFA concentration (plasma FFAs in healthy men 292 ± 58 µmol/l) (41). All patients received at least one nucleoside analogue, seven received at least one protease inhibitor, and five a nonnucleoside analogue. All patients except one had suppressed HIV RNA.

Effect of acipimox on metabolic biochemistry. Basal plasma glucose and insulin concentrations were not different in the placebo and acipimox trials (Fig. 2A and B).

Basal plasma FFA concentration was 51.6% (95% CI 43.0–58.9) lower during the acipimox than during the placebo trial. During stage 1 of the clamp, plasma FFA concentration was reduced 61.3% (53.4–67.8) in the acipimox trial compared with a 46.2% (35.9–54.9) reduction in




FIG. 2. Plasma glucose (A), insulin (B), FFAs (C), and glycerol (D) levels during the basal stage, low-dose insulin infusion (stage 1), and high-dose insulin infusion (stage 2) after overnight treatment with placebo (○) and acipimox (●). Data are presented as means \pm SE for glucose and as geometric means \pm geometric SE for insulin, FFAs, and glycerol.

the placebo trial. During stage 2, the clamp plasma FFA concentrations were not reduced further ($P < 0.0001$ for treatment effect, $P < 0.0001$ for low-dose insulin effect during placebo, $P = 0.035$ for low-dose insulin effect during acipimox, and $P = 0.27$ for high-dose insulin effect) (Fig. 2C).

Basal plasma glycerol concentration was 36.5% (95% CI 29.2–43.1) lower during the acipimox than during the placebo trial. During stage 1, plasma glycerol concentration was reduced 58.7% (53.6–63.3) in the acipimox trial compared with a 43.4% (36.3–49.7) reduction in the placebo trial. During stage 2, plasma glycerol concentrations were not reduced further ($P < 0.0001$ for treatment effect, low-dose insulin effect during placebo, and low-dose insulin effect during acipimox and $P = 0.89$ for high-dose insulin effect) (Fig. 2D).

Effect of acipimox on palmitate R_a . Palmitate R_a decreased by 68.9% (95% CI 52.6–79.5) in the basal stage in the acipimox trial compared with placebo (acipimox: $0.65 \mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ [95% CI 0.29–1.45] versus placebo: $2.08 \mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ [1.41–3.06]; $P < 0.0001$). The decrease in palmitate R_a during stage 1 was greater during acipimox treatment compared with placebo (acipimox: $0.42 \mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ [0.18–0.95] versus placebo: $0.74 \mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ [0.32–1.70] or 79.9 vs. 64.6% compared with baseline; $P = 0.0086$). Palmitate R_a did not differ significantly between the acipimox and placebo trials during stage 2 (acipimox: $0.31 \mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ [0.14–0.71] versus placebo: 0.39 [0.17–0.9] or 85.0 vs. 81.1% compared with baseline; $P = 0.27$). The effect of insulin infusion was significant for both the acipimox ($P = 0.002$) and the placebo ($P < 0.0001$) trials (Fig. 3).

Effect of acipimox on glucose R_a and R_d . During basal conditions, glucose R_a and glucose R_d were the same for the placebo and the acipimox trials (acipimox: $14.77 \mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ [95% CI 8.85–24.66] versus placebo: $14.29 \mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ [11.38–17.93]). Glucose R_a was also not influenced by acipimox treatment during the clamp (stage 1: acipimox: $8.53 \mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ [5.06–14.40] versus placebo: $7.78 \mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ [4.61–13.12] or 41.3 vs.




FIG. 3. Histogram shows effect of treatment (placebo [□] versus acipimox [■]) and time (insulin) on palmitate rate of appearance. Basal = no insulin infusion, clamp stage 1 = 20 mU/m^2 per min insulin infusion, and clamp stage 2 = 50 mU/m^2 per min insulin infusion. * $P < 0.0001$, placebo vs. acipimox during basal stage; † $P < 0.01$, placebo vs. acipimox during clamp stage 1. Time effect = the effect of insulin for both placebo treatment and acipimox treatment. Values are geometric means \pm geometric SE.




FIG. 4. Histograms show effect of treatment (placebo (\square) versus acipimox (\blacksquare)) and time (insulin) on glucose rate of appearance (A) and disappearance (B). Basal = no insulin infusion, clamp stage 1 = 20 mU/m^2 per min insulin infusion, and clamp stage 2 = 50 mU/m^2 per min insulin infusion. * $P = 0.01$, placebo vs. acipimox during clamp stage 1. Time effect = the effect of insulin for placebo treatment and acipimox treatment. Values are geometric means \pm geometric SE. Glucose rates of appearance and disappearance are measured as $\mu\text{mol} \cdot \text{kg body wt}^{-1} \cdot \text{min}^{-1}$.

46.5% compared with baseline. Stage 2: acipimox: 4.96 $\mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ [2.94–8.37] versus placebo: 4.40 $\mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ [2.58–7.51] or 65.9 vs. 69.7% compared with baseline). The effect of insulin infusion on R_a was significant for both the acipimox and the placebo trials ($P = 0.0003$) (Fig. 4A).

Glucose R_d increased to a greater extent in the acipimox than in the placebo trial during stage 1 (acipimox: 26.85 $\mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ [95% CI 18.09–39.86] versus placebo: 20.30 $\mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ [13.67–30.13] or 84.1 vs. 39.2% compared with baseline; $P < 0.01$). Glucose R_d did not differ significantly in the acipimox and placebo trials during high-dose insulin infusion (acipimox: 47.00 $\mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ [31.66–69.77] versus placebo: 40.57 $\mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ [27.09–60.77] or 222.2 vs. 178.3% compared with baseline; $P = 0.16$). The effect of insulin infusion was significant for both the acipimox and the placebo trials ($P < 0.0001$) (Fig. 4B).

Effects of acipimox on regulation of glycogen synthase activity and insulin signaling. Both low and high insulin infusion increased GS activity (I-form) ($P < 0.0001$), decreased phosphorylation on GS site 3a + b ($P <$

0.01), and increased phosphorylation on both GSK-3 β -Ser⁹ ($P < 0.001$) and Akt Thr³⁰⁸ ($P < 0.0001$) (Fig. 5A–D) measured in muscle tissue preparations. In response to acipimox treatment, phosphorylation of GS site 3a + b decreased (treatment effect, $P = 0.017$) (Fig. 5B) and the percent I-form activity of GS increased (treatment effect, $P = 0.01$ [log transformed] and $P = 0.04$ [nontransformed]) (Fig. 5A). Phosphorylation of GSK-3 β and Akt did not change in response to acipimox treatment (GSK-3 β : $P = 0.06$ [nontransformed]; $P = 0.05$ [log transformed] and Akt Thr³⁰⁸: $P = 0.42$) (Fig. 5C and D, respectively). Total GS activity was unchanged by both treatment and insulin (data not shown). The effect of acipimox on R_d appeared partly explained by increased dephosphorylation of GS 3a + b, as 24% of the effect of GS site 3a + b on R_d was due to acipimox treatment.

DISCUSSION

The present study demonstrates that decreasing plasma FFA concentrations and R_a palmitate by acipimox treatment increase peripheral insulin-stimulated glucose uptake in HIV lipodystrophy without any influence on hepatic glucose production.

Similar to the observation of Hadigan et al. (12), who treated patients with HIV lipodystrophy with acipimox for 3 months, we found a decrease of $\sim 50\%$ in basal levels of circulating FFAs and R_a FFAs and an $\sim 30\%$ increase in glucose uptake in response to acipimox. The fact that acute and chronic acipimox treatment elicited the same responses with regard to plasma FFA and insulin sensitivity suggests that chronic lipid-induced insulin resistance can be reversed equally by acute and chronic intervention. The present study adds to former studies by pinpointing the effect specifically to glucose uptake and on muscle insulin signaling. During stage 1 (low insulin doses), there was a significant difference between the placebo and the acipimox trials with regard to plasma FFAs, R_a palmitate, and R_d glucose. During stage 2 (high insulin dose), lipolysis was totally blocked by insulin, which is most likely the reason for no detectable effect of acipimox on glucose metabolism. The response of R_a FFAs to acute acipimox during high-dose insulin is similar to the response of R_a FFAs to chronic acipimox (12).

It is, to some extent, surprising that we did not find any effect of acipimox on hepatic glucose production, even during low insulin dose. This finding conflicts with some previous observations (31,42,43) using acipimox to suppress lipolysis. Three other studies have demonstrated that acute acipimox treatment decreases basal hepatic glucose (31) and increases hepatic sensitivity to insulin in type 2 diabetic subjects (31,42,43). One explanation for the discrepancy could be that the basal levels of plasma FFAs were higher in the diabetic patients. Taken together, our findings suggest that the nature of hepatic insulin resistance in patients with nondiabetic HIV lipodystrophy may, to some extent, differ from that seen in type 2 diabetes. It is still unknown whether increased hepatic glucose production in HIV lipodystrophy is a result of increased gluconeogenesis or increased glycogenolysis or a combination of both. It could be argued that the patients we selected for the present study did not demonstrate hepatic insulin resistance. However, the subjects had increased R_a glucose (8.1 $\mu\text{mol} \cdot \text{kg body wt}^{-1} \cdot \text{min}^{-1}$; $\sim 6.2 \mu\text{mol} \cdot \text{kg fat-free mass}^{-1} \cdot \text{min}^{-1}$) compared with R_a glucose in healthy men ($\sim 2.1 \mu\text{mol} \cdot \text{kg fat-free mass}^{-1} \cdot \text{min}^{-1}$)




FIG. 5. Histograms show effect of treatment (placebo versus acipimox) and time (insulin) on glycogen synthase activity given as percent I-form activity (A) and on phosphorylation of GS at the COOH-terminal site 3a + b (B), of GSK-3 β Ser⁹ (C), and of Akt Thr³⁰⁸ (D) in skeletal muscle. Basal (▨) = no insulin infusion, clamp stage 1 (■) = 20 mU insulin infusion, and clamp stage 2 (□) = 50 mU insulin infusion. Placebo: $n = 8, 7, 8$; and Acipimox: $n = 7, 6, 6$ for basal, clamp stage 1, and clamp stage 2, respectively. * $P < 0.05$; ** $P < 0.01$. Time effect = the effect of insulin for both placebo treatment and acipimox treatment. Values are means \pm SE.

during low insulin dose (3). Inhibition of insulin-stimulated glucose uptake by FFAs is supported by other studies, including experimental designs, where plasma FFA levels were lowered by acipimox (19,30,44,45) or increased by lipid infusions (14,46–48). In these studies, the effect on glucose uptake was observed at a high insulin dose (40–80 mU/m² per min). Our study shows that the effect of suppression of FFA levels may, in fact, be more easily detected at lower insulin concentrations (20 mU/m² per min).

Patients with HIV lipodystrophy are characterized by having decreased oxidative and nonoxidative glucose disposal (25–27). During insulin stimulation, nonoxidative glucose metabolism is thought to primarily represent muscle glycogen synthesis (49). As expected, insulin administration induced increased phosphorylation of Akt Thr³⁰⁸ and GSK-3 β Ser⁹, decreased phosphorylation on GS (site 3a + b), and subsequently increased the activity of GS in both trials. Interestingly, in the response to acipimox treatment, GS activity (%I-form) was greater and phosphorylation of GS (site 3a + b) decreased compared with before treatment. Moreover, acipimox-induced increase in glucose uptake was associated with acipimox-induced improvement of the insulin signaling pathway, although this does not prove causality.

Regulation of GS activity by manipulating FFA concentration is in accordance with previous studies (13–15,17–19). GSK-3 is a serine/threonine kinase that inhibits GS activity by phosphorylation of GS (50) at sites 4, 3c, 3b, and 3a (51). However, in our study, we did not find a significant effect of acipimox treatment on phosphorylation of GSK-3 β Ser⁹. This finding is surprising but seems consistent with the observation in rodents that insulin-

stimulated glycogen synthase activity is inhibited by infusion of lipids independent of deactivation of GSK-3 (24). In addition, Skurat et al. (50) have previously, in an in vitro study, demonstrated that GS (site 3a + b) can be phosphorylated by unidentified protein kinases, suggesting regulation of GS activity by non-GSK-3-mediated pathways (24,52).

Upstream of GSK-3, acipimox treatment correspondingly did not effect phosphorylation of Akt Thr³⁰⁸, a marker of Akt activity. Conflicting data on regulation of phosphorylation of Akt by lipids exist. In a study by Kryszynska et al. (23), intralipid infusion did not change phosphorylation of Akt Thr³⁰⁸ and Akt Ser⁴⁷³ despite reduced whole-body insulin-stimulated glucose disposal. In contrast, two other in vivo studies (humans [21] and rats [24]) find impaired phosphorylation of Akt Thr³⁰⁸ and Ser⁴⁷³ in response to lipid infusion. These discrepancies may indicate that the signaling mechanism is different when inducing acute insulin resistance by lipid infusion compared with reversing chronic insulin resistance by acipimox treatment. Furthermore, it cannot be excluded that acipimox itself may directly stimulate glycogen synthesis in skeletal muscle, although no receptors for nicotinic acid in skeletal muscle have been identified (53). Finally, it is possible that acipimox treatment may lead to improved activation of protein phosphatases regulating GS activity during insulin stimulation. This, however, remains to be investigated.

The present study provides direct evidence that suppression of lipolysis in patients with HIV lipodystrophy improves insulin-stimulated peripheral glucose uptake without influencing hepatic insulin resistance. The increased glucose uptake may in part be explained by increased dephosphorylation of GS (site 3a + 3b), result-

ing in increased GS activity independently of phosphorylation of Akt Thr³⁰⁸ and GSK-3 β Ser⁹.

ACKNOWLEDGMENTS

The Centre of Inflammation and Metabolism is supported by a grant from the Danish National Research Foundation (grant no. 02-512-55). The Copenhagen Muscle Research Centre is supported by grants from the Copenhagen Hospital Corporation, Faculties of Science and of Health Sciences, University of Copenhagen. The study was further supported by the Danish Medical Research Council (grant no. 22-01-009 and 22-003-0177), the Danish AIDS Foundation, the Bikuben Foundation, the Illum Foundation, the Aase and Einar Danielsens Foundation, and National Institutes of Health Grants AR 49869 and DK 56341 (clinical nutrition research unit). J.F.P.W. is the recipient of a Hallas Møller fellowship from the Novo Nordisk Foundation.

We thank the patients for their participation in this study. We are grateful to Ruth Rousing, Hanne Willumsen, Nina Pluszek, Carsten Nielson, and Flemming Jessen for excellent technical help.

REFERENCES

- Carr A, Samaras K, Burton S, Law M, Freund J, Chisholm DJ, Cooper DA: A syndrome of peripheral lipodystrophy, hyperlipidaemia and insulin resistance in patients receiving HIV protease inhibitors. *AIDS* 12:F51–F58, 1998
- Haugaard SB, Andersen O, Dela F, Holst JJ, Storgaard H, Fenger M, Iversen J, Madsbad S: Defective glucose and lipid metabolism in human immunodeficiency virus-infected patients with lipodystrophy involve liver, muscle tissue and pancreatic beta-cells. *Eur J Endocrinol* 152:103–112, 2005
- Reeds DN, Yarasheski KE, Fontana L, Cade WT, Lacity E, DeMoss A, Patterson BW, Powderly WG, Klein S: Alterations in liver, muscle, and adipose tissue insulin sensitivity in men with HIV infection and dyslipidemia. *Am J Physiol Endocrinol Metab* 290:E47–E53, 2006
- Meininger G, Hadigan C, Laposata M, Brown J, Rabe J, Louca J, Aliabadi N, Grinspoon S: Elevated concentrations of free fatty acids are associated with increased insulin response to standard glucose challenge in human immunodeficiency virus-infected subjects with fat redistribution. *Metabolism* 51:260–266, 2002
- Sekhar RV, Jahoor F, White AC, Pownall HJ, Visnegarwala F, Rodriguez-Barradas MC, Sharma M, Reeds PJ, Balasubramanyam A: Metabolic basis of HIV-lipodystrophy syndrome. *Am J Physiol Endocrinol Metab* 283:E332–E337, 2002
- Hadigan C, Borgonha S, Rabe J, Young V, Grinspoon S: Increased rates of lipolysis among human immunodeficiency virus-infected men receiving highly active antiretroviral therapy. *Metabolism* 51:1143–1147, 2002
- Reeds DN, Mittendorfer B, Patterson BW, Powderly WG, Yarasheski KE, Klein S: Alterations in lipid kinetics in men with HIV-dyslipidemia. *Am J Physiol Endocrinol Metab* 285:E490–E497, 2003
- Boden G: Role of fatty acids in the pathogenesis of insulin resistance and NIDDM. *Diabetes* 46:3–10, 1997
- Boden G: Interaction between free fatty acids and glucose metabolism. *Curr Opin Clin Nutr Metab Care* 5:545–549, 2002
- Avramoglou RK, Basciano H, Adeli K: Lipid and lipoprotein dysregulation in insulin resistant states. *Clin Chim Acta* 368:1–19, 2006
- Hadigan C, Rabe J, Meininger G, Aliabadi N, Breu J, Grinspoon S: Inhibition of lipolysis improves insulin sensitivity in protease inhibitor-treated HIV-infected men with fat redistribution. *Am J Clin Nutr* 77:490–494, 2003
- Hadigan C, Liebaw J, Torriani M, Andersen R, Grinspoon S: Improved triglycerides and insulin sensitivity with 3 months of acipimox in human immunodeficiency virus-infected patients with hypertriglyceridemia. *J Clin Endocrinol Metab* 91:4438–4444, 2006
- Dresner A, Laurent D, Marcucci M, Griffin ME, Dufour S, Cline GW, Slezak LA, Andersen DK, Hundal RS, Rothman DL, Petersen KF, Shulman GI: Effects of free fatty acids on glucose transport and IRS-1-associated phosphatidylinositol 3-kinase activity. *J Clin Invest* 103:253–259, 1999
- Boden G, Chen X, Ruiz J, White JV, Rossetti L: Mechanisms of fatty acid-induced inhibition of glucose uptake. *J Clin Invest* 93:2438–2446, 1994
- Roden M, Price TB, Perseghin G, Petersen KF, Rothman DL, Cline GW, Shulman GI: Mechanism of free fatty acid-induced insulin resistance in humans. *J Clin Invest* 97:2859–2865, 1996
- Griffin ME, Marcucci MJ, Cline GW, Bell K, Barucci N, Lee D, Goodyear LJ, Kraegen EW, White MF, Shulman GI: Free fatty acid-induced insulin resistance is associated with activation of protein kinase C θ and alterations in the insulin signaling cascade. *Diabetes* 48:1270–1274, 1999
- Boden G, Jadali F, White J, Liang Y, Mozzoli M, Chen X, Coleman E, Smith C: Effects of fat on insulin-stimulated carbohydrate metabolism in normal men. *J Clin Invest* 88:960–966, 1991
- Kelley DE, Mookan M, Simoneau JA, Mandarino LJ: Interaction between glucose and free fatty acid metabolism in human skeletal muscle. *J Clin Invest* 92:91–98, 1993
- Vaag A, Skott P, Damsbo P, Gall MA, Richter EA, Beck-Nielsen H: Effect of the antilipolytic nicotinic acid analogue acipimox on whole-body and skeletal muscle glucose metabolism in patients with non-insulin-dependent diabetes mellitus. *J Clin Invest* 88:1282–1290, 1991
- Boden G, Cheung P, Stein TP, Kresge K, Mozzoli M: FFA cause hepatic insulin resistance by inhibiting insulin suppression of glycogenolysis. *Am J Physiol Endocrinol Metab* 283:E12–E19, 2002
- Belfort R, Mandarino L, Kashyap S, Wirfel K, Pratipanawat T, Berria R, DeFronzo RA, Cusi K: Dose-response effect of elevated plasma free fatty acid on insulin signaling. *Diabetes* 54:1640–1648, 2005
- Yu C, Chen Y, Cline GW, Zhang D, Zong H, Wang Y, Bergeron R, Kim JK, Cushman SW, Cooney GJ, Atcheson B, White MF, Kraegen EW, Shulman GI: Mechanism by which fatty acids inhibit insulin activation of insulin receptor substrate-1 (IRS-1)-associated phosphatidylinositol 3-kinase activity in muscle. *J Biol Chem* 277:50230–50236, 2002
- Kruszynska YT, Worrall DS, Ofrecio J, Frias JP, Macaraeg G, Olefsky JM: Fatty acid-induced insulin resistance: decreased muscle PI3K activation but unchanged Akt phosphorylation. *J Clin Endocrinol Metab* 87:226–234, 2002
- Kim YB, Shulman GI, Kahn BB: Fatty acid infusion selectively impairs insulin action on Akt1 and protein kinase C λ 2 but not on glycogen synthase kinase-3. *J Biol Chem* 277:32915–32922, 2002
- van d, V, Bisschop PH, Romijn JA, Ackermans MT, Lange JM, Endert E, Reiss P, Sauerwein HP: Lipodystrophy in HIV-1-positive patients is associated with insulin resistance in multiple metabolic pathways. *AIDS* 15:2093–2100, 2001
- Behrens GM, Boerner AR, Weber K, van den HJ, Ockenga J, Brabant G, Schmidt RE: Impaired glucose phosphorylation and transport in skeletal muscle cause insulin resistance in HIV-1-infected patients with lipodystrophy. *J Clin Invest* 110:1319–1327, 2002
- Haugaard SB, Andersen O, Madsbad S, Frosig C, Iversen J, Nielsen JO, Wojtaszewski JF: Skeletal muscle insulin signaling defects downstream of phosphatidylinositol 3-kinase at the level of Akt are associated with impaired nonoxidative glucose disposal in HIV lipodystrophy. *Diabetes* 54:3474–3483, 2005
- Driscoll SD, Meininger GE, Lareau MT, Dolan SE, Killilea KM, Hadigan CM, Lloyd-Jones DM, Klibanski A, Frontera WR, Grinspoon SK: Effects of exercise training and metformin on body composition and cardiovascular indices in HIV-infected patients. *AIDS* 18:465–473, 2004
- Carr A, Emery S, Law M, Puls R, Lundgren JD, Powderly WG: An objective case definition of lipodystrophy in HIV-infected adults: a case-control study. *Lancet* 361:726–735, 2003
- Santomauro AT, Boden G, Silva ME, Rocha DM, Santos RF, Ursich MJ, Strassmann PG, Wajchenberg BL: Overnight lowering of free fatty acids with Acipimox improves insulin resistance and glucose tolerance in obese diabetic and nondiabetic subjects. *Diabetes* 48:1836–1841, 1999
- Fulcher GR, Walker M, Catalano C, Agius L, Alberti KG: Metabolic effects of suppression of nonesterified fatty acid levels with acipimox in obese NIDDM subjects. *Diabetes* 41:1400–1408, 1992
- Wolfe RR, Chinkes D: *Isotope Tracers in Metabolic Research: Principles and Practice of Kinetic Analysis*. Hoboken, NJ, Wiley and Sons, 2005
- Galster AD, Clutter WE, Cryer PE, Collins JA, Bier DM: Epinephrine plasma thresholds for lipolytic effects in man: measurements of fatty acid transport with [1-13C]palmitic acid. *J Clin Invest* 67:1729–1738, 1981
- DeFronzo RA, Tobin JD, Andres R: Glucose clamp technique: a method for quantifying insulin secretion and resistance. *Am J Physiol* 237:E214–E223, 1979
- Plomgaard P, Bouzakri K, Krogh-Madsen R, Mittendorfer B, Zierath JR, Pedersen BK: Tumor necrosis factor- α induces skeletal muscle insulin resistance in healthy human subjects via inhibition of Akt substrate 160 phosphorylation. *Diabetes* 54:2939–2945, 2005
- Patterson BW, Zhao G, Elias N, Hachey DL, Klein S: Validation of a new procedure to determine plasma fatty acid concentration and isotopic enrichment. *J Lipid Res* 40:2118–2124, 1999

37. Hojlund K, Staehr P, Hansen BF, Green KA, Hardie DG, Richter EA, Beck-Nielsen H, Wojtaszewski JF: Increased phosphorylation of skeletal muscle glycogen synthase at NH₂-terminal sites during physiological hyperinsulinemia in type 2 diabetes. *Diabetes* 52:1393–1402, 2003
38. Thomas JA, Schlender KK, Lerner J: A rapid filter paper assay for UDP glucose-glycogen glucosyltransferase, including an improved biosynthesis of UDP-14C-glucose. *Anal Biochem* 25:486–499, 1968
39. R Development Core Team. R: *A Language and Environment for Statistical Computing*. Vienna, Austria, R Foundation for Statistical Computing, 2003
40. National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III): Third report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. *Circulation* 106:3143–3421, 2002
41. Mittendorfer B, Horowitz JF, Klein S: Gender differences in lipid and glucose kinetics during short-term fasting. *Am J Physiol Endocrinol Metab* 281:E1333–E1339, 2001
42. Saloranta C, Franssila-Kallunki A, Ekstrand A, Taskinen MR, Groop L: Modulation of hepatic glucose production by non-esterified fatty acids in type 2 (non-insulin-dependent) diabetes mellitus. *Diabetologia* 34:409–415, 1991
43. Walker M, Agius L, Orskov H, Alberti KG: Peripheral and hepatic insulin sensitivity in non-insulin-dependent diabetes mellitus: effect of nonesterified fatty acids. *Metabolism* 42:601–608, 1993
44. Saloranta C, Groop L, Ekstrand A, Franssila-Kallunki A, Eriksson J, Taskinen MR: Different acute and chronic effects of acipimox treatment on glucose and lipid metabolism in patients with type 2 diabetes. *Diabet Med* 10:950–957, 1993
45. Bajaj M, Suraamornkul S, Romanelli A, Cline GW, Mandarino LJ, Shulman GI, DeFronzo RA: Effect of a sustained reduction in plasma free fatty acid concentration on intramuscular long-chain fatty acyl-CoAs and insulin action in type 2 diabetic patients. *Diabetes* 54:3148–3153, 2005
46. Ferrannini E, Barrett EJ, Bevilacqua S, DeFronzo RA: Effect of fatty acids on glucose production and utilization in man. *J Clin Invest* 72:1737–1747, 1983
47. Bevilacqua S, Bonadonna R, Buzzigoli G, Boni C, Ciociaro D, Maccari F, Giorico MA, Ferrannini E: Acute elevation of free fatty acid levels leads to hepatic insulin resistance in obese subjects. *Metabolism* 36:502–506, 1987
48. Saloranta C, Koivisto V, Widen E, Falholt K, DeFronzo RA, Harkonen M, Groop L: Contribution of muscle and liver to glucose-fatty acid cycle in humans. *Am J Physiol* 264:E599–E605, 1993
49. Shulman GI, Rothman DL, Jue T, Stein P, DeFronzo RA, Shulman RG: Quantitation of muscle glycogen synthesis in normal subjects and subjects with non-insulin-dependent diabetes by ¹³C nuclear magnetic resonance spectroscopy. *N Engl J Med* 322:223–228, 1990
50. Skurat AV, Roach PJ: Phosphorylation of sites 3a and 3b (Ser640 and Ser644) in the control of rabbit muscle glycogen synthase. *J Biol Chem* 270:12491–12497, 1995
51. Lawrence JC Jr, Roach PJ: New insights into the role and mechanism of glycogen synthase activation by insulin. *Diabetes* 46:541–547, 1997
52. Skurat AV, Dietrich AD, Roach PJ: Glycogen synthase sensitivity to insulin and glucose-6-phosphate is mediated by both NH₂- and COOH-terminal phosphorylation sites. *Diabetes* 49:1096–1100, 2000
53. Tunaru S, Kero J, Schaub A, Wufka C, Blaukat A, Pfeffer K, Offermanns S: PUMA-G and HM74 are receptors for nicotinic acid and mediate its antilipolytic effect. *Nat Med* 9:352–355, 2003