

## Type 1 Diabetes

# Evidence for Susceptibility Loci from Four Genome-Wide Linkage Scans in 1,435 Multiplex Families

Patrick Concannon,<sup>1</sup> Henry A. Erlich,<sup>2</sup> Cecile Julier,<sup>3</sup> Grant Morahan,<sup>4</sup> Jørn Nerup,<sup>5</sup> Flemming Pociot,<sup>5</sup> John A. Todd,<sup>6</sup> Stephen S. Rich,<sup>7</sup> and the Type 1 Diabetes Genetics Consortium\*

Type 1 diabetes is a common, multifactorial disease with strong familial clustering (genetic risk ratio [ $\lambda_S$ ]  $\sim 15$ ). Approximately 40% of the familial aggregation of type 1 diabetes can be attributed to allelic variation of HLA loci in the major histocompatibility complex on chromosome 6p21 (locus-specific  $\lambda_S \sim 3$ ). Three other disease susceptibility loci have been clearly demonstrated based on their direct effect on risk, *INS* (chromosome 11p15, allelic odds ratio [OR]  $\sim 1.9$ ), *CTLA4* (chromosome 2q33, allelic OR  $\sim 1.2$ ), and *PTPN22* (chromosome 1p13, allelic OR  $\sim 1.7$ ). However, a large proportion of type 1 diabetes clustering remains unexplained. We report here on a combined linkage analysis of four datasets, three previously published genome scans, and one new genome scan of 254 families, which were consolidated through an international consortium for type 1 diabetes genetic studies ([www.t1dgc.org](http://www.t1dgc.org)) and provided a total sample of 1,435 families with 1,636 affected sibpairs. In addition to the HLA region (nominal  $P = 2.0 \times 10^{-52}$ ), nine non-HLA-linked regions showed some evidence of linkage to type 1 diabetes (nominal  $P < 0.01$ ), including three at (or near) genome-wide significance ( $P < 0.05$ ): 2q31-q33, 10p14-q11, and 16q22-q24. In addition, after taking into account the linkage at the 6p21 (HLA) region, there was evidence supporting linkage for the 6q21 region (empiric  $P < 10^{-4}$ ). More than 80% of the genome could be excluded as harboring type 1 diabetes susceptibility genes of modest effect ( $\lambda_S \geq 1.3$ ) that could be detected by linkage. This study represents one of the largest linkage studies ever performed for any common disease. The results demonstrate some consistency emerging for the existence of susceptibility loci on chromosomes 2q31-q33, 6q21, 10p14-q11, and 16q22-q24 but diminished

support for some previously reported locations. *Diabetes* 54:2995–3001, 2005

Type 1 diabetes is the third most prevalent chronic disease of childhood, affecting up to 0.4% of children in some populations by age 30 years, with an overall lifetime risk of nearly 1% (1,2). It is believed that a large proportion of cases of type 1 diabetes result from the autoimmune destruction of the pancreatic  $\beta$  cells, leading to complete dependence on exogenous insulin to regulate blood glucose levels (3). The etiology of type 1 diabetes is only partially characterized, but it is recognized that both genetic and environmental determinants are important in defining disease risk. Type 1 diabetes clusters in families, based on population-based twin and family studies (4) but does not segregate with a known mode of inheritance (5). The incidence and the age at onset of type 1 diabetes in some populations have changed dramatically since 1950 (6–8). These data, coupled with the incomplete concordance for the phenotype in monozygotic twins (30–70%), suggest that the penetrance of type 1 diabetes susceptibility alleles is strongly influenced by environmental factors (4).

Type 1 diabetes is strongly clustered in families with an overall genetic risk ratio ( $\lambda_S$ ) of  $\sim 15$  (9). At least one locus that contributes strongly to this familial clustering resides within the major histocompatibility complex (MHC) on chromosome 6p21. Genetic, functional, structural, and model studies all suggest that the HLA class II genes (*HLA-DRB1* and *-DQB1*) likely represent the primary determinants of *IDDM1*. The frequency of HLA class II susceptibility alleles also correlates well with the population incidence of type 1 diabetes (10). These studies suggest that the MHC (*IDDM1*) may account for nearly 40% of the observed familial clustering of type 1 diabetes, with a locus-specific  $\lambda_S$  of  $\sim 3$  (11).

Given that HLA alone cannot explain the familial clustering of type 1 diabetes, several, perhaps many, genes remain to be identified. First, in the general population, individuals who carry the high-risk haplotypic combination *DRB1\*04-DQB1\*0302/DRB1\*03-DQB1\*0201* have  $\sim 5\%$  absolute risk of type 1 diabetes. However, within affected sibpair families, this genotype has  $\sim 20\%$  risk (5,12). Second, three non-HLA loci have been identified based on genetic association studies: *IDDM2* [*INS*, 11p15 (13–16)], *IDDM12* [*CTLA4*, 2q33 (17)], and *LYP/PTPN22* [1p13 (18–20)]. Finally, the observed risk of type 1 diabetes in

From the <sup>1</sup>Benaroya Research Institute, Seattle, Washington; <sup>2</sup>Roche Molecular Systems, Alameda, California; <sup>3</sup>Pasteur Institute, Paris, France; <sup>4</sup>The Western Australian Institute of Medical Research, Perth, Australia; the <sup>5</sup>Steno Diabetes Center, Gentofte, Denmark; the <sup>6</sup>Juvenile Diabetes Research Foundation/Wellcome Trust Diabetes Inflammation Laboratory, Cambridge University, Cambridge, U.K.; and <sup>7</sup>Wake Forest University School of Medicine, Winston-Salem, North Carolina.

Address correspondence and reprint requests to Stephen S. Rich, PhD, Department of Public Health Sciences, Wake Forest University School of Medicine, Winston-Salem, NC 27157. E-mail: [srich@wfubmc.edu](mailto:srich@wfubmc.edu).

Received for publication 12 October 2004 and accepted in revised form 29 June 2005.

\*A complete list of the members of the Type 1 Diabetes Genetics Consortium can be found in the APPENDIX.

Additional information for this article can be found in an online appendix at <http://diabetes.diabetesjournals.org>.

IBD, identity-by-descent; LOD, logarithm of odds; MHC, major histocompatibility complex; SNP, single nucleotide polymorphism.

© 2005 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

first- and second-degree relatives declines in a pattern consistent with multiplicative effects of multiple loci (11).

The first type 1 diabetes genome-wide scans for linkage, using fewer than 100 affected sibpair families, identified chromosome 6p21 (*IDDM1*) as the major type 1 diabetes risk locus (21,22). Subsequent studies supported non-HLA loci on chromosomes 11q13 (*IDDM4*) and 6q25 (*IDDM5*) in families from the U.K., the U.S., and France and on chromosome 15q26 (*IDDM3*) in families from Canada. Using both linkage and association approaches, other putative type 1 diabetes susceptibility loci were identified on chromosomes 18q12-q21 (*IDDM6*), 2q33 (*IDDM7*), 6q27 (*IDDM8*), 3q22-q25 (*IDDM9*), 10p11-q11 (*IDDM10*), 14q24-q31 (*IDDM11*), 2q31-q33 (*IDDM12*), 2q34-q35 (*IDDM13*), 6q21 (*IDDM15*), 14q32 (*IDDM16*), 10q25 (*IDDM17*), 5q33 (*IDDM18*), 7p15-p13 (*GCK*), 1q42, 16q22-q24, Xp11 (conditional on HLA-DR genotype), and 8q22-q24 (8–10,12–17). Even though statistical evidence supporting linkage for some of these regions was strong in the initial reports, most regions have not been clearly established in multiple populations (9,23).

A major barrier to type 1 diabetes gene identification, given the likely small locus-specific contribution (low  $\lambda_S$ ) for non-HLA genes, is the relatively small number of newly ascertained affected sibpair families with type 1 diabetes. In addition, previous studies have used available samples from a variety of collections, making the compilation of linkage results difficult because of apparent overlap in families analyzed and uncertainty in the equivalence of allele coding. To facilitate the genetic analysis of type 1 diabetes, results from two previous genome scans of type 1 diabetes were merged, comprising 767 families and 831 affected sibpairs from the U.K. and U.S. (24). The combined analyses supported linkage to at least six non-HLA regions to type 1 diabetes, including *IDDM2* (*INS*, nominal  $P = 6.5 \times 10^{-4}$ ), 2q31-q33 ( $P = 5.1 \times 10^{-4}$ ), and 10p11 ( $P = 3.2 \times 10^{-4}$ ). A third genome scan of 424 type 1 diabetic families with at least two affected relative pairs (464 affected pairs) from Scandinavia (25) found no evidence for linkage at these latter three loci but did support linkage on chromosomes 5q11.2 (nominal  $P = 8.1 \times 10^{-4}$ ) and 16p13 ( $P = 1.6 \times 10^{-4}$ ). The *IDDM15* region on chromosome 6q21 supported linkage ( $P = 7.0 \times 10^{-7}$ ) when HLA was taken into consideration in a combined analysis of U.S., French, and Scandinavian families.

We present here a joint analysis of data from these three prior genome-wide scans (U.S., U.K., and Scandinavia) as well as 254 new families collected for this study, a total of 1,435 multiplex families, for linkage to type 1 diabetes. With an average map information content of 67% (from ~400 polymorphic microsatellite markers in each scan), this family collection provides ~95% power to detect a locus with locus-specific  $\lambda_S \geq 1.3$  and  $P = 10^{-4}$ . In addition to HLA, there was nominal evidence for linkage of type 1 diabetes to 10 other chromosome regions, including 6q21 (*IDDM15*) and 3 that reached genome-wide levels of significance, 2q31-q33 (*IDDM12* and *IDDM7*), 10p11-q14 (*IDDM10*), and 16p12-q24. These data support the existence of non-HLA susceptibility loci for type 1 diabetes and strengthen support for a subset of loci previously proposed to contribute to type 1 diabetes risk.

## RESEARCH DESIGN AND METHODS

Four sets of Caucasian families provided genome scan data for the combined analyses. Three sets of families have been previously published—U.K. (21,26), U.S. (24,26), and Scandinavia (25)—and one set of 254 families that were

newly assembled for this study. The new collection of DNA samples from 254 families was obtained from several sources. DNA samples from 47 U.K. families were identified from the Diabetes U.K. Warren repository (28) that had not been genotyped previously. Families not previously used in published genome scans from the U.S. were contributed by the Joslin Diabetes Center (121 families) and from the Human Biological Data Interchange (76 families). Ten families from Australia were collected by investigators at the Walter and Eliza Hall Institute as previously described (29). In total, there were 1,435 families containing 6,899 individuals (6,358 with genome scan data). A total of 3,109 individuals were affected (with type 1 diabetes), and of these, 3,072 had genotype data (for details of the samples, see table in online appendix [available at <http://diabetes.diabetesjournals.org>]).

**Genotyping.** Microsatellite marker genotyping technologies and allele scoring conventions varied between the different laboratories providing the data for previously published type 1 diabetic families. Details of genotyping of the U.K. (21), U.S. (27), and Scandinavian (25) families have been previously described. The 254 new families were genotyped by the Center for the Inheritance of Disease Research (<http://www.cidr.jhmi.edu/>) using a panel of 405 microsatellite markers. Because direct merging of genotypes (by standardized allele size) was not possible, within-family recoded genotype data from all four sources were merged into a single database. Family-naming conventions between the samples were normalized, and individual marker names were modified to indicate the laboratory of origin for the genotyping. After elimination of marker inconsistencies (see below), genetic markers were selected to form the analysis panel. Multiple independent genotyping occurred for some markers on a subset of individuals by different laboratories. In these cases, only one marker was used for the current analysis. Unless a marker showed inconsistencies in identity-by-descent (IBD) sharing, the marker that was included for analysis was the one scored in the most samples. A total of 1,190 markers were included in the combined analyses. Markers that previously had been added to maps because of association with type 1 diabetes were excluded to avoid bias in the multipoint linkage results. The excluded markers were located on chromosome 2 (*alpha4*, *ND1*, *D2S152*, *CTLA4*, and *IGFBP5*) and chromosome 11 (*INS*, *TH*).

**Statistical analysis.** Before integration of the genetic data, marker error detection and pedigree structure within each dataset were made using PREST software (30). This method uses the genome scan data to determine the likelihood of each specified relationship given the genetic data. Unlikely marker genotypes were resolved by recoding the specific genotype to “unknown.” Occurrences of nonpaternity were resolved by changing the pedigree structure to that which was most likely and then repeating the analysis to confirm appropriate relationships. An integrated marker map was developed by using public databases (Mammalian Genotyping Service, [http://research.marshfieldclinic.org/genetics/Genotyping\\_Service/mgsver2.htm](http://research.marshfieldclinic.org/genetics/Genotyping_Service/mgsver2.htm); Southampton, [http://cedar.genetics.soton.ac.uk/public\\_html/](http://cedar.genetics.soton.ac.uk/public_html/); Cooperative Human Linkage Center, <http://gai.nci.nih.gov/CHLC/>; deCODE, <http://www.decode.is>) as well as physical map and genome sequence information from the University of California at Santa Cruz (<http://genome.ucsc.edu/>) using primer sequences in BLAST searches against the genome sequence. All analyses were based on this “consensus” map. Single and multipoint linkage analyses (based on the consensus map order and distances) were performed using GeneHunter-plus [ $S_{\text{pairs}}$  option (31–33)]. Examination of double recombinants was performed using Merlin software (34). Information content was estimated using Allegro (35).

Estimation of IBD statistics and resulting likelihoods under the null and alternative hypotheses were computed within each dataset. These dataset-specific likelihoods were then combined for the combined linkage analyses. Multipoint linkage analyses were performed, and maximized logarithm of odds (LOD) scores were calculated under an exponential model with  $\delta$  constrained between 0 and 2 (32). Genome-wide empirical  $P$  values were determined by simulating Mendelian transmission with families maintaining the patterns of missing data observed in the sample (36). The fraction of LOD scores observed greater than the nominal value, based on simulations of 10,000 replicates across the genome, provided the estimated genome-wide  $P$  value. Exclusion mapping was performed using the MapMaker/SIBS program (37).

Previous studies have identified a potential type 1 diabetes susceptibility locus near (but not within) the HLA complex (*IDDM15*) (25,27,38). In the presence of strong support for linkage due to *IDDM1*, determining the support for *IDDM15* is complex because of the positive correlation among the IBD proportions for linked loci. Without accounting for this positive correlation, statistical tests are biased toward inferring an epistatic relationship. A simple approach when IBD is known is to compare the observed correlation,  $r$ , between the IBD estimates at two loci (i.e., *IDDM1* and *IDDM15*) with the theoretical correlation,  $\rho = (1-2\theta)^2$ , for two loci separated by recombination fraction,  $\theta$ . The statistic  $t = (z - \zeta)\sqrt{n-3}$ , where  $z = \frac{1}{2} \ln \left( \frac{1+r}{1-r} \right)$  and




FIG. 1. Genome-wide linkage analysis of type 1 diabetes in 1,435 multiplex families.

$\zeta = \frac{1}{2} \ln \left( \frac{1+\rho}{1-\rho} \right)$  allows for the test of “interaction” that contrasts the observed IBD at *IDDM15* based on the distance of *IDDM15* from *IDDM1* and the expected IBD given that distance. Using a sex-averaged map, IBD estimates for each sibpair in the data were computed, and a single IBD estimate was selected from each pedigree. Thus, each of these IBD estimates is independent. Under the null hypothesis of no interaction between these loci,  $t$  approximately has a standard normal distribution. Observed correlations greater than  $\rho$  reflect increased sharing at *IDDM15* over that expected from IBD at *IDDM1* and the hypothesized genetic distance of *IDDM15* from *IDDM1*. A series of 10,000 simulations were performed as described above (36), with the correlation in IBD computed between *IDDM1* and *IDDM15*. The empirical  $P$  value for significance of *IDDM15* was based on the number of simulated correlations greater than that observed in the original data.

## RESULTS

**Linkage analysis.** Analysis of 1,190 genetic markers in 1,435 families revealed that the strongest evidence for linkage to type 1 diabetes was on chromosome 6p21 (nominal  $P = 2.0 \times 10^{-52}$ ) in the MHC (Fig. 1). There were nine non-HLA-linked regions with nominal evidence supporting linkage to type 1 diabetes ( $P < 0.01$ ), including 2q31-q33 (*IDDM7* and *IDDM12*;  $P = 9.0 \times 10^{-5}$ ; genome-wide  $P = 0.016$ ), 10p14-q11 (*IDDM10*,  $P = 1.2 \times 10^{-4}$ ; genome-wide  $P = 0.021$ ), and 16q22-q24 ( $P = 4.9 \times 10^{-4}$ ; genome-wide  $P = 0.075$ ). The estimates of genetic relative varied by region, with *IDDM1* having the largest locus-specific effect ( $\lambda_S \sim 3.3$ ) and the other sites having

low locus-specific effects ( $\lambda_S \sim 1.1$ – $1.2$ ). Individual linkage plots by chromosome are provided in the online appendix (supplementary figures, upper curves).

**Exclusion mapping.** Each of the four datasets had the equivalent of an  $\sim 9$ -centiMorgan (cM) map, providing an average marker information content of  $\sim 66\%$ . The range of information from the markers was 62% for chromosome 15 to 73% for chromosome 16. In the combined data, 82% of the genome could be excluded at  $\text{LOD} < -2$  for loci of effect size  $\lambda_S \geq 1.3$  (supplementary figures, lower curves), and  $>95\%$  could be excluded for  $\lambda_S \geq 1.5$ . Several entire chromosomes could be excluded (chromosomes 7, 8, 18, 20, 21, and 22). The majority of chromosome 6 (due to the strongly linked 6p21/MHC region) and  $<50\%$  of chromosomes 16, 19, and X could be excluded for  $\lambda_S \geq 1.3$ . For effects of  $\lambda_S \geq 1.1$ , only 6% of the genome could be excluded. The extent of exclusion for each chromosome is shown in Table 1.

**IDDM15.** The  $t$ -statistic was computed to determine the increase in sharing at *IDDM15*, beyond that expected from sharing at *IDDM1* and expected decay in sharing due to genetic distance. In the current data, the test statistic was computed for *IDDM15* at a position 37 cM from *IDDM1* (at 47 cM on chromosome 6), using a sex-averaged map. The test statistic was also computed for a range of map positions within 5 cM (32–42 cM) with similar results. At

TABLE 1  
Exclusion mapping of the type 1 diabetes multiplex family data

Chromosome	Excluded $\lambda_S$ (%)			
	~1.1	~1.3	~1.5	~1.7
1	3.52	82.17	100	100
2	1.15	73.76	94.68	100
3	8.93	84.96	99.12	100
4	6.90	95.61	100	100
5	9.74	96.45	100	100
6	0.00	30.77	39.49	41.54
7	0.00	100	100	100
8	0.00	100	100	100
9	12.27	84.76	96.34	100
10	0.00	75.88	100	100
11	4.14	81.63	97.96	100
12	0.00	98.24	100	100
13	0.00	73.33	100	100
14	13.67	82.27	100	100
15	0.86	95.76	100	100
16	0.00	24.24	87.12	100
17	0.00	79.53	100	100
18	23.81	100	100	100
19	0.00	47.83	100	100
20	56.7	100	100	100
21	0.00	100	100	100
22	0.00	100	100	100
X	0.00	39.00	83.00	97.00
Total	5.57	81.64	95.34	96.68

the 37-cM distance from *IDDM1* and *IDDM15*, the expected correlation coefficient between the IBD estimates under the null hypothesis was estimated as  $\rho = 0.2276$ . The observed Pearson's correlation coefficient between the IBD estimates at *IDDM1* and *IDDM15*, using 1,401 informative pedigrees, was  $r = 0.3132$  (empirical  $P < 1.0 \times 10^{-4}$ ). These results support an HLA-independent effect in the *IDDM15* region.

## DISCUSSION

In a previous combined analysis of U.K. and U.S. families (24), it was concluded that an effort to merge and jointly analyze existing families would be required to clarify the role of non-HLA-linked loci in type 1 diabetes. In the present study, this effort has been achieved under the auspices of the Type 1 Diabetes Genetics Consortium (<http://www.t1dgc.org>). We have assembled families and

merged data from three large genome scans and added new data from 254 families not previously scanned. This increased sample size has allowed the exclusion of >80% of the human genome for locus-specific, but population-independent, effects of  $\lambda_S \geq 1.3$ . In addition to continued support for type 1 diabetes susceptibility related to the MHC (*IDDM1*) and *INS* (*IDDM2*), we identified eight regions that supported non-HLA-linked susceptibility. Furthermore, we identified three chromosomes that contained extensive areas for which linkage could not be excluded—chromosomes 16, 19, and X—that could benefit from further genotyping to increase information content and the analysis of additional families.

Three non-HLA-linked regions provided support for linkage at, or near, the genome-wide level of significance ( $P < 0.05$ ): chromosome 2q31-q33 (*IDDM7* and *IDDM12*), 10p14-q11 (*IDDM10*), and 16q22-q24. These locations were unlikely to have occurred by chance, based on our simulations. Together with the six other non-HLA-linked regions that exhibited evidence of linkage at nominal  $P < 0.01$  (Table 2) and the 10-cM map, the data indicate a strong non-HLA genetic effect for type 1 diabetes (39). Furthermore, none of the three most strongly linked regions exhibited support for linkage in the 408 families from Scandinavia (25).

Strong support for linkage (nominal  $P = 7.0 \times 10^{-7}$ ), after taking into account linkage to HLA) to the *IDDM15* locus (6q21), was observed previously in a combined analysis of French, U.S., and Scandinavian families (25,38). In the present study, we have obtained support for *IDDM15* (empirical  $P < 1.0 \times 10^{-4}$ ). The ability to further define the effects of this locus will be facilitated by increased information content in the HLA region and in the region surrounding *IDDM15* to better estimate the observed IBD sharing and model the residual linkage to the HLA region, including taking into account sex-specific genetic map differences.

The *IDDM12* locus lies within the 2q31-q33 region and has been attributed to single nucleotide polymorphisms (SNPs) in the 3'-untranslated region of *CTLA4* (15); however, the modest  $\lambda_S$  value predicted from the odds ratios (ORs) (1.1–1.2) of the disease-associated SNPs at *CTLA4* ( $\lambda_S \sim 1.01$ ) in type 1 diabetes seems unlikely to fully account for the magnitude of the observed evidence for linkage (regional  $\lambda_S \sim 1.19$ ). This result suggests the presence of other loci in the 2q31-q33 region, if this linkage is confirmed in other future studies. Originally, *IDDM7* at

TABLE 2  
Multipoint linkage analysis of four genome scans for type 1 diabetes ( $P < 0.01$ )

Chromosome	Position (cM)	Closest marker	Nominal LOD	Nominal $P$ value	$\lambda_S$	Genome-wide $P$ value*
2q31-q33	192	<i>D2S2167</i>	3.34	$9.0 \times 10^{-5}$	1.19	0.016
3p13-p14	98	<i>D3S1261</i>	1.52	$8.2 \times 10^{-3}$	1.15	0.649
6p21	47	<i>TNFA</i>	116.30	$4.9 \times 10^{-52}$	3.35	$<1.0 \times 10^{-4}$
6q21	80	<i>D6S283</i>	22.39	$7.0 \times 10^{-7}$	1.56	†
9q33-q34	150	<i>D9S260</i>	2.20	$1.5 \times 10^{-3}$	1.13	0.191
10p14-q11	61	<i>D10S1426</i>	3.21	$1.2 \times 10^{-4}$	1.12	0.021
11p15	2	<i>D11S922</i>	1.87	$3.4 \times 10^{-3}$	1.16	0.371
12q14-q12	81	<i>D12S375</i>	1.66	$5.8 \times 10^{-3}$	1.10	0.528
16p12-q11.1	56	<i>D16S3131</i>	1.88	$3.3 \times 10^{-3}$	1.17	0.363
16q22-q24	108	<i>D16S504</i>	2.64	$4.9 \times 10^{-4}$	1.19	0.075
19p13.3-p13.2	25	<i>INSR</i>	1.92	$3.0 \times 10^{-3}$	1.15	0.338

\*All genome-wide empiric  $P$  values are based on simulations of 10,000 replicates. †Empirical  $P$  value determined by simulation and estimated correlation of IBD between HLA and *IDDM15*, not genome-wide (see RESEARCH DESIGN AND METHODS); empirical  $P < 1.0 \times 10^{-4}$ .

chromosome 2q33 was assigned on the basis of evidence of allelic association of the *D2S152* microsatellite marker, but this association has not been substantiated. The evidence supporting linkage in the current study does not include the putative *IDDM13* locus at chromosome 2q34-q35 (29). Linkage of type 1 diabetes to 10p14-q13 (*IDDM10*) is well supported by the current and past studies (24,26); however, there has been little follow-up other than association analyses of the functional candidate gene *GAD2*, suggesting that this gene is not a type 1 diabetes susceptibility locus (40,41). The observed locus-specific effects for 2q31-q33 ( $\lambda_S \sim 1.19$ ) and 10p14-q11 ( $\lambda_S \sim 1.12$ ) suggest that a single common susceptibility allele would have an allelic association (OR)  $\sim 3$ . This effect should be identifiable in a fine-mapping association study using dense SNP maps across the regions.

Support for a type 1 diabetes susceptibility locus on chromosome 16p12-q11.1, which was observed independently in both the combined U.K. and U.S. families (nominal  $P = 4.5 \times 10^{-3}$ ) and in the Scandinavian families ( $P = 2 \times 10^{-4}$ ), remained in the present study ( $P = 3.3 \times 10^{-3}$ ). A recent analysis of four rheumatoid arthritis genome scans (42) reported evidence for linkage at chromosomes 6p21 (HLA;  $P = 2 \times 10^{-5}$ ) and 16p-cen ( $P = 0.004$ ). Because rheumatoid arthritis, antithyroid autoimmune disease, and type 1 diabetes cluster in families more often than expected by chance (43), evidence for linkage for any one of these autoimmune diseases could be informative for others. Evidence for linkage in U.K. families with early-onset rheumatoid arthritis (44) to chromosome 16p has previously been demonstrated ( $P = 3.2 \times 10^{-4}$ ). The comparison between linkage scan results for type 1 diabetes and rheumatoid arthritis provides other interesting similarities. The largest, single, combined scan of rheumatoid arthritis families (45) reported significant linkage of rheumatoid arthritis to chromosome 6p21 (HLA;  $P = 5 \times 10^{-12}$ ), and some evidence ( $P < 0.005$ ) of rheumatoid arthritis linked to six other regions (1q43, 6q21, 10q21, 12q12, 17p13, and 18q21). This overlap with potential type 1 diabetes susceptibility at 6q21, 12q12, and 16p-cen may not be coincidental in the etiology of these autoimmune diseases.

Recently, evidence for association of type 1 diabetes with alleles in the *PTPN22* locus (chromosome 1p13) has been reported (18), and this association has been confirmed (19,20). *PTPN22* encodes a lymphoid-specific tyrosine phosphatase (LYP) and is also associated with autoimmune thyroid disease, rheumatoid arthritis, and SLE (46). The absence of evidence supporting linkage of type 1 diabetes to chromosome 1p13 (*DIS206*) in the 1,435 families studied here is not surprising given the magnitude of the *PTPN22* association with type 1 diabetes. The OR of *PTPN22* is large ( $\sim 1.7$ ) but the  $\lambda_S$  is  $\sim 1.05$ . Thus, to detect linkage at  $P < 0.001$  with 50% power, a sample of  $>8,000$  affected sibpair families would be required, using a fully informative genetic map. Assuming a multiplicative model, the contribution of *PTPN22* to type 1 diabetes (based on the observed OR) is  $\sim 2\%$ , much lower than HLA (40–50%). Nevertheless, the knowledge that *PTPN22* may be involved in risk to type 1 diabetes, as well as in other autoimmune diseases, is significant and could provide insight into modulating T-cell activity for disease prevention.

Several previously supported regions of linkage have diminished support in the current analyses. Type 1 diabetes susceptibility locus on chromosome 1q42 was strongly

supported (nominal  $P = 9.8 \times 10^{-5}$ ) in a study of 679 U.K. and U.S. families (24,47) but exhibited decreasing linkage support in a follow-up analysis of 616 families using a denser map in the region ( $P = 4.0 \times 10^{-4}$ ). The evaluation of previously supported regions is difficult, even in the present study with over 1,600 affected sibpairs, particularly for regions with low  $\lambda_S$  (e.g.,  $\lambda_S \sim 1.1$ ). For example, the region on 1q42 was originally supported with  $MLS = 3.31$  and  $\lambda_S \sim 1.5$  (27). Although the current sample excludes this region at the reported  $\lambda_S \sim 1.5$ , support for linkage in this region has decreased with increasing sample size from a  $LOD = 2.20$  (24), to the current  $LOD = 0.87$  (nominal  $P = 1.4 \times 10^{-2}$ ) with  $\lambda_S \sim 1.05$ . At the current estimated magnitude of genetic effect, the 1q42 region could not be excluded.

In a combined analysis of animal model and human linkage data from a number of autoimmune diseases, chromosome 18q12-q21 demonstrated evidence of linkage, which has now been supported by the analysis of congenic strains in NOD mice (48). There was no support for loci on chromosome 18 at  $P < 0.05$  in the current study. Three independent studies of type 1 diabetes have reported linkage to chromosome 8q (49–51), but there was no support for 8q in this study. Additional previously reported loci with relatively little support for linkage in the current study include *IDDM4* (11q13), *IDDM6* (18q12-q21), *IDDM9* (3q22-q25), *IDDM11* (14q24-q31), *IDDM16* (14q32), *IDDM17* (10q25), and *IDDM18* (5q33). These data suggest that these putative type 1 diabetes susceptibility loci represent either false-positive results or have very small effects that may be more readily detected in certain populations because of variation in allele frequencies or other factors, including the possibility of population-specific genetic or environmental effects.

Linkage and fine mapping studies in mouse models of SLE (52) and of type 1 diabetes (53) have demonstrated that a single linkage peak may be composed of several susceptibility loci. In human populations, a linkage signal may be observed by the chance clustering of several disease loci, each with relatively weak locus-specific effects. The presence of multiple susceptibility loci may also account, in part, for broad linkage peaks often observed in studies of complex, common diseases. This underlying complexity would also increase the difficulty to obtain convincing results in future fine-mapping association studies. Both development of novel analytical approaches and increased sample size will be necessary to resolve this apparent complexity (54,55). Through the efforts of consortia (such as our international effort), it will be possible to increase the number of families for type 1 diabetes (<http://www.t1dgc.org>), which would increase power and allow exclusion of loci  $\lambda_S \geq 1.2$ , as well as provide standardized samples and reagents for future fine-mapping studies.

These results suggest two parallel tracks for the identification of type 1 diabetes susceptibility loci. First, systematic fine mapping of all the variants responsible for the HLA linkage to type 1 diabetes is justified, especially within the 4-Mb HLA region. Second, further exploration of potential non-HLA regions described here is now justified, including chromosomes 2q31-q33, 6q21, 10p14-q11, and 16q22-q24. Because sample sizes in linkage and association studies have historically been small and few genes (out of  $\sim 25,000$ ) have been studied in depth, future collaborative efforts and establishment of accessible re-

sources for study should increase the yield of true disease susceptibility loci.

#### ACKNOWLEDGMENTS

P.C. has received support from the National Institute of Diabetes and Digestive and Kidney Diseases and Juvenile Diabetes Research Foundation. S.S.R. has received support from the National Institute of Diabetes and Digestive and Kidney Diseases and the JDRF. J.A.T. has received support from the JDRF and the Wellcome Trust.

We acknowledge the assistance of Mark Brown, Mathew Barber, Heather Cordell, Neil Walker, Carl Langefeld and Nancy Cox.

#### APPENDIX

##### Members of the Type 1 Diabetes Genetics Consortium

Asia-Pacific Network: Eesh Bhatia, Francois Bonnici, Pik To Cheung, Peter Colman, Andrew Cotterill, Jenny Couper, Ric Cutfield, Elizabeth Davis, Tim Davis, Paul Dixon, Kim Donaghue, Paul Drury, Mark Harris, Len Harrison, Tim Jones, Uma Kanga, Alok Kanungo, Betty Kek, Jeremy Krebs, Yann-Jinn Lee, Margaret Lloyd, Amanda Loth, Narinder Mehra, Grant Morahan, C.B. Sanjeevi, Brian Tait, Mike Varney, Jinny Willis, Loke Kah Yin, Goh Siok Ying.

European Network: Francisco J. Ampudia-Blasco, Jesus Argente, Magdalena Avbelj, Gulja Babadjanova, Klaus Badenhoop, Lubomir Barak, Christos Bartsocas, Tadej Batelino, Emilia Belda, Polly Bingley, Bernhard O. Boehm, Eulalia Bragues, Raffaella Buzzetti, Joyce Carlson, Luis Castano, Anna Casu, Ondrej Cinek, Raquel Corripio, Alberto de Leiva, V.A. Ruiz Esquide, Ana Fagulha, Marta Hernandez Garcia, Cristian Guja, Alona Hamou, Erifli Hatziagelaki, Simon Heath, Kaire Heilman, Nora Hosszafalusi, Constantin Ionescu-Tirgoviste, Cecile Julier, Ida Kinalska, Ingrid Kockum, Kalinka Koprivarova, Adam Kretowski, Dora Krikovszky, Nebojsa Lalic, Nicole Lambracht, Merce Lara, Mark Lathrop, Katharina Laubner, Claire Levy-Marchal, Johnny Ludvigsson, Laszlo Madacsy, Mara Marga, Didac Mauricio, Jorn Nerup, Antanas Norkus, Anna Okruszko, Xavier Palomer, Teresa Pedro, Moshe Phillip, Valdis Pirags, Flemming Pociot, Galina Popova, Paolo Pozzilli, Bart O. Roep, Silke Rosinger, Ana Maria Varela Sande, Ilhan Satman, Edith Schober, Jochen Seufert, Jan Skrha, Gyula Soltesz, Giatgen Spinass, Juraj Stanik, Tanya Szendeffy, Vallo Tillmann, Dag Undlien, Vaidotas Urbanavicius, Luciana Valente, Bart Van der Auwera, Andriani Vazeou-Gerasimidi, Dzilda Velickiene, Ana Wagner, Markus Walter, Alistair Williams, Miroslav Wurzbürger, Anette-G Ziegler.

North American Network: Alan Aldrich, Mark Anderson, Christophe Benoist, Noureddine Berka, Patrick Concannon, Mark Daly, Jayne Danska, Larry Dolan, David Donaldson, Alessandro Doria, Janice Dorman, George Eisenbarth, Henry Erlich, Pamela Fain, Rosanna Fiallo-Scharer, Kenneth Gabbay, Daniel Geraghty, Soumitra Ghosh, Steven Gitelman, Nat Goodman, Gregory Goodwin, Carla Greenbaum, William Hagopian, John Hansen, Joel Hirschhorn, Leroy Hood, Kevin Kaiserman, Jean Lawrence, Victoria Magnuson, Jennifer Marks, John Mayberry, Elizabeth Mayer-Davis, Richard McIndoe, Brad McNeney, Eric Mickelson, Antoinette Moran, Gerald Nepom, Janelle Noble, Jill Norris, Tihamer Orban, David Owerbach, Andrew Paterson, Catherine Pihoker, Constantin Polychronakos, Al-

berto Pugliese, Philip Raskin, Marian Rewers, Henry Rodriguez, Jerry Rotter, Monique Roy, Desmond Schatz, Gary Schoch, Jin-Xiong She, Richard Spielman, Andrea Steck, Kent Taylor, Jay Tischfield, Ellen Toth, Diane Wherrett, Stephen Willi, Darrell Wilson, Lue Ping Zhao.

U.K. Network: Francesco Cucca, David Dunger, Simon Howell, Sarah Nutland, Helen Rance, Luc Smink, John Todd, Neil Walker, Barry Widmer, Heather Withers.

Coordinating Center: Don Babcock, Stephanie Beck, Mark Brown, Cralen Davis, Mark Espeland, Clara Goro-dezky, Jason Griffin, Mark Hall, Teresa Harnish, Laura Henrick, John Hepler, Joan Hilner, Letitia Howard, Ethan Lange, Carl Langefeld, Josyf Mychaleckyj, June Pierce, David Reboussin, Stephen Rich, Scott Rushing, Michele Sale, Elizabeth Sides, Beverly Snively, Michael Steffes, Augy Thiel, Lynne Wagenknecht, Dustin Williams, Jianzhao Xu, Belinda Youngdahl.

#### REFERENCES

- Rewers M, LaPorte RE, King H, Tuomilehto J: Trends in the prevalence and incidence of diabetes: insulin-dependent diabetes mellitus in childhood. *World Health Stat Q* 41:179–189, 1988
- Rewers MJ: The changing face of the epidemiology of insulin-dependent diabetes mellitus (IDDM): research designs and models of disease causation. *Ann Med* 23:419–426, 1991
- Genuth S, Alberti KG, Bennett P, Buse J, Defronzo R, Kahn R, Kitzmiller J, Knowler WC, Lebovitz H, Lemmark A, Nathan D, Palmer J, Rizza R, Saudek C, Shaw J, Steffes M, Stern M, Tuomilehto J, Zimmet P; Expert Committee on the Diagnosis and Classification of Diabetes Mellitus: Follow-up report on the diagnosis of diabetes mellitus. *Diabetes Care* 31:3160–3167, 2003
- Hirshhorn JN: Genetic epidemiology of type 1 diabetes. *Pediatr Diabetes* 4:87–100, 2003
- Noble JA, Valdes AM, Cook M, Klitz W, Thomson G, Erlich HA: The role of HLA class II genes in insulin-dependent diabetes mellitus: molecular analysis of 180 Caucasian, multiplex families. *Am J Hum Genet* 59:1134–1148, 1996
- Hyttinen V, Kaprio J, Kinnunen L, Koskenvuo M, Tuomilehto J: Genetic liability of type 1 diabetes and the onset age among 22,650 young Finnish twin pairs: a nationwide follow-up study. *Diabetes* 52:1052–1055, 2003
- Svensson J, Carstensen B, Molbak A, Christau B, Mortensen HB, Nerup J, Borch-Johnsen K: Increased risk of childhood type 1 diabetes in children born after 1985. *Diabetes Care* 25:2197–2201, 2002
- Onkamo P, Vaananen S, Karvonen M, Tuomilehto J: Worldwide increase in incidence of type 1 diabetes: the analysis of the data on published incidence trends. *Diabetologia* 42:1395–1403, 1999
- Pociot F, McDermott MF: Genetics of type 1 diabetes mellitus (Review). *Genes Immun* 3:235–249, 2002
- Ronningen KS, Keiding N, Green A, EURODIAB ACE Study Group: Europe and diabetes: correlations between the incidence of childhood-onset type 1 diabetes in Europe and HLA genotypes. *Diabetologia* 44 (Suppl. 3):B51–B59, 2001
- Rich SS: Mapping genes in diabetes: a genetic epidemiological perspective. *Diabetes* 39:1315–1319, 1990
- Valdes AM, Noble JA, Genin E, Clerget-Darpoux F, Erlich HA, Thomson G: Modeling of HLA class II susceptibility to type 1 diabetes reveals an effect associated with DPB1. *Genet Epidemiol* 21:212–223, 2001
- Bell GI, Horita S, Karam JH: A polymorphic locus near the human insulin gene is associated with insulin-dependent diabetes mellitus. *Diabetes* 33:176–183, 1984
- Julier C, Hyer RN, Davies J, Merlin F, Soularue P, Briant L, Cathelineau G, Deschamps I, Rotter JJ, Froguel P: Insulin-IGF2 region on chromosome 11p encodes a gene implicated in HLA-DR4-dependent diabetes susceptibility. *Nature* 354:155–159, 1991
- Julier C, Lucassen A, Villedieu P, Delepine M, Levy-Marchal C, Danze PM, Bianchi F, Boitard C, Froguel P, Bell J: Multiple DNA variant association analysis: application to the insulin gene region in type I diabetes. *Am J Hum Genet* 55:1247–1254, 1994
- Barratt BJ, Payne F, Lowe CE, Hermann R, Healy BC, Harold D, Concannon P, Gharani N, McCarthy MI, Olavesen MG, McCormack R, Guja C, Ionescu-Tirgoviste C, Undlien DE, Ronningen KS, Gillespie KM, Tuomilehto-Wolf E, Tuomilehto J, Bennett ST, Clayton DG, Cordell HJ, Todd JA: Remapping the insulin gene/IDDM2 locus in type 1 diabetes. *Diabetes* 53:1884–1889, 2004

17. Ueda H, Howson JM, Esposito L, Heward J, Snook H, Chamberlain G, Rainbow DB, Hunter KM, Smith AN, Di Genova G, Herr MH, Dahlman I, Payne F, Smyth D, Lowe C, Twells RC, Howlett S, Healy B, Nutland S, Rance HE, Everett V, Smink LJ, Lam AC, Cordell HJ, Walker NM, Bordin C, Hulme J, Motzo C, Cucca F, Hess JF, Metzker ML, Rogers J, Gregory S, Allahabadi A, Nithiyanthan R, Tuomilehto-Wolf E, Tuomilehto J, Bingley P, Gillespie KM, Undlien DE, Ronningen KS, Guja C, Ionescu-Tirgoviste C, Savage DA, Maxwell AP, Carson DJ, Patterson CC, Franklyn JA, Clayton DG, Peterson LB, Wicker LS, Todd JA, Gough SC: Association of the T-cell regulatory gene CTLA4 with susceptibility to autoimmune disease. *Nature* 423:506–511, 2003
18. Bottini N, Musumeci L, Alonso A, Rahmouni S, Nika K, Rostamkhani M, MacMurray J, Meloni GF, Lucarelli P, Pellicchia M, Eisenbarth GS, Comings D, Mustelin T: A functional variant of lymphoid tyrosine phosphatase is associated with type I diabetes. *Nat Genet* 36:337–338, 2004
19. Smyth DJ, Cooper JD, Collins JE, Heward JM, Franklyn JA, Howson JMM, Walker N, Vella A, Nutland S, Rance HE, Maier LM, Barratt BJ, Guja C, Ionescu-Tirgoviste CA, Savage D, Dunger DB, Widmer B, Strachan DP, Ring SM, Clayton DG, Twells RCJ, Gough SCL, Todd JA: Replication of an association between the lymphoid tyrosine phosphatase locus LYP/PTPN22 with type 1 diabetes, and evidence for its role as a general autoimmunity locus. *Diabetes* 53:3020–3023, 2004
20. Onengut-Gumuscu S, Ewens KG, Spielman RS, Concannon P: A functional polymorphism (1858C/T) in the PTPN22 gene is linked and associated with type 1 diabetes in multiplex families. *Genes Immun* 5:678–680, 2004
21. Davies JL, Kawaguchi Y, Bennett ST, Copeman JB, Cordell HJ, Pritchard LE, Reed PW, Gough SCL, Jenkins SC, Palmer SM, Balfour KM, Row B, Farrall M, Barnett AH, Bain SC, Todd JA: A genome-wide search for human type 1 diabetes susceptibility genes. *Nature* 371:130–136, 1994
22. Hashimoto L, Habita C, Beressi J, Delepine M, Besse C, Cambon-Thomsen A, Deschamps I, Rotter J, Djoulah S, James M, Froguel P, Weissenbach J, Lathrop GM, Julier C: Genetic mapping of a susceptibility locus for insulin-dependent diabetes mellitus on chromosome 11q. *Nature* 371:161–164, 1994
23. Melanitou E, Fain P, Eisenbarth GS: Genetics of type 1A (immune mediated) diabetes. *J Autoimmun* 21:93–98, 2003
24. Cox NJ, Wapelhorst B, Morrison VA, Johnson L, Pinchuk L, Spielman RS, Todd JA, Concannon P: Seven regions of the genome show evidence of linkage to type 1 diabetes in a consensus analysis of 767 multiplex families. *Am J Hum Genet* 69:820–830, 2001
25. Mein CA, Esposito L, Dunn MG, Johnson GC, Timms AE, Goy JV, Smith AN, Sebag-Montefiore L, Merriman ME, Wilson AJ, Pritchard LE, Cucca F, Barnett AH, Bain SC, Todd JA: A search for type 1 diabetes susceptibility genes in families from the United Kingdom. *Nat Genet* 19:297–300, 1998
26. Concannon P, Gogolin-Ewens KJ, Hinds D, Wapelhorst B, Morrison VA, Stirling B, Mitra M, Farmer J, Williams SR, Cox NJ, Bell GI, Risch N, Spielman RS: A second-generation screen of the human genome for susceptibility to insulin-dependent diabetes mellitus (IDDM). *Nat Genet* 19:292–296, 1998
27. European Consortium for IDDM Genome Studies: A genomewide scan for type 1-diabetes susceptibility in Scandinavian families: identification of new loci with evidence of interactions. *Am J Hum Genet* 69:1301–1313, 2001
28. Bain SC, Todd JA, Barnett AH: The British Diabetic Association: Warren repository. *Autoimmunity* 7:83–85, 1990
29. Morahan G, Huang D, Tait B, Colman PG, Harrison LC: Markers on distal chromosome 2q linked to insulin-dependent diabetes mellitus. *Science* 272:1811–1813, 1996
30. McPeck MS, Sun L: Statistical tests for detection of misspecified relationships by use of genome screen data. *Am J Hum Genet* 66:1076–1094, 2000
31. Kruglyak L, Daly MJ, Reeve-Daly MP, Lander ES: Parametric and non-parametric linkage analysis: a unified multipoint approach. *Am J Hum Genet* 58:1347–1363, 1996
32. Kong A, Cox NJ: Allele-sharing models: LOD scores and accurate linkage tests. *Am J Hum Genet* 61:1179–1188, 1997
33. Langefeld CD, Boehnke M: Multiple trait nonparametric linkage regression. *Am J Hum Genet* 65:A258, 1999
34. Abecasis GR, Cherny SS, Cookson WOC, Cardon LR: MERLIN: rapid analysis of dense genetic maps using sparse gene flow trees. *Nat Genet* 30:97–101, 2002
35. Gudbjartsson DF, Jonasson K, Frigge M, Kong A: Allegro, a new computer program for multipoint linkage analysis. *Nat Genet* 25:12–13, 2000
36. Hauser ER, Boehnke M: Genetic linkage analysis of complex genetic traits by using affected sibling pairs. *Biometrics* 54:1238–1246, 1998
37. Kruglyak L, Lander ES: Complete multipoint sib-pair analysis of qualitative and quantitative traits. *Am J Hum Genet* 57:439–454, 1995
38. Delepine M, Pociot F, Habita C, Hashimoto L, Froguel P, Rotter J, Combon-Thomsen A, Deschamps I, Djoulah S, Weissenbach J, Nerup J, Lathrop M, Julier C: Evidence of a non-MHC susceptibility locus in type 1 diabetes linked to HLA on chromosome 6. *Am J Hum Genet* 60:174–187, 1997
39. Cucca F, Goy JV, Kawaguchi Y, Esposito L, Merriman ME, Wilson AJ, Cordell HJ, Bain SC, Todd JA: A male-female bias in type 1 diabetes and linkage to chromosome Xp in MHC HLA-DR3-positive patients. *Nat Genet* 3:301–302, 1998
40. Wiltshire S, Cardon LR, McCarthy MI: Evaluating the results of genome-wide linkage scans of complex traits by locus counting. *Am J Hum Genet* 71:1175–1182, 2002
41. Wapelhorst B, Bell GI, Risch N, Spielman RS, Concannon P: Linkage and association studies in insulin-dependent diabetes with a new dinucleotide repeat polymorphism at the GAD<sub>65</sub> locus. *Autoimmunity* 21:127–130, 1995
42. Johnson GC, Payne F, Nutland S, Stevens H, Tuomilehto-Wolf E, Tuomilehto J, Todd JA: A comprehensive, statistically powered analysis of GAD2 in type 1 diabetes. *Diabetes* 51:2866–2870, 2002
43. Fisher SA, Lanchbury JS, Lewis CM: Meta-analysis of four rheumatoid arthritis genome-wide linkage studies: confirmation of a susceptibility locus on chromosome 16. *Arthritis Rheum* 48:1200–1206, 2003
44. Tait KF, Marshall T, Berman J, Carr-Smith J, Rowe B, Todd JA, Bain SC, Barnett AH, Gough SC: Clustering of autoimmune disease in parents of siblings from the type 1 diabetes Warren repository. *Diabet Med* 21:358–362, 2004
45. MacKay K, Eyre S, Myerscough A, Millicic A, Barton A, Laval S, Barrett J, Lee D, While S, Jhon S, Brown MA, Bell J, Silman A, Ollier W, Wordsworth P, Worthington J: Whole-genome linkage analysis of rheumatoid arthritis susceptibility loci in 252 affected sibling pairs in the United Kingdom. *Arthritis Rheum* 46:632–639, 2002
46. Jawaheer D, Seldin MF, Amos CI, Chen WV, Shigeta R, Etzel C, Damle A, Xiao X, Chen D, Lum RF, Monteiro J, Kern M, Criswell LA, Albani S, Nelson JL, Clegg DO, Pope R, Schroeder HW Jr, Bridges SL Jr, Pisetsky DS, Ward R, Kastner DL, Wilder RL, Pincus T, Callahan LF, Flemming D, Wener MH, Gregersen PK, North American Rheumatoid Arthritis Consortium: Screening the genome for rheumatoid arthritis susceptibility genes: a replication study and combined analysis of 512 multicase families. *Arthritis Rheum* 48:906–916, 2003
47. Begovich AB, Carlton VE, Honigberg LA, Schrodi SJ, Chokkalingam AP, Alexander HC, Ardlie KG, Huang Q, Smith AM, Spoerke JM, Conn MT, Chang M, Chang SY, Saiki RK, Catanese JJ, Leong DU, Garcia VE, McAllister LB, Jeffery DA, Lee AT, Batliwalla F, Remmers E, Criswell LA, Seldin MF, Kastner DL, Amos CI, Sninsky JJ, Gregersen PK: A missense single-nucleotide polymorphism in a gene encoding a protein pyrosine phosphatase (PTPN22) is associated with rheumatoid arthritis. *Am J Hum Genet* 75:330–337, 2004
48. Ewens KG, Johnson LN, Wapelhorst B, O'Brien K, Gutin S, Morrison VA, Street C, Gregory SG, Spielman RS, Concannon P: Linkage and association with type 1 diabetes on chromosome 1q42. *Diabetes* 51:3318–3325, 2002
49. Merriman T, Twells R, Merriman M, Eaves I, Cox R, Cucca F, McKinney P, Shield J, Baum D, Bosi E, Pozzilli P, Nistico L, Buzzetti R, Joner G, Ronningen K, Thorsby E, Undlien D, Pociot F, Nerup J, Bain S, Barnett A, Todd J: Evidence by allelic association-dependent methods for a type 1 diabetes polygene (IDDM6) on chromosome 18q21. *Hum Mol Genet* 6:1003–1010, 1997
50. Luo DF, Buzzetti R, Rotter JI, Maclaren NK, Raffel LJ, Nistico L, Govannini C, Pozzilli P, Thomson G, She JX: Confirmation of three susceptibility genes to insulin-dependent diabetes mellitus: IDDM4, IDDM5 and IDDM8. *Hum Mol Genet* 5:693–698, 1996
51. Sale MM, FitzGerald LM, Charlesworth JC, Bowden DW, Rich SS: Evidence for a novel type 1 diabetes susceptibility locus on chromosome 8. *Diabetes* 51 (Suppl. 3):S316–S319, 2002
52. Vaessen N, Heutink P, Houwing-Duistermaat JJ, Snijders PJ, Rademaker T, Testers L, Batstra MR, Sandkuijl LA, van Duijn CM, Oostra BA: A genome-wide search for linkage disequilibrium with type 1 diabetes in a recent genetically isolated population from the Netherlands. *Diabetes* 51:856–859, 2002
53. Nguyen C, Limaye N, Wakeland EK: Susceptibility genes in the pathogenesis of murine lupus. *Arthritis Res* 4 (Suppl. 3):S255–S263, 2002
54. Todd JA, Wicker LS: Genetic protection from the inflammatory disease type 1 diabetes in humans and animal models. *Immunity* 15:387–395, 2001
55. Rich SS, Concannon P: Challenges and strategies for investigating the genetic complexity of common human diseases. *Diabetes* 51 (Suppl. 3):S288–S294, 2002
56. Pociot F, Karlens AE, Pedersen CB, Aalund M, Nerup J for the European Consortium for IDDM Genome Studies: Novel analytical methods applied to type 1 diabetes genome-scan data. *Am J Hum Genet* 74:647–660, 2004