

Common Missense Variant in the Glucokinase Regulatory Protein Gene Is Associated With Increased Plasma Triglyceride and C-Reactive Protein but Lower Fasting Glucose Concentrations

Marju Orho-Melander,¹ Olle Melander,¹ Candace Guiducci,² Pablo Perez-Martinez,^{3,4,5} Dolores Corella,⁵ Charlotta Roos,¹ Ryan Tewhey,² Mark J. Rieder,⁶ Jennifer Hall,⁷ Goncalo Abecasis,⁸ E. Shyong Tai,⁹ Cullan Welch,⁷ Donna K. Arnett,¹⁰ Valeriya Lyssenko,¹ Eero Lindholm,¹ Richa Saxena,² Paul I.W. de Bakker,² Noel Burt,² Benjamin F. Voight,² Joel N. Hirschhorn,² Katherine L. Tucker,¹¹ Thomas Hedner,¹² Tiinamaija Tuomi,^{13,14} Bo Isomaa,¹⁴ Karl-Fredrik Eriksson,¹ Marja-Riitta Taskinen,¹³ Björn Wahlstrand,¹² Thomas E. Hughes,¹⁵ Laurence D. Parnell,⁴ Chao-Qiang Lai,⁴ Göran Berglund,¹⁶ Leena Peltonen,¹⁷ Erkki Vartiainen,¹⁸ Pekka Jousilahti,¹⁸ Aki S. Havulinna,¹⁸ Veikko Salomaa,¹⁸ Peter Nilsson,¹ Leif Groop,^{1,13} David Altshuler,^{2,19,20} Jose M. Ordovas,⁴ and Sekar Kathiresan^{2,21}

OBJECTIVE—Using the genome-wide association approach, we recently identified the glucokinase regulatory protein gene (*GCKR*, rs780094) region as a novel quantitative trait locus for plasma triglyceride concentration in Europeans. Here, we sought to study the association of *GCKR* variants with metabolic phe-

notypes, including measures of glucose homeostasis, to evaluate the *GCKR* locus in samples of non-European ancestry and to fine-map across the associated genomic interval.

RESEARCH DESIGN AND METHODS—We performed association studies in 12 independent cohorts comprising >45,000 individuals representing several ancestral groups (whites from Northern and Southern Europe, whites from the U.S., African Americans from the U.S., Hispanics of Caribbean origin, and Chinese, Malays, and Asian Indians from Singapore). We conducted genetic fine-mapping across the ~417-kb region of linkage disequilibrium spanning *GCKR* and 16 other genes on chromosome 2p23 by imputing untyped HapMap single nucleotide polymorphisms (SNPs) and genotyping 104 SNPs across the associated genomic interval.

RESULTS—We provide comprehensive evidence that *GCKR* rs780094 is associated with opposite effects on fasting plasma triglyceride ($P_{\text{meta}} = 3 \times 10^{-56}$) and glucose ($P_{\text{meta}} = 1 \times 10^{-13}$) concentrations. In addition, we confirmed recent reports that the same SNP is associated with C-reactive protein (CRP) level ($P = 5 \times 10^{-5}$). Both fine-mapping approaches revealed a common missense *GCKR* variant (rs1260326, Pro446Leu, 34% frequency, $r^2 = 0.93$ with rs780094) as the strongest association signal in the region.

CONCLUSIONS—These findings point to a molecular mechanism in humans by which higher triglycerides and CRP can be coupled with lower plasma glucose concentrations and position *GCKR* in central pathways regulating both hepatic triglyceride and glucose metabolism. *Diabetes* 57:3112–3121, 2008

From the ¹Department of Clinical Sciences, University Hospital Malmö, Clinical Research Center, Lund University, Malmö, Sweden; the ²Program in Medical and Population Genetics, Broad Institute of the Massachusetts Institute of Technology and Harvard University, Cambridge, Massachusetts; the ³Lipids and Atherosclerosis Research Unit, Reina Sofia University Hospital, University of Cordoba, Cordoba, Spain; the ⁴Nutrition and Genomics Laboratory, Jean Mayer-U.S. Department of Agriculture Human Nutrition Research Center on Aging at Tufts University, Boston, Massachusetts; the ⁵Genetic and Molecular Epidemiology Unit and CIBER Fisiopatología de la Obesidad y Nutrición, School of Medicine University of Valencia, Valencia, Spain; the ⁶Department of Genome Sciences, University of Washington, Seattle, Washington; the ⁷Lillehei Heart Institute, Department of Medicine, University of Minnesota, Minneapolis, Minnesota; the ⁸Center for Statistical Genetics, Department of Biostatistics, University of Michigan School of Public Health, Ann Arbor, Michigan; the ⁹Department of Endocrinology, Singapore General Hospital, Singapore; the ¹⁰Dietary Assessment and Epidemiology Research Program, Jean Mayer-U.S. Department of Agriculture Human Nutrition Research Center on Aging, Tufts University, Boston, Massachusetts; the ¹¹Department of Epidemiology, University of Alabama, Birmingham, Alabama; the ¹²Department of Clinical Pharmacology, Sahlgrenska Academy, Göteborg, Sweden; the ¹³Department of Medicine, Helsinki University Hospital, University of Helsinki, Helsinki, Finland; the ¹⁴Folkhälsan Research Center, Helsinki, Finland; the ¹⁵Novartis Institutes for BioMedical Research, Cambridge, Massachusetts; the ¹⁶Department of Clinical Sciences, Medicine, Lund University, Malmö, Sweden; the ¹⁷Department of Molecular Medicine, National Public Health Institute, Biomedicum, Helsinki, Finland; the ¹⁸Department of Epidemiology and Health Promotion, National Public Health Institute, Helsinki, Finland; the ¹⁹Center for Human Genetic Research and Department of Molecular Biology, Massachusetts General Hospital, Boston, Massachusetts; the ²⁰Department of Genetics, Harvard Medical School, Boston, Massachusetts; and the ²¹Cardiovascular Disease Prevention Center, Cardiology Division, Department of Medicine, Massachusetts General Hospital, Harvard Medical School, Boston, Massachusetts.

Corresponding author: Marju Orho-Melander, marju.ortho-melander@med.lu.se.

Received 17 April 2008 and accepted 29 July 2008.

Published ahead of print at <http://diabetes.diabetesjournals.org> on 4 August 2008. DOI: 10.2337/db08-0516.

© 2008 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Recently, in the genome-wide association Diabetes Genetics Initiative (DGI) Study for 19 traits, including plasma lipids, we provided evidence that the glucokinase (*GCK*) regulatory protein gene (*GCKR*) region was a novel quantitative trait locus associated with plasma triglyceride concentration (1). Of all single nucleotide polymorphisms (SNPs) tested, an intronic SNP at *GCKR* (rs780094) explained the greatest proportion of interindividual variability in plasma triglycerides (1).

GCKR regulates *GCK*, which functions as a glucose sensor responsible for glucose phosphorylation in the first step of glycolysis. The discoveries that inactivating muta-

tions in GCK cause maturity onset diabetes of the young type 2 (2) and activating *GCK* mutations lead to permanent hyperinsulinemic hypoglycemia (3) emphasize that GCK plays a major role in glucose metabolism. *GCKR*-deficient mice have reduced GCK expression but maintain nearly normal GCK activity and show impaired glucose clearance (4). Furthermore, adenoviral-mediated overexpression of *GCKR* in mouse liver increased GCK activity and lowered fasting blood glucose (5) and overexpression of *GCK* in liver led to lowered blood glucose and increased triglyceride concentrations (6,7). Thus, experimental evidence suggests that perturbation of the GCKR pathway has opposing effects of triglyceride and glucose metabolism.

In our original report, SNP rs780094 in *GCKR* was associated with fasting triglyceride levels in two independent samples, each of Northern European ancestry ($P = 3.7 \times 10^{-8}$ and 8.7×10^{-8} , respectively) (1). After initial identification and replication of a chromosomal region associated with a trait, key next steps include extension of the association finding to related phenotypes, validation of the association finding in different ethnicities, and fine-mapping to identify the putative causal variant. Recently, our initial finding was replicated in a Danish study in which a strong association was found between the rs780094 T allele and elevated fasting triglyceride levels but also lower insulin levels, better insulin sensitivity, and a moderately decreased risk of type 2 diabetes (8). In addition, recent genome-wide association studies identified an association between the same *GCKR* intronic SNP and C-reactive protein (CRP) levels (9,10).

Hereby, we sought to examine the effect of SNP rs780094 on triglycerides and related metabolic traits, including fasting glucose concentrations, in 12 samples representing a range of ancestral groups and including a large prospective study with a mean follow-up time of 23 years. In addition, we performed fine-mapping in one of these samples to identify the strongest association signal in the region.

RESEARCH DESIGN AND METHODS

The genetic association studies were performed in 12 study samples as shown in Table 1. For all studies, informed consent was obtained from the study subjects, and the study protocols were approved by local ethics committees. All study cohorts genotyped for GCKR as part of this experiment are included in this report.

The DGI Study material consisting of 2,931 individuals from Finland and Sweden (1,464 patients with type 2 diabetes and 1,467 nondiabetic control subjects) was ascertained as previously described (1,11,12). DGI samples were genotyped on the Affymetrix 500K chip (1) and were used in the present study for the in silico fine-mapping. In addition, the DGI samples were used for the genotype fine-mapping of the *GCKR* locus and for the analyses of apolipoprotein B (apoB) and free fatty acids (FFAs) according to rs780094.

The Malmö Diet and Cancer Study–Cardiovascular Cohort (MDC-CC) represents 6,103 people that were randomly selected to participate in a study of the epidemiology of carotid artery disease from a large, community-based prospective epidemiological cohort of 28,449 people recruited for a baseline examination between 1991 and 1996 (13). Of the MDC-CC participants, 597 did not provide a baseline plasma sample or did not have triglyceride data available, leaving 5,506 individuals available for the analyses of lipid traits. Of these, 5,023 nondiabetic individuals had data on fasting blood glucose, and 4,867 had data on fasting insulin. In MDC-CC, cardiovascular events were ascertained through linkage of the 10-digit personal identification number of each Swedish citizen with three registries: the Swedish Hospital Discharge Register, the Swedish Cause of Death Register, and the Stroke Register of Malmö. The prespecified composite end point of cardiovascular events was defined as myocardial infarction (according to codes 410 and 121 in ICD-9 and ICD-10), ischemic stroke (according to codes 434 and 436 in ICD-9 and I63 or I64 in ICD-10), and death from coronary heart disease (codes 412 and 414 in ICD-9 or I22–I23 and I25 in ICD-10 in the Swedish Cause of Death Register). We excluded cases of intracerebral or subarachnoid hemorrhage. The status of

cigarette smoking was elicited by a self-administered questionnaire and was coded as never, former, or current. A parental or sibling history of myocardial infarction was also elicited by a questionnaire. CRP was measured by a high-sensitivity assay (Tina-quant CRP; Roche Diagnostics). Study participants underwent a B-mode ultrasonography of the right carotid artery. Intima-media thickness (IMT) of the common carotid artery was measured according to standardized protocol by a trained certified sonographer as previously described (14,15).

In the Malmö Preventive Project (MPP), 33,346 citizens from Malmö in southern Sweden participated in a health screening during 1974–1992 (16). Of individuals participating in the initial screening, 4,931 had died, and 551 were lost from follow-up. Of the eligible individuals, 25,000 were invited to a rescreening visit during 2002–2006, which included a physical examination and fasting blood samples for measurements of plasma glucose and lipids. Of the invited subjects, 17,284 people participated in the rescreening, and of these, 17,037 individuals had DNA, 16,998 individuals had fasting triglyceride data, and 16,976 individuals had fasting glucose data available for the study. The mean follow-up time was 23.4 ± 4.6 years. Of the 17,037 individuals, 5,946 were also participants of the MDC-CC, bringing the number of unique individuals in the MPP (i.e., not included in MDC-CC) to 11,091. Because of the overlap between MDC-CC and MPP materials, all of the phenotype association analyses in MPP included in the meta-analysis were performed in the unique cohort of 11,091 participants, whereas the prospective analyses were performed in the whole MPP cohort with DNA and available phenotypes. Individuals on lipid-lowering medication ($n = 3,510$) were excluded from the triglyceride analyses, and individuals with diabetes either at baseline or follow-up ($n = 3,029$) were excluded from the analyses of fasting blood glucose. For the analysis of progression to type 2 diabetes in MPP, 16,061 nondiabetic subjects, 2,063 of whom developed type 2 diabetes, were included. Diagnosis of diabetes was confirmed from patient records or based on a fasting plasma glucose concentration >7.0 mmol/L.

During the follow-up visit, a subgroup from the MPP study participated in more extensive metabolic studies, including a euglycemic-hyperinsulinemic clamp combined with indirect calorimetry to measure hepatic glucose output. All individuals underwent a physical examination, and a subgroup of 199 men (Malmö men, MPP-MM) with impaired glucose tolerance (IGT) underwent more extensive metabolic studies, including a new oral glucose tolerance test (OGTT) and a euglycemic-hyperinsulinemic clamp combined with infusion of [$^3\text{-H}$]glucose to measure hepatic glucose output (17). After the follow-up time, 66 of the 199 men with IGT at baseline had normal glucose tolerance, 52 had impaired fasting glucose and/or IGT, and 81 had type 2 diabetes. Type 2 diabetic patients were treated either with diet alone (42%) or with oral hypoglycemic agents, which were withheld the day before the clamp.

The Nordic Diltiazem (NORDIL) Study is an intervention study in 10,881 patients with hypertension from Sweden and Norway (diastolic blood pressure ≥ 100 mmHg at least twice, mean age 60 ± 7 years) of whom 5,152 Swedish patients provided DNA and were included in the present study (18). The study participants had been randomized to antihypertensive treatment with either the calcium-antagonist diltiazem or β -blocker/thiazide diuretic to compare efficacy of the two drug therapies to prevent cardiovascular end points (18). Lipoprotein and lipid measurements at the baseline examination (before the initiation of antihypertensive therapy) were studied in this report.

The Diabetes Registry sample consists of 2,777 type 2 diabetic patients from the Skania Diabetes 2000 Registry, a local diabetes registry in southern Sweden (19). The majority of registered patients came from the city of Malmö in southern Sweden. Type 2 diabetes was classified according to 1997 World Health Organization (WHO) criteria (20).

The Botnia-Prevalence, Prediction, and Prevention of Diabetes (Botnia-PPP) Study is a population-based study from the Botnia region of western Finland. The current study was initiated in 2004 in a population comprising $\sim 135,000$ individuals. Using a population registry, a random sample of subjects aged 18–75 years was selected: In age-groups 18–29 and 60–74 years, 1 of 10 individuals was randomly selected; and in age-group 30–59 years, 1 of 15 individuals was randomly selected. Altogether, 6,075 individuals were invited to participate in the study, and 3,621 took part. Of these individuals, 3,495 have data and DNA available for the current study.

The FINRISK97 study is a population-based cross-sectional survey which consists of 8,191 people aged 25–74 years from five geographical areas of Finland (the Helsinki, southwestern Finland, North Karelia, Oulu, and Kuopio regions) (21). The study followed the protocol of the WHO Multinational Monitoring of Trends and Determinants in Cardiovascular Disease (MONICA) Project, an international study conducted under the auspices of the WHO.

The Valencia study population comprised 1,608 individuals randomly selected from the Valencia region on the eastern Mediterranean coast of Spain and examined between 1999 and 2003 (22). This sample comprised randomly

TABLE 1
Clinical characteristics of the study cohorts

	Finland and Sweden		Sweden			Finland	
	DGI (ref. 1)	MDC-CC (ref. 13)	MPP (ref. 16)	NORDIL (ref. 18)	Skania Diabetes 2000 Registry (ref. 19)	Botnia-PPP	FINRISK97
<i>n</i>	2,930	5,506	17,037	5,152	2,777	3,495	8,191
<i>n</i> (men/women)	1,448/1,482	2,282/3,224	10,927/6,110	2,567/2,585	1,583/1,075	1,663/1,841	4,082/4,109
Age (years)	62 ± 11	58 ± 6	46 ± 7	60 ± 7	63 ± 11	49 ± 16	48 ± 13
BMI (kg/m ²)	28 ± 4	26 ± 4	24 ± 3	28 ± 4	30 ± 5	26 ± 4	27 ± 5
Triglycerides (mg/dl)	144 ± 101	122 ± 71	117 ± 71	159 ± 108	231 ± 275	116 ± 71	133 ± 92
HDL (mg/dl)	49 ± 15	53 ± 14	—	53 ± 21	44 ± 13	—	54 ± 14
LDL (mg/dl)	145 ± 51	161 ± 38	—	160 ± 43	137 ± 40	—	134 ± 36
fb-glucose (mg/dl)	121 ± 49	93 ± 25	88 ± 13	96 ± 28	189 ± 69	84 ± 15	—
HOMA (mmol × mU)	1.7 ± 1.4	2.0 ± 2.7	—	—	—	—	—
Type 2 diabetes (%)	49.9	8.4	0.0	8.7	100.0	3.8	5.4

Continued on facing page

selected workers, using a continuously updated computerized population register, and subjects randomly selected from the general population.

The Dallas Heart Study is a multiethnic, probability-based sample of Dallas County, weighted such that 50% of the study population was black (23). Ethnicity was self-reported and consisted of non-Hispanic blacks, non-Hispanic whites, and Hispanics. The study population included 3,469 individuals from one of these three ethnic groups with fasting venous blood samples.

The Genetics of Lipid-Lowering Drugs and Diet Network (GOLDN) Study sample consists of 1,062 individuals and is part of the Program for Genetic Interactions Network (24). The majority of participants in the GOLDN Study were re-recruited from three-generational pedigrees from two National Heart, Lung, and Blood Institute (NHLBI) Family Heart Study (FHS) field centers (Minneapolis, MN, and Salt Lake City, UT) (24). The NHLBI FHS is a multicenter, population-based study of genetic and environmental determinants of cardiovascular disease (CVD) and associated risk factors. Nearly all subjects were of European ancestry.

The longitudinal Boston Puerto Rican Health Study includes 837 Puerto Rican (Hispanics of Caribbean origin) men and women aged 45–75 years in the greater Boston area (25,26). As one of eight nationally funded National Institutes of Health (NIH) Centers on Population Health and Health Disparities, the study is investigating health disparities in the Puerto Rican population. Participants were recruited from the Boston area through door-to-door enumeration, following a sampling scheme based on identification of the 2000 U.S. Census blocks containing Hispanics, and in partnership with community organizations.

The Singaporean National Health Survey 98 (Singaporean NHS-98) study was an initiative to determine the risk factors for the major noncommunicable diseases in Singapore (27,28). A total of 3,973 subjects who participated in the Singaporean NHS-98 and had the data needed for the current study were included in this study. In brief, 11,200 individuals from addresses representing the house-type (a proxy for socioeconomic status) distribution of the entire Singapore housing population were selected from the National Database on Dwellings. From these individuals, a random sample was selected by disproportionate stratified and systematic sampling. The Malays and Indians were oversampled to ensure that prevalence estimates for these minority groups were reliable.

Lipid and glucose phenotypes. Plasma total cholesterol, HDL cholesterol, and triglycerides in each study were measured using standard enzymatic methods from fasting blood samples, with the exception of FINRISK97, in which the lipid measurements were performed from blood samples collected in a “semifasting” state; i.e., the participants were instructed to fast for 4 h and to avoid fatty meals earlier during the day. ApoB in the DGI was measured using an immunochemical assay (Orion Diagnostica, Espoo, Finland), and FFA was measured by an enzymatic colorimetric ACS-ACOD-MEHA method (Wako Chemicals, Neus, Germany).

Fasting blood or plasma glucose was measured by glucose oxidase methods as previously described (1,16–18, 22–27), and blood glucose was converted to plasma glucose using a correction factor of 1.13. Fasting serum insulin was measured using radioimmunoassay in DGI, MDC-CC, MPP-MM, GOLDN, and Singaporean NHS-98 samples (1,11–13,21,24,25). Homeostasis

model assessment (HOMA) insulin resistance index was calculated using the following formula: fasting plasma glucose × fasting insulin/22.5. The insulinogenic index was calculated as [(insulin 30 – fasting insulin)/(glucose 30 – fasting glucose)].

Genotyping. Genotyping was performed either by matrix-assisted laser desorption ionization-time of flight mass spectrometry on the Sequenom MassARRAY platform (San Diego, CA) or by allelic discrimination method on the ABI 7900 instrument (Applied Biosystems, Foster City, CA). The studied SNPs were in Hardy-Weinberg equilibrium in all studied populations ($P > 0.01$) except rs780094 in the Singaporean Chinese population ($P = 0.000063$). Because of deviation from Hardy-Weinberg equilibrium, a random sample of 12% of the Chinese samples was reanalyzed in a separate assay, and the genotyping error rate was 0.3%.

Genotype fine-mapping. *GCKR* rs780094 lies in a large region of linkage disequilibrium on chromosome 2. We defined the associated interval to be ~417 kb based on linkage disequilibrium between the index SNP (rs780094) and SNPs upstream and downstream of the index SNP. rs1049817 was furthest upstream with an $r^2 \geq 0.25$ with the index SNP, and rs13023194 was the furthest downstream with an $r^2 \geq 0.25$ with the index SNP. The interval between rs1049817 and rs13023194 spans 416,543 bases (National Center for Biotechnology Information human genome sequence Build 35). In this interval, there are 17 annotated genes: *MPV17*, *GTF3C2*, *EIF2B4*, *SNX17*, *ZNF513*, *PPM1G*, *NRBP1*, *KRTCAP3*, *IFT172*, *FNDC4*, *GCKR*, *C2orf16*, *ZNF512*, *CCDC121*, *XABI*, *SUPT7L*, and *SLC4A1AP*. To fine-map the association signal across this interval, we selected 120 SNPs for genotyping based on the following criteria: 1) tag SNPs ($n = 33$) that captured all common SNPs (minor allele frequency > 0.05) across the ~417 kb at an $r^2 > 0.8$; 2) all coding SNPs ($n = 83$) present in HapMap CEU for these genes; and 3) a set of SNPs predicted to be microRNA binding sites ($n = 4$). Of these 120 SNPs, 104 were successfully designed for genotyping assays and were genotyped in the DGI sample.

In silico fine-mapping. We also conducted fine-mapping using a second approach: imputation of untyped SNPs. We imputed untyped SNPs across the region using a recently developed Markov Chain Haplotyping algorithm (MACH 1.0) (29). This method predicts genotypes for untyped SNPs in a given study using two inputs: genotypes at typed SNPs in the study sample and the entire set of genotypes in HapMap (www.hapmap.org) for a given reference sample. Here, the inputs were the following: genotypes from the Affymetrix 500K array in the DGI study and ~2.2 million SNPs in the HapMap CEU samples (a reference sample of European ancestry).

Expression studies. To evaluate whether the transcript level of *GCKR* or *GCK* varied by genotype, we studied 101 human liver samples with both measured transcript levels and genotypes. Transcript levels for 60 samples were assessed by the Human Ref8 v.2 Illumina chip, and genotypes were measured by the Illumina 500K array (30). For 41 additional samples, *GCKR* and *GCK* transcript levels were measured in human liver tissue obtained from the University of Minnesota Tissue Procurement Center (Minneapolis, MN) following institutional review board guidelines. DNA was extracted using a Qiagen extraction protocol according to the manufacturer's directions. Samples were genotyped by primer extension with detection by matrix-assisted

TABLE 1
Continued

Spain		U.S.				Singapore		
		Dallas Heart Study				NHS-98 (refs. 27, 28)		
Valencia	Blacks	Hispanics	Whites	GOLDN (ref. 24)	BPRHS (refs. 25,26)	Chinese	Malays	Asian Indians
1,608	1,825	601	1,043	1,062	837	2,691	734	548
760/848	770/1,055	251/350	500/543	506/556	225/612	1,227/1,464	353/381	262/286
42 ± 14	45 ± 10	40 ± 9	45 ± 10	49 ± 16	58 ± 7	38 ± 12	39 ± 13	41 ± 12
26 ± 5	32 ± 8	31 ± 7	29 ± 7	28 ± 6	32 ± 8	23 ± 4	26 ± 5	25 ± 5
112 ± 66	107 ± 95	151 ± 130	139 ± 107	135 ± 85	159 ± 99	120 ± 72	143 ± 87	145 ± 83
51 ± 11	52 ± 15	46 ± 11	48 ± 15	47 ± 13	45 ± 12	55 ± 14	50 ± 13	44 ± 12
131 ± 36	105 ± 37	107 ± 33	108 ± 34	131 ± 34	122 ± 31	108 ± 35	150 ± 42	143 ± 40
96 ± 24	105 ± 49	106 ± 43	98 ± 34	101 ± 18	124 ± 54	101 ± 24	109 ± 39	111 ± 38
—	—	—	—	3.6 ± 2.6	—	1.8 ± 1.3	2.4 ± 2.2	2.8 ± 2.5
3.9	14.2	12.0	6.6	7.6	35.7	2.2	4.5	8.4

Data are means ± SD (continuous measures). To convert the values to millimoles per liter, multiply triglycerides by 0.01129, HDL and LDL by 0.02586, and fasting blood glucose (fB-glucose) by 0.0556. BPRHS, Boston Puerto Rican Health Study.

laser desorption ionization-time of flight mass spectroscopy using a Sequenom platform. RNA was extracted using the Qiagen RNeasy mini kit on pulverized tissue according to the manufacturer's protocols (Qiagen). RNA was reverse transcribed to cDNA using the Superscript III first-strand synthesis from Invitrogen. RT-PCR was performed on an ABI 7900 using ABI TaqMan primer probe sets (HS01564551_G1, *GCK*; and HS01100274_M1, *GCKR*) and normalized to cyclophilin. Data are expressed as Δ Ct (cycle threshold) values.

Statistical analyses. SNP-phenotype association analyses were performed by multivariate linear regression using an additive genetic model. Because of deviation from normal distribution, fasting triglyceride, blood glucose, and CRP concentrations, and carotid IMT measurements were log transformed before analyses. For each participant, residual values were created for log-triglycerides, HDL cholesterol, and LDL cholesterol after adjustment for age, sex, and diabetes status. Residuals were created for BMI after adjustment for age and sex. For analyses of lipid traits, individuals on lipid-lowering medication were excluded (except in NORDIL and Diabetes Registry, where this information was not available). Log-fasting glucose, HOMA insulin resistance index, and insulinogenic index were studied only in nondiabetic individuals, and residuals were created after adjustment for age and sex. To limit the undue influence of outliers in the regression analysis, for each trait we excluded the bottom and top 0.5% of the trait-level distribution in each study sample. We tested the null hypothesis that the trait residuals do not differ by the analyzed genotypes.

To summarize the statistical evidence across the multiple cohorts, we conducted a fixed-effects variance-weighted meta-analysis. We computed a weighted average of the β -coefficient estimates and SEs (from the linear regression models) using the inverse of the variance in each cohort as weights. Heterogeneity between the studies was tested using the Cochran test. One-way ANOVA was used to compare expression levels of *GCK* and *GCKR* in the human liver samples according to genotype. Survival analyses were performed using Cox regression analysis with either age and sex or age, sex, LDL cholesterol, HDL cholesterol, triglycerides, BMI, systolic blood pressure (sBP), diastolic blood pressure (dBP), smoking, family history of myocardial infarction, lipid-lowering medication, antihypertensive medication, and log-CRP as independent predictors.

Analysis of association between log-carotid IMT and *GCKR* was performed by linear regression analysis in individuals without prevalent myocardial infarction or type 2 diabetes after adjustment for age and sex. Because our sample included a modest number of CVD events (321 incident case subjects and 4,781 event-free control subjects), we assessed power to detect an association between *GCKR* SNP genotype and CVD. We approximated hazards ratios using case-control design for discrete traits (31). At 5% significance level, we had a power of 0.58 for a SNP genotype with an odds ratio (OR) of 1.2 per risk allele (that is, OR 1.00, 1.20, and 1.44 for carriers of 0, 1, and 2 risk alleles, respectively), power 0.88 for an OR of 1.30, and power 0.98 for an OR of 1.4 per risk allele, assuming a risk allele frequency of 34% (the Leu allele of SNP rs1260326 has an allele frequency of 34%). For the association of Pro446Leu with carotid IMT ($n = 4,859$ individuals analyzed), at 5% signifi-

cance level, we had a power of 0.88 to detect an association for a SNP genotype that explained 0.2% of the variance in carotid IMT.

In the MPP prospective study, the ORs for risk of developing type 2 diabetes were calculated using logistic regression analyses adjusted for age at inclusion and time to last follow-up, BMI, and sex. Differences in change of fasting triglyceride by genotype during the follow-up time were tested by linear regression analysis adjusted for age at inclusion, sex, time to last follow-up, and diabetes status. After excluding individuals with diabetes at baseline and on follow-up, differences in change of fasting glucose were tested using linear regression analyses with adjustment for age at inclusion, sex, and time to last follow-up.

All statistical analyses were conducted using either SPSS version 14.0 or PLINK (32). All nominal P values of <0.05 were considered significant. All reported P values are two sided.

RESULTS

Association with plasma triglyceride concentrations. The initial association of rs780094 with triglyceride concentrations was studied in each of 12 study samples, representing a range of ancestral groups (Table 2). In all but one sample, the T allele was associated with higher triglycerides (Table 2; P for association ranging from 0.29 to 6×10^{-10}); for example, in the MDC-CC, each copy of the T allele was associated with ~ 5.5 mg/dl higher triglycerides. Across the studies, SNP rs780094 explained between 0.1 and 1.2% of triglyceride variance (after accounting for age, sex, and diabetes status).

Combining the data from the studied 46,549 individuals provided robust evidence for association between the minor T allele at rs780094 and higher triglyceride levels (meta-analysis $P = 3 \times 10^{-56}$; Table 2). The minor T-allele frequency varied from 16.1% in U.S. blacks to 47.6% in the Spanish from Valencia, and the effect size per T allele ranged from 0.6 mg/dl in Dallas Heart Study Hispanics to ~ 6.2 mg/dl in the MPP (Table 2). Mean effect size was ~ 4.2 mg/dl, and we did not observe significant heterogeneity between the studies ($P = 0.15$). We found that the T allele was associated with higher triglycerides in population-based samples and in cohorts of patients with diabetes and hypertension. Furthermore, the T allele was associated with higher triglycerides regardless of the mean triglycer-

TABLE 2
Triglyceride concentrations according to genotype at *GCKR* rs780094* in 12 studies comprising 46,549 individuals

Country	Study	CC (mg/dl)	CT (mg/dl)	TT (mg/dl)	Minor allele frequency	Z score	P value
Finland and Sweden	DGI*	136 ± 89 (1,142)	144 ± 103 (1,194)	164 ± 114 (300)	0.34	-5.76	3.7 × 10 ⁻⁸
Sweden	MDC-CC*	117 ± 70 (2,207)	123 ± 71 (2,457)	128 ± 74 (639)	0.35	-5.45	1.7 × 10 ⁻⁷
Sweden	MPP	107 ± 70 (4,059)	113 ± 77 (4,616)	117 ± 72 (1,425)	0.37	-6.17	1.3 × 10 ⁻⁹
Sweden	NORDIL	151 ± 78 (2,223)	157 ± 79 (2,220)	172 ± 87 (572)	0.34	-6.14	7.4 × 10 ⁻⁹
Sweden	Skania Diabetes 2000 Registry	205 ± 148 (1,076)	224 ± 97 (1,069)	253 ± 104 (259)	0.33	-4.87	1.8 × 10 ⁻⁶
Finland	Botnia-PPP	106 ± 58 (1,273)	115 ± 69 (1,429)	125 ± 69 (409)	0.36	-5.62	4.8 × 10 ⁻⁸
Finland	FINRISK	128 ± 86 (3,009)	130 ± 89 (3,398)	142 ± 105 (931)	0.36	-5.00	8.0 × 10 ⁻⁷
Spain	Valencia	108 ± 66 (446)	112 ± 64 (792)	118 ± 71 (370)	0.48	-2.26	0.02
U.S.	Dallas Heart Study blacks	103 ± 90 (1,163)	108 ± 78 (444)	110 ± 59 (44)	0.16	-2.53	0.01
U.S.	Dallas Heart Study Hispanics	149 ± 138 (244)	151 ± 124 (263)	161 ± 155 (58)	0.34	-0.56	0.29
U.S.	Dallas Heart Study whites	131 ± 115 (342)	135 ± 97 (455)	164 ± 124 (145)	0.40	-4.04	4.4 × 10 ⁻⁵
U.S.	GOLDN	109 ± 75 (378)	143 ± 90 (538)	133 ± 87 (146)	0.39	-2.24	0.03
U.S.	BPRHS	153 ± 104 (423)	165 ± 92 (330)	174 ± 118 (84)	0.30	-2.96	0.003
Singapore	Singapore NHS-98 Chinese	112 ± 64 (842)	120 ± 71 (1233)	131 ± 82 (616)	0.46	-5.41	3.0 × 10 ⁻⁵
Singapore	Singapore NHS-98 Malays	132 ± 75 (268)	150 ± 91 (332)	153 ± 97 (134)	0.41		0.02
Singapore	Singapore NHS-98 Asian Indians	140 ± 79 (332)	151 ± 89 (188)	163 ± 109 (28)	0.22		0.08
Total n = 46,549		19,427	20,958	6,164	Meta-analysis P value 3 × 10 ⁻⁵⁶		

Data means ± SD (*n*) (continuous raw measures). Association analyses were conducted with an outcome variable of residual log-triglyceride concentration after adjustment for age, sex, and diabetes status. For ease of interpretation, unadjusted triglyceride concentrations are presented in the table. To convert the values to millimoles per liter, multiply triglycerides by 0.01129. All *P* values are two sided. *DGI and MDC-CC results have been reported in ref. 1. BPRHS, Boston Puerto Rican Health Study.

ide level in the sample (e.g., the mean triglyceride concentration is considerably lower in a population-based sample, such as MDC-CC [at 122 mg/dl] compared with the Diabetes Registry cohort [at 231 mg/dl] comprising entirely individuals with type 2 diabetes).

Association with plasma triglyceride-related metabolic traits. We next explored the relationship between SNP rs780094 and related metabolic traits, including plasma LDL cholesterol, HDL cholesterol, apolipoprotein concentrations, FFA concentrations, and BMI. SNP rs780094 was not associated with LDL cholesterol or HDL cholesterol in any of the samples (supplementary Table 1, which is available in an online appendix at <http://dx.doi.org/10.2337/db08-0516>). As expected, given the correlation between triglyceride and apoB concentration ($r = 0.42$ [$P < 0.0001$] and 0.57 [$P < 0.0001$] in DGI and FINRISK97, respectively), rs780094 was associated with apoB concentration in a meta-analysis of the DGI and FINRISK97 cohorts ($P = 7.5 \times 10^{-5}$), with the T-allele carriers having the highest apoB concentration. Fasting FFA and triglyceride concentrations are weakly correlated ($r = 0.25$ and 0.20 in DGI and MPP-MM, respectively), and we did not observe any association between rs780094 and fasting FFA ($P = 0.39$ and 0.90 in DGI and MPP-MM cohorts, respectively) or suppression of FFA levels at 2 h in an OGTT ($P = 0.70$ and 0.98 in DGI and MPP-MM cohorts, respectively). *GCKR* rs780094 was nominally associated with BMI in the Singaporean NHS-98 study ($P = 0.04$) but not in any of the other samples (supplementary Table 2).

Because two genome-wide association studies recently reported association between CRP levels and rs780094 and

rs1260326 (9,10), we tested for association between *GCKR* rs1260326 and rs780094 and CRP in MDC-CC. Both SNPs were strongly associated with CRP levels with the T-allele carriers having significantly higher levels (CC 2.5 ± 4.7 , CT 2.6 ± 4.2 , and TT 2.9 ± 4.3 mg/l, $P = 4.5 \times 10^{-5}$ in linear regression analysis of rs1260326 adjusted for age and sex).

Association with measures of glucose metabolism. We studied the association of SNP rs780094 with fasting glucose concentrations in 33,995 nondiabetic individuals and HOMA estimates of insulin resistance in 11,084 nondiabetic individuals. Despite the fact that the T allele was consistently associated with higher triglycerides, T-allele carriers had significantly lower fasting plasma glucose levels in six of the studied populations, and a similar trend was observed in the other populations (Table 3, $P_{\text{meta}} = 1 \times 10^{-13}$); for example, in the MDC-CC, each copy of the T allele was associated with ~0.5 mg/dl lower fasting blood glucose. In addition, T-allele carriers were more insulin sensitive as estimated by the HOMA insulin resistance index ($P_{\text{meta}} = 5.0 \times 10^{-5}$). In DGI and Botnia PPP cohorts, we could calculate the insulinogenic index during an OGTT in 999 and 3,184 nondiabetic individuals, respectively. Insulin secretion capacity calculated as the insulinogenic index did not differ significantly between the different *GCKR* genotype carriers ($P = 0.72$ and 0.27 , respectively).

Encouraged by these results for intermediate traits, we also tested association of rs780094 with type 2 diabetes in our Nordic cohorts with similar minor allele frequency of the SNP (DGI, MDC-CC, NORDIL, Diabetes Registry, and FINRISK97). Of 24,034 individuals, 5,578 had type 2 diabe-

TABLE 3
Measures of glucose tolerance according to genotype at GCKR rs780094 among nondiabetic individuals within the study populations

Country	Study	Fasting plasma glucose			P value	HOMA (insulin resistance)			P value
		CC (mg/dl)	CT (mg/dl)	TT (mg/dl)		CC (mmol × mU)	CT (mmol × mU)	TT (mmol × mU)	
Finland and Sweden	DGI	101 ± 31 (609)	99 ± 32 (675)	99 ± 31 (170)	0.25	1.8 ± 1.2 (571)	1.8 ± 1.0 (623)	1.7 ± 1.0 (161)	0.03
Sweden	MDC-CC	99 ± 9 (2,043)	99 ± 9 (2,301)	98 ± 17 (610)	0.02	1.8 ± 1.7 (1,982)	1.8 ± 1.2 (2,226)	1.7 ± 1.2 (592)	0.04
Sweden	MPP	87 ± 9 (3,893)	87 ± 8 (4,435)	86 ± 8 (1,364)	0.00048	—	—	—	—
Sweden	NORDIL	102 ± 16 (2,024)	102 ± 16 (2,025)	99 ± 16 (538)	0.004	—	—	—	—
Finland	Botnia-PPP	83 ± 10 (1,383)	82 ± 10 (1,526)	81 ± 12 (444)	0.00022	—	—	—	—
Spain	Valencia	94 ± 17 (433)	92 ± 16 (754)	92 ± 19 (359)	0.01	—	—	—	—
U.S.	Dallas Heart Study blacks	91 ± 13 (1,102)	92 ± 16 (414)	89 ± 10 (42)	0.23	—	—	—	—
U.S.	Dallas Heart Study Hispanics	94 ± 10 (222)	94 ± 11 (246)	92 ± 10 (59)	0.22	—	—	—	—
U.S.	Dallas Heart Study whites	92 ± 12 (347)	91 ± 11 (475)	92 ± 13 (148)	0.32	—	—	—	—
U.S.	GOLDN	99 ± 13 (353)	100 ± 14 (492)	97 ± 9 (141)	0.47	3.3 ± 2.3 (352)	3.4 ± 2.4 (490)	3.3 ± 1.9 (141)	0.75
U.S.	BPRHS	104 ± 31 (262)	103 ± 18 (218)	99 ± 16 (55)	0.16	—	—	—	—
Singapore	Singapore NHS-98 Chinese	101 ± 19 (823)	99 ± 16 (1,203)	98 ± 19 (604)	0.001	1.9 ± 1.3 (823)	1.8 ± 1.3 (1,203)	1.7 ± 1.3 (604)	0.01
Singapore	Singapore NHS-98 Malays	104 ± 25 (256)	105 ± 30 (317)	101 ± 14 (128)	0.40	2.5 ± 2.5 (256)	2.3 ± 2.0 (317)	2.3 ± 2.3 (128)	0.32
Singapore	Singapore NHS-98 Asian Indians	107 ± 33 (298)	102 ± 22 (177)	106 ± 19 (27)	0.22	3.0 ± 2.7 (298)	2.6 ± 2.1 (177)	2.5 ± 2.8 (27)	0.11
Total: 33,995		14,048	15,258	4,689	Meta-analysis P value 7 × 10 ⁻¹⁴	4,328	5,090	1,666	Meta-analysis P value 5.0 × 10 ⁻⁵

Data are means ± SD (*n*) or *n*. Continuous raw measures or blood glucose converted to plasma glucose using a correction factor of 1.13. Association analyses were conducted with an outcome variable of residual log-glucose concentration after adjustment for age and sex. For ease of interpretation, unadjusted glucose concentrations are presented in the table. HOMA insulin resistance index was calculated using the following formula: fasting plasma glucose × fasting insulin/22.5. Two-sided *P* values are shown for all cohorts. To convert the fasting plasma glucose values to millimoles per liter, multiply by 0.055.




FIG. 1. In silico and genotype fine-mapping of the *GCKR* locus. To define the strongest signal for the association on chromosome 2p23 for triglycerides, a region spanning ~417 kb and containing 17 annotated genes was fine-mapped by two different approaches, imputation of untyped SNPs (29) (or so-called in silico fine-mapping) (A) and genotyping tagging SNPs across the region (B). Both in silico and genotype fine-mapping methods indicated the Pro446Leu as the variant with the strongest association with triglyceride levels. The genotype consensus rate between the imputed genotypes and genotyped genotypes was ~95.7%.

tes, and the frequency of the rs780094 T-allele was 35.2% among nondiabetic individuals compared with 33.4% in type 2 diabetic patients (OR 0.93 [95% CI 0.88–0.97], $P = 0.0006$). Of the nondiabetic individuals, 2.3% were homozygous for the T allele compared with 10.9% of type 2 diabetic patients, and the TT genotype was associated with a protection from type 2 diabetes when compared with CC-genotype carriers (OR 0.84 [0.76–0.93], $P = 0.0008$).

We next evaluated whether the rs780094 genotype was associated with hepatic glucose output in 125 men who had undergone a hyperinsulinemic-euglycemic clamp with infusion of [^3H]glucose and assessment of basal and clamp hepatic glucose production. *GCKR* rs780094 T-allele carriers did not have a significantly lower basal rate of hepatic glucose production (CC 2.26 ± 0.27 , CT 2.18 ± 0.20 , and TT $2.16 \pm 0.09 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$, $P = 0.10$), but their hepatic glucose output during the hyperinsulinemic state was slightly lower compared with that of C-allele carriers (CC 0.26 ± 0.59 , CT 0.20 ± 0.40 , and TT $0.09 \pm 0.22 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$, $P = 0.01$).

Fine-mapping of the *GCKR* locus. We fine-mapped the *GCKR* region with two different approaches, by genotyping tag and coding SNPs across the region and by imputing untyped SNPs (or so-called in silico fine-mapping) (Fig. 1). We genotyped 104 SNPs across the ~417-kb region and studied the association of these SNPs with triglyceride concentrations (supplementary Table 3). A common *GCKR* coding SNP rs1260326 (Pro446Leu) gave the strongest signal for association with triglycerides ($P = 9.4 \times 10^{-10}$).

Fine-mapping by imputation also revealed that *GCKR* coding SNP rs1260326 (Pro446Leu) gave the strongest signal for triglyceride concentrations ($P = 1.5 \times 10^{-9}$) in the associated interval on chromosome 2p23. In HapMap CEU, *GCKR* coding SNP rs1260326 shows strong linkage disequilibrium to the intronic SNP rs780094 ($r^2 = 0.93$). We performed regression analysis, including both rs1260326 and rs780094 as predictors of triglyceride levels in MDC-CC, but because of the strong correlation between the SNPs, none of the two were significant in this analysis ($P = 0.18$ and 0.80 for rs1260326 and rs780094, respectively).

Figure 1 summarizes the results of both fine-mapping approaches. Both the genotyping and the in silico fine-

mapping methods indicated the Pro446Leu as the variant with strongest association with triglyceride levels. To evaluate the fidelity of the MACH imputation algorithm (24), we compared the genotypes generated by Sequenom genotyping for 57 SNPs with that predicted by imputation. The genotype consensus rate was 95.7%.

Longitudinal changes in fasting triglyceride and glucose stratified by *GCKR* Pro446Leu genotype. In the MPP cohort, the Pro446Leu was strongly associated with higher triglycerides and lower fasting blood glucose both at baseline ($P = 6 \times 10^{-22}$ and 0.0005 , respectively) and after the mean follow-up period of 23.4 years ($P = 3 \times 10^{-29}$ and 0.004 , respectively) (Fig. 2). In addition, the triglyceride levels of the Leu446 carriers increased more over time compared with those of homozygous Pro446 carriers ($P = 8 \times 10^{-5}$), whereas change in fasting glucose over time did not differ by genotype status (Fig. 2).

In the MPP study, among 17,037 individuals free of type 2 diabetes at baseline, 2,063 (12.1%) individuals developed type 2 diabetes during the follow-up period. Carriage of the Leu allele trended to protect from development of type 2 diabetes (OR 0.96 [95% CI 0.91–1.02], $P = 0.27$).

Association of *GCKR* variation with CVD and carotid IMT. In MDC-CC, 321 individuals experienced the first CVD end point during the mean follow-up time of 10.5 ± 1.8 years. Neither rs780094 nor rs1260326 predicted CVD ($P = 0.85$ and 0.45 , respectively). The results were similar when age, sex, LDL cholesterol, HDL cholesterol, triglycerides, BMI, sBP, dBP, smoking, family history of myocardial infarction, lipid-lowering medication, antihypertensive medication, and CRP were included as covariates. *GCKR* variants were also not associated with carotid IMT in MDC-CC. No association was detected between *GCKR* variants and common carotid artery IMT ($P = 0.94$ and 0.63 for rs780094 and rs1260326, respectively).

Hepatic expression of *GCK* and *GCKR* according to *GCKR* genotypes. We next examined whether rs780094 or Pro446Leu was associated with transcript levels of *GCKR* and/or *GCK* in human liver. In a modest number of liver samples ($n = 60$), neither rs780094 nor Pro446Leu genotype was associated with transcript levels of *GCK* or *GCKR* (supplementary Table 4).




FIG. 2. Fasting triglycerides and blood glucose levels in MPP participants before and after a mean follow-up time of 23.4 years according to *GCKR* Pro446Leu. In a large prospective study, the Pro446Leu was strongly associated with higher triglycerides and lower fasting blood glucose both at baseline and after the follow-up. In addition, among 12,528 individuals not on lipid-lowering medication, fasting triglyceride levels increased more during the follow-up time among Leu446 allele carriers compared with homozygous Pro446 carriers ($P = 8 \times 10^{-5}$), whereas the change in fasting blood glucose during the follow-up time was not significant among 12,964 individuals without diabetes either at baseline or at follow-up.

DISCUSSION

In line with the opposite effects of *GCKR*-pathway manipulation on glucose and triglyceride concentrations in rodent models and a recent association study in Danes (8), our study provides compelling evidence that common DNA sequence variants in *GCKR* are associated with opposite effects on fasting triglyceride and glucose concentrations in humans and modest protection from type 2 diabetes. Both imputation and genotype fine-mapping of the *GCKR* locus yielded a nonsynonymous coding SNP (Pro446Leu) as the strongest association signal, suggesting the hypothesis that this nonsynonymous coding SNP is the causal variant for the observed associations. We also provide evidence that the Leu446 allele carriers increase their triglyceride levels more over time compared with noncarriers. In addition, our data suggest that, at least within regions of high linkage disequilibrium, genotypes predicted by imputation are highly accurate and may provide a good starting point for genotype fine-mapping.

The exact mechanism for the effect of the *GCKR* variant on blood glucose, triglycerides, and CRP remains to be defined. A potential explanation is the opposite and overriding effects of increased glucose utilization and glycolytic flux on liver glucose and lipid metabolism. With increased glucose utilization and glycolytic flux, PEPCK and glucose-6-phosphatase are downregulated, whereas G6P, phosphofruktokinase, and fatty acid synthase are upregulated. These changes increase glycogen synthesis and malonyl CoA concentration and direct fatty-acyl-CoA into de novo lipogenesis and VLDL triglyceride production (33). However, the consequence of in vivo glucose metabolism is enhanced suppression of hepatic glucose output (33). Our observation that the *GCKR* variant T allele associates with higher triglycerides, lower fasting glucose, and lower hepatic glucose output during a euglycemic-hyperinsulinemic clamp agrees with this hypothesis. Finally, the T-allele carriers had insulin secretion capacity similar to that of noncarriers. Thus, the association is

similar to that of *GCKR*-30G/A, which affects the glucose levels needed to induce insulin secretion.

Our human studies propose the hypothesis that *GCKR* Pro446Leu may mimic the consequences of *GCKR* overexpression in rodent models with upregulation of glucose utilization and VLDL-triglyceride synthesis and downregulation of gluconeogenesis. The Pro446Leu variant has been introduced into rat cDNA but was not found to affect the functional properties of the rat protein when prepared by overexpression in *Escherichia coli* (34). Unfortunately, the human protein could not be produced using that expression system (32). Although human and rat *GCKR* share 88% identity, the human protein is importantly different from rat *GCKR*: human *GCKR* is a more potent inhibitor of G6P than rat *GCKR* in the absence of fructose-6-phosphate, and human *GCKR* has higher affinity for fructose-6-phosphate (35). Thus, the potential impact of the amino acid difference on overall structure and function of human *GCKR* remains to be defined.

There are conflicting data on the association between circulating triglyceride concentrations and risk of CVD (36,37). Our data combined with data for other common variants suggest a potential explanation for the varying risk associated with high triglycerides. DNA sequence variants in some genes (e.g., *APOB*) have been associated with both increased triglycerides and markers of increased atherosclerosis risk, such as elevated LDL cholesterol (38). Similarly, common genetic variations in both lipoprotein lipase (*LPL*) and apoA5 (*APOA5*) genes (rs328 and rs3133506, respectively) are associated with both increased triglycerides and markers of increased atherosclerosis risk, such as decreased HDL cholesterol (39,40). However, at *GCKR* Pro446Leu, the variant allele is associated with higher triglycerides and higher CRP levels but also with a favorable metabolic marker, namely decreased glucose. Our finding of no association between the *GCKR* variant and CVD events or carotid IMT is thus not surprising but instead proposes that the risk of CVD associated with

higher triglycerides may vary based on the specific profile of genetic variants in different genes contributing to an increased triglyceride concentration. However, given the limited number of CVD events in our study, this result needs further confirmation in other studies.

We provide convincing evidence that common variation in *GCKR* is associated with opposite effects on fasting plasma triglyceride and glucose concentrations in multiple human populations and demonstrate that the strongest association signal resides at coding SNP rs1260326 (Pro446Leu) in *GCKR*. Taken together, the data position *GCKR* in central pathways regulating both hepatic triglyceride and glucose metabolism in humans.

ACKNOWLEDGMENTS

M.O.-M. is supported by the Diabetes program at the Lund University and Novo Nordic Foundations, the Swedish Medical Research Council, the Swedish Heart and Lung Foundation, the Region Skåne, the Medical Faculty of Lund University, the Malmö University Hospital, the Albert Pålsson Research Foundation, and the Crafoord Foundation. O.M. is supported by the Swedish Medical Research Council, the Swedish Heart and Lung Foundation, the Region Skåne, the Medical Faculty of Lund University, the Malmö University Hospital, the Albert Pålsson Research Foundation, the Crafoord Foundation, the Swedish Medical Society, the Ernhöld Lundströms Research Foundation, the Mossfelt Foundation, and the King Gustav V and Queen Victoria Foundation. M.J.R. is supported by NIH Grant R01-NS-053646. V.L. is supported by the Sigrid Juselius Foundation. K.L.T. is supported by U.S. Department of Agriculture Contract 58-1950-9-001, NIH Grant P01-AG-023394-03, NIH Contract 53-K06-5-10, and NIH-NHLBI Grants U01-HL-72524 and HL-54776. M.-R.T. is supported by the Sigrid Juselius Foundation, the Finnish Heart Foundation, and the Clinical Research Institute, Helsinki University Central Hospital. L.G. is supported by the Sigrid Juselius Foundation, the Finnish Diabetes Research Foundation, The Folkhalsan Research Foundation, Knut and Alice Wallenberg Stiftelse, the Clinical Research Institute HUCH, and a Linné grant from the Swedish Research Council. J.M.O. is supported by U.S. Department of Agriculture Contract 58-1950-9-001, NIH Grant P01-AG-023394-03, NIH Contract 53-K06-5-10, and NIH-NHLBI grants U01-HL-72524 and HL-54776. S.K. is supported by a Doris Duke Charitable Foundation Clinical Scientist Development Award, a charitable gift from the Fannie E. Rippel Foundation, and NIH Grant K23-HL-083102. The University of Washington School of Pharmacy Human Liver Bank is supported in part by NIH Grant PO1-GM-32165. Liver expression studies are supported in part by the University of Washington School of Pharmacy Drug Metabolism, Transport and Pharmacogenomics Research program (funded by unrestricted gifts from Abbott, Allergan, Amgen, Bend Research, Bristol-Myers Squibb, Eli Lilly, Johnson & Johnson, Merck, Roche, and Pfizer).

We thank Dr. Helen Hobbs and Dr. Jonathan Cohen for their contribution of the Dallas Heart Study data and the Donald W. Reynolds Foundation for its support of the Dallas Heart Study. We are indebted to the staff and participants of all of the study populations for their important contributions. We thank Malin Svensson for technical assistance in Malmö and the RSKC2 (Region Skania) genotyping facility for help with genotyping of the NORDIL sample.

REFERENCES

1. The Diabetes Genetics Initiative of the Broad Institute of MIT and Harvard, Lund University, and Novartis Institutes for BioMedical Research: Genome-wide association analysis identifies loci for type 2 diabetes and triglyceride levels. *Science* 316:1331–1336, 2007
2. Vionnet N, Stoffel M, Takeda J, Yasuda K, Bell GI, Zouali H, Lesage S, Velho G, Iris F, Passa P, Froguel P: Nonsense mutation in the glucokinase gene causes early-onset non-insulin-dependent diabetes mellitus. *Nature* 356: 721–722, 1992
3. Glaser B, Chiu KC, Anker R, Nestorowicz A, Landau H, Ben-Bassat H, Shlomai Z, Kaiser N, Thornton PS, Stanley CA: Familial hyperinsulinism caused by an activating glucokinase mutation. *N Engl J Med* 338:226–230, 1998
4. Grimsby J, Coffey JW, Dvorozniak MT, Magram J, Li G, Matschinsky FM, Shiota C, Kaur S, Magnuson MA, Grippo JF: Characterization of glucokinase regulatory protein-deficient mice. *J Biol Chem* 275:7826–7831, 2000
5. Slosberg ED, Desai UJ, Fanelli B, St Denny I, Connelly S, Kaleko M, Boettcher BR, Caplan SL: Treatment of type 2 diabetes by adenoviral-mediated overexpression of the glucokinase regulatory protein. *Diabetes* 50:1813–1820, 2001
6. O'Doherty RM, Lehman DL, Telemaque-Potts S, Newgard CB: Metabolic impact of glucokinase overexpression in liver: lowering of blood glucose in fed rats is accompanied by hyperlipidemia. *Diabetes* 48:2022–2027, 1999
7. Ferre T, Riu E, Bosch F, Valera A: Evidence from transgenic mice that glucokinase is rate limiting for glucose utilization in the liver. *FASEB J* 10:1213–1218, 1996
8. Sparsø T, Andersen G, Nielsen T, Burgdorf KS, Gjesing AP, Nielsen AL, Albrechtsen A, Rasmussen SS, Jørgensen T, Borch-Johnsen K, Sandbæk A, Lauritzen T, Madsbad S, Hansen T, Pedersen O: The *GCKR* rs780094 polymorphism is associated with elevated fasting serum triacylglycerol, reduced fasting and OGTT-related insulinaemia, and reduced risk of type 2 diabetes. *Diabetologia* 51:70–75, 2008
9. Ridker PM, Pare G, Parker A, Zee RY, Danik JS, Buring JE, Kwiatkowski D, Cook NR, Miletich JP, Chasman DI: Loci related to metabolic-syndrome pathways including *LEPR*, *HNF1A*, *IL6R*, and *GCKR* associate with plasma C-reactive protein: the Women's Genome Health Study. *Am J Hum Genet* 82:1185–1192, 2008
10. Reiner AP, Barber MJ, Guan Y, Ridker PM, Lange LA, Chasman DI, Walston JD, Cooper GM, Jenny NS, Rieder MJ, Durda JP, Smith JD, Novembre J, Tracy RP, Rotter JJ, Stephens M, Nickerson DA, Krauss RM: Polymorphisms of the *HNF1A* gene encoding hepatocyte nuclear factor-1 alpha are associated with C-reactive protein. *Am J Hum Genet* 82:1193–1201, 2008
11. Groop L, Forsblom C, Lehtovirta M, Tuomi T, Karanko S, Nissen M, Ehrnström BO, Forsen B, Isomaa B, Snickars B, Taskiran M-R: Metabolic consequences of a family history of NIDDM (the Botnia study): evidence for sex-specific parental effects. *Diabetes* 45:1585–1593, 1996
12. Bøg-Hansen E, Lindblad U, Bengtsson K, Ransta J, Melander A, Råstam L: Risk factor clustering in patients with hypertension and non-insulin-dependent diabetes mellitus: The Skaraborg Hypertension Project. *J Intern Med* 243:223–232, 1998
13. Berglund G, Elmstahl S, Janzon L, Larsson SA: The Malmö Diet and Cancer Study: design and feasibility. *J Intern Med* 233:45–51, 1993
14. Rosvall M, Janzon L, Berglund G, Engström G, Hedblad B: Incident stroke is related to carotid IMT even in the absence of plaque. *Atherosclerosis* 179:325–331, 2005
15. Rosvall M, Janzon L, Berglund G, Hedblad B: Incident coronary events and case fatality in relation to common carotid intima-media thickness. *J Intern Med* 257:430–437, 2005
16. Berglund G, Nilsson P, Eriksson KF, Nilsson JA, Hedblad B, Kristenson H, Lindgärde F: Long-term outcome of the Malmö Preventive Project: mortality and cardiovascular morbidity. *J Intern Med* 247:19–29, 2000
17. Tripathy D, Eriksson KF, Orho-Melander M, Fredriksson J, Ahlqvist G, Groop L: Parallel manifestation of insulin resistance and beta cell decompensation is compatible with a common defect in type 2 diabetes. *Diabetologia* 47:782–793, 2004
18. Hansson L, Hedner T, Lund-Johansen P, Kjeldsen SE, Lindholm LH, Syvertsen JO, Lanke J, de Faire U, Dahlöf B, Karlberg BE: Randomised trial of effects of calcium antagonists compared with diuretics and beta-blockers on cardiovascular morbidity and mortality in hypertension: the Nordic Diltiazem (NORDIL) study. *Lancet* 356:359–365, 2000
19. Lindholm E, Agardh E, Tuomi T, Groop L, Agardh CD: Classifying diabetes according to the new WHO clinical stages. *Eur J Epidemiol* 17:983–989, 2001
20. World Health Organization: *Definition, Diagnosis, and Classification of Diabetes Mellitus and Its Complications. Report of a WHO Consultation.*

- Part 1: Diagnosis and Classification of Diabetes Mellitus*. Geneva, World Health Organization, 1999
21. Vartiainen E, Jousilahti P, Alftan G, Sundvall J, Pietinen P, Puska P: Cardiovascular risk factor changes in Finland 1972–1997. *Int J Epidemiol* 29:49–56, 2000
 22. Corella D, Guillén M, Sáiz C, Portolés O, Sabater A, Cortina S, Folch J, González JI, Ordovas JM: Environmental factors modulate the effect of the APOE genetic polymorphism on plasma lipid concentrations: ecogenetic studies in a Mediterranean Spanish population. *Metabolism* 50:936–944, 2001
 23. Victor RG, Haley RW, Willett DL, Peshock RM, Vaeth PC, Leonard D, Basit M, Cooper RS, Iannacchione VG, Visscher WA, Staab JM, Hobbs HH, Dallas Heart Study Investigators: The Dallas Heart Study: a population-based probability sample for the multidisciplinary study of ethnic differences in cardiovascular health. *Am J Cardiol* 93:1473–1480, 2004
 24. Higgins M, Province M, Heiss G, Eckfeldt J, Ellison RC, Folsom AR, Rao DC, Sprafka JM, Williams R: NHLBI Family Heart Study: objectives and design. *Am J Epidemiol* 143:1219–1228, 1996
 25. Tucker KL: Stress and nutrition in relation to excess development of chronic disease in Puerto Rican adults living in the Northeastern USA. *J Med Invest* 52:252–258, 2005
 26. Tucker KL, Bianchi L, Maras J, Bermudez O: Adaptation of a food frequency questionnaire to assess diets of Puerto Rican and non-Hispanic adults. *Am J Epidemiol* 148:507–518, 1998
 27. Cutter J, Tan BY, Chew SK: Levels of cardiovascular disease risk factors in Singapore following a national intervention programme. *Bull World Health Organ* 79:908–915, 2001
 28. Deurenberg-Yap M, Li T, Tan WL, van Staveren WA, Chew SK, Deurenberg P: Can dietary factors explain differences in serum cholesterol profiles among different ethnic groups (Chinese, Malays and Indians) in Singapore? *Asia Pac J Clin Nutr* 10:39–45, 2001
 29. Scott LJ, Mohlke KL, Bonnycastle LL, Willer CJ, Li Y, Duren WL, Erdos MR, Stringham HM, Chines PS, Jackson AU, Prokunina-Olsson L, Ding CJ, Swift AJ, Narisu N, Hu T, Pruim R, Xiao R, Li XY, Conneely KN, Riebow NL, Sprau AG, Tong M, White PP, Hetrick KN, Barnhart MW, Bark CW, Goldstein JL, Watkins L, Xiang F, Saramies J, Buchanan TA, Watanabe RM, Valle TT, Kinnunen L, Abecasis GR, Pugh EW, Doheny KF, Bergman RN, Tuomilehto J, Collins FS, Boehnke M: A genome-wide association study of type 2 diabetes in Finns detects multiple susceptibility variants. *Science* 316:1341–1345, 2007
 30. Rieder MJ, Reiner AP, Gage BF, Nickerson DA, Eby CS, McLeod HL, Blough DK, Thummel KE, Veenstra DL, Rettie AE: Effect of VKORC1 haplotypes on transcriptional regulation and warfarin dos. *N Engl J Med* 352:2285–2293, 2005
 31. Sham PC, Cherny SS, Purcell S, Hewitt JK: Power of linkage versus association analysis of quantitative traits, by use of variance-components models, for sibship data. *Am J Hum Genet* 66:1616–1630, 2000
 32. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P, Bakker PIW, Daly MJ, Sham PC: PLINK: A tool set for whole genome association and population based linkage analyses. *Am J Hum Genet* 81:559–575, 2007
 33. Gerich JE: Control of glycaemia. *Baillieres Endocrinol Metab* 7:551–586, 1993
 34. Veiga-da-Cunha M, Delplanque J, Gillain A, Bonthron DT, Boutin P, Van Schaftingen E, Froguel P: Mutations in the glucokinase regulatory protein gene in 2p23 in obese French Caucasians. *Diabetologia* 46:704–711, 2003
 35. Brocklehurst KJ, Davies RA, Agius L: Differences in regulatory properties between human and rat glucokinase regulatory protein *Biochem J* 378: 693–697, 2004
 36. Sarwar N, Danesh J, Eiriksdottir G, Sigurdsson G, Wareham N, Bingham S, Boekholdt SM, Khaw KT, Gudnason V: Triglycerides and the risk of coronary heart disease: 10,158 incident cases among 262,525 participants in 29 Western prospective studies. *Circulation* 115:450–458, 2007
 37. McBride PE: Triglycerides and risk for coronary heart disease. *JAMA* 298:336–338, 2007
 38. Sposito AC, Gonbert S, Turbin G, Chapman MJ, Thillet J: Common promoter C516T polymorphism in the ApoB gene is an independent predictor of carotid atherosclerotic disease in subjects presenting a broad range of plasma cholesterol levels. *Arterioscler Thromb Vasc Biol* 24:2192–2195, 2004
 39. Rip J, Nieman MC, Wareham NJ, Luben R, Bingham SA, Day NE, van Miert JN, Hutten BA, Kastelein JJ, Kuivenhoven JA, Khaw KT, Boekholdt SM: Serum lipoprotein lipase concentration and risk for future coronary artery disease: the EPIC-Norfolk prospective population study. *Arterioscler Thromb Vasc Biol* 26:637–642, 2006
 40. Lai CQ, Demissie S, Cupples LA, Zhu Y, Adiconis Y, Parnell LD, Corella D, Ordovas JM: Influence of the APOA5 locus on plasma triglyceride, lipoprotein subclasses, and CVD risk in the Framingham Heart Study. *J Lipid Res* 45:2096–2105, 2004