

Glucagon Supports Postabsorptive Plasma Glucose Concentrations in Humans With Biologically Optimal Insulin Levels

Benjamin A. Cooperberg and Philip E. Cryer

OBJECTIVE—Based on the premise that postabsorptive patients with type 1 diabetes receiving intravenous insulin in a dose that maintains stable euglycemia are receiving biologically optimal insulin replacement, we tested the hypothesis that glucagon supports postabsorptive plasma glucose concentrations in humans.

RESEARCH DESIGN AND METHODS—Fourteen patients with type 1 diabetes were studied after an overnight fast on up to five occasions. Insulin was infused intravenously to hold plasma glucose concentrations at ~ 100 mg/dl (5.6 mmol/l) overnight and fixed from -60 to 240 min the following morning. From 0 through 180 min the patients also received 1) saline, 2) octreotide 30 ng \cdot kg $^{-1}$ \cdot min $^{-1}$ with growth hormone replacement or octreotide with growth hormone, plus 3) glucagon in doses of 0.5 ng \cdot kg $^{-1}$ \cdot min $^{-1}$, 4) 1.0 ng \cdot kg $^{-1}$ \cdot min $^{-1}$, and 5) 2.0 ng \cdot kg $^{-1}$ \cdot min $^{-1}$.

RESULTS—Compared with a mean \pm SE of 98 ± 5 mg/dl (5.4 ± 0.3 mmol/l) at 180 min during saline, mean plasma glucose concentrations declined to 58 ± 1 mg/dl (3.2 ± 0.1 mmol/l) ($P < 0.001$) at 180 min during octreotide plus saline and were 104 ± 16 mg/dl (5.8 ± 0.9 mmol/l) (NS), 143 ± 13 mg/dl (7.9 ± 0.7 mmol/l) ($P = 0.004$), and 160 ± 15 mg/dl (8.9 ± 0.8 mmol/l) ($P < 0.001$) at 180 min during octreotide plus glucagon in doses of 0.5 , 1.0 , and 2.0 ng \cdot kg $^{-1}$ \cdot min $^{-1}$, respectively.

CONCLUSIONS—In the setting of biologically optimal insulin replacement, suppression of glucagon secretion with octreotide caused a progressive fall in plasma glucose concentrations that was prevented by glucagon replacement. These data document that glucagon supports postabsorptive glucose concentrations in humans. *Diabetes* 59:2941–2944, 2010

The current interest in the development of drugs that block the action or secretion of glucagon for the treatment of diabetes rests on the premise that glucagon supports the plasma glucose concentration and, therefore, that glucagon, in concert with insulin deficiency, may play a role in the pathogenesis of hyperglycemia in diabetes (1–4). That premise is based largely on studies with somatostatin including those with the pancreatic clamp technique (4,5). That technique involves infusion of somatostatin (or of the somatostatin analog octreotide) (6), which suppresses insulin and glu-

From the Division of Endocrinology, Metabolism and Lipid Research, Washington University School of Medicine, St. Louis, Missouri.

Corresponding author: Philip E. Cryer, pcrayer@wustl.edu.

Received 26 May 2010 and accepted 29 July 2010. Published ahead of print at <http://diabetes.diabetesjournals.org> on 10 August 2010. DOI: 10.2337/db10-0750.

© 2010 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

cagon secretion among other actions, alone and with insulin replacement, glucagon replacement, and both insulin and glucagon replacement to document the roles of suppression of those hormones in the changes in glycemia (or in glucose kinetics when glucose concentrations are clamped) that result from administration of somatostatin (7–10).

Obviously, the biological appropriateness of the putative replacement doses of insulin and glucagon are critical to the interpretation of pancreatic clamp data (11,12). Excessive insulin replacement (or insufficient glucagon replacement) would confound the data. Insulin has been infused peripherally in doses of 0.14 (13), 0.15 (7), 0.20 (14), and 0.24 (15) mU \cdot kg $^{-1}$ \cdot min $^{-1}$ in humans to replace basal insulin levels during infusion of somatostatin. However, we found that intravenous insulin doses as low as 0.15 mU \cdot kg $^{-1}$ \cdot min $^{-1}$ are excessive; when infused alone in healthy humans they drove plasma glucose concentrations down to subnormal levels and thus activated glucose counterregulatory systems (nearly complete suppression of insulin secretion and stimulation of glucagon and epinephrine secretion) (11). Although it also lowered plasma glucose concentrations, insulin infused in a dose of 0.10 mU \cdot kg $^{-1}$ \cdot min $^{-1}$ did not drive glucose down to subnormal levels and therefore did not activate glucose counterregulatory systems, at least over 2 h (11). Accordingly, we used the latter lower insulin replacement dose in additional pancreatic clamp studies in healthy adults (12). Octreotide infusion caused plasma glucose concentrations to decrease and then increase as expected (6–10); octreotide plus insulin in a dose of 0.10 mU \cdot kg $^{-1}$ \cdot min $^{-1}$ caused a sustained decrease in plasma glucose consistent with the interpretation that glucagon, in concert with insulin, supports the postabsorptive plasma glucose concentration. However, the addition of glucagon in a dose of 1.0 ng \cdot kg $^{-1}$ \cdot min $^{-1}$, a putative replacement dose (16), to octreotide and insulin did not raise glucose levels to those observed during infusion of octreotide alone. Therefore, the insulin dose was still too high, the glucagon dose was too low, or both.

To clarify this issue, we tested the hypothesis that glucagon supports the postabsorptive plasma glucose concentration in insulin-sufficient patients with type 1 diabetes. Our premise is that demonstrably endogenous insulin-deficient patients with type 1 diabetes infused intravenously with insulin in a dose that maintains stable euglycemia are receiving a biologically optimal insulin replacement dose.

RESEARCH DESIGN AND METHODS

Fourteen patients with type 1 diabetes—ten women and four men with a mean \pm SD age of 33 ± 9 years, weight of 78 ± 17 kg, BMI of 26.7 ± 4.3 kg/m 2 , duration of diabetes of 16 ± 11 years, and A1C of $7.8 \pm 0.8\%$ —gave their




FIG. 1. Mean \pm SE plasma glucose concentrations (A) and plasma insulin (B) and glucagon (C) concentrations in patients with type 1 diabetes infused with insulin in doses that maintain euglycemia from -60 to 240 min with infusions of 1) saline (shaded area) ($n = 14$), 2) octreotide (with growth hormone replacement) (\bullet) ($n = 14$) (glucose was infused if necessary to prevent glucose levels <55 mg/dl [3.0 mmol/l]), 3) octreotide (with growth hormone) plus glucagon 0.5 ng \cdot kg $^{-1}$ \cdot min $^{-1}$ (\circ) ($n = 6$), 4) plus glucagon 1.0 ng \cdot kg $^{-1}$ \cdot min $^{-1}$ (\blacktriangle) ($n = 14$), and 5) plus glucagon 2.0 ng \cdot kg $^{-1}$ \cdot min $^{-1}$ (\triangle) ($n = 10$). Oct., octreotide.

written consent to participate in this study, which was approved by the Washington University Human Research Protection Office and conducted at the Washington University Clinical Research Unit (CRU). Six patients were using a multiple daily injection insulin regimen with a long-acting insulin analog as the basal insulin and a rapid-acting insulin analog as the prandial insulin. Eight patients were using a continuous subcutaneous insulin infusion regimen with a rapid acting insulin analog. With values less than the detection unit of 0.1 ng/ml assigned that value, their mean \pm SD fasting plasma C-peptide concentrations were 0.2 ± 0.2 ng/dl ($<0.1 \pm 0.1$ nmol/l).

Subjects were studied, after an overnight fast, on up to five occasions. They took their last dose of long-acting insulin, if used, the morning of the day prior to study and managed their diabetes with a rapid-acting insulin that day. They reported to the CRU the evening prior to the study. An intravenous line was inserted, and the plasma glucose concentration was held at ~ 100 mg/dl (5.6 mmol/l) overnight with intravenous infusion of variable doses of regular insulin (Novolin R; Novo Nordisk, Princeton, NJ). The following morning, that insulin dose was adjusted, if necessary, and infused from -60 to 240 min. Along with insulin, infused on all study occasions, initially the subjects received intravenous 1) saline ($n = 14$), 2) octreotide 30 ng \cdot kg $^{-1}$ \cdot min $^{-1}$ ($n = 14$), 3) octreotide plus glucagon 1.0 ng \cdot kg $^{-1}$ \cdot min $^{-1}$ ($n = 14$), or 4) octreotide plus glucagon 2.0 ng \cdot kg $^{-1}$ \cdot min $^{-1}$ ($n = 10$), in random sequence, from 0 – 180 min. After 10 studies, the glucagon dose of 2.0 ng \cdot kg $^{-1}$ \cdot min $^{-1}$ was discontinued and six patients received octreotide plus the glucagon dose of 0.5 ng \cdot kg $^{-1}$ \cdot min $^{-1}$ from 0 – 180 min. Growth hormone (Genotropin; Pharmacia and Upjohn, Kalamazoo, MI) was infused, in a dose of 4.7 ng \cdot kg $^{-1}$ \cdot min $^{-1}$ (15), along with octreotide. Glucose was infused intravenously, if necessary to prevent plasma glucose concentrations <55 mg/dl (3.0 mmol/l). Blood samples were drawn through an indwelling line in a hand, with that hand kept in a $\sim 55^\circ\text{C}$ plexiglas box to provide arterialized venous samples, every 15 min for plasma glucose determinations and every 30 min for plasma insulin, glucagon, growth hormone, and epinephrine and norepinephrine measurements from -60 to 240 min. Blood pressure and heart rate were recorded (Dash 3000; GE Healthcare, Fairfield, CT) at 30 -min intervals, and the electrocardiogram was monitored throughout.

Analytical methods. Plasma glucose concentrations were measured with a glucose oxidase method (YSI Glucose Analyzer; Yellow Springs Instruments, Yellow Springs, OH). Plasma insulin, C-peptide, and growth hormone were measured with two-site chemiluminescent assays (Immulite 1000; Siemens,

Los Angeles, CA), plasma glucagon with the Linco radioimmunoassay (Millipore, Temecula, CA), and plasma epinephrine and norepinephrine with a single isotope derivative (radioenzymatic) method (17).

Statistical methods. Except where the SD is specified, data are reported as the mean \pm SE. The time- and condition-related data were analyzed by repeated-measures mixed model ANOVA. P values <0.05 were considered to indicate statistically significant differences.

RESULTS

Plasma glucose concentrations. Plasma glucose concentrations (Fig. 1A) were stable during saline infusion; they were 101 ± 3 mg/dl (5.6 ± 0.2 mmol/l) at 0 min and 98 ± 3 mg/dl (5.4 ± 0.2 mmol/l) at 180 min. Glucose levels declined, from 102 ± 3 mg/dl (5.7 ± 0.2 mmol/l) to 58 ± 1 mg/dl (3.2 ± 0.1 mmol/l) (P vs. saline <0.001) during octreotide infusion. Indeed, to prevent glucose levels <55 mg/dl (3.0 mmol/l), glucose was infused (0.9 ± 0.2 mg \cdot kg $^{-1}$ \cdot min $^{-1}$) in the 3rd h in 10 of the 14 patients. Glucose levels were stable during octreotide plus glucagon 0.5 ng \cdot kg $^{-1}$ \cdot min $^{-1}$ (98 ± 2 mg/dl [5.4 ± 0.1 mmol/l]) to 104 ± 15 mg/dl [5.8 ± 0.9 mmol/l]), rose from 104 ± 3 mg/dl (5.8 ± 0.2 mmol/l) to 143 ± 13 mg/dl (7.9 ± 0.7 mmol/l) (P vs. saline = 0.004) during octreotide plus glucagon 1.0 ng \cdot kg $^{-1}$ \cdot min $^{-1}$ and rose from 97 ± 3 mg/dl (5.4 ± 0.2 mmol/l) to 160 ± 15 mg/dl (8.9 ± 0.8 mmol/l) (P vs. saline <0.001) during octreotide plus glucagon 2.0 ng \cdot kg $^{-1}$ \cdot min $^{-1}$. The insulin infusion rates were 0.12 ± 0.02 , 0.14 ± 0.02 , 0.13 ± 0.02 , 0.16 ± 0.02 , and 0.13 ± 0.02 mU \cdot kg $^{-1}$ \cdot min $^{-1}$, respectively, in the five studies.

Plasma insulin and glucagon concentrations. Plasma insulin concentrations (Fig. 1B) were stable (13 ± 2 $\mu\text{U/ml}$ [78 ± 12 pmol/l] at 0 min and 12 ± 2 $\mu\text{U/ml}$ [72 ± 12 pmol/l] at 180 min) during saline infusion and the other

study conditions. Plasma glucagon concentrations (Fig. 1C) were also stable (50 ± 3 pg/ml [14.4 ± 0.9 pmol/l] at 0 min and 52 ± 4 pg/ml [14.9 ± 1.1 pmol/l] at 180 min) during saline infusion. They decreased to 43 ± 3 pg/ml (12.3 ± 0.9 pmol/l) during infusion of octreotide alone and increased to 51 ± 5 pg/ml (14.6 ± 1.4 pmol/l), 105 ± 10 pg/ml (30.1 ± 2.9 pmol/l), and 166 ± 19 pg/ml (47.7 ± 5.5 pmol/l) during infusion of octreotide plus glucagon in doses of 0.5, 1.0, and 2.0 ng · kg⁻¹ · min⁻¹, respectively.

Other data. Because growth hormone was infused, plasma growth hormone concentrations were stable during all octreotide infusions (e.g., 2.2 ± 0.6 ng/ml [97 ± 26 pmol/l] at 0 min and 1.7 ± 0.2 ng/ml [75 ± 9 pmol/l] at 180 min during octreotide with growth hormone without glucagon replacement). Plasma epinephrine concentrations tended to increase when plasma glucose levels fell during infusion of octreotide with growth hormone without glucagon replacement (data not shown). Heart rate, systolic blood pressure, and diastolic blood pressure were similar under the five study conditions (data not shown).

DISCUSSION

The optimal insulin replacement dose during pancreatic clamps in humans remains to be determined (11,12). There is evidence that insulin doses of 0.14–0.24 mU · kg⁻¹ · min⁻¹, used earlier (7,13–15), are excessive; indeed, an insulin dose of 0.10 mU · kg⁻¹ · min⁻¹ lowered plasma glucose concentration in healthy humans (11). When that dose was infused with octreotide, the addition of a putative replacement dose (16) of glucagon did not recreate the glycemic pattern observed during infusion of octreotide alone (12). The latter finding suggests that the insulin dose was still too high, the glucagon dose was too low, or both. The biologically optimal insulin replacement dose would be one that maintains euglycemia in the absence of endogenous insulin secretion. That condition is met in patients with type 1 diabetes, who have virtually no endogenous insulin secretion, receiving infusions of insulin in doses that maintain stable euglycemia over time.

In such patients receiving biologically optimal insulin replacement in fixed doses, we tested the hypothesis that glucagon supports the postabsorptive plasma glucose concentration by comparing the plasma glucose concentrations during saline infusion with those during infusion of the somatostatin analog octreotide (with growth hormone replacement) alone and with three doses of glucagon. During infusion of saline, plasma glucose levels were stable. During infusion of octreotide without glucagon, plasma glucose levels declined progressively to subnormal levels. Indeed, glucose infusions were required to prevent glucose levels <55 mg/dl (3.0 mol/l) in 10 of the 14 patients. Remarkably, that decline in plasma glucose was prevented, and euglycemia was maintained, by coinfusion of only 0.5 ng · kg⁻¹ · min⁻¹ of glucagon with octreotide. The latter finding provides additional evidence that such low glucagon doses are biologically active (4,16,18,19). Thus, these data indicate that even the lower insulin dose used in our previous pancreatic clamp study (12) was still too high and that the glucagon dose was not too low. They convincingly confirm that glucagon supports the postabsorptive plasma glucose concentration in humans. Indeed, the glycemic response to the lowest dose of glucagon and the hyperglycemic responses to the higher doses suggest a substantial impact of glucagon.

Limitations of the present study include the fact that

although four limbs—saline, octreotide without glucagon, octreotide plus glucagon 1.0 ng · kg⁻¹ · min⁻¹, and octreotide plus glucagon 2.0 ng · kg⁻¹ · min⁻¹—were conducted in random sequence, the octreotide plus glucagon 0.5 ng · kg⁻¹ · min⁻¹ limb was performed toward the end of the study in a subset of the patients. In addition, the glucagon radioimmunoassay that was used cross-reacts with molecular species in addition to 3,500 Da glucagon (4,5). Thus, despite clear effects on plasma glucose concentrations, the decrement in measured plasma glucagon during infusion of octreotide and the increment during infusion of octreotide with the lowest dose of glucagon were small. Finally, other neuroendocrine and cardiovascular effects of octreotide were not studied. However, complete reversal of the glucose lowering effect of octreotide by coinfusion of a relatively low dose of glucagon suggests that these other effects play little, if any, role in the changes in plasma glucose levels.

Interestingly, the insulin infusion doses that maintained euglycemia in these patients were slightly greater than 0.10 mU · kg⁻¹ · min⁻¹, a dose we used earlier (12) and now judge excessive in nondiabetic individuals. However, patients with type 1 diabetes are known to be somewhat resistant to the actions of insulin (20–22).

These data document that glucagon, in concert with insulin, supports the postabsorptive plasma glucose concentrations in humans. Thus, they support the notion that glucagon, in concert with insulin deficiency, plays a role in the pathogenesis of hyperglycemia in diabetes (1–4) and the interest in the development of drugs that block the action or secretion of glucagon for the treatment of diabetes (4).

ACKNOWLEDGMENTS

This study was supported, in part, by National Institutes of Health grants R37-DK27085 and UL1-RR24992 and by a fellowship award from the American Diabetes Association.

P.E.C. has served as a consultant to MannKind Corporation, Marcadia Biotech, and Merck & Company in the past year. B.A.C. has nothing to declare. No other potential conflicts of interest relevant to this article were reported.

B.A.C. and P.E.C. planned the study. B.A.C. performed the study. B.A.C. and P.E.C. wrote the manuscript.

Parts of this study were presented in abstract form at the 70th Scientific Sessions of the American Diabetes Association, Orlando, Florida, 25–29 June 2010.

The authors acknowledge the assistance of the staff of the Washington University Clinical Research Unit and the technical assistance of Krishan Jethi and of the staff of the Core Laboratory for Clinical Studies: Licia Rowe, Laura Karsteter, Nalima Parikh, Tanya Eden, Shirley Frei, Janice Bathon, Paula Wood, and David Gibson. Janet Dedede, assistant to P.E.C., prepared this manuscript.

REFERENCES

1. Unger RH, Orci L. Glucagon and the A cell: physiology and pathophysiology (first of two parts). *N Engl J Med* 1981;304:1518–1524
2. Unger RH, Orci L. Glucagon and the A cell: physiology and pathophysiology (second of two parts). *N Engl J Med* 1981;304:1575–1580
3. Shah P, Vella A, Basu A, Basu R, Schwenk WF, Rizza RA. Lack of suppression of glucagon contributes to postprandial hyperglycemia in subjects with type 2 diabetes mellitus. *J Clin Endocrinol Metab* 2000;85:4053–4059
4. Dunning BE, Gerich JE. The role of alpha-cell dysregulation in fasting and postprandial hyperglycemia in type 2 diabetes and therapeutic implications. *Endocr Rev* 2007;28:253–283

5. Cherrington AD. Control of glucose production in vivo by insulin and glucagon. In *Handbook of Physiology*. Sect. 7, The Endocrine System, volume 2, The Endocrine Pancreas and the Regulation of Metabolism. Jefferson LS, Cherrington AD, Eds. New York, Oxford University Press 2001, p. 759–785
6. Krentz AJ, Boyle PJ, Macdonald LM, Schade DS. Octreotide: a long-acting inhibitor of endogenous hormone secretion for human metabolic investigations. *Metabolism* 1994;43:24–31
7. Liljenquist JE, Mueller GL, Cherrington AD, Keller U, Chiasson J-L, Perry JM, Lacy WW, Rabinowitz D. Evidence for an important role of glucagon in the regulation of hepatic glucose production in normal man. *J Clin Invest* 1977;59:369–374
8. Sherwin RS, Hendler R, DeFronzo R, Wahren J, Felic P. Glucose homeostasis during prolonged suppression of glucagon and insulin secretion by somatostatin. *Proc Natl Acad Sci U S A* 1977;74:348–352
9. Lins PE, Efendić S. Hyperglycemia induced by somatostatin in normal subjects. *Horm Metab Res* 1976;8:497–498
10. Rosenz SG, Clutter WE, Shah SD, Miller JP, Bier DM, Cryer PE. Direct alpha-adrenergic stimulation of hepatic glucose production in human subjects. *Am J Physiol* 1983;245:E616–E626
11. Breckenridge SM, Raju B, Arbelaez AM, Patterson BW, Cooperberg BA, Cryer PE. Basal insulin, glucagon, and growth hormone replacement. *Am J Physiol Endocrinol Metab* 2007;293:E1303–E1310
12. Breckenridge SM, Cooperberg BA, Arbelaez AM, Patterson BW, Cryer PE. Glucagon, in concert with insulin, supports the postabsorptive plasma glucose concentration in humans. *Diabetes* 2007;56:2442–2448
13. Nielsen MF, Nyholm B, Caumo A, Chandramouli V, Schumann WC, Cobelli C, Landau BR, Rizza RA, Schmitz O. Prandial glucose effectiveness and fasting gluconeogenesis in insulin-resistant first-degree relatives of patients with type 2 diabetes. *Diabetes* 2000;49:2135–2141
14. Rosen SG, Clutter WE, Berk MA, Shah SD, Cryer PE. Epinephrine supports the postabsorptive plasma glucose concentration and prevents hypoglycemia when glucagon secretion is deficient in man. *J Clin Invest* 1984;73:405–411
15. Nielsen MF, Basu R, Wise S, Caumo A, Cobelli C, Rizza RA. Normal glucose-induced suppression of glucose production but impaired stimulation of glucose disposal in type 2 diabetes: evidence for a concentration-dependent defect in uptake. *Diabetes* 1998;47:1735–1747
16. Gerich JE, Lorenzi M, Bier DM, Tsalikian E, Schneider V, Karam JH, Forsham PH. Effects of physiologic levels of glucagon and growth hormone on human carbohydrate and lipid metabolism. Studies involving administration of exogenous hormone during suppression of endogenous hormone secretion with somatostatin. *J Clin Invest* 1976;57:875–884
17. Shah SD, Clutter WE, Cryer PE. External and internal standards in the single-isotope derivative (radioenzymatic) measurement of plasma norepinephrine and epinephrine. *J Lab Clin Med* 1985;106:624–629
18. Lins PE, Wajngot A, Adamson U, Vranic M, Efendić S. Minimal increases in glucagon levels enhance glucose production in man with partial hypoinulinemia. *Diabetes* 1983;32:633–636
19. Ward WK, Best JD, Halter JB, Porte D Jr. Prolonged infusion of somatostatin with glucagon replacement increases plasma glucose and glucose turnover in man. *J Clin Endocrinol Metab* 1984;58:449–453
20. DeFronzo RA, Hendler R, Simonson D. Insulin resistance is a prominent feature of insulin-dependent diabetes. *Diabetes* 1982;31:795–801
21. DeFronzo RA, Simonson D, Ferrannini E. Hepatic and peripheral insulin resistance: a common feature of type 2 (non-insulin-dependent) and type 1 (insulin-dependent) diabetes mellitus. *Diabetologia* 1982;23:313–319
22. Pang TT, Narendran P. Addressing insulin resistance in type 1 diabetes. *Diabet Med* 2008;25:1015–1024