

Response to Comment on: Cani et al. (2007) Metabolic Endotoxemia Initiates Obesity and Insulin Resistance: *Diabetes* 56:1761–1772

Rémy Burcelin,¹ Patrice D. Cani,¹ and Jacques Amar²

We thank Saito and Reiko (1), who point out an important fact with regard to the potential sources of circulating lipopolysaccharide (LPS) in metabolic endotoxemia and its correlation with obesity. This relationship has been shown in humans and animal models (2,3). In the light of our recent data (4), which causally demonstrate the role of intestinal bacterial LPS on the onset of metabolic diseases, Saito and Reiko remind us that changes in plasma LPS, driven by periodontitis, may be a causative factor of the relationship between periodontitis and metabolic diseases. We agree that, in addition to intestinal microflora, periodontitis could be an important source of endotoxemia. Interestingly, regarding Saito and Reiko's hypothesis, our group has previously shown an interaction between the soluble CD14 receptor (the main LPS receptor), a CD14 gene polymorphism, and cigarette smoking (5), in which the deleterious role of buccal microbiota and periodontitis

have been well established (6). The next step to pursue these hypotheses will be to demonstrate whether a reduction of metabolic endotoxemia, by means of a dietary intervention or treatment of periodontal disease, impact the subsequent risk for metabolic disease.

REFERENCES

1. Saito T, Reiko H: Comment on: Cani et al. (2007) Metabolic endotoxemia initiates obesity and insulin resistance: *Diabetes* 56:1761–1772. *Diabetes* 456:e20, 2007. DOI: 10.2337/db07-1181
2. Saito T, Shimazaki Y, Sakamoto M: Obesity and periodontitis. *N Engl J Med* 339:482–483, 1998
3. Tomofuji T, Ekni D, Yamanak R, Kusano H, Azuma T, Sanbe T, Tamaki N, Yamamoto T, Watanabe T, Miyauch M, Takat T: Chronic administration of lipopolysaccharide and proteases induces periodontal inflammation and hepatic steatosis in rats. *J Periodontol* 78:1999–2006, 2007
4. Cani PD, Amar J, Iglesias MA, Poggi M, Knauf C, Bastelica D, Neyrinck AM, Fava F, Tuohy KM, Chabo C, Waget A, Delmee E, Cousin B, Sulpice T, Chamontin B, Ferrieres J, Tanti JF, Gibson GR, Casteilla L, Delzenne NM, Alessi MC, Burcelin R: Metabolic endotoxemia initiates obesity and insulin resistance. *Diabetes* 56:1761–1772, 2007
5. Gomes SC, Piccinin FB, Oppermann RV, Susin C, Nonnenmacher CI, Mutters R, Marcantonio RA: Periodontal status in smokers and never-smokers: clinical findings and real-time polymerase chain reaction quantification of putative periodontal pathogens. *J Periodontol* 77:1483–1490, 2006
6. Amar J, Ruidavets JB, Bal dit Sollier C, Bongard V, Boccalon H, Chamontin B, Drouet L, Ferrieres J: CD14 C(-260)T gene polymorphism, circulating soluble CD14 levels and arteriosclerosis. *J Hypertens* 8:1523–1528, 2004

From ¹INSERM U858, Institute of Molecular Medicine I²MR, Toulouse, France; and ²INSERM U558, Toulouse, France

Address correspondence and reprint requests to Rémy Burcelin, I²MR U858, IFR 31, Hôpital Rangueil, BP 84225, Toulouse 31432 Cedex 4, France. E-mail: burcelin@toulouse.inserm.fr.

Received and accepted for publication 5 September 2007.

DOI: 10.2337/db07-1262

© 2007 by the American Diabetes Association.