

Urologic Complications of Diabetes

JEANETTE S. BROWN, MD¹
 HUNTER WESSELLS, MD²
 MICHAEL B. CHANCELLOR, MD³
 STUART S. HOWARDS, MD⁴
 WALTER E. STAMM, MD²

ANN E. STAPLETON, MD²
 WILLIAM D. STEERS, MD⁴
 STEPHEN K. VAN DEN EEDEN, PHD⁵
 KEVIN T. MCVARY, MD, FACS⁶

Diabetes and urologic diseases are very common health problems that markedly increase in prevalence and incidence with advancing age (1–3). Diabetes is associated with an earlier onset and increased severity of urologic diseases, resulting in costly and debilitating urologic complications. Urologic complications, including bladder dysfunction, sexual and erectile dysfunction, as well as urinary tract infections (UTIs), have a profound effect on the quality of life of men and women with diabetes. This review presents a comprehensive overview of the current understanding of clinical and basic research on urologic complications of diabetes and recommendations for future directions for research and clinical care.

BLADDER DYSFUNCTION—Over 50% of men and women with diabetes have bladder dysfunction (4,5). Current understanding of bladder dysfunction reflects a progressive condition encompassing a broad spectrum of lower urinary tract symptoms including urinary urgency, frequency, nocturia, and incontinence. Previously, the dysfunction has been classically described as diminished

bladder sensation, poor contractility, and increased postvoid residual urine, termed bladder cystopathy (6). However, bladder cystopathy most likely represents end-stage bladder failure with symptoms of infrequent voiding, difficulty initiating voiding, and postvoid fullness and is relatively uncommon.

A number of clinical studies in men and women with diabetes have reported bladder instability or hypersensitivity as the most frequent finding, ranging from 39–61% of subjects (5,7). Diminished bladder contractility or sensation has been found less often (5), and an acontractile bladder appears to be quite uncommon.

Bladder dysfunction in women

In women, urinary incontinence is estimated to affect nearly 50% of middle aged and older women, leading to significant distress, limitations in daily functioning, and poorer quality of life (8,9). Diabetes has been identified as an important independent risk factor for incontinence in several large observational studies, including the Nurses' Health Study, and is associated with 30–100% increased risk (9–12). This suggests that interventions

that prevent or delay onset of diabetes may also prevent urinary incontinence.

The Diabetes Prevention Program (DPP) randomized trial demonstrated that an intensive lifestyle intervention involving weight loss and exercise reduced the incidence of diabetes among women with impaired glucose tolerance (IGT) (13). Prevalence of weekly stress incontinence was also substantially decreased by the DPP intensive lifestyle intervention. Metformin had no effect on incontinence (14). Weight loss, as has been shown in other studies among women without IGT (15), was the most likely mechanism for the effect of the lifestyle intervention on stress incontinence. Differential diabetes incidence among the treatment groups appeared to play almost no mediating role. Importantly, reducing incontinence may be a powerful motivator for women with IGT to choose lifestyle modification to prevent diabetes. The DPP-Outcomes Study, a follow-up study of the DPP cohort, will examine longer-term effects of the intervention on incontinence.

Obesity is a significant and important risk factor for incontinence and type 2 diabetes. The Action for Health in Diabetes (Look AHEAD) is an ongoing randomized trial examining the effects of interventions designed to produce sustained weight loss on cardiovascular events in 5,145 obese and overweight participants ages 45–75 with type 2 diabetes. An ancillary study will evaluate incontinence annually throughout the trial and postintervention follow-up. The Look AHEAD study offers a unique opportunity to determine whether intentional weight loss also decreases incidence or severity of incontinence and the degree to which this may be mediated by effects on diabetes severity. Additionally, oversampling of racial/ethnic groups at elevated risk of diabetes will help to elucidate differential incontinence patterns observed in earlier studies (11).

Recent large observational studies have identified urge incontinence, an involuntary loss of urine with a feeling of urgency, as increased among women with diabetes, while there was no increased risk for stress incontinence, involuntary

From the ¹University of California, San Francisco, California; the ²University of Washington, Seattle, Washington; the ³University of Pittsburgh, Pittsburgh, Pennsylvania; the ⁴University of Virginia, Charlottesville, Virginia; the ⁵Division of Research, Kaiser Permanente, Oakland, California; and ⁶Northwestern University, Chicago, Illinois.

Address correspondence and reprint requests to Jeanette S. Brown, MD, University of California, San Francisco, 1635 Divisadero St., Suite 600, San Francisco, CA 94115. E-mail: brownj@obgyn.ucsf.edu.

Received for publication 22 March 2004 and accepted in revised form 21 September 2004.

H.W. holds stock in and has received honoraria from Pfizer. W.E.S. has received honoraria or consulting fees from Corixa, Proctor/Gamble, Bayer, GlaxoSmithKline, ActivBiotics, Medimmune, Chiron, Aventis/Pasteur, Preventis, and Osel and has received grant or research support from ActivBiotics, Proctor/Gamble, Medimmune, and Bayer. K.T.M. is a member on an advisory panel for and has received consulting fees from Pfizer, Eli Lilly, ICOS/Lilly, Merck, and GlaxoSmithKline.

Abbreviations: AGE, advanced glycation end product; ASB, asymptomatic bacteriuria; BPH, benign prostatic hyperplasia; DPP, Diabetes Prevention Program; ED, erectile dysfunction; EDIC, Epidemiology of Diabetes Interventions and Complications Study; ET-1, endothelin-1; IGT, impaired glucose tolerance; IL, interleukin; LUTS, lower urinary tract symptoms; NGF, nerve growth factor; NOS, nitric oxide synthase; PKC, protein kinase C; PMN, polymorphonuclear leukocyte; STZ, streptozotocin.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

© 2005 by the American Diabetes Association.

loss of urine with physical activity (16). Diabetic women treated with insulin are at considerably higher risk of urge incontinence than those treated with oral medications or diet (16). Other aspects of diabetes severity, including glycemic control and microvascular complications resulting in damage to innervation of the bladder, have been suggested as possible mechanisms for increasing incontinence (6). However, most previous studies of incontinence in women with diabetes were conducted in patients attending specialty clinics, did not differentiate between type 1 and type 2 diabetes, and did not adjust for other established incontinence risk factors, including age, parity, and obesity. As a result, little is known about how diabetes may contribute to the incidence or severity of urinary incontinence.

Common treatments for urinary incontinence in women include conservative management (e.g., pelvic muscle training or bladder training), medications, and surgery (17). Because the safety and efficacy of these treatments have not been determined in women with diabetes, conservative management should be the first line of treatment in this population.

Bladder dysfunction in men

In men, lower urinary tract symptoms (LUTS) are common, age-related complaints that are often attributed to benign prostatic hyperplasia (BPH). LUTS and BPH increase rapidly with age starting at about age 50 years (18). Straining, intermittency, postvoid dribbling, and weak stream may signify urethral obstruction from BPH. However, among men with diabetes, similar symptoms may also result from bladder dysfunction due to denervation and poor detrusor contractility. Other complex associations of LUTS and BPH among diabetic men include symptoms of urgency, frequency, and nocturia that may occur from detrusor overactivity, resulting from BPH, and/or microvascular complications associated with diabetes, increasing hyperactivity of the detrusor. The failure to differentiate LUTS from BPH in studies of diabetic men has contributed to the confusing evidence now seen in the literature (19). Other important factors that contribute to this confusion are concerns such as selected or different study populations, low response rates, inadequate control of potential confounders, and small sample sizes in many studies.

The effect of diabetes on the development or presence of LUTS and BPH remains controversial. As noted, this is due in part to a lack of uniformity in the definition of the outcome. Recent evidence suggests that LUTS may occur more frequently among men with diabetes, with an estimated 25% to nearly twofold increased risk of LUTS in men with diabetes (20–23). Additionally, among men with BPH, diabetes is associated with more LUTS symptoms compared with nondiabetic men (23). BPH may or may not be associated with diabetes. Early studies suggested that diabetes increases prostate size consistent with BPH (24). However, studies using surgery for BPH as the outcome have reported conflicting results, with some reporting that diabetic men were at higher risk (25) and others reporting high blood glucose levels to be associated with a lower risk (26). Two more recent cohort studies found conflicting results. In a study of BPH with LUTS or surgery, a nonsignificant increase was reported (27), whereas a large cohort study using cross-sectional data reported a decreased risk of BPH and increased risk of LUTS (22). In a prospective study of BPH progression, diabetes was found to increase the risk of LUTS but was not associated with an increase in prostate volume (28).

While there is clinical overlap between the presence of BPH and LUTS, they can be manifestations of different pathophysiological pathways mediated through hormonal, environmental, genetic, neuropathic, and (micro)vascular influences, particularly in the diabetic patient. A growing body of experimental evidence indicates that diabetes and obstruction affect different populations of visceral afferents supplying the bladder. In both conditions early alterations in sodium and potassium channels occur similar to neuropathic models. These changes trigger altered excitability, leading to detrusor overactivity and urinary frequency. With time, impaired contractility due to a myopathy can lead to incomplete emptying. Thus, a combination of several factors with differing time courses lead to LUTS and known urodynamic findings, making discerning an etiology and distinguishing classic diabetic cystopathy from neural plasticity accompanying obstruction due to BPH problematic.

Many of the difficulties in reconciling the literature are the broad variation in

definitions used to define BPH and/or LUTS; often they are used interchangeably. Future studies need to take as much care as possible to delineate each condition and to examine each of the underlying pathways before we are able to better understand how diabetes affects BPH and LUTS. Whether outcomes for treatment of LUTS and/or BPH differ between diabetic and nondiabetic men has not been adequately examined in randomized trials.

Pathophysiology

The biology of diabetes-associated bladder complications can be due to an alteration in the detrusor smooth muscle, neuronal dysfunction, and urothelial dysfunction. The experimental model most often used to assess bladder complications is the streptozotocin (STZ) rat model. Pharmacological studies of smooth muscle dysfunction on isolated diabetic rat bladder strips have generated a considerable amount of controversy. Bladder smooth muscle contraction is mediated by acetylcholine released by the pelvic nerve acting on muscarinic receptors. Responsiveness of diabetic bladder strips to externally applied muscarinic agonists has been reported as increased, decreased, or described as little or no change in responsiveness (29,30). Even when an increased responsiveness is found, the mechanism is unclear. An increase in muscarinic receptor density has been found at both 2 and 8 weeks of STZ-induced diabetes (31). A recent study found an increase in β_1 -receptor-mediated relaxation response in isolated detrusor smooth muscle strips from 8- to 10-week STZ-induced diabetic rats (32).

One diabetes-related change that most experts agree on is an increased responsiveness of isolated rat bladder strips to electrical field stimulation (33). Explanations for this change include an increased diabetes-related alteration in membrane lipid composition or increased neurotransmitter release. It has been suggested that changes are related to increased calcium channel activity or enhanced calcium sensitivity (33). Most recently, it has been reported that decreased function was more notable in strips from diabetic rats with enlarged bladders (34). An increased depolarization of myocytes in STZ-induced diabetic rat bladder strips to the application of acetylcholine has been found, indicating en-

hanced muscarinic sensitivity in the diabetic bladder (35).

Neuronal dysfunction may reflect the deficiency of axonal transport of nerve growth factor (NGF) and play an important role in inducing diabetic neuropathy. It has been recently reported in STZ-induced diabetic rats that decreased NGF levels in the bladder and lumbosacral dorsal root ganglia are associated with bladder dysfunction, suggesting the feasibility of NGF gene therapy for the treatment of diabetic cystopathy (36).

The bladder urothelium is important for the regulation of permeability, transport, and endocytosis across the bladder wall. It has become increasingly clear that the urothelium is not only a passive barrier against urea and ion diffusion, but that it can also function as a sensor, controlling bladder function and dysfunction. The effects of diabetes on urothelial receptors and urothelial signaling mechanisms have not been extensively studied. It has been well established that in the STZ-induced diabetic rat model, there are progressive increases in total bladder tissue with hypertrophy of the bladder wall and dilation of the bladder. Both smooth muscle and urothelium (percentage of total tissue) have been shown to increase significantly in a time-dependent manner. Thus, it has been found that the epithelium from STZ-induced diabetic rat bladders was at least twice as thick and heavy than that from controls (37).

SEXUAL DYSFUNCTION—Erectile dysfunction, reduced libido, orgasmic dysfunction, and retrograde ejaculation are established complications found with variable prevalence in men with diabetes. Reduced desire, decreased arousal, and painful intercourse are increased sexual problems in women with diabetes, with a reported prevalence of 18–42% (38,39). Whether the same pathophysiological mechanisms are involved in male and female sexual dysfunction and in patients with type 1 and type 2 diabetes has not been determined.

Male sexual dysfunction

Erectile dysfunction (ED) occurs in a substantial number of men with diabetes, with prevalence estimates ranging from 20 to 71% (2,40,41). ED in diabetic men significantly impacts their quality of life (41,42). The prevalence of disorders of

desire, orgasm, and ejaculation has not been determined.

A number of longitudinal studies have estimated the disease risk of ED in men with diabetes (2,43,44). The 10-year incidence of ED was 25% in a population-based cohort of individuals with type 1 diabetes followed in the Wisconsin Epidemiological Study of Diabetic Retinopathy (43). The Massachusetts Male Aging Study (44) found that the crude incidence of ED in the men with diabetes was 50.7 per 1,000 population-years versus 24.8 in those without diabetes. This figure was similar to results in a convenience sample of 1,010 men with diabetes (68 per 1,000 population-years) (2).

Risk factors associated with an increased risk of ED include hypertension, lipid disorders, coronary heart disease, LUTS, older age, higher BMI, and cigarette smoking. In men with diabetes, the relative risk for ED increases with poor glycemic control, duration of diabetes, and the number of other nonurologic complications of diabetes (i.e., retinopathy, nephropathy, and limb loss) (40,45–47). Studies that focus on the impact of race/ethnicity, modifiable risk factors (e.g., weight loss), and comorbidities on ED are still needed. Recently investigators reported a decrease in relative risk of ED with increased physical activity (48). Whether this reduced risk applies to diabetic men is not known.

Randomized trials demonstrated the success of pharmacological treatment of ED. Phosphodiesterase-5 inhibition leads to significant improvements in function in 50–70% of type 1 and type 2 diabetic patients with ED and controlled hyperglycemia (49,50). However, the efficacy is reduced compared with nondiabetic populations (51). Nonresponders to oral treatment benefit from intracavernosal injections of prostaglandin-E1 and related agents, as >80% of men with diabetes develop adequate penile rigidity (52).

As a result of diabetes, progressive replacement of cavernosal smooth muscle by fibrosis may lead to complete erectile failure (53). Effective surgical interventions in such cases are limited to penile implants. The risk of periprosthetic infection after implantation in diabetic men ranges from 3.2–15% (54). The relationship between glycemic control and risk of infection was not confirmed in a prospective trial (54), although the study may have been underpowered.

The reduced efficacy of all treatments makes diabetic men a population for whom novel prevention and early intervention strategies are needed. Pharmacological approaches targeting smooth muscle cells and neuronal tissue are most relevant (55,56). Examples include administration of vascular endothelial growth factor (57) and immunophilin ligands (58) in animal models of ED. Approaches such as gene therapy that deal with the impaired delivery of nitric oxide (NO) to cavernosal smooth muscle cells may be most important for men with diabetes who develop severe ED (59,60).

Pathophysiology. NO from nonadrenergic noncholinergic neurons and the endothelium is required to induce smooth muscle relaxation in the corpus cavernosum, resulting in sinusoidal filling and penile erection (61). When compared with control subjects, diabetic men with ED have impaired neurogenic and endothelium-mediated relaxation of smooth muscle, increased accumulation of advanced glycation end products (AGEs), and altered expression of arginase, a competitor with NO synthase (NOS) for its substrate L-arginine (62,63).

Human studies. Neuropathy and arterial disease have all been implicated in the pathogenesis of diabetic ED (64,65). Complications of diabetes that are associated with an increased risk of ED include peripheral or autonomic neuropathy, nephropathy, and retinopathy (2).

Nocturnal penile tumescence testing has been used to infer parasympathetic damage of the penis in diabetic men without vascular disease (65). Neurodiagnostic tests have confirmed that diabetic men with ED have altered conduction of both unmyelinated lower-extremity sensory fibers (66,67) and myelinated pudendal somatosensory fibers. However, diagnostic tests that measure autonomic nerve function and the integrity of cavernosal innervation require development and validation.

Adequate arterial inflow is necessary for erection, and cavernosal arterial morphology, flow (68), and diameter differ between diabetic and nondiabetic populations with ED. Overall, the preponderance of human studies suggests that neuropathy is the most important risk factor associated with ED in diabetes, leading to impaired NO signaling. Whether microangiopathy directly impairs erectile function or merely mediates autonomic neuropathy is unknown.

Animal studies. The proposed mechanisms for ED in experimental models of diabetes include central and/or autonomic neuropathy, smooth muscle cellular dysfunction, endothelial dysfunction, and tissue remodeling (69–72).

Downregulation of the neuronal isoform of NOS has been demonstrated in rats with both type 1 and 2 diabetes (73). A recent study suggests that penile nitrergic nerves undergo a selective degenerative process in two phases (74). In the first phase, nNOS protein is depleted in the axons with resultant impairment of erectile function. In the second phase, a synergistic action of endogenous NO and accumulated AGEs in the autonomic ganglia results in selective nitrergic apoptosis. Mechanisms responsible for this neural degeneration and other vascular complications of diabetes may include increased polyol pathway flux, intracellular AGE accumulation, activation of protein kinase C (PKC), and increased flux through the hexosamine pathway (75).

Studies in diabetic animals have identified abnormalities in the central nervous system, spinal cord, autonomic ganglia and nerves, and in the penis. The result of neural degeneration may be to induce apoptosis, cell turnover, and tissue remodeling in the cavernosal spaces. Extensive smooth muscle and endothelial degradation in the corpora cavernosa of diabetic penis has been noted in a model of type 1 diabetes (69). This degradation was accompanied by profound ED, significantly decreased expression of the morphogenic protein sonic hedgehog (Shh) in the smooth muscle of the corpora cavernosa, and increased Shh mRNA expression in the nerves, corpora, and urethra (76). Taken together, these results suggest that in diabetes, ED is the result of a neuro-pathic process leading to denervation of the major pelvic ganglia and cavernous nerves, which in turn results in extensive but truncated tissue remodeling.

Vasoconstrictor signaling pathways are important physiological modulators of the erectile process, and perturbations in these pathways may contribute to the pathophysiology of ED (77). In experimental diabetes, penile smooth muscle has augmented force responses to two important vasoconstrictors, endothelin-1 (ET-1) and the α -adrenergic agonist phenylephrine (78). Investigators hypothesized that a Ca^{2+} -sensitization pathway regulates smooth muscle con-

tractile tone, and ρ -kinase and PKC have been implicated as proteins that promote penile flaccidity (79,80). Diabetic Zucker rats with impaired erections demonstrated increased phenylephrine and ET-1 constrictive activity along with elevated expression of PKC isozymes and RhoA and ρ -kinase (C. Wingard, unpublished data). Other studies have shown increased sensitivity to ET-1 and increased ρ -kinase expression using diabetic rabbits (78). The conclusion drawn from these studies is that in diabetes, elements of a Ca^{2+} -sensitization mechanism participate in impairment of smooth muscle relaxation in the penis.

Delineation of the interaction among PKC, RhoA/ ρ -kinase, and NO/cGMP signaling, the relative contribution of each to neurally generated penile erection, and their potential alterations in diabetes requires further study.

Female sexual dysfunction

Female sexual dysfunction is associated with biological, psychological, and social determinants (3). It includes disorders of desire/libido, arousal, inhibited orgasm, and sexual pain (81). Prevalence increases with age, cardiovascular disease, diabetes, cancer, hysterectomy, and neurological conditions (38,82).

Studies of female sexual dysfunction in women with diabetes are limited. Prevalence of dysfunction has been reported in 18–27% of women with type 1 diabetes and in 42% of women with type 2 diabetes (83,84). Women with type 1 diabetes have a nearly twofold greater prevalence of sexual dysfunction than women without diabetes (83). Women with more diabetic complications have more sexual dysfunction. Psychological factors including depression, adaptation to diabetes, and quality of partner relationship are likely to contribute to the sexual dysfunction of women with diabetes and require quantification in future studies. As female sexual complaints may be ignored by caregivers and patients alike, clinicians should be encouraged to query patients about these issues.

In vitro and in vivo approaches may provide insights into disease mechanisms (85–87). Hormonal determinants and possible therapies in women remain controversial (88). The safety and efficacy of testosterone (89) or sildenafil citrate (90) have not been studied in women with diabetes.

ASYMPTOMATIC BACTERIURIA AND SYMPTOMATIC UTIS

— Over the years, evidence from many epidemiological studies have suggested that asymptomatic bacteriuria (ASB) and symptomatic UTIs occur more commonly in women with diabetes than in those without diabetes (91). Most of these studies, however, were not prospective cohort designs and are thus subject to multiple biases characteristically associated with case-control, retrospective, or cross-sectional studies. Further, the majority of the data has been collected in patients with type 2 diabetes and in women; therefore, data regarding these relationships in type 1 diabetes and in men are less available. Many of these studies also failed to adjust for relevant confounding variables or comorbidities, and many were done before widespread use of improved methods of glycemic control.

Recent studies have focused on the relationship of ASB to diabetes (92–94). In women without diabetes, ASB is relatively uncommon and increases risk of UTI but does not lead to serious sequelae (95). Diabetic women have a two- to threefold higher prevalence of ASB and are at risk for developing more serious consequences (92,93). For women with diabetes and ASB, those with type 2 diabetes have an increased risk for development of a symptomatic UTI (96), and women with type 1 diabetes are at an increased risk for pyelonephritis and subsequent impairment of renal function (94).

Assuming that ASB is more common and that the consequences of ASB may be deleterious among women with diabetes, the question as to whether to attempt to eradicate ASB in women with diabetes is of considerable relevance. In a randomized controlled trial of type 1 and type 2 diabetic women with ASB, women were randomized to treatment with antimicrobials or no treatment for episodes of ASB >3 years (97). Importantly, the study demonstrated that screening and treatment of episodes of ASB had no impact on overall occurrence of symptomatic UTI or hospitalizations.

Recent studies have demonstrated an increased risk of symptomatic UTIs primarily in women with type 2 diabetes. In postmenopausal women, type 2 diabetes has been associated with a twofold increased risk for UTIs (98–100). Additionally, women on treatment with oral

hypoglycemics or insulin have a three- to fourfold higher risk of UTI, possibly indicating an association with severity of diabetes (98).

Knowledge of risk factors for UTIs in diabetic women is important to identify women in need of therapy to prevent serious complications. Currently, risk factors for UTI in diabetic women are not well defined and may differ by type of diabetes. ASB increases risk for UTI among women with type 2 diabetes (96). However, as previously stated, screening and treatment of ASB have not shown benefit (97). Among women with type 1 diabetes, sexual activity has been identified as the most important risk factor for the development of UTI, similar to women without diabetes (96,101). Continuous or postcoital prophylaxis with low-dose antimicrobial agents as well as intermittent self-treatment with antimicrobials are recommended strategies to prevent recurrent UTIs in women without diabetes (102). However, trials have not been performed in women with diabetes. Although previously suggested as possible risk factors, duration of diabetes or elevated HbA_{1c} levels have not been shown to increase the risk of UTI in recent studies (96,98).

Among the most striking effects of diabetes with regard to UTI appears to be the associated risk of infections progressing to complications, severe outcomes, or being due to unusual infecting organisms. For example, emphysematous cystitis and pyelonephritis, relatively rare infections, occur almost exclusively in diabetic patients (91). Other clinical manifestations that are unique or strongly associated with diabetes include abscess formation and renal papillary necrosis. Bacteremia secondary to UTI and pyelonephritis may also be more common in patients with diabetes. Also of importance is the fact that many diabetic patients are infected with non-*Escherichia coli* species, in particular *Klebsiella*, other gram-negative rods, enterococci, and group B streptococci (96,103). Additionally, urinary infections and ASB with *Candida albicans* occur commonly in diabetic women but infrequently in other women (91). Thus, although not documented in prospective studies, there appears to be a clear association of unusual infecting organisms and unusually severe clinical manifestations in patients with both type 1 and type 2 diabetes, primarily women. These obser-

vations should be reevaluated in the light of recent concepts of the pathogenesis of UTI (104).

Pathogenesis. Clinical observations have suggested various pathogenic mechanisms of UTI in patients without diabetes, generating extensive laboratory studies of concepts such as the importance of pathogen virulence in the urinary tract. Several recent studies have applied principles of uropathogenesis developed for uncomplicated UTI to understanding the pathogenesis of UTI among patients with diabetes.

The development of UTI in women is preceded by colonization of the vaginal and periurethral epithelium by the infecting organism (105). Ascension to the bladder and/or bacteremia may then ensue. *E. coli* causes the overwhelming majority of UTIs among most groups of patients without diabetes and is also the most common cause of UTI among men or women with diabetes in community-acquired or nosocomial UTI. However, *E. coli* among diabetic women causes a significantly lower proportion of UTIs than among nondiabetic men or women (91).

Bacterial pathogenesis in the urinary tract has been extensively investigated for *E. coli* infections of patients without diabetes (106). Uropathogenic *E. coli* are specialized for success in the urinary tract, elaborating virulence determinants such as adhesins (type 1, P and S fimbriae, and afimbrial adhesin), which bind to specific molecules in the uroepithelium, such as glycosphingolipids and uroplakins (106). Additional virulence factors of *E. coli* associated with UTI and other extraintestinal infections include hemolysin, lipopolysaccharide, cytotoxic necrotizing factor, siderophores, and others (106).

Several older studies of *E. coli* virulence determinants among organisms cultured from patients with diabetes and other medically compromising conditions demonstrated a decreased prevalence of these urovirulence factors (107). However, a recent study examined the prevalence of virulence factors of *E. coli* isolated from episodes of ASB from diabetic women, correlated these with the patients' clinical characteristics, and then compared the results with those of historical control subjects (108). In contrast with the prior finding, the prevalence of virulence factors among the ASB isolates from women with diabetes was nearly identical to findings among ASB isolates

from patients with no medically compromising condition, except for a reduced frequency of specific O:K:H serotypes associated with UTI (108). Three virulence determinants were associated with a decline in renal function (108).

Another recent study examined the ability of three representative clinical isolates of uropathogenic *E. coli* to adhere to uroepithelial cells collected from urine of women with and without diabetes (109). Uropathogenic *E. coli* expressing type 1 fimbriae but no P fimbriae were twice as adherent to cells from women with diabetes as compared with cells collected from the women without diabetes (109). No differences in adherence were seen using the *E. coli* strain expressing no known uropathogenic adhesins or for the isolate expressing only P fimbriae (109). These data suggest that uroepithelial cells from women with diabetes may have intrinsic differences in nature and/or amount and/or affinity of binding receptors for type 1 fimbriae, such as uroplakins.

Because of known alterations in polymorphonuclear leukocyte (PMN) function in high-glucose states, it has been suggested that leukocyte dysfunction occurs in patients with diabetes and increases risk of UTI. However, it has been difficult to correlate in vitro findings with clinical effects. A recent, well-designed study of diabetic women found no differences in five standard measures of PMN function among PMNs isolated from women with diabetes, with or without ASB, as compared with cells isolated from control subjects without diabetes (110). Adaptive immune functions appear to be intact in diabetes, based on findings of normal antibody production and responses to vaccines.

Abnormalities in innate immune functions may also occur in diabetes, but these have not been systematically investigated. Some studies have shown increased baseline levels of cytokines (e.g., interleukin [IL]-6 and IL-8) in serum from patients with diabetes (111). Results of studies of stimulating immune cells from diabetic patients have been conflicting (112). One recent study found that monocytes from women with type 1 diabetes produced lower amounts of proinflammatory cytokines upon stimulation with lipopolysaccharide when compared with cells collected from women without diabetes (113). Women with diabetes who developed bacteriuria also produced

lower urinary IL-6 concentrations, as compared with specimens from bacteriuric control subjects without diabetes (113).

Antimicrobial peptides, an important component of host defense in infections occurring at mucosal surface, such as UTI, have been scarcely investigated in diabetes. There is evidence that AGEs, found in serum of diabetic subjects, may inhibit the function of components of the innate immune system, such as lactoferrin and lysozyme (114). It is not clear what role lactoferrin and/or lysozyme might play in the defense against UTI. On the other hand, human PMNs produce α -defensins, and β -defensins are synthesized at epithelial surfaces including the upper respiratory tract, nasal mucosa, tongue, kidney, pancreas, colon, female reproductive tract, and conjunctiva (115). β -Defensins have also been detected in urine (116). Additional studies of potential adaptive and innate immune dysfunction in diabetes and its potential effect on the risk of UTI are warranted.

CONCLUSIONS — Although urologic complications are common and major health problems in men and women with diabetes, data to define expected prevalence, incidence, and risk factors as well as interventions to reduce the risk of developing these complications are limited. Intensive glycemic control delays the onset and progression of microvascular complications of diabetes in both type 1 and type 2 diabetes (117,118). If microvascular complications also damage the vascular and neurologic innervation of the urethral sphincter, bladder, and corpora cavernosa, then intensive glycemic control may prevent or improve severity of urologic complications.

Data on lower urinary tract symptoms and incontinence, sexual and erectile dysfunction, as well as UTIs, are currently being collected in the Epidemiology of Diabetes Interventions and Complications Study (EDIC), a prospective cohort of participants from the Diabetes Control and Complications Trial (DCCT). The EDIC offers a unique opportunity to further understand the epidemiology of urologic complications among a well-characterized cohort of men and women with type 1 diabetes. Most importantly, analysis of EDIC urologic complications data will determine whether intensive glycemic control during the DCCT that re-

duced the development of retinopathy, nephropathy, and neuropathy in participants of EDIC also reduce the risk of developing urologic complications. Similar studies would be useful among men and women with type 2 diabetes.

Clinical outcomes of common treatments for urologic diseases among diabetic men and women have not been critically examined. Large randomized controlled trials are needed to assess the efficacy and safety of conservative, pharmacologic, and surgical treatments of urologic complications.

Additionally, the pathophysiology and possible mechanisms by which diabetes might promote urologic complications are not clear. The multicellular composition of bladder and erectile tissue make it unlikely that a single mechanism underlies both voiding and sexual dysfunction in diabetes. The complex impact of hyperglycemia on nerve, epithelium (or endothelium), and mesenchymal components of genitourinary organs points toward multiple future therapeutic classes that may be relevant during specific phases of diabetes.

In summary, new research initiatives are needed to further understand the dimensions, including possible basic disease mechanisms, and burden of urologic complications as a health outcome in men and women with diabetes. The paucity of knowledge has been a barrier to developing the best methods of prevention and treatment of urologic complications.

References

- Harris MI, Flegal KM, Cowie CC, Eberhardt MS, Goldstein DE, Little RR, Wiedmeyer HM, Byrd-Holt DD: Prevalence of diabetes, impaired fasting glucose, and impaired glucose tolerance in U.S. adults: the Third National Health and Nutrition Examination Survey, 1988–1994. *Diabetes Care* 21:518–524, 1998
- Fedeles D, Bortolotti A, Coscelli C, Santusanio F, Chatenoud L, Colli E, Lavezari M, Landoni M, Parazzini F: Erectile dysfunction in type 1 and type 2 diabetics in Italy: on behalf of Gruppo Italiano Studio Deficit Erettile nei Diabetici. *Int J Epidemiol* 29:524–531, 2000
- Laumann EO, Paik A, Rosen RC: Sexual dysfunction in the United States: prevalence and predictors. *JAMA* 281:537–544, 1999
- Goldman HB, Appell RA: Voiding dysfunction in women with diabetes mellitus. *Int Urogynecol J Pelvic Floor Dysfunct* 10:130–133, 1999
- Kaplan SA, Te AE, Blaivas JG: Urodynamic findings in patients with diabetic cystopathy. *J Urol* 153:342–344, 1995
- Frimodt-Moller C: Diabetic cystopathy: epidemiology and related disorders. *Ann Intern Med* 92:318–321, 1980
- Menendez V, Cofan F, Talbot-Wright R, Ricart MJ, Gutierrez R, Carretero P: Urodynamic evaluation in simultaneous insulin-dependent diabetes mellitus and end stage renal disease (see comments). *J Urol* 155:2001–2004, 1996
- Perry S, Shaw C, Assassa P, Dallosso H, Williams K, Brittain KR, Mensah F, Smith N, Clarke M, Jagger C, Mayne C, Castleden CM, Jones J, McGrother C: An epidemiological study to establish the prevalence of urinary symptoms and felt need in the community: the Leicestershire MRC Incontinence Study: Leicestershire MRC Incontinence Study Team. *J Public Health Med* 22:427–434, 2000
- Wetle T, Scherr P, Branch LG, Resnick NM, Harris T, Evans D, Taylor JO: Difficulty with holding urine among older persons in a geographically defined community: prevalence and correlates. *J Am Geriatr Soc* 43:349–355, 1995
- Brown J, Grady D, Ouslander J, Herzog A, Varner R, Posner S: Prevalence of urinary incontinence and associated risk factors in postmenopausal women: Heart & Estrogen/Progestin Replacement Study (HERS) Research Group. *Obstet Gynecol* 94:66–70, 1999
- Sampsel CM, Harlow SD, Skurnick J, Brubaker L, Bondarenko I: Urinary incontinence predictors and life impact in ethnically diverse perimenopausal women. *Obstet Gynecol* 100:1230–1238, 2002
- Lifford K, Curhan G, Grodstein F: Type 2 diabetes and risk of developing urinary incontinence in the Nurses' Health Study (Abstract). In *Urologic Complications of Diabetes*. Bethesda, MD, National Institutes of Health, 2003, p. 31
- Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM: Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 346:393–403, 2002
- Brown JS, Barrett-Connor E, Nyberg LM, JW. K, TJ O, Ma Y, Vittinghoff E: Incontinence in women with impaired glucose tolerance: results of the Diabetes Prevention Program (Abstract). *J Urol* 171:325–326, 2004
- Subak LL, Johnson C, Whitcomb E, Boban D, Saxton J, Brown JS: Does weight loss improve incontinence in moderately obese women? *Int J Urogyn* 13:40–43, 2002

16. Jackson RA, Vittinghoff E, Kanaya AM, Resnick HE, Kritchevsky S, Miles T, Simonsick E, Brown JS, for the Health, Aging and Body Composition (Health ABC) Research Group: Urinary incontinence in elderly women: finding from the Health, Aging, and Body Composition Study. *Obstet Gynecol* 104:301–307, 2004
17. Holroyd-Leduc JM, Straus SE: Management of urinary incontinence in women: scientific review. *JAMA* 291:986–995, 2004
18. Boyle P, Robertson C, Mazzetta C, Keech M, Hobbs FD, Fourcade R, Kiemeny L, Lee C: The prevalence of lower urinary tract symptoms in men and women in four centres: the UrEpik study. *BJU Int* 92:409–414, 2003
19. Blaivas JG: Cause and effect of lower urinary tract symptoms (Editorial). *J Urol* 168: 1464, 2002
20. Klein BE, Klein R, Lee KE, Bruskewitz RC: Correlates of urinary symptom scores in men. *Am J Public Health* 89:1745–1748, 1999
21. Joseph MA, Harlow SD, Wei JT, Sarma AV, Dunn RL, Taylor JM, James SA, Cooney KA, Doerr KM, Montie JE, Schottenfeld D: Risk factors for lower urinary tract symptoms in a population-based sample of African-American men. *Am J Epidemiol* 157:906–914, 2003
22. Van Den Eeden SK, Enger S, Caan BJ, Quesenberry CPJ, Sternfeld B, Ferrara A, Loo R, Rowell S, Habel LA: Urologic conditions and diabetes in the California Men's Health Study (Abstract). In *Urologic Complications of Diabetes*. Bethesda, Maryland, National Institutes of Health, 2003, p. 49
23. Michel MC, Mehlburger L, Schumacher H, Bressel HU, Goepel M: Effect of diabetes on lower urinary tract symptoms in patients with benign prostatic hyperplasia. *J Urol* 163:1725–1729, 2000
24. Bourke JB, Griffin JP: Diabetes mellitus and prostatic hyperplasia. *Br Med J* 4:832–833, 1968
25. Glynn RJ, Campion EW, Bouchard GR, Silbert JE: The development of benign prostatic hyperplasia among volunteers in the Normative Aging Study. *Am J Epidemiol* 121:78–90, 1985
26. Sidney S, Quesenberry C Jr, Sadler MC, Lydick EG, Guess HA, Cattolica EV: Risk factors for surgically treated benign prostatic hyperplasia in a prepaid health care plan. *Urology* 38:13–19, 1991
27. Meigs JB, Mohr B, Barry MJ, Collins MM, McKinlay JB: Risk factors for clinical benign prostatic hyperplasia in a community-based population of healthy aging men. *J Clin Epidemiol* 54:935–944, 2001
28. Burke JP, Jacobson DJ, Roberts RO, Girman CJ, Lieber MM, Jacobsen SJ: BPH progression and diabetes mellitus in Olmsted County, Minnesota (Poster Abstract). In *Urologic Complications of Diabetes*. Bethesda, Maryland, National Institutes of Health, 2003, p. 61
29. Kanda M, Eto K, Tanabe N, Sugiyama A, Hashimoto K, Ueno A: Effects of ONO-2235, an aldose reductase inhibitor, on muscarinic receptors and contractile response of the urinary bladder in rats with streptozotocin-induced diabetes. *Jpn J Pharmacol* 73:221–228, 1997
30. Longhurst PA, Belis JA: Abnormalities of rat bladder contractility in streptozotocin-induced diabetes mellitus. *J Pharmacol Exp Ther* 238:773–777, 1986
31. Tong YC, Cheng JT, Wan WC: Effects of Ba-Wei-Die-Huang-Wan on the cholinergic function and protein expression of M2 muscarinic receptor of the urinary bladder in diabetic rats. *Neurosci Lett* 330:21–24, 2002
32. Kubota Y, Nakahara T, Mitani A, Maruko T, Sakamoto K, Ishii K: Augmentation of rat urinary bladder relaxation mediated by beta1-adrenoceptors in experimental diabetes. *Eur J Pharmacol* 467:191–195, 2003
33. Waring JV, Wendt IR: Effects of streptozotocin-induced diabetes mellitus on intracellular calcium and contraction of longitudinal smooth muscle from rat urinary bladder. *J Urol* 163:323–330, 2000
34. Bezuijen MW, Levendusky MC, Longhurst PA: Functional response of bladder strips from streptozotocin diabetic rats depends on bladder mass. *J Urol* 169:2397–2401, 2003
35. Hashitani H, Suzuki H: Altered electrical properties of bladder smooth muscle in streptozotocin-induced diabetic rats. *Br J Urol* 77:798–804, 1996
36. Sasaki K, Chancellor MB, Phelan MW, Yokoyama T, Fraser MO, Seki S, Kubo K, Kumon H, Groat WC, Yoshimura N: Diabetic cystopathy correlates with a long-term decrease in nerve growth factor levels in the bladder and lumbosacral dorsal root ganglia. *J Urol* 168:1259–1264, 2002
37. Pitre DA, Ma T, Wallace LJ, Bauer JA: Time-dependent urinary bladder remodeling in the streptozotocin-induced diabetic rat model. *Acta Diabetol* 39:23–27, 2002
38. Enzlin P, Mathieu C, Vanderschueren D, Demyttenaere K: Diabetes mellitus and female sexuality: a review of 25 years' research. *Diabet Med* 15:809–815, 1998
39. Webster L: Management of sexual problems in diabetic patients. *Br J Hosp Med* 51:465–468, 1994
40. Klein R, Klein BE, Lee KE, Moss SE, Cruickshanks KJ: Prevalence of self-reported erectile dysfunction in people with long-term IDDM. *Diabetes Care* 19: 135–141, 1996
41. De Berardis G, Franciosi M, Belfiglio M, Di Nardo B, Greenfield S, Kaplan SH, Pellegrini F, Sacco M, Tognoni G, Valentini M, Nicolucci A: Erectile dysfunction and quality of life in type 2 diabetic patients: a serious problem too often overlooked. *Diabetes Care* 25:284–291, 2002
42. Penson DF, Latini DM, Lubeck DP, Wallace KL, Henning JM, Lue TF: Do impotent men with diabetes have more severe erectile dysfunction and worse quality of life than the general population of impotent patients? Results from the Exploratory Comprehensive Evaluation of Erectile Dysfunction (ExCEED) database. *Diabetes Care* 26:1093–1099, 2003
43. Klein BE, Klein R: Ten year incidence of self-reported erectile dysfunction in people with long-term type 1 diabetes. *J Diabetes Complications*. In press
44. Johannes CB, Araujo AB, Feldman HA, Derby CA, Kleinman KP, McKinlay JB: Incidence of erectile dysfunction in men 40 to 69 years old: longitudinal results from the Massachusetts Male Aging Study. *J Urol* 163:460–463, 2000
45. Romeo JH, Sefitel AD, Madhun ZT, Aron DC: Sexual function in men with diabetes type 2: association with glycemic control. *J Urol* 163:788–791, 2000
46. Fedele D, Coscelli C, Santeusano F, Bortolotti A, Chatenoud L, Colli E, Landoni M, Parazzini F: Erectile dysfunction in diabetic subjects in Italy: Gruppo Italiano Studio Deficit Eretille nei Diabetici. *Diabetes Care* 21:1973–1977, 1998
47. Bortolotti A, Fedele D, Chatenoud L, Colli E, Coscelli C, Landoni M, Lavezzari M, Santeusano F, Parazzini F: Cigarette smoking: a risk factor for erectile dysfunction in diabetics. *Eur Urol* 40:392–396 (discussion 397), 2001
48. Bacon CG, Mittleman MA, Kawachi I, Giovannucci E, Glasser DB, Rimm EB: Sexual function in men older than 50 years of age: results from the Health Professionals Follow-Up Study. *Ann Intern Med* 139:161–168, 2003
49. Saenz de Tejada I, Anglin G, Knight JR, Emmick JT: Effects of tadalafil on erectile dysfunction in men with diabetes. *Diabetes Care* 25:2159–2164, 2002
50. Goldstein I, Young JM, Fischer J, Bangerter K, Segerson T, Taylor T: Vardenafil, a new phosphodiesterase type 5 inhibitor, in the treatment of erectile dysfunction in men with diabetes: a multicenter double-blind placebo-controlled fixed-dose study. *Diabetes Care* 26:777–783, 2003
51. Goldstein I, Lue TF, Padma-Nathan H, Rosen RC, Steers WD, Wicker PA: Oral

- sildenafil in the treatment of erectile dysfunction: Sildenafil Study Group. *N Engl J Med* 338:1397–1404, 1998
52. Lepore G, Nosari I: Efficacy of oral sildenafil in the treatment of erectile dysfunction in diabetic men with positive response to intracavernosal injection of alprostadil. *Diabetes Care* 24:409–411, 2001
 53. Nehra A, Goldstein I, Pabby A, Nugent M, Huang YH, de las Morenas A, Krane RJ, Udelson D, Saenz de Tejada I, Moreland RB: Mechanisms of venous leakage: a prospective clinicopathological correlation of corporeal function and structure. *J Urol* 156:1320–1329, 1996
 54. Carson CC, Mulcahy JJ, Govier FE: Efficacy, safety and patient satisfaction outcomes of the AMS 700CX inflatable penile prosthesis: results of a long-term multicenter study: AMS 700CX Study Group. *J Urol* 164:376–380, 2000
 55. Lin CS, Lau A, Tu R, Lue TF: Expression of three isoforms of cGMP-binding cGMP-specific phosphodiesterase (PDE5) in human penile cavernosum. *Biochem Biophys Res Commun* 268:628–635, 2000
 56. Mills TM, Chitale K, Lewis RW, Webb RC: Nitric oxide inhibits RhoA/Rho-kinase signaling to cause penile erection. *Eur J Pharmacol* 439:173–174, 2002
 57. Byrne RR, Henry GD, Rao DS, Huynh TT, Pippen AM, Annex BH, Hagen PO, Donatucci CF: Vascular endothelial growth factor restores corporeal smooth muscle function in vitro. *J Urol* 165:1310–1315, 2001
 58. Burnett AL, Becker RE: Immunophilin ligands promote penile neurogenesis and erection recovery after cavernous nerve injury. *J Urol* 171:495–500, 2004
 59. Magee TR, Ferrini M, Garban HJ, Vernet D, Mitani K, Rajfer J, Gonzalez-Cadavid NF: Gene therapy of erectile dysfunction in the rat with penile neuronal nitric oxide synthase. *Biol Reprod* 67:20–28, 2002
 60. Christ GJ, Rehman J, Day N, Salkoff L, Valcic M, Melman A, Geliebter J: Intra-corporal injection of hSlo cDNA in rats produces physiologically relevant alterations in penile function. *Am J Physiol* 275:H600–H608, 1998
 61. Rajfer J, Aronson WJ, Bush PA, Dorey FJ, Ignarro LJ: Nitric oxide as a mediator of relaxation of the corpus cavernosum in response to nonadrenergic, noncholinergic neurotransmission. *N Engl J Med* 326:90–94, 1992
 62. Seftel AD, Vaziri ND, Ni Z, Razmjouei K, Fogarty J, Hampel N, Polak J, Wang RZ, Ferguson K, Block C, Haas C: Advanced glycation end products in human penis: elevation in diabetic tissue, site of deposition, and possible effect through iNOS or eNOS. *Urology* 50:1016–1026, 1997
 63. Bivalacqua TJ, Hellstrom WJ, Kadowitz PJ, Champion HC: Increased expression of arginase II in human diabetic corpus cavernosum: in diabetic-associated erectile dysfunction. *Biochem Biophys Res Commun* 283:923–927, 2001
 64. Autieri MV, Melman A, Christ GJ: Identification of a down-regulated mRNA transcript in corpus cavernosum from diabetic patients with erectile dysfunction. *Int J Impot Res* 8:69–73, 1996
 65. Bax G, Marin N, Piarulli F, Lamonica M, Bellio F, Fedele D: Rigiscan evaluation of specific nervous impairment in patients with diabetes and erectile disorders. *Diabetes Care* 21:1159–1161, 1998
 66. Vardi Y, Sprecher E, Kanter Y, Livne PM, Hemli JA, Yarnitsky D: Polyneuropathy in impotence. *Int J Impot Res* 8:65–68, 1996
 67. Wellmer A, Sharief MK, Knowles CH, Misra VP, Kopelman P, Ralph D, Anand P: Quantitative sensory and autonomic testing in male diabetic patients with erectile dysfunction. *BJU Int* 83:66–70, 1999
 68. Jevtic MJ, Edson M, Jarman WD, Herrera HH: Vascular factor in erectile failure among diabetics. *Urology* 19:163–168, 1982
 69. McVary KT, Rathnau CH, McKenna KE: Sexual dysfunction in the diabetic BB/WOR rat: a role of central neuropathy. *Am J Physiol* 272:R259–R267, 1997
 70. Rehman J, Chenven E, Brink P, Peterson B, Walcott B, Wen YP, Melman A, Christ G: Diminished neurogenic but not pharmacological erections in the 2- to 3-month experimentally diabetic F-344 rat. *Am J Physiol* 272:H1960–H1971, 1997
 71. El-Sakka AI, Lin CS, Chui RM, Dahiya R, Lue TF: Effects of diabetes on nitric oxide synthase and growth factor genes and protein expression in an animal model. *Int J Impot Res* 11:123–132, 1999
 72. Escrig A, Marin R, Abreu P, Gonzalez-Mora JL, Mas M: Changes in mating behavior, erectile function, and nitric oxide levels in penile corpora cavernosa in streptozotocin-diabetic rats. *Biol Reprod* 66:185–189, 2002
 73. Podlasek CA, Zelter DJ, Bervig TR, Gonzalez CM, McKenna KE, McVary KT: Characterization and localization of nitric oxide synthase isoforms in the BB/WOR diabetic rat. *J Urol* 166:746–755, 2001
 74. Celtek S, Foxwell NA, Moncada S: Two phases of nitroergic neuropathy in streptozotocin-induced diabetic rats. *Diabetes* 52:2353–2362, 2003
 75. Brownlee M: Biochemistry and molecular cell biology of diabetic complications. *Nature* 414:813–820, 2001
 76. Podlasek CA, Zelter DJ, Harris JD, Meroz CL, Tang Y, McKenna KE, McVary KT: Altered Sonic hedgehog signaling is associated with morphological abnormalities in the penis of the BB/WOR diabetic rat. *Biol Reprod* 69:816–827, 2003
 77. Wingard CJ, Husain S, Williams J, James S: RhoA-Rho kinase mediates synergistic ET-1 and phenylephrine contraction of rat corpus cavernosum. *Am J Physiol Regul Integr Comp Physiol* 285:R1145–R1152, 2003
 78. Chang S, Hypolite JA, Changolkar A, Wein AJ, Chacko S, DiSanto ME: Increased contractility of diabetic rabbit corpora smooth muscle in response to endothelin is mediated via Rho-kinase beta. *Int J Impot Res* 15:53–62, 2003
 79. Wingard CJ, Johnson JA, Holmes A, Prikosh A: Improved erectile function after Rho-kinase inhibition in a rat castrate model of erectile dysfunction. *Am J Physiol Regul Integr Comp Physiol* 284:R1572–R1579, 2003
 80. Chitale K, Wingard CJ, Clinton Webb R, Branam H, Stopper VS, Lewis RW, Mills TM: Antagonism of Rho-kinase stimulates rat penile erection via a nitric oxide-independent pathway. *Nat Med* 7:119–122, 2001
 81. Basson R, Berman J, Burnett A, Derogatis L, Ferguson D, Fourcroy J, Goldstein I, Graziottin A, Heiman J, Laan E, Leiblum S, Padma-Nathan H, Rosen R, Segraves K, Segraves RT, Shabsigh R, Sipski M, Wagner G, Whipple B: Report of the international consensus development conference on female sexual dysfunction: definitions and classifications. *J Urol* 163:888–893, 2000
 82. Wincze JP, Albert A, Bansal S: Sexual arousal in diabetic females: physiological and self-report measures. *Arch Sex Behav* 22:587–601, 1993
 83. Enzlin P, Mathieu C, Van den Bruel A, Bosteels J, Vanderschueren D, Demyttenaere K: Sexual dysfunction in women with type 1 diabetes: a controlled study. *Diabetes Care* 25:672–677, 2002
 84. Schiel R, Muller UA: Prevalence of sexual disorders in a selection-free diabetic population (JEVIN). *Diabetes Res Clin Pract* 44:115–121, 1999
 85. Munarriz R, Kim NN, Goldstein I, Traish AM: Biology of female sexual function. *Urol Clin North Am* 29:685–693, 2002
 86. Hale TM, Heaton JP, Adams MA: A framework for the present and future development of experimental models of female sexual dysfunction. *Int J Impot* 15 (Suppl. 5):S75–S79, 2003
 87. Bechara A, Bertolino MV, Casabe A, Munarriz R, Goldstein I, Morin A, Secin F, Literat B, Pesaresi M, Fredotovich N: Duplex Doppler ultrasound assessment

- of clitoral hemodynamics after topical administration of alprostadil in women with arousal and orgasmic disorders. *J Sex Marital Ther* 29 (Suppl. 1):1–10, 2003
88. Bachmann G, Bancroft J, Braunstein G, Burger H, Davis S, Dennerstein L, Goldstein I, Guay A, Leiblum S, Lobo R, Nottelovitz M, Rosen R, Sarrel P, Sherwin B, Simon J, Simpson E, Shifren J, Spark R, Traish A: Female androgen insufficiency: the Princeton consensus statement on definition, classification, and assessment. *Fertil Steril* 77:660–665, 2002
 89. Shifren JL, Braunstein GD, Simon JA, Casson PR, Buster JE, Redmond GP, Burki RE, Ginsburg ES, Rosen RC, Leiblum SR, Caramelli KE, Mazer NA: Transdermal testosterone treatment in women with impaired sexual function after oophorectomy. *N Engl J Med* 343:682–688, 2000
 90. Berman JR, Berman LA, Toler SM, Gill J, Haughie S: Safety and efficacy of sildenafil citrate for the treatment of female sexual arousal disorder: a double-blind, placebo controlled study. *J Urol* 170:2333–2338, 2003
 91. Stapleton A: Urinary tract infections in patients with diabetes. *Am J Med* 113 (Suppl. 1A):80S–84S, 2002
 92. Zhanel GG, Nicolle LE, Harding GK: Prevalence of asymptomatic bacteriuria and associated host factors in women with diabetes mellitus: the Manitoba Diabetic Urinary Infection Study Group. *Clin Infect Dis* 21:316–322, 1995
 93. Geerlings SE, Stolk RP, Camps MJ, Netten PM, Hoekstra JB, Bouter PK, Bravboer B, Collet TJ, Jansz AR, Hoepelman AM: Asymptomatic bacteriuria can be considered a diabetic complication in women with diabetes mellitus. *Adv Exp Med Biol* 485:309–314, 2000
 94. Geerlings SE, Stolk RP, Camps MJ, Netten PM, Collet JT, Schneeberger PM, Hoepelman AI: Consequences of asymptomatic bacteriuria in women with diabetes mellitus. *Arch Intern Med* 161:1421–1427, 2001
 95. Hooton TM, Scholes D, Stapleton AE, Roberts PL, Winter C, Gupta K, Samadpour M, Stamm WE: A prospective study of asymptomatic bacteriuria in sexually active young women. *N Engl J Med* 343:992–997, 2000
 96. Geerlings SE, Stolk RP, Camps MJ, Netten PM, Collet TJ, Hoepelman AI: Risk factors for symptomatic urinary tract infection in women with diabetes. *Diabetes Care* 23:1737–1741, 2000
 97. Harding GK, Zhanel GG, Nicolle LE, Cheang M: Antimicrobial treatment in diabetic women with asymptomatic bacteriuria. *N Engl J Med* 347:1576–1583, 2002
 98. Boyko EJ, Fihn SD, Scholes D, Chen CL, Normand EH, Yarbro P: Diabetes and the risk of acute urinary tract infection among postmenopausal women. *Diabetes Care* 25:1778–1783, 2002
 99. Fihn SD, Boyko EJ, Scholes D, Jackson S, Hu K, Abraham L, Gupta K: Diabetes and the risk of urinary tract infection in community dwelling, post-menopausal women (Abstract). In *Urologic Complications of Diabetes*. Bethesda, MD, National Institutes of Health, 2003, p. 29–30
 100. Brown JS, Vittinghoff E, Kanaya AM, Agarwal SK, Hulley S, Foxman B: Urinary tract infections in postmenopausal women: effect of hormone therapy and risk factors. *Obstet Gynecol* 98:1045–1052, 2001
 101. Hooton TM, Scholes D, Hughes JP, Winter C, Roberts PL, Stapleton AE, Stergachis A, Stamm WE: A prospective study of risk factors for symptomatic urinary tract infection in young women. *N Engl J Med* 335:468–474, 1996
 102. Fihn SD: Clinical practice: acute uncomplicated urinary tract infection in women. *N Engl J Med* 349:259–266, 2003
 103. Hansen DS, Gottschau A, Kolmos HJ: Epidemiology of Klebsiella bacteraemia: a case control study using *Escherichia coli* bacteraemia as control. *J Hosp Infect* 38:119–132, 1998
 104. Stamm WE, Norrby SR: Urinary tract infections: disease panorama and challenges. *J Infect Dis* 183 (Suppl. 1):S1–S4, 2001
 105. Wullt B, Bergsten G, Fischer H, Godaly G, Karpman D, Leijonhufvud I, Lundstedt AC, Samuelsson P, Samuelsson M, Svensson ML, Svanborg C: The host response to urinary tract infection. *Infect Dis Clin North Am* 17:279–301, 2003
 106. Johnson JR: Microbial virulence determinants and the pathogenesis of urinary tract infection. *Infect Dis Clin North Am* 17:261–278, viii, 2003
 107. Brauner A, Katouli M, Ostenson CG: P-fimbriation and haemolysin production are the most important virulence factors in diabetic patients with *Escherichia coli* bacteraemia: a multivariate statistical analysis of seven bacterial virulence factors. *J Infect* 31:27–31, 1995
 108. Geerlings SE, Brouwer EC, Gaastra W, Stolk R, Diepersloot RJ, Hoepelman AI: Virulence factors of *Escherichia coli* isolated from urine of diabetic women with asymptomatic bacteriuria: correlation with clinical characteristics. *Antonie Van Leeuwenhoek* 80:119–127, 2001
 109. Geerlings SE, Meiland R, van Lith EC, Brouwer EC, Gaastra W, Hoepelman AI: Adherence of type 1-fimbriated *Escherichia coli* to uroepithelial cells: more in diabetic women than in control subjects. *Diabetes Care* 25:1405–1409, 2002
 110. Balasoiu D, van Kessel KC, van Kats-Renaud HJ, Collet TJ, Hoepelman AI: Granulocyte function in women with diabetes and asymptomatic bacteriuria. *Diabetes Care* 20:392–395, 1997
 111. Zozulinska D, Majchrzak A, Sobieska M, Wiktorowicz K, Wierusz-Wysocka B: Serum interleukin-8 level is increased in diabetic patients. *Diabetologia* 42:117–118, 1999
 112. Geerlings SE, Hoepelman AI: Immune dysfunction in patients with diabetes mellitus (DM). *FEMS Immunol Med Microbiol* 26:259–265, 1999
 113. Geerlings SE, Brouwer EC, Van Kessel KC, Gaastra W, Stolk RP, Hoepelman AI: Cytokine secretion is impaired in women with diabetes mellitus. *Eur J Clin Invest* 30:995–1001, 2000
 114. Li YM, Tan AX, Vlassara H: Antibacterial activity of lysozyme and lactoferrin is inhibited by binding of advanced glycation-modified proteins to a conserved motif. *Nat Med* 1:1057–1061, 1995
 115. Huttner KM, Bevins CL: Antimicrobial peptides as mediators of epithelial host defense. *Pediatr Res* 45:785–794, 1999
 116. Valore EV, Park CH, Quayle AJ, Wiles KR, McCray PB Jr, Ganz T: Human beta-defensin-1: an antimicrobial peptide of urogenital tissues. *J Clin Invest* 101:1633–1642, 1998
 117. UK Prospective Diabetes Study (UKPDS) Group: Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* 352:837–853, 1998
 118. Wan P, Lau J, Chalmore T: Meta-analysis of effects of intensive blood glucose control on the late diabetic complications of type 1 diabetes. *Lancet* 341:877–890, 1993