

# Histological Course of Nonalcoholic Fatty Liver Disease in Japanese Patients

Tight glycemic control, rather than weight reduction, ameliorates liver fibrosis

ERIKA HAMAGUCHI, MD, PHD<sup>1</sup>  
TOSHINARI TAKAMURA, MD, PHD<sup>1</sup>  
MASARU SAKURAI, MD, PHD<sup>2</sup>  
EISHIRO MIZUKOSHI, MD, PHD<sup>3</sup>  
YOH ZEN, MD, PHD<sup>4</sup>  
YUMIE TAKESHITA, MD, PHD<sup>1</sup>

SEIICHIRO KURITA, MD, PHD<sup>1</sup>  
KUNIAKI ARAI, MD, PHD<sup>3</sup>  
TATSUYA YAMASHITA, MD, PHD<sup>3</sup>  
MOTOKO SASAKI, MD, PHD<sup>5</sup>  
YASUNI NAKANUMA, MD, PHD<sup>5</sup>  
SHUICHI KANEKO, MD, PHD<sup>3</sup>

**OBJECTIVE** — The goal of this study was to examine whether metabolic abnormalities are responsible for the histological changes observed in Japanese patients with nonalcoholic fatty liver disease (NAFLD) who have undergone serial liver biopsies.

**RESEARCH DESIGN AND METHODS** — In total, 39 patients had undergone consecutive liver biopsies. Changes in their clinical data were analyzed, and biopsy specimens were scored histologically for stage.

**RESULTS** — The median follow-up time was 2.4 years (range 1.0–8.5). Liver fibrosis had improved in 12 patients (30.7%), progressed in 11 patients (28.2%), and remained unchanged in 16 patients (41%). In a Cox proportional hazard model, decrease in A1C and use of insulin were associated with improvement of liver fibrosis independent of age, sex, and BMI. However,  $\Delta$ A1C was more strongly associated with the improvement of liver fibrosis than use of insulin after adjustment for each other ( $\chi^2$ ; 7.97 vs. 4.58, respectively).

**CONCLUSIONS** — Tight glycemic control may prevent histological progression in Japanese patients with NAFLD.

*Diabetes Care* 33:284–286, 2010

Accumulating trans-sectional evidence suggests that the presence of multiple metabolic disorders, including obesity, diabetes, dyslipidemia, hypertension, and ultimately metabolic syndrome, are associated with nonalcoholic fatty liver disease (NAFLD) (1). However, it remains unclear which metabolic abnormalities are responsible for the pathological progression of NAFLD, especially in Japanese patients, who generally are not severely obese compared with Western patients.

We retrospectively compared clinical features with the histological changes in the livers of Japanese patients with NAFLD who had undergone serial liver biopsies.

## RESEARCH DESIGN AND METHODS

We recruited 195 patients with clinically suspected NAFLD who had undergone liver biopsies at Kanazawa University Hospital from 1997 through 2008. For details about the study subjects and the exclusion criteria, see supplementary Fig. 1 in the online appendix, available at <http://care.diabetesjournals.org/cgi/content/full/dc09-0148/DC1>. Of 178 patients diagnosed histologically as having NAFLD, 39 had undergone serial liver biopsies.

ria, see supplementary Fig. 1 in the online appendix, available at <http://care.diabetesjournals.org/cgi/content/full/dc09-0148/DC1>. Of 178 patients diagnosed histologically as having NAFLD, 39 had undergone serial liver biopsies.

## Data collection

Clinical information, including age, sex, body measurements, and prevalence of metabolic abnormalities, was obtained for each patient. Venous blood samples drawn for laboratory testing before the liver biopsies were obtained. All subjects had been administered a 75-g oral glucose tolerance test at baseline and at follow-up.

## Liver biopsies

Biopsies were obtained after a thorough clinical evaluation and receipt of signed informed consent from each patient. All biopsies were analyzed twice and at separate times randomly by a single pathologist who was blinded to the clinical information and the order in which the biopsies were obtained. The biopsied tissues were scored for steatosis, stage, and grade as described (2), according to the standard criteria for grading and staging of nonalcoholic steatohepatitis proposed by Brunt et al. (3).

For additional details on subjects, data collection methods, liver pathology, and statistical analyses, see supplementary Methods in the online appendix.

**RESULTS** — The basal clinical and biochemical data from 39 patients with NAFLD are described in supplementary Table 1. Prevalence of type 2 diabetes, hypertension, and dyslipidemia were 77, 36, and 64%, respectively. The median follow-up period was 2.4 years (range 1.0–8.5). Medications for diabetes and medication changes during the follow-up period are described in supplementary Table 2. Seventeen patients treated with oral diabetic agents were switched to insulin therapy after the initial biopsy. No patients initiated pioglitazone during follow-up.

From the <sup>1</sup>Department of Disease Control and Homeostasis, Kanazawa University Graduate School of Medical Science, Ishikawa, Japan; the <sup>2</sup>Department of Epidemiology and Public Health, Kanazawa Medical University, Ishikawa, Japan; the <sup>3</sup>Department of Gastroenterology, Kanazawa University Hospital, Ishikawa, Japan; the <sup>4</sup>Division of Pathology, Kanazawa University Hospital, Ishikawa, Japan; and the <sup>5</sup>Department of Human Pathology, Kanazawa University Graduate School of Medical Science, Ishikawa, Japan.

Corresponding author: Toshinari Takamura, [ttakamura@m-kanazawa.jp](mailto:ttakamura@m-kanazawa.jp).

Received 8 February 2009 and accepted 20 October 2009. Published ahead of print at <http://care.diabetesjournals.org> on 30 October 2009. DOI: 10.2337/dc09-0148.

© 2010 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Table 1—Baseline and follow-up clinical features and gradients of laboratory markers associated with changes in liver fibrosis in 39 patients with NAFLD

	Baseline			P	Follow-up			P
	Improved	Stable	Progressed		Improved	Stable	Progressed	
n	12	16	11		12	16	11	—
Simple fatty liver:nonalcoholic steatohepatitis (n)	3:9	9:7	10:1		10:2	9:7	6:5	
Age (years)	51.5 (29–66) 5:7	48.5 (20–79) 12:4	51.5 (29–66) 5:7	0.97 0.17	26.9 (22.8–31.2)	29.1 (24.3–44.8)	30.7 (24.1–36.3)	0.13
Sex (M:F)	5:7	12:4	5:7	0.17	23 (11–28)	26 (15–71)	24 (14–164)	0.20
BMI (kg/m <sup>2</sup> )	27.5 (23.2–34.1)	27.7 (22.5–44.4)	30.9 (23.4–37.7)	0.74	21 (11–53)	36 (21–66)	31 (12–202)	0.10
Aspartate transaminase (IU/l)	70 (11–106)	29 (14–86)	32 (13–83)	0.05	103 (93–220)	121 (83–198)	116 (88–199)	0.51
Alanine transaminase (IU/l)	71 (10–209)	48 (23–81)	40 (11–162)	0.13	6.0 (5.0–9.0)	6.2 (5.0–10.0)	7.0 (6.0–11.0)	0.10
Fasting plasma glucose (mg/dl)	133 (96–207)	143 (87–414)	111 (76–167)	0.20	3.1 (1.5–8.5)	3.4 (1.9–7.7)	3.9 (1.6–11.1)	0.76
A1C (%)	8.2 (4.7–11.6)	8.0 (4.9–13.6)	6.2 (5.1–9.5)	0.27	0.33 (0.28–0.37)	0.32 (0.30–0.35)	0.31 (0.29–0.34)	0.82
HOMA-IR	3.9 (0.7–5.5)	3.4 (1.9–7.7)	3.9 (1.6–11.1)	0.91	2.4 (1.6–4.5)	2.4 (1.6–4.5)	1.9 (1.3–4.5)	0.80
QUICKI	0.32 (0.29–0.40)	0.31 (0.27–0.34)	0.31 (0.29–0.35)	0.32	2.0 (1.3–5.9)	4.3 (1.9–15.9)	4.5 (2.3–8.8)	0.75
Muscle insulin resistance	2.1 (1.5–4.0)	1.7 (0.3–3.3)	3.0 (2.1–4.4)	0.20	192 (114–224)	115 (57–241)	131 (36–173)	0.68
Hepatic insulin resistance (×10 <sup>6</sup> )	5.3 (2.3–10.2)	5.0 (2.3–10.0)	3.7 (1.4–10.6)	0.66	53 (40–71)	52 (39–64)	52 (36–79)	0.92
Total cholesterol (mg/dl)	191 (128–276)	187 (129–252)	206 (163–244)	0.57	21.5 (6.3–31.8)	162 (110–614)	62 (10–171)	0.05
Triglycerides (mg/dl)	111 (28–224)	114 (36–204)	96 (36–521)	0.87	0.10 (0.00–0.24)	0.10 (0.00–0.24)	0.09 (0.00–0.89)	0.89
HDL cholesterol (mg/dl)	47 (35–82)	51 (31–73)	48 (20–74)	0.68	8.3 (3.2–14.0)	24.5 (0.0–57.0)	24.3 (0.0–140.3)	0.63
Platelets (×10 <sup>4</sup> /μl)	21.1 (9.4–30.8)	23.0 (7.0–38.2)	24.3 (20.2–41.2)	0.29	0.6 (0.4–0.8)	0.5 (0.5–233.0)	0.6 (0.4–1.0)	0.96
Ferritin (μg/dl)	185 (13–452)	397 (190–604)	46 (10–347)	0.14	21.5 (6.3–31.8)	162 (110–614)	62 (10–171)	0.05
hs-CRP	0.40 (0.08–7.53)	0.14 (0.02–0.61)	0.06 (0.00–0.30)	0.23	8.3 (3.2–14.0)	24.5 (0.0–57.0)	24.3 (0.0–140.3)	0.63
Type IV collagen 7S (ng/dl)	5.1 (2.7–10.0)	4.1 (3.1–7.2)	3.7 (3.3–4.5)	0.27	0.5 (0.5–233.0)	0.5 (0.5–233.0)	0.6 (0.4–1.0)	0.96
HA (ng/dl)	20.6 (0.0–144.7)	25.5 (11.5–299)	30.4 (0.0–61.7)	0.66	0.3 (–1.8 to 7.1)	0.3 (–1.8 to 7.1)	0.3 (–1.8 to 7.1)	0.02
P-III-P (U/ml)	0.6 (0.5–1.2)	0.6 (0.4–45.0)	0.5 (0.4–0.6)	0.07	–4.7 (–10.6 to 10.2)	2.2 (–9.4 to 13.4)	–0.9 (–12.7 to 9.6)	0.04
Diabetes (%)	82	69	64	0.59	–1.3 (–4.4 to 1.2)	–0.3 (–4.3 to 3.3)	–0.7 (–6.1 to 1.8)	0.81
Dyslipidemia (%)	73	63	73	0.95	–1.9 (–6.0 to 0.4)	–1.2 (–6.1 to 4.4)	0.3 (–1.8 to 7.1)	0.02
Hypertension (%)	64	18	36	0.03	–4.7 (–10.6 to 10.2)	2.2 (–9.4 to 13.4)	–0.9 (–12.7 to 9.6)	0.04
Metabolic syndrome (%)	73	38	27	0.18	–1.3 (–4.4 to 1.2)	–0.3 (–4.3 to 3.3)	–0.7 (–6.1 to 1.8)	0.81
AAIC					–1.9 (–6.0 to 0.4)	–1.2 (–6.1 to 4.4)	0.3 (–1.8 to 7.1)	0.02
Δbody weight					–4.7 (–10.6 to 10.2)	2.2 (–9.4 to 13.4)	–0.9 (–12.7 to 9.6)	0.04
AHOMA-IR					–1.3 (–4.4 to 1.2)	–0.3 (–4.3 to 3.3)	–0.7 (–6.1 to 1.8)	0.81

Data are medians (range) or %. A Kruskal-Wallis test and a  $\chi^2$  test were used to compare the continuous and categorical variables among three groups. HA, hyaluronic acid; hs-CRP, high-sensitivity C-reactive protein; P-III-P, procollagen III peptide.

Liver fibrosis improved in 12 patients (30.7%), progressed in 11 patients (28.2%), and remained unchanged in 16 patients (41%). As shown in Table 1, fasting plasma glucose, A1C, insulin resistance indicators, and prevalence of metabolic disorders were not significantly different among the three liver fibrosis groups. In the Cox proportional hazard model (supplementary Table 3), although some of the confidence intervals were very wide because of the small sample size, improvement of liver fibrosis was significantly associated with changes in A1C between the initial and final liver biopsies ( $\Delta$ A1C) ( $P = 0.005$ ) and use of insulin for the treatment of diabetes ( $P = 0.019$ ). Both  $\Delta$ A1C and use of insulin were independently associated with the improvement of liver fibrosis after adjusted for each other. However,  $\Delta$ A1C was more strongly associated with the improvement of liver fibrosis than use of insulin ( $\chi^2$ ; 7.97 vs. 4.58, respectively; supplementary Table 3).

**CONCLUSIONS**— In the present study, we showed that a change in glycemic control ( $\Delta$ A1C), but not changes in insulin resistance indicators, was an independent predictor of the progression of liver fibrosis in Japanese patients with NAFLD. This is the first report identifying a change in A1C as a predictor of the histological course in the liver of patients with NAFLD. Two of five previous longitudinal studies have identified obesity, higher BMI, and homeostasis model assessment of insulin resistance (HOMA-IR) as predictors of liver fibrosis progression in Western populations (4,5). The difference between those results and the results of the present study may be due in part to differences in the assessed severity of obesity and insulin resistance between the populations. We previously demonstrated that diabetes is an independent risk factor for the progression of liver fibrosis in hepatitis C (6) and that diabetes accelerates the pathology of nonalcoholic steatohepatitis in the type 2 diabetic rat model OLETF (7).

Liver fibrosis is closely associated with two regulators of fibrosis: transforming growth factor (TGF)- $\beta$  (8,9) and plasminogen activator inhibitor type 1 (PAI-1) (8,10). High glucose levels induce the expression of TGF- $\beta$  (11) and PAI-1 (12). We previously reported that the expression of TGF family member

genes, PAI-1, and genes involved in fibrogenesis are upregulated in the livers of patients with type 2 diabetes (13,14), suggesting that a diabetic state increases the risk for liver fibrosis.

In the present study, only  $\Delta$ A1C was associated with the progression of liver fibrosis, but not liver inflammation (data not shown). We speculate that the reduction of A1C inhibits the expression of master genes such as TGF- $\beta$  and PAI-1 that are involved in the regulation of fibrogenesis, rather than genes involved in liver inflammation, and thereby improves liver fibrosis in NAFLD.

The major limitation of this study was small population size. We could not evaluate the changes of liver histology according to the difference in detail characteristics such as treatment of diabetes. Lower statistical power of this study should be considered when we evaluate the results.

In conclusion, our study suggested that  $\Delta$ A1C could predict liver fibrosis progression in Japanese patients with NAFLD, and tight glycemic control may ameliorate liver fibrosis. Future large-scale prospective studies are needed to confirm our results.

**Acknowledgments**— This study was supported in part by a grant-in-aid from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

No potential conflicts of interest relevant to this article were reported.

We thank Dr. Akiko Shimizu, Dr. Tsuguhito Ota, and Dr. Hirofumi Misu for recruiting the patients.

## References

- Duvnjak M, Lerotic I, Barsic N, Tomasic V, Virovic Jukic L, Velagic V. Pathogenesis and management issues for non-alcoholic fatty liver disease. *World J Gastroenterol* 2007;13:4539–4550
- Sakurai M, Takamura T, Ota T, Ando H, Akahori H, Kaji K, Sasaki M, Nakanuma Y, Miura K, Kaneko S. Liver steatosis, but not fibrosis, is associated with insulin resistance in nonalcoholic fatty liver disease. *J Gastroenterol* 2007;42:312–317
- Brunt EM, Janney CG, Di Bisceglie AM, Neuschwander-Tetri BA, Bacon BR. Non-alcoholic steatohepatitis: a proposal for grading and staging the histological lesions. *Am J Gastroenterol* 1999;94:2467–2474
- Ekstedt M, Franzen LE, Mathiesen UL, Thorelius L, Holmqvist M, Bodemar G, Kechagias S. Long-term follow-up of pa-

- tients with NAFLD and elevated liver enzymes. *Hepatology* 2006;44:865–873
- Fassio E, Alvarez E, Dominguez N, Landeira G, Longo C. Natural history of non-alcoholic steatohepatitis: a longitudinal study of repeat liver biopsies. *Hepatology* 2004;40:820–826
- Kita Y, Mizukoshi E, Takamura T, Sakurai M, Takata Y, Arai K, Yamashita T, Nakamoto Y, Kaneko S. Impact of diabetes mellitus on prognosis of patients infected with hepatitis C virus. *Metabolism* 2007;56:1682–1688
- Ota T, Takamura T, Kurita S, Matsuzawa N, Kita Y, Uno M, Akahori H, Misu H, Sakurai M, Zen Y, Nakanuma Y, Kaneko S. Insulin resistance accelerates a dietary rat model of nonalcoholic steatohepatitis. *Gastroenterology* 2007;132:282–293
- Matsuzawa N, Takamura T, Kurita S, Misu H, Ota T, Ando H, Yokoyama M, Honda M, Zen Y, Nakanuma Y, Miyamoto K, Kaneko S. Lipid-induced oxidative stress causes steatohepatitis in mice fed an atherogenic diet. *Hepatology* 2007;46:1392–1403
- Uno M, Kurita S, Misu H, Ando H, Ota T, Matsuzawa-Nagata N, Kita Y, Nabemoto S, Akahori H, Zen Y, Nakanuma Y, Kaneko S, Takamura T. Tranilast, an antifibrogenic agent, ameliorates a dietary rat model of nonalcoholic steatohepatitis. *Hepatology* 2008;48:109–118
- Bergheim I, Guo L, Davis MA, Lambert JC, Beier JI, Duveau I, Luyendyk JP, Roth RA, Arteel GE. Metformin prevents alcohol-induced liver injury in the mouse: critical role of plasminogen activator inhibitor-1. *Gastroenterology* 2006;130:2099–2112
- Sugimoto R, Enjoji M, Kohjima M, Tsuruta S, Fukushima M, Iwao M, Sonta T, Kotoh K, Inoguchi T, Nakamura M. High glucose stimulates hepatic stellate cells to proliferate and to produce collagen through free radical production and activation of mitogen-activated protein kinase. *Liver Int* 2005;25:1018–1026
- Suzuki M, Akimoto K, Hattori Y. Glucose upregulates plasminogen activator inhibitor-1 gene expression in vascular smooth muscle cells. *Life Sci* 2002;72:59–66
- Takamura T, Sakurai M, Ota T, Ando H, Honda M, Kaneko S. Genes for systemic vascular complications are differentially expressed in the livers of type 2 diabetic patients. *Diabetologia* 2004;47:638–647
- Takeshita Y, Takamura T, Hamaguchi E, Shimizu A, Ota T, Sakurai M, Kaneko S. Tumor necrosis factor-alpha-induced production of plasminogen activator inhibitor 1 and its regulation by pioglitazone and cerivastatin in a nonmalignant human hepatocyte cell line. *Metabolism* 2006;55:1464–1472