

# Relationship of Glycemic Control, Exogenous Insulin, and C-Peptide Levels to Ischemic Heart Disease Mortality Over a 16-Year Period in People With Older-Onset Diabetes

The Wisconsin Epidemiologic Study of Diabetic Retinopathy (WESDR)

FLAVIO E. HIRAI, MD, MPH<sup>1,2</sup>  
 SCOT E. MOSS, MA<sup>1</sup>  
 BARBARA E.K. KLEIN, MD, MPH<sup>1</sup>  
 RONALD KLEIN, MD, MPH<sup>1</sup>

**OBJECTIVE** — The purpose of this study was to examine the relationship of glycemic control and exogenous and endogenous insulin levels with all-cause and cause-specific mortality (ischemic heart disease and stroke) in an older-onset diabetic population.

**RESEARCH DESIGN AND METHODS** — The Wisconsin Epidemiologic Study of Diabetic Retinopathy (WESDR) is an ongoing, prospective, population-based cohort study of individuals with diabetes first examined in 1980–1982. A stratified sample of all individuals with diabetes diagnosed at 30 years of age or older was labeled “older-onset” ( $n = 1,370$ ). Those participating in the 1984–1986 examination phase ( $n = 1,007$ ) were included in the analysis. Endogenous insulin was determined by measurements of plasma C-peptide (in nanomoles per liter), and exogenous insulin was calculated in units per kilogram per day. Glycemic control was determined by levels of glycosylated hemoglobin ( $HbA_1$ ).

**RESULTS** — After 16 years of follow-up, 824 individuals died (all-cause mortality); 358 deaths involved ischemic heart disease and 137 involved stroke. C-peptide and  $HbA_1$  were significantly associated with all-cause and ischemic heart disease mortality in our study. The hazard ratio (95% CI) values for all-cause mortality were 1.12 (1.07–1.17) per 1% increase in  $HbA_1$ , 1.20 (0.85–1.69) per 1 unit  $\cdot$  kg<sup>-1</sup>  $\cdot$  day<sup>-1</sup> increase in exogenous insulin, and 1.15 (1.04–1.29) per 1 nmol/l increase in C-peptide and for ischemic heart disease mortality were 1.14 (1.06–1.22), 1.50 (0.92–2.46), and 1.19 (1.02–1.39) for  $HbA_1$ , exogenous insulin, and C-peptide, respectively, after adjusting for relevant confounders. C-peptide was associated with stroke mortality only among men (1.65 [1.07–2.53]).

**CONCLUSIONS** — Our results show that individuals with higher endogenous insulin levels are at higher risk of all-cause, ischemic heart disease, and stroke mortality.

*Diabetes Care* 31:493–497, 2008

From the <sup>1</sup>Department of Ophthalmology and Visual Sciences, University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin; and the <sup>2</sup>Ophthalmology Federal University of São Paulo, São Paulo, Brazil.

Address correspondence to Ronald Klein, MD, MPH, University of Wisconsin Madison, Ophthalmology and Visual Sciences, 610 N. Walnut St., 4th Floor WARF, Madison, Wisconsin 53726. E-mail: kleinr@epi.ophth.wisc.edu.

Received for publication 19 June 2007 and accepted in revised form 14 November 2007.

Published ahead of print at <http://care.diabetesjournals.org> on 19 November 2007. DOI: 10.2337/dc07-1161.

**Abbreviations:** CVD, cardiovascular disease; WESDR, Wisconsin Epidemiologic Study of Diabetic Retinopathy.

© 2008 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

There has been an interest in investigating the association of high levels of endogenous and exogenous insulin with cardiovascular morbidity and mortality (1–9), particularly after the results of the Diabetes Control and Complications Trial (DCCT), which recommended tight glycemic control in order to decrease the risk of microvascular complications in diabetic individuals (10). The rationale has been based on results from clinical and experimental studies that demonstrated the influence of exogenous insulin on arterial wall changes that ultimately contributed to the development of atherosclerosis (11–13). However, increased levels of endogenous insulin and the use of exogenous insulin have been associated with increased cardiovascular morbidity and mortality in some studies (3,5–7,9) but not in others (2,14,15). The purpose of this study was to investigate the relationship of glycemic control and exogenous and endogenous insulin levels with all-cause and cause-specific mortality (ischemic heart disease and stroke) in an older-onset diabetic population.

## RESEARCH DESIGN AND METHODS

The Wisconsin Epidemiologic Study of Diabetic Retinopathy (WESDR) is an ongoing, prospective, population-based cohort study of individuals who were receiving care for diabetes in 11 counties in Wisconsin in 1978–1979. A stratified sample of all individuals with diabetes diagnosed at 30 years of age or older was included and was labeled “older-onset” ( $n = 1,370$ ). The first examination phase was in 1980–1982, and follow-up examinations were performed every 4–6 years. Plasma C-peptide was first measured at the 1984–1986 follow-up visit, which, for the purposes of this report, was considered to be the baseline examination ( $n = 1,007$ ).

**Table 1—Baseline characteristics of individuals with older-onset diabetes in the 1984–1986 WESDR**

	n	Mean ± SD or %
Age (years)	1,007	68.6 ± 11.0
Duration of diabetes (years)	1,007	15.3 ± 7.9
HbA <sub>1c</sub> (%)	980	9.3 ± 1.9
C-peptide (nmol/l)	960	1.0 ± 0.9
Systolic blood pressure (mmHg)	979	143.1 ± 23.3
Diastolic blood pressure (mmHg)	978	75.4 ± 12.3
BMI (kg/m <sup>2</sup> )	884	29.1 ± 5.9
Sex (percent male)	1,006	44.9
Insulin use (yes)	1,007	56.4
Cardiovascular disease (present)	990	34.1
Hypertension (present)	991	57.0
Smoking status (current)	1,001	12.9
Diabetic retinopathy (present)	992	66.0

Detailed protocols used in WESDR have been published elsewhere (16,17). Briefly, baseline and all follow-up examinations included detailed medical history with information about cardiovascular disease (CVD) and cigarette smoking and measurements of blood pressure, height, and weight. Nonfasting endogenous insulin was determined by measurement of plasma C-peptide (in nanomoles per liter), and exogenous insulin was calculated in units per kilogram per day. Glycemic control was determined by levels of glycosylated hemoglobin (HbA<sub>1c</sub>). Stereoscopic seven-field 30° color retinal photographs were taken, and diabetic retinopathy was classified according to a modified Airlie House Classification Scheme (18).

Deaths were ascertained by contacting family members and physicians, review of newspaper obituaries, and use of vital status records. For cause-specific analyses, any mention on the death certificate was considered an event. Causes of death were defined according to the ICD 9th and 10th revisions. Ischemic heart disease mortality was defined according to codes 410.0–414.9 (ICD-9) and I20.0–I25.9 (ICD-10), and stroke mortality was defined according to codes 430.0–438.9 (ICD-9) and I60.0–I69.9 (ICD-10). The survival interval considered in the current analysis was a 16-year period from baseline until death or the date of last contact by 31 December 2002. Of the 1,007 examined, 824 had died, 151 were living as of 31 December 2002, and 32 had their last living contact date before 31 December 2002. The survival interval for the latter group was the period

between the baseline examination and the last contact date.

Diabetic retinopathy was defined as the presence of mild (i.e., microaneurysms only, hard or soft exudates, or hemorrhages with or without microaneurysms) or worse retinopathy. A positive history of CVD was defined as having history of angina, heart attack, or stroke. A person was considered hypertensive if systolic or diastolic blood pressure was ≥140 or 90 mmHg, respectively, or using antihypertensive medication. Current smokers were those who had smoked more than 100 cigarettes and continued to smoke until the day of the examination.

#### Statistical analysis

Age- and sex-adjusted and multivariable analyses were performed. The use of exogenous insulin was analyzed as a binary variable (yes/no) and as a continuous variable (units · kg<sup>-1</sup> · day<sup>-1</sup>), and plasma C-peptide (nmol/l) and HbA<sub>1c</sub> (%) were analyzed as continuous variables in the whole population. To analyze the influence of the use of exogenous insulin on mortality, we calculated a propensity score that was a conditional probability of receiving insulin given some individual covariates (age, duration of diabetes, levels of HbA<sub>1c</sub>, BMI, history of CVD, and presence of diabetic retinopathy) in order to decrease the chances of bias by indication (i.e., physicians tend to prescribe insulin to those with poorer glycemic control and therefore at higher risk for developing comorbidities and for death). Propensity score was then added to our regression models, and multivariable analyses were performed with the Cox

proportional hazards model. Hazard ratios (HRs) and 95% CIs were estimated, and *P* values <0.05 were considered significant.

In addition, HbA<sub>1c</sub>, plasma C-peptide, and exogenous insulin were categorized into tertiles or quartiles. Dummy variables were created and added into multivariable models. It was observed that the magnitude of HRs remained constant for each category increment in all these variables. For example, C-peptide was categorized into tertiles, and changes from one tertile to another showed similar HR values (~1.05); thus, those variables were kept as a continuous variable in our analyses. Analyses were performed using SAS version 9 (SAS Institute, Cary, NC).

The institutional review board approved the study, and consent forms were obtained from all participants. This research was conducted in accordance with the principles of the Declaration of Helsinki.

**RESULTS**— After 16 years of follow-up, 824 individuals died (all-cause mortality); 358 deaths involved ischemic heart disease, and 137 involved stroke. Table 1 shows the characteristics of individuals with older-onset diabetes at baseline. The mean age was 68.6 ± 11.0 years, and 44.9% of the individuals were male.

Age- and sex-adjusted HR showed that HbA<sub>1c</sub> was significantly associated with increased all-cause (HR 1.13 [95% CI 1.09–1.17], per 1% increase) and ischemic heart disease mortality (1.14 [1.08–1.21], per 1% increase) (Table 2). Use of exogenous insulin was also associated with increased age- and sex-adjusted all-cause mortality (1.62 [1.41–1.87] comparing those using vs. not using exogenous insulin and 1.70 [1.23–2.34] per 1 unit · kg<sup>-1</sup> · day<sup>-1</sup> increase) and ischemic heart disease mortality (1.74 [1.40–2.16] comparing those using vs. not using exogenous insulin and 2.10 [1.32–3.36] per 1 per 1 unit · kg<sup>-1</sup> · day<sup>-1</sup> increase). Plasma C-peptide was not significantly associated with all-cause mortality (0.95 [0.87–1.03] per 1 nmol/l increase) and ischemic heart disease mortality (0.96 [0.85–1.09] per 1 nmol/l increase) in the age- and sex-adjusted models.

In multivariable analyses, HbA<sub>1c</sub> and C-peptide were significantly associated with all-cause and ischemic heart disease mortality after controlling for potential confounders (age, sex, BMI, duration of diabetes, systolic blood pressure, history

of CVD, diabetic retinopathy status, and smoking) (Table 2). We also assessed the association of HbA<sub>1c</sub>, plasma C-peptide, and amount of insulin used with mortality stratified by sex in our multivariable analysis. HbA<sub>1c</sub> was associated with increased all-cause mortality among both men (HR 1.16 [95% CI 1.08–1.25]) and women (1.09 [1.03–1.15] per 1% increase), with ischemic heart disease mortality among both men (1.24 [1.11–1.38]) and women (1.09 [1.00–1.19]), and with stroke mortality among men only (1.26 [1.10–1.53]). Higher levels of plasma C-peptide were also associated with increased all-cause mortality among women (1.18 [1.03–1.36] per 1 nmol/l increase) and ischemic heart disease mortality (1.43 [1.13–1.83]) and stroke mortality (1.65 [1.07–2.53]) among men (Table 3).

Analysis of a subsample ( $n = 369$ ) showed that the inclusion of serum lipids in our multivariable models did not change the positive association of endogenous insulin (HR 1.12 [95% CI 0.96–1.32] for all-cause and 1.09 [0.87–1.38] for ischemic heart disease) or exogenous insulin (1.17 [0.87–1.57] for all-cause and 1.18 [0.75–1.84] for ischemic heart disease mortality).

**CONCLUSIONS**— The WESDR provides a unique opportunity to investigate long-term longitudinal associations of risk factors due to its population-based design and length of follow-up. Our results showed that higher levels of endogenous insulin, measured by plasma levels of C-peptide at baseline, and HbA<sub>1c</sub> were associated with increased risk of all-cause and CVD mortality among individuals with older-onset diabetes.

In our study, exogenous insulin was associated with all-cause and ischemic heart disease mortality independently of HbA<sub>1c</sub> level. Our results are consistent with data from previous population-based studies investigating the association of endogenous insulin and mortality (6,7,19). However, most of these studies were done in the general population. In a population-based study in Finland, hyperinsulinemia had a modest association with increased cardiovascular mortality in middle-aged men (7). In a meta-analysis, Hu et al. (6) evaluated 11 European studies and concluded that plasma fasting insulin levels were associated with a 1.5-times increased chance of CVD mortality in men and ~2.7 in women after controlling for potential confounders.

In the Atherosclerosis Risk in Communities (ARIC) study, higher levels of endogenous insulin were significantly associated with increased thickness of the carotid artery wall in both men and women (20). It is believed that insulin stimulates smooth muscle cell proliferation in arterial walls and increases lipid synthesis and, as a consequence, leads to the formation of lipid lesions in arterial tissues (12). Insulin is also associated with the activation of plasminogen activator inhibitor-1, which is involved in the development of thrombosis (21).

In our study, higher levels of plasma C-peptide were also associated with increased risk of death involving stroke among men. In the ARIC study (22), elevated fasting insulin concentrations were also associated with up to a 19% increase in ischemic stroke incidence.

We found positive relationships between exogenous insulin use and mortality in our age- and sex-adjusted models. However, the use of exogenous insulin could have been targeted more to those at higher risk of developing complications and possibly with higher risk of death. Thus, the significant associations observed in the age- and sex-adjusted models were no longer statistically significant after the adjustment for several confounders and the propensity score. Although some studies showed positive associations between the use of exogenous insulin and the development of CVD, findings from randomized controlled clinical trials showed that the benefits of tight glycemic control in reducing the risk of long-term complications of diabetes (i.e., retinopathy, nephropathy, and neuropathy) outweigh the possible harmful effects of exogenous insulin treatment (13,23). The University Group Diabetes Program showed no evidence that insulin treatment influenced the risk of CVD or mortality (24) in individuals with type 2 diabetes. The Epidemiology of Diabetes Interventions and Complications (EDIC) study, a long-term follow-up of the DCCT, reported a slower rate of progression of arterial wall thickness (25) and a 42% reduced risk of cardiovascular events (26) in the intensive therapy group compared with the group under conventional treatment among type 1 diabetic patients. In the UK Prospective Diabetes Study (UKPDS), there was no increase in rates of myocardial infarction among participants assigned to receive either insulin therapy or sulfonylurea compared with those under conventional therapy (diet)

**Table 2—Association of glycemic control, C-peptide, and exogenous insulin with all-cause and cause-specific mortality**

	All-cause mortality		Ischemic heart disease mortality		Stroke	
	Age-sex adjusted	Multivariable	Age-sex adjusted	Multivariable	Age-sex adjusted	Multivariable
HbA <sub>1c</sub> (per 1%)	1.13 (1.09–1.17)	1.12 (1.07–1.17)	1.14 (1.08–1.21)	1.14 (1.06–1.22)	1.08 (0.98–1.18)	1.11 (0.99–1.23)
C-peptide (per 1 nmol/l)	0.95 (0.87–1.03)	1.15 (1.04–1.29)	0.96 (0.85–1.09)	1.19 (1.02–1.39)	0.96 (0.79–1.16)	1.09 (0.85–1.40)
Exogenous insulin (yes/no)	1.62 (1.41–1.87)	1.12 (0.93–1.35)	1.74 (1.40–2.16)	1.06 (0.80–1.42)	1.61 (1.13–2.28)	1.28 (0.81–2.02)
Exogenous insulin (per 1 unit · kg <sup>-1</sup> · day <sup>-1</sup> )	1.70 (1.23–2.34)	1.20 (0.85–1.69)	2.10 (1.32–3.36)	1.50 (0.92–2.46)	0.90 (0.37–2.14)	0.73 (0.29–1.79)

Data are HR (95% CI). Multivariable models included 1) for HbA<sub>1c</sub>: age, sex, BMI, diabetes duration, systolic blood pressure, history of CVD, presence of diabetic retinopathy, cigarette smoking, plasma C-peptide, and exogenous insulin use; 2) for exogenous insulin: sex, systolic blood pressure, smoking status, plasma C-peptide, and propensity score; 3) for plasma C-peptide: age, sex, BMI, diabetes duration, systolic blood pressure, history of CVD, presence of diabetic retinopathy, cigarette smoking, time since last meal, exogenous insulin use, and HbA<sub>1c</sub>.

Table 3—Association of glycemic control, C-peptide, and exogenous insulin with all-cause and cause-specific mortality stratified by sex

	All-cause mortality		Ischemic heart disease mortality		Stroke	
	Men	Women	Men	Women	Men	Women
HbA <sub>1c</sub> (per 1%)	1.16 (1.08–1.25)	1.09 (1.03–1.15)	1.24 (1.11–1.38)	1.09 (1.00–1.19)	1.26 (1.04–1.53)	1.02 (0.89–1.18)
C-peptide (per 1 nmol/l)	1.19 (0.99–1.41)	1.18 (1.03–1.36)	1.43 (1.13–1.83)	1.09 (0.88–1.36)	1.65 (1.07–2.53)	0.99 (0.71–1.39)
Exogenous insulin (yes/no)	0.92 (0.70–1.21)	1.35 (1.04–1.75)	0.94 (0.62–1.43)	1.23 (0.83–1.84)	0.95 (0.44–2.02)	1.52 (0.85–2.73)
Exogenous insulin (per 1 unit • kg <sup>-1</sup> • day <sup>-1</sup> )	1.05 (0.53–2.09)	1.25 (0.84–1.85)	1.21 (0.45–3.23)	1.62 (0.92–2.86)	0.10 (0.01–1.01)	1.05 (0.42–2.63)

Data are HR (95% CI). Multivariable models included 1) for HbA<sub>1c</sub>: age, sex, BMI, diabetes duration, systolic blood pressure, history of CVD, presence of diabetic retinopathy, cigarette smoking, plasma C-peptide, and exogenous insulin use; 2) for exogenous insulin: sex, systolic blood pressure, smoking status, plasma C-peptide, and propensity score; 3) for plasma C-peptide: age, sex, BMI, diabetes duration, systolic blood pressure, history of CVD, presence of diabetic retinopathy, cigarette smoking, time since last meal, exogenous insulin use, and HbA<sub>1c</sub>.

(27). However, hyperglycemia was associated with increased risk of myocardial infarction in that study (28), providing further evidence of the benefits of tight glycemic control.

One of the major concerns regarding the role of endogenous or exogenous insulin as a predictor of CVD has been their close association with other important risk factors for CVD such as obesity, dyslipidemia, and hypertension (29). Després et al. (19) reported that hyperinsulinemia was an independent risk factor for incident ischemic heart disease in the Quebec Cardiovascular Study, after controlling for these variables. In our study, adjustments for potential confounders, as well as the role of glycemic control, were addressed in our multivariable analysis. However, residual confounding is always a concern. We did not adjust for levels of serum lipids in our analysis because serum levels of cholesterol and HDL were only measured in a subsample of this cohort (n = 369) during this second examination phase. Analysis of this subsample showed that the inclusion of lipid profile in our multivariable models did not change the positive association between endogenous or exogenous with all-cause and ischemic heart disease mortality. These associations were not statistically significant possibly due to the smaller sample size. In addition, the need to have insulin therapy and the higher levels of endogenous insulin might reflect underlying insulin resistance, which is also known to be associated with some CVD risk factors such as atherosclerosis, dyslipidemia, and hypertension (8,30).

Another possible limitation might be due to loss to follow-up in the 4-year interval between the between the first and second examination phases. Because we used the second follow-up visit as baseline, if death (our main cause of loss to follow-up) was associated with insulin levels or use of exogenous insulin, we might have underestimated the strength of the association reported. In addition, women had a lower number of events (i.e., deaths) than men, and therefore low power might have also influenced our ability to detect associations among female participants.

Future reports from clinical trials such as the Bypass Angioplasty Revascularization Investigation 2 Diabetes (BARI 2D) Trial (31), with longer follow-up, may lead to a better understanding of the influence of insulin in cardiovascular morbidity

and mortality. Our results support the idea that individuals with higher endogenous insulin levels should be closely followed due to their higher risk of all-cause and ischemic heart disease mortality.

**Acknowledgments**—This work was supported, in part, by research grants EY016379 (to R.K. and B.E.K.K.) from the National Institutes of Health, Bethesda, Maryland, and by the Mentor-Based Postdoctoral Fellowship Award to R.K. from the American Diabetes Association, Alexandria, VA.

This work was partially presented at the 66th Annual Meeting of the American Diabetes Association, Washington, DC, 9–13 June 2006.

We are indebted to the WESDR cohort participants and physicians and their staffs for their continued support and participation since 1980.

**References**

1. Ruige JB, Assendelft WJ, Dekker JM, Kostense PJ, Heine RJ, Bouter LM: Insulin and risk of cardiovascular disease: a meta-analysis. *Circulation* 97:996–1001, 1998
2. Welborn TA, Knuijan MW, Ward N, Whittall DE: Serum insulin is a risk marker for coronary heart disease mortality in men but not in women. *Diabetes Res Clin Pract* 26:51–59, 1994
3. Perry IJ, Wannamethee SG, Whincup PH, Shaper AG, Walker MK, Alberti KG: Serum insulin and incident coronary heart disease in middle-aged British men. *Am J Epidemiol* 144:224–234, 1996
4. Pyorala M, Miettinen H, Laakso M, Pyorala K: Hyperinsulinemia and the risk of stroke in healthy middle-aged men: the 22-year follow-up results of the Helsinki Policemen Study. *Stroke* 29:1860–1866, 1998
5. Pyorala M, Miettinen H, Laakso M, Pyorala K: Plasma insulin and all-cause, cardiovascular, and noncardiovascular mortality: the 22-year follow-up results of the Helsinki Policemen Study. *Diabetes Care* 23:1097–1102, 2000
6. Hu G, Qiao Q, Tuomilehto J, Eliasson M, Feskens EJ, Pyorala K: Plasma insulin and cardiovascular mortality in non-diabetic European men and women: a meta-analysis of data from eleven prospective studies. *Diabetologia* 47:1245–1256, 2004
7. Lakka HM, Lakka TA, Tuomilehto J, Sivenius J, Salonen JT: Hyperinsulinemia and the risk of cardiovascular death and acute coronary and cerebrovascular events in men: the Kuopio Ischaemic Heart Disease Risk Factor Study. *Arch Intern Med* 160:1160–1168, 2000
8. Rewers M, Zaccaro D, D'Agostino R, Haffner S, Saad MF, Selby JV, Bergman R,

- Savage P: Insulin sensitivity, insulinemia, and coronary artery disease: the Insulin Resistance Atherosclerosis Study. *Diabetes Care* 27:781–787, 2004
9. Folsom AR, Szklo M, Stevens J, Liao F, Smith R, Eckfeldt JH: A prospective study of coronary heart disease in relation to fasting insulin, glucose, and diabetes: the Atherosclerosis Risk in Communities (ARIC) Study. *Diabetes Care* 20:935–942, 1997
  10. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus: the Diabetes Control and Complications Trial Research Group. *N Engl J Med* 329:977–986, 1993
  11. Nordestgaard BG, Agerholm-Larsen B, Stender S: Effect of exogenous hyperinsulinaemia on atherogenesis in cholesterol-fed rabbits. *Diabetologia* 40:512–520, 1997
  12. Stout RW: Insulin and atheroma: 20-yr perspective. *Diabetes Care* 13:631–654, 1990
  13. Muis MJ, Bots ML, Bilo HJ, Hoogma RP, Hoekstra JB, Grobbee DE, Stolk RP: High cumulative insulin exposure: a risk factor of atherosclerosis in type 1 diabetes? *Atherosclerosis* 181:185–192, 2005
  14. Ferrara A, Barrett-Connor EL, Edelstein SL: Hyperinsulinemia does not increase the risk of fatal cardiovascular disease in elderly men or women without diabetes: the Rancho Bernardo Study, 1984–1991. *Am J Epidemiol* 140:857–869, 1994
  15. Welin L, Eriksson H, Larsson B, Ohlson LO, Svardsudd K, Tibblin G: Hyperinsulinaemia is not a major coronary risk factor in elderly men: the study of men born in 1913. *Diabetologia* 35:766–770, 1992
  16. Klein R, Klein BE, Moss SE, Cruickshanks KJ: Association of ocular disease and mortality in a diabetic population. *Arch Ophthalmol* 117:1487–1495, 1999
  17. Klein R, Klein BE, Moss SE, Davis MD, DeMets DL: The Wisconsin Epidemiologic Study of Diabetic Retinopathy. III. Prevalence and risk of diabetic retinopathy when age at diagnosis is 30 or more years. *Arch Ophthalmol* 102:527–532, 1984
  18. Grading diabetic retinopathy from stereoscopic color fundus photographs—an extension of the modified Airlie House classification: ETDRS report number 10: Early Treatment Diabetic Retinopathy Study Research Group. *Ophthalmology* 98 (Suppl. 5):786–806, 1991
  19. Despres JP, Moorjani S, Lupien PJ, Tremblay A, Nadeau A, Bouchard C: Regional distribution of body fat, plasma lipoproteins, and cardiovascular disease. *Arteriosclerosis* 10:497–511, 1990
  20. Folsom AR, Eckfeldt JH, Weitzman S, Ma J, Chambless LE, Barnes RW, Cram KB, Hutchinson RG: Relation of carotid artery wall thickness to diabetes mellitus, fasting glucose and insulin, body size, and physical activity: Atherosclerosis Risk in Communities (ARIC) Study Investigators. *Stroke* 25:66–73, 1994
  21. Schneider DJ, Nordt TK, Sobel BE: Attenuated fibrinolysis and accelerated atherogenesis in type II diabetic patients. *Diabetes* 42:1–7, 1993
  22. Folsom AR, Rasmussen ML, Chambless LE, Howard G, Cooper LS, Schmidt MI, Heiss G: Prospective associations of fasting insulin, body fat distribution, and diabetes with risk of ischemic stroke: the Atherosclerosis Risk in Communities (ARIC) Study Investigators. *Diabetes Care* 22:1077–1083, 1999
  23. Tseng CH: Exogenous insulin use and hypertension in adult patients with type 2 diabetes mellitus. *Arch Intern Med* 166:1184–1189, 2006
  24. Genuth S: Exogenous insulin administration and cardiovascular risk in non-insulin-dependent and insulin-dependent diabetes mellitus. *Ann Intern Med* 124:104–109, 1996
  25. Nathan DM, Lachin J, Cleary P, Orchard T, Brillon DJ, Backlund JY, O'Leary DH, Genuth S: Intensive diabetes therapy and carotid intima-media thickness in type 1 diabetes mellitus. *N Engl J Med* 348:2294–2303, 2003
  26. Nathan DM, Cleary PA, Backlund JY, Genuth SM, Lachin JM, Orchard TJ, Raskin P, Zinman B: Intensive diabetes treatment and cardiovascular disease in patients with type 1 diabetes. *N Engl J Med* 353:2643–2653, 2005
  27. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33): UK Prospective Diabetes Study (UKPDS) Group. *Lancet* 352:837–853, 1998
  28. Adler AI, Neil HA, Manley SE, Holman RR, Turner RC: Hyperglycemia and hyperinsulinemia at diagnosis of diabetes and their association with subsequent cardiovascular disease in the United Kingdom prospective diabetes study (UKPDS 47). *Am Heart J* 138:S353–S359, 1999
  29. Vinik A, Flemmer M: Diabetes and macrovascular disease. *J Diabetes Complications* 16:235–245, 2002
  30. Reaven GM, Laws A: Insulin resistance, compensatory hyperinsulinaemia, and coronary heart disease. *Diabetologia* 37:948–952, 1994
  31. Magee MF, Isley WL: Rationale, design, and methods for glycemic control in the Bypass Angioplasty Revascularization Investigation 2 Diabetes (BARI 2D) Trial. *Am J Cardiol* 97:20G–30G, 2006