

Moving to an A1C-Based Diagnosis of Diabetes Has a Different Impact on Prevalence in Different Ethnic Groups

DIRK L. CHRISTENSEN, MSc, PhD^{1,2}
 DANIEL R. WITTE, MD, PhD²
 LYDIA KADUKA, MSc³
 MARIT E. JØRGENSEN, MD, PhD⁴
 KNUT BORCH-JOHNSEN, MD, DMSc^{2,5}

VISWANATHAN MOHAN, MD, FRCP, PhD, DSc⁶
 JONATHAN E. SHAW, MD, MRCP⁷
 ADAM G. TABÁK, MD, PhD^{8,9}
 DORTE VISTISEN, MSc, PhD²

RESEARCH DESIGN AND METHODS

Six studies including populations from different ethnic origins were included in the analysis (6–11). Populations from Denmark (Inter99 study), the U.K. (Whitehall II study, phase 7), Australia (The Australian Diabetes, Obesity and Lifestyle Study [AusDiab]), Greenland (Inuit Health in Transition Study), Kenya, and India (Chennai Urban Rural Epidemiology Study [CURES]) were included.

Data were collected during the period 1999–2009. Participants were excluded if they had missing OGTT or A1C measurements or known diabetes (self-reported). In the Inter99 study, 5.6% were not of Danish nationality and were excluded from the analyses. In the Whitehall II study, whites were included in the main analysis, whereas south Asian (4.2%) and black (1.9%) participants were analyzed in a subsidiary analysis. In the AusDiab study, only individuals born in Australia or New Zealand who spoke English at home and were not of Aboriginal/Torres Strait Islander origin were included (76.2%). In the Inuit Health in Transition Study, only Inuit participants were included in the analysis (95.5%). The participants in the study from Kenya were all black, and participants in the CURES study were all of Indian origin. A total of 23,094 participants were included in this analysis.

Participants were categorized into four groups based on their OGTT results (diabetes or no diabetes) and A1C levels (<6.5% or ≥6.5%). Exact 95% CIs were calculated for proportions (12). The probability of an A1C ≥6.5% among diabetic case subjects based on an OGTT was calculated. This probability is effectively the sensitivity of an A1C cut point of 6.5% with the WHO criteria as the gold standard. The magnitude of the difference in probability between centers was analyzed using logistic regression analysis adjusted for relevant confounders (age, sex, BMI, waist circumference, and smoking). A1C assays were aligned to the Diabetes Control and Complications Trial assay at each study center according to local labora-

OBJECTIVE — To compare screen-detected diabetes prevalence and the degree of diagnostic agreement by ethnicity with the current oral glucose tolerance test (OGTT)-based and newly proposed A1C-based diagnostic criteria.

RESEARCH DESIGN AND METHODS — Six studies (1999–2009) from Denmark, the U.K., Australia, Greenland, Kenya, and India were tested for the probability of an A1C ≥6.5% among diabetic case subjects based on an OGTT. The difference in probability between centers was analyzed by logistic regression adjusting for relevant confounders.

RESULTS — Diabetes prevalence was lower with the A1C-based diagnostic criteria in four of six studies. The probability of an A1C ≥6.5% among OGTT-diagnosed case subjects ranged widely (17.0–78.0%) by study center. Differences in diagnostic agreement between ethnic subgroups in the U.K. study were of the same magnitude as between-country comparisons.

CONCLUSIONS — A shift to an A1C-based diagnosis for diabetes will have substantially different consequences for diabetes prevalence across ethnic groups and populations.

Diabetes Care 33:580–582, 2010

Recently, an international expert committee report recommended a shift in the diagnostic tool for diabetes from the 75-g oral glucose tolerance test (OGTT) to A1C (1), thereby proposing replacement of the current World Health Organization (WHO) criteria (2). More specifically, an A1C threshold of ≥6.5% was recommended, as this value has been shown to be strongly related to retinopathy (1). In their report, the international expert committee emphasizes that it is premature to establish separate diagnostic thresholds based on

race/ethnicity and that the new diagnostic criterion is likely to identify different individuals than those identified by the WHO criteria (1). Previous studies have shown A1C levels in individuals with impaired glucose tolerance or diabetes to differ by race and ethnicity (3–5). We aimed to compare diabetes prevalence and the degree of diagnostic agreement between the OGTT- and A1C-based definitions by race/ethnicity in six different countries. Below, the term diabetes is referring to diabetes assessed by one OGTT or one A1C at screening.

From the ¹Department of International Health, Immunology and Microbiology, University of Copenhagen, Copenhagen, Denmark; the ²Steno Diabetes Center, Gentofte, Denmark; the ³Centre for Public Health Research, KEMRI, Nairobi, Kenya; the ⁴Centre for Health Research in Greenland, Directorate of Health and National Institute of Public Health, University of Southern Denmark, Copenhagen, Denmark; the ⁵Research Centre for Prevention and Health, Glostrup Hospital, Copenhagen, Denmark; the ⁶Madras Diabetes Research Foundation and Dr. Mohan's Diabetes Specialities Centre, World Health Organization Collaborating Centre for Noncommunicable Diseases Prevention and Control, Chennai, India; the ⁷Baker IDI Heart and Diabetes Institute, Caulfield, Victoria, Australia; the ⁸Department of Epidemiology and Public Health, University College London, London, U.K.; and the ⁹1st Department of Internal Medicine, Semmelweis University Faculty of Medicine, Budapest, Hungary.

Corresponding author: Dorte Vistisen, dtvs@steno.dk.

Received 5 October 2009 and accepted 4 December 2009. Published ahead of print at <http://care.diabetesjournals.org> on 15 December 2009. DOI: 10.2337/dc09-1843.

© 2010 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Table 1—Background characteristics and diabetes prevalence by OGTT and A1C diagnostic criteria in different ethnic groups

	Denmark	U.K.	Australia	Greenland	Kenya	India
	Inter99	Whitehall II (phase 7)	AusDiab	Inuit Health in Transition		CURES
Study period	1999–2001	2002–2004	1999–2000	2005–2009	2005–2006	2001–2004
<i>n</i>	5,932	4,563	7,800	2,321	296	2,182
Age (years)	46.2 ± 7.9	60.5 ± 5.9	50.9 ± 14.4	44.1 ± 14.6	37.6 ± 10.6	38.8 ± 12.6
Male subjects (%)	49.7 (48.4–51.0)	73.9 (72.6–75.2)	44.4 (43.3–45.5)	43.4 (41.4–45.5)	44.6 (38.8–50.5)	46.0 (43.9–48.1)
BMI (kg/m ²)	26.2 ± 4.5	26.5 ± 4.2	26.9 ± 4.9	26.4 ± 5.1	22.1 ± 4.6	23.0 ± 4.0
Waist circumference (cm)	86.5 ± 13.2	93.2 ± 12.0	90.6 ± 13.8	91.9 ± 13.3	79.9 ± 12.2	83.0 ± 11.4
Current smoker (%)	36.0 (34.8–37.2)	6.8 (6.1–7.6)	16.3 (15.5–17.2)	66.1 (64.1–68.0)	10.5 (7.3–14.6)	18.6 (17.0–20.3)
Fasting plasma glucose (mmol/l)	5.5 ± 0.8	5.3 ± 0.7	5.4 ± 0.7	5.7 ± 0.8	4.5 ± 0.9	5.1 ± 1.7
2-h plasma glucose (mmol/l)	6.2 ± 2.1	6.5 ± 2.0	6.2 ± 2.2	5.9 ± 2.4	5.6 ± 1.7	7.0 ± 3.5
A1C (%)	5.8 ± 0.5	5.2 ± 0.5	5.1 ± 0.4	5.7 ± 0.4	5.0 ± 0.6	5.9 ± 1.2
Diabetes by OGTT (%)	4.2 (3.7–4.8)	3.7 (3.2–4.3)	4.0 (3.6–4.4)	7.0 (6.0–8.1)	3.4 (1.6–6.1)	10.2 (9.0–11.6)
Diabetes by A1C (%)	6.7 (6.1–7.3)	1.0 (0.7–1.3)	0.7 (0.5–0.9)	3.9 (3.1–4.7)	1.4 (0.4–3.4)	12.9 (11.5–14.4)
A1C ≥6.5% given						
diabetes by OGTT (%)	42.6 (36.4–49)	25.0 (18.7–32.3)	17.0 (13.0–21.7)	29.6 (22.7–37.3)	20.0 (2.5–55.6)	78.0 (72.0–83.3)
Diabetes by OGTT given						
A1C ≥6.5% (%)	27.0 (22.7–31.7)	91.3 (79.2–97.6)	98.1 (90.1–100)	53.3 (42.5–63.9)	50.0 (6.8–93.2)	61.9 (56.0–67.6)

Data are means ± SD and proportions (95% CI). Diabetes by OGTT: fasting plasma glucose ≥7.0 mmol/l or 2-h plasma glucose ≥11.1 mmol/l. Diabetes by A1C: A1C ≥6.5%.

tory guidelines (assay details in the online appendix at <http://care.diabetesjournals.org/cgi/content/full/dc09-1843/DC1>).

RESULTS— The prevalence of diabetes was lower in four of six studies (Whitehall II, AusDiab, Inuit Health in Transition, and Kenya) with the A1C diagnostic criterion than with the OGTT (Table 1). The probability of a person having an A1C ≥6.5% given the presence of diabetes according to the OGTT differed by study center (range 17.0–78.0%). Overall, the magnitude of this difference between centers was independent of differences in age and sex distributions. Further adjustment for BMI, waist circumference, and smoking reduced the magnitude of the difference between some centers, but the overall difference remained significant ($P < 0.0001$). Pairwise comparisons between centers on this difference in probability were significant. Exceptions were the contrasts between Whitehall II and Greenland, and the comparisons between Kenya on the one hand and Inter99, Whitehall II, AusDiab, and Greenland on the other. These results did not change when adjusting for age, sex, BMI, waist circumference, and smoking.

We also performed a subsidiary analysis on the south Asian ($n = 204$) and

black ($n = 91$) minority groups in the Whitehall II study. The differences in agreement between the two diagnostic criteria for diabetes between these ethnic subgroups within Whitehall matched those observed between populations in the main analysis (online appendix). Disregarding differences in study size, the overall prevalence of diabetes was 18% lower with an A1C-based diagnostic test for diabetes. The corresponding probability of A1C ≥6.5% among diabetic case subjects based on an OGTT was 43.5%.

CONCLUSIONS— The diabetes prevalence was more likely to be lower than higher when replacing the OGTT diagnostic criteria with A1C. The rate was 63% higher in the Inter99 study, while it was 82% lower in the AusDiab study. These differences are quite substantial and may in part be due to methodological differences.

There was also a significant discrepancy in the magnitude of the OGTT and A1C diabetes diagnosis overlap between study populations of different ethnic origins, even after adjusting for age, sex, BMI, waist circumference, and smoking. However, the differences between the white populations of Inter99, Whitehall II, and AusDiab were also significant and of the same magnitude, suggesting that

part of the discrepancy in overlap can be ascribed to difference in study methodology such as the A1C assay method. On the other hand, the subsidiary analysis of the south Asian and black minorities compared with the white majority group of Whitehall II indicates that discrepancies are at least partly due to ethnic differences.

The lack of a significant difference in the pairwise comparisons between the Kenyan population and four of five other studies does not rule out a true difference in the probability of A1C ≥6.5% among OGTT-diagnosed diabetic case subjects but may be due to the limited number of individuals in the Kenyan data.

Although we cannot dismiss the possibility that part of the observed diagnostic inconsistency is due to methodological differences between studies, we can conclude that the proposed shift to A1C as the diagnostic tool for diabetes is likely to have a substantially different impact on diabetes prevalence in different populations, partly due to differences in race/ethnicity. However, future analyses on ethnic differences between studies using the same methodology are needed.

Acknowledgments— The CURES field studies were supported by the Chennai Willingdon

Corporate Foundation, Chennai, India. This is the 86th publication from Chennai Urban Rural Epidemiology Study (CURES 86). The Whitehall II study is supported by the following bodies in the U.K.: the Medical Research Council, the Economic and Social Research Council, the British Heart Foundation, the Health and Safety Executive, the Department of Health; and in the U.S.: the National Heart, Lung, and Blood Institute (HL36310), the National Institutes of Health/National Institute on Aging (AG13196), the Agency for Health Care Policy Research (HS06516), and The John D. and Catherine T. MacArthur Foundation. The Inter99 study was supported by the Danish Medical Research Council, the Danish Center for Evaluation and Health Technology Assessment, Novo Nordisk, Copenhagen County, the Danish Heart Foundation, the Danish Diabetes Association, the Danish Pharmaceutical Association, the Augustinus Foundation, the Ib Henriksen Foundation, and the Beckett Foundation. The Kenya study was supported by the Danish International Development Agency (DANIDA) (91202), the University of Copenhagen (Cluster of International Health), the Steno Diabetes Center, the Beckett Foundation, the Brødrene Hartmann Foundation, the Dagmar Marshall Foundation, the Dr. Thorvald Madsen's Grant, and the Kong Christian den Tiende's Foundation.

K.B.-J. is head of the Steno Diabetes Center, a hospital integrated in the Danish National Health Care Service but owned by Novo Nordisk. K.B.-J. holds shares in Novo Nordisk. K.B.-J. and J.E.S. were on the expert committee that published the new diagnostic recommendations. The remaining authors have no

relevant conflict of interest to declare. No other potential conflicts of interest relevant to this article were reported.

We thank all study participants for their cooperation. Likewise, we thank all coauthors of the original papers of diabetes prevalence studies on which this paper is based.

References

1. The International Expert Committee. International Expert Committee Report on the role of the A1C assay in the diagnosis of diabetes. *Diabetes Care* 2009;32:1327–1334
2. World Health Organization. *Definition, Diagnosis, and Classification of Diabetes Mellitus and its Complications: Report of a WHO Consultation. Part 1. Diagnosis and Classification of Diabetes Mellitus*. Geneva, World Health Org., 1999, p. 1–59
3. Herman WH, Dungan KM, Wolffenbuttel BHR, Buse JB, Fahrback JL, Jiang H, Martin S. Racial and ethnic differences in mean plasma glucose, hemoglobin A1c, and 1,5-anhydroglucitol in over 2000 patients with type 2 diabetes. *J Clin Endocrinol Metab* 2009;94:1689–1694
4. Herman WH, Ma Y, Uwaifo G, Haffner S, Kahn SE, Horton ES, Lachin JM, Montez MG, Brenneman T, Barrett-Connor E. Differences in A1C by race and ethnicity among patients with impaired glucose tolerance in the diabetes prevention program. *Diabetes Care* 2007;30:2453–2457
5. Cohen RM. A1C: does one size fit all? *Diabetes Care* 2007;30:2756–2758
6. Inuit Health in Transition. *Greenland Survey 2005–2008: Population Sample and Survey Methods*. Bjerregaard, Denmark, University of Southern Denmark, National Institute of Public Health, 2009, p. 3–13
7. Christensen DL, Friis H, Mwaniki DL, Kilonzo B, Tetens I, Boit MK, Omondi B, Kaduka L, Borch-Johnsen K. Prevalence of glucose intolerance and associated risk factors in rural and urban populations of different ethnic groups in Kenya. *Diabetes Res Clin Pract* 2009;84:303–310
8. Deepa M, Pradeepa R, Rema M, Mohan A, Deepa R, Shanthirani S, Mohan V. The Chennai Urban Rural Epidemiology Study (CURES): study design and methodology (urban component) (CURES-I). *J Assoc Physicians India* 2003;51:863–870
9. Dunstan DW, Zimmet PZ, Welborn TA, Cameron AJ, Shaw J, de Courten M, Jolley D, McCarty DJ. The Australian Diabetes, Obesity and Lifestyle Study (AusDiab): methods and response rates. *Diabetes Res Clin Pract* 2002;57:119–129
10. Glumer C, Jorgensen T, Borch-Johnsen K. Prevalences of diabetes and impaired glucose regulation in a Danish population: the Inter99 study. *Diabetes Care* 2003;26:2335–2340
11. Marmot MG, Smith GD, Stansfeld S, Patel C, North F, Head J, White I, Brunner E, Feeney A. Health inequalities among British civil servants: the Whitehall II study. *Lancet* 1991;337:1387–1393
12. Harte D. Non asymptotic binomial confidence intervals [article online], 2002. Available from <http://www.statsresearch.co.nz/pdf/confint.pdf>. Accessed 1 September 2009