

# Lipotoxicity and Decreased Islet Graft Survival

CRISTIANE B. LEITÃO, MD, PHD<sup>1,2</sup>  
 KARINA BERNETTI, MD<sup>1</sup>  
 THIPAPORN THARAVANIJ, MD<sup>1</sup>  
 PABLO CURE, MD<sup>1</sup>

VINCENZO LAURIOLA, EPC<sup>1</sup>  
 PER-OLOF BERGGREN, PHD<sup>3</sup>  
 CAMILLO RICORDI, MD<sup>1,4,5,6</sup>  
 RODOLFO ALEJANDRO, MD<sup>1,5</sup>

**OBJECTIVE** — To evaluate if baseline serum lipids are associated with islet graft survival in type 1 diabetes islet transplant (ITx) recipients.

**RESEARCH DESIGN AND METHODS** — Baseline fasting lipid profile was collected from 44 ITx recipients. Comparisons were performed between subjects below and above the median values of each lipid fraction. Differences in outcomes were compared by Kaplan-Meier curves and Cox regression analysis.

**RESULTS** — Subjects with baseline fasting plasma triglycerides and VLDL cholesterol above the median had shorter islet graft survival (triglycerides:  $39.7 \pm 6.1$  vs.  $61.3 \pm 6.6$  months,  $P = 0.029$ , and VLDL:  $41.5 \pm 5.7$  vs.  $62.8 \pm 7.3$  months,  $P = 0.032$ ). Total, LDL, and HDL cholesterol did not influence islet function. Triglycerides (odds ratio 2.97 [95% CI 1.03–8.52],  $P = 0.044$ ) maintained its association with graft failure after adjustments for confounders.

**CONCLUSIONS** — Higher baseline triglycerides are associated with earlier decline in islet graft function. Prospective clinical trials should address whether it is directly caused by lipotoxicity and if strategies focusing on lowering serum lipids may prolong islet graft survival.

*Diabetes Care* 33:658–660, 2010

Increased free fatty acids (FFAs) cause  $\beta$ -cell dysfunction and death (1–3).  $\beta$ -Cell lipid accumulation is mediated by defective intracellular lipid oxidation associated with leptin resistance (4). This abnormality can be corrected by insulin sensitizers or leptin therapy (5,6). In addition, FFA-induced endoplasmic reticulum stress has been implicated in  $\beta$ -cell apoptosis (7), which could be minimized by glucagon-like peptide-1 agonists (8). Islet grafts infused directly into the liver receive lipid-rich postprandial blood. Insulin secreted by islet grafts promote triglyceride deposition in surrounding hepatocytes, and multifocal steatosis has been reported in ~20% of islet transplant (ITx) recipients (9,10). Moreover, an insulin resistance phenotype and tendency

for higher serum triglycerides, factors associated with steatosis (11), were predictors of shorter graft survival (12). The aim of this study was to determine whether the lipid profile of type 1 diabetic ITx recipients is associated with islet graft survival.

## RESEARCH DESIGN AND METHODS

A retrospective cohort study was conducted in 44 type 1 diabetic subjects (37 ITx alone subjects; 7 islet after kidney [IAK] subjects), post-allogeneic ITx between 2000 and 2007 (follow-up  $40.9 \pm 23.5$  months). All patients have achieved the goal of glucose stability and avoidance of hypoglycemia, and 28 (64%) achieved insulin independence. ITx-related procedures were previously

described (13). Immunosuppressive regimen consisted of tacrolimus and sirolimus. Three IAK recipients were on corticosteroid maintenance doses. Fourteen subjects were converted to mycophenolate mofetil or mycophenolic acid, as per protocol ( $n = 6$ ) or due to side effects ( $n = 8$ ). Protocol procedures were approved by the University of Miami Health Research Ethics Board, and informed consent was obtained.

Clinical variables (demography, anthropometry, and family history of type 2 diabetes), insulin dosage per kilogram, islet autoantibodies, number of infusions and islet equivalents infused, exenatide use, and immunosuppressive medication were recorded. Outcomes were graft dysfunction (positive C-peptide, fasting glucose  $>140$  mg/dl and/or postprandial glucose  $>180$  mg/dl more than three times in a 1-week period, and/or A1C  $>6.5\%$  in two consecutive measurements) and graft failure (fasting C-peptide  $\leq 0.10$  ng/ml [two consecutive measurements in absence of hypoglycemia] or stimulated C-peptide  $\leq 0.3$  ng/ml).

Fasting lipids (total cholesterol, HDL cholesterol, VLDL cholesterol, and triglycerides) were measured by enzymatic method, and LDL cholesterol was calculated (Friedewald equation). Medians of serum lipids were calculated (total cholesterol: 177 mg/dl, LDL: 96 mg/dl, HDL: 67 mg/dl, VLDL: 13 mg/dl, and triglycerides: 65 mg/dl). Fasting glucose (hexokinase), A1C (high-performance liquid chromatography; BioRad, Richmond, CA), and autoantibodies (radioimmunoassay) were obtained. C-peptide was measured by double antibody radioimmunoassay at fasting and during a mixed-meal test (Boost high protein; Novartis/Sandoz, Nestle Nutrition). Kaplan-Meier curves (log-rank [Mantel-Cox] test) were used to compare time to outcomes (graft dysfunction and failure) between subjects with lipids below and above their median value. Adjustments for confounders were performed with Cox regression analysis.  $P$  values of  $<0.05$  (two-tailed) were significant (SSPS version 16.0).

**RESULTS** — Age at first ITx was  $43.0 \pm 8.6$  years, and diabetes duration was  $30.5 \pm 11.7$  years. Eighteen (41%) recip-


From the <sup>1</sup>Diabetes Research Institute, Miami, Florida; the <sup>2</sup>Endocrine Division of Hospital de Clínicas de Porto Alegre, Porto Alegre, Brazil; the <sup>3</sup>Rolf Luft Research Center for Diabetes and Endocrinology, Karolinska Institutet, Stockholm, Sweden; the <sup>4</sup>DeWitt Daughtry Family Department of Surgery, University of Miami-Miller School of Medicine, Miami, Florida; the <sup>5</sup>Department of Medicine, University of Miami-Miller School of Medicine, Miami, Florida; and the <sup>6</sup>Jackson Memorial Hospital, University of Miami Transplant Institute, University of Miami-Miller School of Medicine, Miami, Florida.

Corresponding author: Rodolfo Alejandro, ralejand@med.miami.edu.

Received 28 July 2009 and accepted 3 December 2009. Published ahead of print at <http://care.diabetesjournals.org> on 15 December 2009. DOI: 10.2337/dc09-1387.

© 2010 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.



**Figure 1**—Islet graft failure according to serum lipids below (black line) and above (gray line) the median values for triglycerides (A) and VLDL cholesterol (B). Comparisons were done with Kaplan-Meier curves and log-rank (Mantel-Cox) test.

ients were male and all were white. Subjects with baseline fasting plasma triglycerides above the median had earlier graft dysfunction ( $6.1 \pm 1.5$  vs.  $17.3 \pm 3.4$  months,  $P < 0.001$ ) and failure ( $39.7 \pm 6.1$  vs.  $61.3 \pm 6.6$  months,  $P = 0.029$ ) (Fig. 1A) in comparison with those with lower values. Similar results were found for VLDL cholesterol (dysfunction:  $6.2 \pm 1.6$  vs.  $16.7 \pm 3.2$  months,  $P = 0.001$ ; failure:  $41.5 \pm 5.7$  vs.  $62.8 \pm 7.3$  months,  $P = 0.032$ ) (Fig. 1B). Total, LDL, and HDL cholesterol were not determinants of islet function (data not shown).

To clarify if variables associated with higher triglycerides and/or VLDL cholesterol were determinants of shorter graft survival, we compared clinical and labo-

ratory characteristics of subjects below and above their median values (data not shown). Patients with triglycerides and/or VLDL cholesterol above median were more frequently male and IAK protocol participants, had positive family history of type 2 diabetes and overweight, were on higher doses of insulin per kilogram pre-ITx, and received a longer period of sirolimus/tacrolimus combination. These variables were included in multivariate analysis with time-to-graft-failure as the dependant variable. Triglycerides (odds ratio 2.97 [95% CI 1.03–8.52],  $P = 0.044$ ) sustained its association with graft failure, while VLDL cholesterol (3.06 [0.99–9.45],  $P = 0.052$ ) attained borderline significance. Other variables were an-

alyzed on separate multivariate models based on their biological relevance (HLA mismatches, cold ischemia duration, age, diabetes duration, BMI, autoantibodies, and immunosuppressant's serum trough levels) without modifying the results.

**CONCLUSIONS**— In ITx recipients, higher baseline triglycerides predict earlier graft dysfunction and failure. VLDL cholesterol produced similar outcomes, probably by the same mechanisms, since VLDL cholesterol is mainly composed by triglycerides.

Lipotoxicity has been pointed as one of the mechanisms responsible for  $\beta$ -cell dysfunction and death in type 2 diabetes (1). Concerns about similar effects in ITx have been raised by posttransplant image studies showing steatosis (9,10). However, the significance of steatosis in humans is not clear, being described either as a marker of good function (9) or dysfunction (10).

Recently, lipid toxicity has been studied in an animal model of ITx (14), and liver triglyceride accumulation was associated with poorer islet graft function and histological appearance (reduced  $\beta$ -cell mass and increased islet fibrosis) (14). Notably, these abnormalities were corrected by therapies targeting lipid supply (restrictive diet) or deposition (leptin gene therapy) (14).

Besides direct toxic effects, another interesting hypothesis connecting higher triglycerides and islet graft survival can be formulated. Lipoprotein apo-C3 provokes human  $\beta$ -cell apoptosis (15). This lipoprotein is a component of triglyceride-rich VLDL cholesterol molecules, raising the possibility of an extra mechanism of  $\beta$ -cell damage.

To the best of our knowledge, this is the first report of serum lipid association with islet graft survival in humans. This finding adds to the understanding of multiple and complex mechanisms of  $\beta$ -cell survival and calls attention to potential new therapies targeting serum lipids, preventing lipotoxicity and possibly leading to longer islet survival.

The study limitations are retrospective analysis and small sample size. We are aware that our results are not definitive and should be confirmed in larger cohorts with more detailed laboratorial analysis, including measurement of FFAs, and quantification of liver lipid deposition by spectroscopy. Our aim was to prove a concept and bring this idea to discussion. Higher baseline triglycerides are

associated with earlier decline in islet graft function. Prospective trials should address whether it is directly caused by lipotoxicity and if strategies reducing serum lipids may prolong islet graft survival.

**Acknowledgments**— This study was supported by the National Institutes of Health (NIH)/National Center for Research Resources (U42 RR016603, M01 RR16587), Juvenile Diabetes Research Foundation International (no. 4-2000-946, 4-2004-361), the NIH/National Institute of Diabetes and Digestive and Kidney Diseases (5 R01 DK55347, 5 R01 DK056953), the state of Florida, and the Diabetes Research Institute Foundation (Hollywood, FL). C.B.L. was the recipient of a scholarship from Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

No potential conflicts of interest relevant to this article were reported.

Parts of this article were presented in abstract form at the 69th Scientific Sessions of the American Diabetes Association, New Orleans, Louisiana, 5–9 June 2009.

#### References

1. Lee Y, Hirose H, Ohneda M, Johnson JH, McGarry JD, Unger RH. Beta-cell lipotoxicity in the pathogenesis of non-insulin-dependent diabetes mellitus of obese rats: impairment in adipocyte-beta-cell relationships. *Proc Natl Acad Sci U S A* 1994; 91:10878–10882
2. Kusminski CM, Shetty S, Orci L, Unger RH, Scherer PE. Diabetes and apoptosis: lipotoxicity. *Apoptosis* 2009;14:1484–1495
3. Zhou YP, Grill VE. Long-term exposure of rat pancreatic islets to fatty acids inhibits glucose-induced insulin secretion and biosynthesis through a glucose fatty acid cycle. *J Clin Invest* 1994;93:870–876
4. Shimabukuro M, Koyama K, Chen G, Wang MY, Trieu F, Lee Y, Newgard CB, Unger RH. Direct antidiabetic effect of leptin through triglyceride depletion of tissues. *Proc Natl Acad Sci U S A* 1997;94: 4637–4641
5. Shimabukuro M, Zhou YT, Lee Y, Unger RH. Troglitazone lowers islet fat and restores beta cell function of Zucker diabetic fatty rats. *J Biol Chem* 1998;273:3547–3550
6. Kakuma T, Lee Y, Higa M, Wang Z, Pan W, Shimomura I, Unger RH. Leptin, troglitazone, and the expression of sterol regulatory element binding proteins in liver and pancreatic islets. *Proc Natl Acad Sci U S A* 2000;97:8536–8541
7. Cunha DA, Hekerman P, Ladriere L, Bazzara-Castro A, Ortis F, Wakeham MC, Moore F, Rasschaert J, Cardozo AK, Bel-lomo E, Overbergh L, Mathieu C, Lupi R, Hai T, Herchuelz A, Marchetti P, Rutter GA, Eizirik DL, Cnop M. Initiation and execution of lipotoxic ER stress in pancreatic beta-cells. *J Cell Sci* 2008;121:2308–2318
8. Cunha DA, Ladrière L, Ortis F, Igoillo-Esteve M, Gurzov EN, Lupi R, Marchetti P, Eizirik DL, Cnop M. GLP-1 agonists protect pancreatic  $\beta$ -cells from lipotoxic endoplasmic reticulum stress through up-regulation of BiP and JunB. *Diabetes* 2009;2851–2862
9. Markmann JF, Rosen M, Siegelman ES, Soulen MC, Deng S, Barker CF, Naji A. Magnetic resonance-defined periportal steatosis following intraportal islet transplantation: a functional footprint of islet graft survival? *Diabetes* 2003;52: 1591–1594
10. Bhargava R, Senior PA, Ackerman TE, Ryan EA, Paty BW, Lakey JR, Shapiro AM. Prevalence of hepatic steatosis after islet transplantation and its relation to graft function. *Diabetes* 2004;53:1311–1317
11. Browning JD, Horton JD. Molecular mediators of hepatic steatosis and liver injury. *J Clin Invest* 2004;114:147–152
12. Leitão CB, Bernetti K, Tharavanij T, Cure P, Ricordi C, Alejandro R. Type 2 diabetes mellitus phenotype and graft survival after islet transplantation. *Transplantation* 2009;88:57–61
13. Froud T, Ricordi C, Baidal DA, Hafiz MM, Ponte G, Cure P, Pileggi A, Poggioli R, Ichii H, Khan A, Ferreira JV, Pugliese A, Esquenazi VV, Kenyon NS, Alejandro R. Islet transplantation in type 1 diabetes mellitus using cultured islets and steroid-free immunosuppression: Miami experience. *Am J Transplant* 2005;5:2037–2046
14. Lee Y, Ravazzola M, Park BH, Bashmakov YK, Orci L, Unger RH. Metabolic mechanisms of failure of intraportally transplanted pancreatic  $\beta$ -cells in rats: role of lipotoxicity and prevention by leptin. *Diabetes* 2007;56:2295–2301
15. Juntti-Berggren L, Refai E, Appelskog I, Andersson M, Imreh G, Dekki N, Uhles S, Yu L, Griffiths WJ, Zaitsev S, Leibiger I, Yang SN, Olivecrona G, Jornvall H, Berggren PO. Apolipoprotein CIII promotes Ca<sup>2+</sup>-dependent beta cell death in type 1 diabetes. *Proc Natl Acad Sci U S A* 2004; 101:10090–10094