

- gies. *Endocr Pract* 12:142–147, 2006
6. Monnier L, Lapinski H, Colette C: Contributions of fasting and postprandial plasma glucose increments to the overall diurnal hyperglycemia of type 2 diabetic patients: variations with increasing levels of HbA_{1c}. *Diabetes Care* 26:881–885, 2003
 7. Haffner SM, D'Agostino R Jr, Mykkanen L, Tracy R, Howard B, Rewers M, Selby J, Savage PJ, Saad MF: Insulin sensitivity in subjects with type 2 diabetes: relationship to cardiovascular risk factors: the Insulin Resistance Atherosclerosis Study. *Diabetes Care* 22:562–568, 1999
 8. Parikh H, Groop L: Candidate genes for type 2 diabetes. *Rev Endocr Metab Disord* 5:151–176, 2004
 9. Esposito K, Giugliano D, Nappo F, Marfella R, the Campanian Postprandial Hyperglycemia Study Group: Regression of carotid atherosclerosis by control of postprandial hyperglycemia in type 2 diabetes mellitus. *Circulation* 110:214–219, 2004
 10. Chiasson JL, Josse RG, Gomis R, Hanefeld M, Karasik A, Laakso M, the STOP-NIDDM Trial Research Group: Acarbose treatment and the risk of cardiovascular disease and hypertension in patients with impaired glucose tolerance: the STOP-NIDDM trial. *JAMA* 290:486–494, 2003