

Subclinical Inflammation and Diabetic Polyneuropathy

MONICA/KORA Survey F3 (Augsburg, Germany)

CHRISTIAN HERDER, PHD¹
 MARK LANKISCH, MD²
 DAN ZIEGLER, MD, FRCPE^{1,3}
 WOLFGANG RATHMANN, MD, MSPH⁴
 WOLFGANG KOENIG, MD⁵
 THOMAS ILLIG, PHD⁶

ANGELA DÖRING, MD⁶
 BARBARA THORAND, PHD, MPH⁶
 ROLF HOLLE, PHD⁷
 GUIDO GIANI, PHD⁴
 STEPHAN MARTIN, MD^{1,8}
 CHRISTA MEISINGER, MD, MPH⁶

OBJECTIVE — Subclinical inflammation represents a risk factor of type 2 diabetes and several diabetes complications, but data on diabetic neuropathies are scarce. Therefore, we investigated whether circulating concentrations of acute-phase proteins, cytokines, and chemokines differ among diabetic patients with or without diabetic polyneuropathy.

RESEARCH DESIGN AND METHODS — We measured 10 markers of subclinical inflammation in 227 type 2 diabetic patients with diabetic polyneuropathy who participated in the population-based MONICA/KORA Survey F3 (2004–2005; Augsburg, Germany). Diabetic polyneuropathy was diagnosed using the Michigan Neuropathy Screening Instrument (MNSI).

RESULTS — After adjustment for multiple confounders, high levels of C-reactive protein and interleukin (IL)-6 were most consistently associated with diabetic polyneuropathy, high MNSI score, and specific neuropathic deficits, whereas some inverse associations were seen for IL-18.

CONCLUSIONS — This study shows that subclinical inflammation is associated with diabetic polyneuropathy and neuropathic impairments. This association appears rather specific because only certain immune mediators and impairments are involved.

Diabetes Care 32:680–682, 2009

Subclinical inflammation is a risk factor not only for type 2 diabetes (1) but also for diabetes complications such as cardiovascular disease, stroke, diabetic nephropathy, and diabetic retinopathy (2–4). However, the association between subclinical inflammation and diabetic polyneuropathy has only been in-

vestigated in small studies without definitive results (5,6).

Therefore, the aim of this study was to investigate systematically whether patients with type 2 diabetes with or without diabetic polyneuropathy exhibit a different immune profile and whether associations between subclinical inflammation

and diabetic polyneuropathy as well as the individual components of diabetic polyneuropathy are independent of anthropometric and metabolic factors.

RESEARCH DESIGN AND METHODS

— Data are based on the MONICA/KORA Survey F3 study (2004–2005). The present study includes 227 participants of the MONICA/KORA Survey F3 with type 2 diabetes. Diabetic polyneuropathy was diagnosed based on the Michigan Neuropathy Screening Instrument (MNSI) (7). A cutoff of >2 points in the continuous MNSI score was used as previously suggested (7–10) and identified 111 individuals with diabetic polyneuropathy. The study design; assessment of neuropathic impairments; collection of information regarding demographic, anthropometric, clinical, sociodemographic, and lifestyle variables and medication; measurement of metabolic and immune mediators markers; and statistical analysis are described in the online appendix (<http://care.diabetesjournals.org/cgi/content/full/dc08-2011/DC1>).

RESULTS — The prevalence of diabetic polyneuropathy as defined by an MNSI score >2 was 48.9% (95% CI 42.4–55.4) in our study population (supplementary Figure A1 and Table A1 in the online appendix). Patients with diabetic polyneuropathy had higher levels of C-reactive protein (CRP) ($P = 0.013$), interleukin (IL)-6 ($P = 0.0091$), and interferon- γ -inducible protein-10 ($P = 0.039$) compared with those in patients without diabetic polyneuropathy, whereas leukocyte count and levels of serum amyloid A, IL-18, tumor necrosis factor- α , adiponectin, IL-8, and monocyte chemoattractant protein-1 did not differ significantly (Table A1).

Both CRP ($r = 0.23$; $P = 0.0006$) and IL-6 ($r = 0.25$; $P = 0.0001$) were highly significantly associated with the continuous MNSI score in univariate analyses and in multiple linear regression models that adjusted for anthropometric, metabolic, and lifestyle factors; anti-inflammatory medication; and recent respiratory infec-

From the ¹Institute for Clinical Diabetology, German Diabetes Center, Leibniz Center for Diabetes Research at Heinrich Heine University, Düsseldorf, Germany; the ²Department of Medicine III, Helios Clinic Wuppertal, Wuppertal, Germany; the ³Department of Medicine/Metabolic Diseases, Heinrich Heine University Düsseldorf, Düsseldorf, Germany; the ⁴Institute of Epidemiology and Biometrics, German Diabetes Center, Leibniz Center for Diabetes Research at Heinrich Heine University, Düsseldorf, Germany; the ⁵Department of Internal Medicine II-Cardiology, University of Ulm Medical Center, Ulm, Germany; the ⁶Institute of Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany; the ⁷Institute of Health Economics and Health Care Management, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany; and ⁸Sana Krankenhaus Gerresheim, Sana Kliniken Düsseldorf, Düsseldorf, Germany.

Corresponding author: Christian Herder, christian.herder@ddz.uni-duesseldorf.de.

Received 7 November 2008 and accepted 3 January 2009.

Published ahead of print at <http://care.diabetesjournals.org> on 8 January 2009. DOI: 10.2337/dc08-2011.

C.H. and M.L. contributed equally to this study.

© 2009 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Table 1—Association among MNSI, individual components of diabetic polyneuropathy, and immunological variables (multivariable-adjusted analysis)

Dependent variable	Model 1		Model 2		Model 3		Model 4	
	β	<i>P</i>	β	<i>P</i>	β	<i>P</i>	β	<i>P</i>
MNSI score								
CRP (mg/l)	0.17	0.0005	0.14	0.0043	0.14	0.0055	0.14	0.0067
IL-6 (pg/ml)	0.09	0.0015	0.08	0.0082	0.08	0.0090	0.08	0.0076
IL-18 (pg/ml)	−0.03	0.13	−0.03	0.15	−0.03	0.094	−0.03	0.11
Ankle reflex score								
CRP (mg/l)	0.14	0.0037	0.10	0.060	0.09	0.076	0.09	0.086
IL-6 (pg/ml)	0.08	0.0058	0.05	0.080	0.05	0.087	0.06	0.079
IL-18 (pg/ml)	−0.05	0.012	−0.04	0.023	−0.05	0.014	−0.05	0.013
Vibration perception score								
CRP (mg/l)	−0.01	0.075	−0.008	0.23	−0.008	0.23	−0.007	0.28
IL-6 (pg/ml)	−0.01	0.0036	−0.008	0.034	−0.008	0.031	−0.008	0.035
IL-18 (pg/ml)	0.003	0.22	0.002	0.34	0.002	0.31	0.003	0.26
Temperature perception score								
CRP (mg/l)	−0.04	0.39	−0.05	0.23	−0.06	0.17	−0.06	0.21
IL-6 (pg/ml)	−0.04	0.18	−0.04	0.17	−0.04	0.15	−0.04	0.13
IL-18 (pg/ml)	0.01	0.35	0.02	0.22	0.02	0.25	0.02	0.31
Appearance of feet								
CRP (mg/l)	0.39	0.010	0.36	0.019	0.34	0.026	0.36	0.024
IL-6 (pg/ml)	0.20	0.037	0.19	0.041	0.19	0.047	0.20	0.040
IL-18 (pg/ml)	0.03	0.65	0.01	0.85	−0.004	0.95	0.004	0.94
Pain perception (pinprick)								
CRP (mg/l)	0.11	0.55	0.003	0.99	0.01	0.94	−0.03	0.88
IL-6 (pg/ml)	0.25	0.023	0.20	0.082	0.20	0.076	0.20	0.085
IL-18 (pg/ml)	−0.15	0.021	−0.14	0.045	−0.13	0.059	−0.15	0.035
Pain or discomfort in the lower limbs								
CRP (mg/l)	−0.06	0.74	−0.04	0.79	−0.05	0.76	−0.11	0.52
IL-6 (pg/ml)	−0.06	0.59	−0.06	0.53	−0.06	0.55	−0.07	0.53
IL-18 (pg/ml)	0.10	0.095	0.11	0.087	0.10	0.10	0.10	0.10

Regression coefficients β and *P* values for MNSI score and individual impairments are from multiple linear regression models with ln-transformed concentrations of immune mediators as dependent variables. Details on the assessment of neuropathic deficits are given in the online appendix. Model 1: adjusted for age and sex; model 2: adjustment for model 1 variables plus waist circumference, duration of diabetes, A1C, hypertension, and total cholesterol; model 3: adjustment for model 2 variables plus smoking, high alcohol intake, and physical activity; model 4: adjustment for model 3 variables plus lipid-lowering medication, nonsteroidal anti-inflammatory drugs, and recent respiratory infections.

tions ($P < 0.01$ in all models; Table 1). The most consistent associations with individual neuropathic deficits were observed for CRP and IL-6, and some associations between IL-18 and neuropathic deficits were found (Table 1). High levels of CRP or IL-6 were associated with impaired ankle reflex, high vibration perception threshold, abnormal appearance of feet, and impaired pain perception (pinprick). In the case of IL-18, low levels of this cytokine were associated with impaired ankle reflex and impaired pain perception (pinprick). Associations for the other immune markers were less pronounced or absent (data not shown).

CONCLUSIONS— The present study demonstrates for the first time at the population level that several immune mediators are associated with diabetic polyneuropathy and that many of these associations

persist when adjusting for multiple potential confounders. CRP and IL-6 were most consistently associated with diabetic polyneuropathy and some neuropathic deficits. Although the strength of the correlations with the MNSI score was moderate ($r \leq 0.25$), it is important that these associations remained statistically significant after adjustment for multiple confounders, including duration of diabetes and A1C, so that immune activation in diabetic polyneuropathy cannot be explained solely as a consequence of hyperglycemia or other metabolic disturbances. Associations of CRP and IL-6 with cardiac autonomic neuropathy have been previously reported in two small studies in patients with type 1 diabetes (11,12). Furthermore, we describe significant associations of subclinical inflammation with some of the individual neuropathic impairments. Interestingly, impaired temperature per-

ception and pain or discomfort in the lower limbs were not associated with any of the measured immune mediators, indicating that the association between subclinical inflammation and diabetic polyneuropathy may only affect certain components of diabetic polyneuropathy, whereas others may be independent of immune activation.

The association between subclinical inflammation and diabetic polyneuropathy appears relatively complex because higher CRP and IL-6 levels were associated with diabetic polyneuropathy, whereas for IL-18, an inverse association was found. CRP is an acute-phase protein that is produced in the liver, and its main inducer is IL-6. This relationship is reflected by the high degree of correlation between these two mediators in the present study ($r = 0.56$; $P < 0.001$) (supplementary Table A2). Because IL-18 is

also widely considered a proinflammatory cytokine (13), the hypothesis of IL-18 being protective against at least some neurological symptoms (14,15) needs to be corroborated by further studies.

The population-based design of the MONICA/KORA Survey F3, the extensive immunological phenotyping, and the multiple regression analyses represent strengths of this study. This study also has limitations. First, MNSI as a diagnostic tool only allows identification of patients with high risk of diabetic polyneuropathy. Although most patients will have diabetic polyneuropathy, in some cases neuropathy may not have been due to diabetes, but an exclusion of other potential causes of neuropathy for this relatively large study sample was not feasible. However, MNSI has been validated in other diabetic populations (7–10). In addition, we did not conduct more objective electrophysiological tests to assess neuropathic deficits, which could have had an impact on our findings. Second, this study is cross-sectional, so one cannot distinguish between true risk factors and associations that could be due to reverse causation or simply coincidental. Third, the study lacked a nondiabetic control group, which would have been necessary to demonstrate unequivocally that the diabetic study participants had increased levels of proinflammatory markers compared with those in healthy individuals without type 2 diabetes. Fourth, the study relied on measurements of immune mediators at a single time point only and at different stages of disease. Fifth, the study was not designed to assess neuropathic pain in detail; thus, further studies are necessary to clarify the association between subclinical inflammation and this specific neuropathic symptom. Sixth, we performed multiple tests in this study, so type I errors cannot be ruled out.

Taken together, the data indicate that diabetic polyneuropathy and some of its individual impairments are significantly associated with subclinical inflammation (in particular with CRP, IL-6, and, inversely, IL-18). Prospective studies will be required to assess the time course and causal relevance of subclinical inflamma-

tion in the development of diabetic polyneuropathy in order to test whether immunomodulation could become a treatment option for patients with diabetic polyneuropathy.

Acknowledgments—The KORA research platform and the MONICA/KORA Augsburg studies are financed by the Helmholtz Zentrum München, German Research Center for Environmental Health, which is funded by the German Federal Ministry of Education, Science, Research, and Technology (Berlin, Germany), and by the State of Bavaria. This study was funded by the Deutsche Diabetes Gesellschaft (German Diabetes Association; grant to C.H.), the Federal Ministry of Health (Berlin, Germany), and the Ministry of Innovation, Science, Research and Technology of the state North Rhine-Westphalia (Düsseldorf, Germany).

No potential conflicts of interest relevant to this article were reported.

We thank all members of the Helmholtz Zentrum München and the field staff in Augsburg who were involved in the conduct of the studies. We thank Ulrike Poschen, Karin Röhrig, Gabi Gornitzka, Astrid Hoffmann (all from the German Diabetes Center), and Gerlinde Trischler (University of Ulm Medical Center) for excellent technical assistance; and Andrea Schneider and Margit Heier (both from Helmholtz Zentrum München) for data management and definition of variables.

References

- Kolb H, Mandrup-Poulsen T: An immune origin of type 2 diabetes? *Diabetologia* 48: 1038–1050, 2005
- Van Gaal LF, Mertens IL, De Block CE: Mechanisms linking obesity with cardiovascular disease. *Nature* 444:875–880, 2006
- Navarro-Gonzalez JF, Mora-Fernandez C: The role of inflammatory cytokines in diabetic nephropathy. *J Am Soc Nephrol* 19: 433–442, 2008
- Adamis AP, Berman AJ: Immunological mechanisms in the pathogenesis of diabetic retinopathy. *Semin Immunopathol* 30:65–84, 2008
- Matsuda M, Kawasaki F, Inoue H, Kanda Y, Yamada K, Harada Y, Saito M, Eto M, Matsuki M, Kaku K: Possible contribution of adipocytokines on diabetic neuropathy. *Diabetes Res Clin Pract* 66 (Suppl. 1): S121–S123, 2004
- Gonzalez-Clemente JM, Mauricio D, Richart C, Broch M, Caixas A, Megia A, Gimenez-Palop O, Simon I, Martinez-Riquelme A, Gimenez-Perez G, Vendrell J: Diabetic neuropathy is associated with activation of the TNF-alpha system in subjects with type 1 diabetes mellitus. *Clin Endocrinol (Oxf)* 63:525–529, 2005
- Feldman EL, Stevens MJ, Thomas PK, Brown MB, Canal N, Greene DA: A practical two-step quantitative clinical and electrophysiological assessment for the diagnosis and staging of diabetic neuropathy. *Diabetes Care* 17:1281–1289, 1994
- Lunetta M, Le Moli R, Grasso G, Sangiorgio L: A simplified diagnostic test for ambulatory screening of peripheral diabetic neuropathy. *Diabetes Res Clin Pract* 39:165–172, 1998
- Jia WP, Shen Q, Bao YQ, Lu JX, Li M, Xiang KS: Evaluation of the four simple methods in the diagnosis of diabetic peripheral neuropathy. *Zhonghua Yi Xue Za Zhi* 86:2707–2710, 2006 [in Chinese]
- Moghtaderi A, Bakhshipour A, Rashidi H: Validation of Michigan neuropathy screening instrument for diabetic peripheral neuropathy. *Clin Neurol Neurosurg* 108:477–481, 2006
- Gonzalez-Clemente JM, Vilardell C, Broch M, Megia A, Caixas A, Gimenez-Palop O, Richart C, Simon I, Martinez-Riquelme A, Arroyo J, Mauricio D, Vendrell J: Lower heart rate variability is associated with higher plasma concentrations of IL-6 in type 1 diabetes. *Eur J Endocrinol* 157:31–38, 2007
- Lanza GA, Pitocco D, Navarese EP, Sestito A, Sgueglia GA, Manto A, Infusino F, Musella T, Ghirlanda G, Crea F: Association between cardiac autonomic dysfunction and inflammation in type 1 diabetic patients: effect of beta-blockade. *Eur Heart J* 28:814–820, 2007
- Felderhoff-Mueser U, Schmidt OI, Oberholzer A, Bührer C, Stahel PF: IL-18: a key player in neuroinflammation and neurodegeneration? *Trends Neurosci* 28: 487–493, 2005
- Zhang XM, Duan RS, Chen Z, Quezada HC, Mix E, Winblad B, Zhu J: IL-18 deficiency aggravates kainic acid-induced hippocampal neurodegeneration in C57BL/6 mice due to an overcompensation by IL-12. *Exp Neurol* 205:64–73, 2007
- Andoh T, Kishi H, Motoki K, Nakanishi K, Kurashiki Y, Muraguchi A: Protective effect of IL-18 on kainate- and IL-1 β -induced cerebellar ataxia in mice. *J Immunol* 180:2322–2328, 2008