

Plasma sRAGE Is Independently Associated With Urinary Albumin Excretion in Type 2 Diabetes

PER M. HUMPERT, MD
STEFAN KOPF
ZDENKA DJURIC, MD
THORALF WENDT, MD

MICHAEL MORCOS, MD
PETER P. NAWROTH, MD
ANGELIKA BIERHAUS, PHD

The receptor for advanced glycation end products (RAGE) has been shown to be involved in the pathogenesis of late diabetes complications (1–5; rev. in 6). However, little is known about the physiologic function of endogenous sRAGE, a splice variant of the full-length receptor found in plasma (7). It was recently reported that plasma sRAGE levels are diminished in type 1 and type 2 diabetes and correlate inversely with intima-media thickness (8,9), suggesting a protective role of high sRAGE levels in the development of late vascular complications. In view of these data, we aimed to decipher an association of plasma sRAGE with albuminuria as a marker of microvascular damage in 90 patients with type 2 diabetes and regular glomerular filtration rate (GFR).

RESEARCH DESIGN AND METHODS

Ninety type 2 diabetic patients were recruited from family practices, being referred to our diabetes outpatient clinic for specialist treatment after giving written consent. The study was approved by the local ethics committee. For eligibility, patients had to test positive for albuminuria in two separate urine samples (>20 mg/dl albumin). Detailed patient characteristics are given in Table 1. Twenty-four-hour urine samples were collected on 3 consecutive days, and the mean of albumin excretion was taken for statistical evaluation. All blood values as well as ambulatory 24-h blood pressure

values (given as mean of 24 h) were taken on the day of study entrance. GFR was estimated using the Cockcroft-Gault formula (10). sRAGE antigen was detected in plasma by enzyme-linked immunosorbent assay (R&D Systems, Wiesbaden, Germany), as suggested by the manufacturer. For statistical evaluation, variables were correlated using Pearson's coefficient, and a *t* test was performed for comparison of sRAGE levels between subgroups of the cohort. A forward and backward stepwise multivariate linear regression model was calculated to detect independent associations of variables with albuminuria and sRAGE levels. $P \geq 0.1$ for *F* values was taken as criterion for exclusion of variables. SPSS 13.0 software was used for all statistical testing (SPSS, Chicago, IL).

RESULTS— sRAGE levels in the cohort of type 2 diabetic patients ($1,191 \pm 496$ pg/ml) were similar to the concentrations previously detected in type 1 diabetic patients using the same assay ($1,320 \pm 459$ pg/ml, [11]). In this cohort, sRAGE positively correlated with the extent of 24-h albumin excretion ($R = 0.27$, $P = 0.01$). sRAGE did not correlate with GFR ($R = -0.04$, $P = \text{NS}$), fasting glucose ($R = 0.12$, $P = \text{NS}$), or HbA_{1c} ($R = 0.05$, $P = \text{NS}$). There was also no significant correlation of mean systolic ($R = -0.02$, $P = \text{NS}$) or diastolic ($R = -0.07$, $P = \text{NS}$) 24-h blood pressure with sRAGE. Neither patients with macrovas-

cular disease nor patients affected by retinopathy or polyneuropathy showed decreased levels of sRAGE (not shown).

In a stepwise linear regression model, 24-h albumin excretion correlated independently with sRAGE ($\beta = 0.29$, $P = 0.008$, $R^2 = 0.13$; Table 1). Although there were no significant differences in sRAGE levels between men and women ($1,150 \pm 482$ vs. $1,343 \pm 531$ pg/ml, $P = \text{NS}$), sex was identified as an additional influence factor in this multivariable model ($\beta = 0.24$, $P = 0.03$). Furthermore, the results of the linear regression model did not change when ACE inhibitor/angiotensin receptor antagonist therapy, as known influence factors on albumin excretion and sRAGE levels (12–14), were added in a larger regression model.

CONCLUSIONS— This is the first study showing an independent positive correlation of plasma sRAGE with albumin excretion in type 2 diabetic patients. Since there was no significant correlation of sRAGE with GFR, filtration deficits can be excluded as an underlying cause of this finding. Hence, plasma sRAGE levels might represent an early marker of microvascular dysfunction and diabetic nephropathy in type 2 diabetes. This finding was unexpected, since previous reports stated an inverse correlation of intima-media thickness with sRAGE levels in type 1 and type 2 diabetes (8,9). In addition, a nonsignificant inverse correlation of sRAGE with albuminuria was reported in a smaller cohort of type 1 diabetic patients, of which only nine were affected by microalbuminuria (9). Previous studies (9,11) also showed contradictory results concerning plasma sRAGE levels comparing healthy control subjects and type 1 diabetic patients affected by retinopathy. In our cohort of 90 type 2 diabetic patients with albuminuria, there was no correlation of plasma sRAGE with HbA_{1c}. While the inverse correlation previously reported in type 1 and type 2 diabetes is likely due to inclusion of both patients and healthy control subjects (8,9), the inconclusive results on plasma sRAGE levels in type 1 and type 2 diabetes could be

From the Department of Medicine 1, University of Heidelberg, Heidelberg, Germany.

Address correspondence and reprint requests to Angelika Bierhaus, PhD, Medizinische Klinik 1, Universitätsklinikum Heidelberg, Im Neuenheimer Feld 410, 69120 Heidelberg, Germany. E-mail: angelika.bierhaus@med.uni-heidelberg.de.

Received for publication 16 December 2005 and accepted in revised form 6 February 2006.

Abbreviations: GFR, glomerular filtration rate; RAGE, receptor for advanced glycation end products.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

DOI: 10.2337/dc05-2478

© 2006 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Table 1—Patient characteristics and correlation between sRAGE and variables in all patients

Characteristics (n = 90)		Univariate*		Multivariate†	
		R	P	β	P
Age (years)	58.1 ± 6.9	0.03	NS	—	—
Sex (women/men)	19/71	—	—	0.24	0.03
Diabetes duration (years)	12.8 ± 7.8	0.15	NS	—	—
BMI (kg/m ²)	33.5 ± 5.9	−0.02	NS	—	—
HbA _{1c} (%)	7.4 ± 1.2	0.05	NS	—	—
Fasting glucose (mmol/l)	8.2 ± 3.0	0.12	NS	—	—
GFR (ml/min)	130.8 ± 51.3	−0.04	NS	—	—
Mean 24-h systolic blood pressure (mmHg)	139.4 ± 16.5	−0.02	NS	—	—
Mean 24-h diastolic blood pressure (mmHg)	80.4 ± 8.0	−0.07	NS	—	—
24-h albumin excretion (mg)	180.5 ± 437.9	0.27	0.01	0.29	0.008
Macrovascular complications (%)	33.3	—	—	—	—
Retinopathy (%)	26.7	—	—	—	—
Polyneuropathy (%)	50.0	—	—	—	—
ACE inhibitors/AT-R antagonists (%)	85.9	—	—	—	—
Insulin therapy (%)	58.8	—	—	—	—
Oral antidiabetic medications (%)	72.9	—	—	—	—
Statins (%)	63.5	—	—	—	—

Data are means ± SD. *Pearson's univariate correlation coefficients. †A stepwise multivariate regression analysis was performed, R² = 0.13. Data are shown for variables included in the model.

due to the composition of patient groups. Since this study identifies a positive correlation of albuminuria and sRAGE levels in type 2 diabetic patients with normal GFR, albuminuria as well as renal function should be included in future studies.

The results of this study do not support the initial hypothesis of a microvascular protection mediated by secretion of endogenous sRAGE, possibly scavenging its inflammatory ligands (6), but points to a role of sRAGE as a marker of microvascular dysfunction. Although application of exogenous sRAGE improved renal function in experimental settings (15,16), it seems questionable whether the plasma concentration of sRAGE in patients (~1,200 pg/ml or <0.1 nmol/l) is sufficient to scavenge its ligands, known to accumulate in diabetes. sRAGE was previously shown to bind proinflammatory AGEs in a saturable and dose-dependant manner at a K_d of ~75 nmol/l (17). However, a function for sRAGE as a potential marker of microvascular dysfunction in diabetic nephropathy seems plausible, since sRAGE is a splice variant of full-length RAGE, which is upregulated in diabetic nephropathy (1,2).

In conclusion, this study points to a possible role of sRAGE as a marker of early diabetic nephropathy in type 2 diabetic patients. Future prospective clinical studies will have to define the value of endogenous sRAGE as a predictive marker for diabetic nephropathy and renal failure.

Acknowledgments—This study was supported by the Lautenschläger Stiftung (to P.P.N.) and the European Foundation for the Study of Diabetes (to A.B.).

We thank K. Schaefer for her dedicated practical help.

References

- Abel M, Ritthaler U, Zhang Y, Deng Y, Schmidt AM, Greten J, Sernau T, Wahl P, Andrassy K, Ritz E, et al.: Expression of receptors for advanced glycosylated end-products in renal disease. *Nephrol Dial Transplant* 10:1662–1667, 1995
- Tanji N, Markowitz GS, Fu C, Kislinger T, Taguchi A, Pischetsrieder M, Stern D, Schmidt AM, D'Agati VD: Expression of advanced glycation end products and their cellular receptor RAGE in diabetic nephropathy and nondiabetic renal disease. *J Am Soc Nephrol* 11:1656–1666, 2000
- Yamamoto Y, Kato I, Doi T, Yonekura H, Ohashi S, Takeuchi M, Watanabe T, Yamagishi S, Sakurai S, Takasawa S, Okamoto H, Yamamoto H: Development and prevention of advanced diabetic nephropathy in RAGE-overexpressing mice. *J Clin Invest* 108:261–268, 2001
- Bucciarelli LG, Wendt T, Qu W, Lu Y, Lalla E, Rong LL, Goova MT, Moser B, Kislinger T, Lee DC, Kashyap Y, Stern DM, Schmidt AM: RAGE blockade stabilizes established atherosclerosis in diabetic apolipoprotein E-null mice. *Circulation* 106:2827–2835, 2002
- Bierhaus A, Haslbeck KM, Humpert PM, Liliensiek B, Dehmer T, Morcos M, Sayed AA, Andrassy M, Schiekofer S, Schneider

JG, Schulz JB, Heuss D, Neundorfer B, Dierl S, Huber J, Tritschler H, Schmidt AM, Schwaninger M, Haering HU, Schleicher E, Kasper M, Stern DM, Arnold B, Nawroth PP: Loss of pain perception in diabetes is dependent on a receptor of the immunoglobulin superfamily. *J Clin Invest* 114:1741–1751, 2004

- Bierhaus A, Humpert PM, Morcos M, Wendt T, Chavakis T, Arnold B, Stern DM, Nawroth PP: Understanding RAGE, the receptor for advanced glycation end products. *J Mol Med* 83:876–886, 2005
- Yonekura H, Yamamoto Y, Sakurai S, Petrova RG, Abedin MJ, Li H, Yasui K, Takeuchi M, Makita Z, Takasawa S, Okamoto H, Watanabe T, Yamamoto H: Novel splice variants of the receptor for advanced glycation end-products expressed in human vascular endothelial cells and pericytes, and their putative roles in diabetes-induced vascular injury. *Biochem J* 370:1097–1109, 2003
- Koyama H, Shoji T, Yokoyama H, Motoyama K, Mori K, Fukumoto S, Emoto M, Shoji T, Tamei H, Matsuki H, Sakurai S, Yamamoto Y, Yonekura H, Watanabe T, Yamamoto H, Nishizawa Y: Plasma level of endogenous secretory RAGE is associated with components of the metabolic syndrome and atherosclerosis. *Arterioscler Thromb Vasc Biol* 25:2587–2593, 2005
- Katakami N, Matsuhisa M, Kaneto H, Matsuoka TA, Sakamoto K, Nakatani Y, Ohtoshi K, Hayaishi-Okano R, Kosugi K, Hori M, Yamasaki Y: Decreased endogenous secretory advanced glycation end product receptor in type 1 diabetic patients: its possible association with dia-

- betic vascular complications. *Diabetes Care* 28:2716–2721, 2005
10. Sampson MJ, Drury PL: Accurate estimation of glomerular filtration rate in diabetic nephropathy from age, body weight, and serum creatinine. *Diabetes Care* 15: 609–612, 1992
 11. Challier M, Jacqueminet S, Benabdes-selam O, Grimaldi A, Beaudoux JL: Increased serum concentrations of soluble receptor for advanced glycation endproducts in patients with type 1 diabetes. *Clin Chem* 51:1749–1750, 2005
 12. Forbes JM, Thorpe SR, Thallas-Bonke V, Pete J, Thomas MC, Deemer ER, Bassal S, El-Osta A, Long DM, Panagiotopoulos S, Jerums G, Osicka TM, Cooper ME: Modulation of soluble receptor for advanced glycation end products by angiotensin-converting enzyme-1 inhibition in diabetic nephropathy. *J Am Soc Nephrol* 16: 2363–2372, 2005
 13. Parving HH, Lehnert H, Brochner-Mortensen J, Gomis R, Andersen S, Arner P: The effect of irbesartan on the development of diabetic nephropathy in patients with type 2 diabetes. *N Engl J Med* 345: 870–878, 2001
 14. Marre M, Lievre M, Vasmant D, Gallois Y, Hadjadj S, Reglier JC, Chatellier G, Mann J, Viberti GC, Passa P: Determinants of elevated urinary albumin in the 4,937 type 2 diabetic subjects recruited for the DIABHYCAR Study in Western Europe and North Africa. *Diabetes Care* 23 (Suppl. 2):B40–B48, 2000
 15. Flyvbjerg A, Denner L, Schrijvers BF, Tilton RG, Mogensen TH, Paludan SR, Rasch R: Long-term renal effects of a neutralizing RAGE antibody in obese type 2 diabetic mice. *Diabetes* 53:166–172, 2004
 16. Wendt TM, Tanji N, Guo J, Kislinger TR, Qu W, Lu Y, Bucciarelli LG, Rong LL, Moser B, Markowitz GS, Stein G, Bierhaus A, Liliensiek B, Arnold B, Nawroth PP, Stern DM, D'Agati VD, Schmidt AM: RAGE drives the development of glomerulosclerosis and implicates podocyte activation in the pathogenesis of diabetic nephropathy. *Am J Pathol* 162:1123–1137, 2003
 17. Kislinger T, Fu C, Huber B, Qu W, Taguchi A, Du Yan S, Hofmann M, Yan SF, Pischetsrieder M, Stern D, Schmidt AM: N(epsilon)-(carboxymethyl)lysine adducts of proteins are ligands for receptor for advanced glycation end products that activate cell signaling pathways and modulate gene expression. *J Biol Chem* 274:31740–31749, 1999