

OBSERVATIONS

Sulfonylurea Treatment in Young Children With Neonatal Diabetes

Dealing with hyperglycemia, hypoglycemia, and sick days

Recently, heterozygous activating mutations in the genes forming the ATP-sensitive K⁺ channel (K_{ATP} channel), *KCNJ11* and *ABCC8*, have been shown to cause neonatal diabetes (1–4). Sulfonylurea treatment restores insulin secretion in these patients (3,5,6), but information on the practical management of children with mutated K_{ATP} channels taking this medication is limited.

We report clinical aspects of the successful transfer to oral treatment in three cases of young children with *KCNJ11* and *ABCC8* mutations (Table 1). All parents gave written consent.

In case 1, a girl was transferred from

insulin to glibenclamide at 17 months (7) and had been on this treatment for 2 years. During this period, blood glucose testing decreased from 5–6 to 2–3 tests/day. As blood glucose levels were not affected by the ingestion of different amounts of carbohydrates, a free diet was initiated. Unexplained episodes of hyperglycemia were occasionally observed, and an appropriate decrease in the blood glucose level was observed with the usual dose of glibenclamide, even in cases of hyperglycemia (blood glucose 350 mg/dl). When the parents missed one dose, blood glucose was 455 mg/dl without ketosis, which was treated at home with lispro insulin dose and administration of the missed sulfonylurea dose. Only one episode of symptomatic hypoglycemia (30 mg/dl) occurred and was successfully treated with fruit juice and a temporary decrease in glibenclamide.

Minor episodes of viral respiratory disease were managed by decreasing the sulfonylurea dose to avoid hypoglycemia. One episode of rotavirus diarrhea was managed in a hospital using insulin and by stopping glibenclamide. Upon discharge, the sulfonylurea was restarted at

the previous dose. Ketones were not detected on any of these acute illnesses.

In case 2, a boy with a *KCNJ11* mutation was successfully transferred from insulin to glibenclamide at 38 months. This patient also had some episodes of unexpected hyperglycemia, which responded to taking the normal glibenclamide dose. An episode of a febrile upper respiratory tract viral illness was managed with a decrease of the glibenclamide dose, and ketones were not detected and insulin was not required.

In case 3, a girl was treated with insulin until the confirmation of a novel mutation in *ABCC8* when aged 3 years. Unexpectedly, a low dose of glibenclamide (0.1 mg · kg⁻¹ · day⁻¹) not only allowed the stopping of insulin but also resulted in episodes of asymptomatic hypoglycemia. The dose was reduced and then discontinued completely for 12 days. However, as hyperglycemia recurred tolbutamide was started, which resulted in good control without hypoglycemia.

These cases show that the use of sulfonylureas in children with K_{ATP} mutations differ from adults with type 2 diabetes. In two cases, glibenclamide was

Table 1—Clinical and molecular study of the three children with early-onset diabetes and information about their sulfonylureas treatment

Case	1	2	3
Molecular study	R201L in <i>KCNJ11</i>	R201H in <i>KCNJ11</i>	Q211K in <i>ABCC8</i>
Age diabetes was diagnosed (months)	4	6	4
Ketoacidosis at onset	+	+	+
Insulin dose before sulfonylurea treatment (units · kg ⁻¹ · day ⁻¹)	0.6	0.7	0.3
Type of insulin used	Glargine/lispro	Isophane/lispro	Glargine/lispro
A1C before sulfonylurea treatment (%)	7.3	8.9	6.7
Sulfonylureas treatment that allowed stopping insulin			
Age sulfonylurea treatment began (years)	1.4	3.1	3
Drug used for successful transfer	Glibenclamide	Glibenclamide	Glibenclamide
Dose (mg · kg ⁻¹ · day ⁻¹)	0.8	0.6	0.3
Doses/day (n)	2	3	2
Dose (mg/dose)	3.75, 3.75	2.5, 3.0, 3.5	1.5 - 1.5
Sulfonylureas treatment at the last medical visit			
Drug used at the last medical visit	Glibenclamide	Glibenclamide	Tolbutamide
Duration of sulfonylureas treatment (months)	26.0	5.0	4.5
Quantity/day (mg · kg ⁻¹ · day ⁻¹)	0.3	0.6	1.4
Doses/day (n)	3	3	1
Quantity/dose (mg/dose)	0.75, 1.25, 2.25	2.25, 2.75, 3.75	15.6
A1C during sulfonylureas treatment (%)	5.0–6.5	5.8	6.8
Medical problems observed during sulfonylureas treatment	Initial diarrhea Unexplained hyperglycemia Hyperglycemia associated with missed dose Sick day management	Sick day management Unexplained occasional hyperglycemia	Hypoglycemia with low doses of glibenclamide Transitory diarrhea with glibenclamide

best given three times a day. These children also required a higher nighttime dose to lower morning glucose, possibly as sulfonylureas act through facilitating the response to incretins in this type of diabetes (5). Sulfonylurea treatment was well tolerated; however, the risk of hypoglycemia persists, and education in their prevention and treatment should be given. Hyperglycemia, even at blood glucose level 350 mg/dl, responded to the usual dose of sulfonylureas, but if these patients consistently miss medications they risk ketoacidosis.

ETHEL CODNER, MD¹
SARAH E. FLANAGAN, BSC²
FRANCISCA UGARTE, MD³
HERNAN GARCÍA, MD⁴
TERESA VIDAL, MD⁵
SIÂN ELLARD, PHD, MRCPATH²
ANDREW T. HATTERSLEY, DM, FRCP²

From the ¹Institute of Maternal and Child Research, School of Medicine, University of Chile, Santiago, Chile; the ²Institute of Biomedical and Clinical Science, Peninsula Medical School, Exeter, U.K.; the ³Hospital Exequiel González Cortés, Santiago, Chile; the ⁴Clínica Santa María, Santiago, Chile; and the ⁵Hospital Dr. Hernán Henríquez, Temuco, Chile.

Address correspondence to Prof. Andrew T. Hattersley, Institute of Biomedical and Clinical Science, Peninsula Medical School, Barrack Road, Exeter EX2 5DW, U.K. E-mail: a.t.hattersley@ex.ac.uk.

DOI: 10.2337/dc06-2134
© 2007 by the American Diabetes Association.

Acknowledgments— This work was supported, in part, by the Fondo Nacional de Desarrollo Científico y Tecnológico, Chile (FONDECYT) Grant 1050452 (to E.C.) and by Wellcome Trust.

We are grateful to Dr. María Isabel Hernández and Alejandra Ávila, BSc, from the Institute of Maternal and Child Research, University of Chile, Santiago, Chile, for their help in the care of these patients.



References

- Gloyn AL, Pearson ER, Antcliff JF, Proks P, Bruining GJ, Slingerland AS, Howard N, Srinivasan S, Silva JMCL, Molnes J, Edghill EL, Frayling TM, Temple IK, Mackay D, Shield JPH, Sumnik Z, van Rhijn A, Wales JKH, Clark P, Gorman S, Aisenberg J, Ellard S, Njolstad PR, Ashcroft FM, Hattersley AT: Activating mutations in the gene encoding the ATP-sensitive potassium-channel subunit Kir6.2 and permanent neonatal diabetes. *N Engl J Med* 350:1838–1849, 2004
- Proks P, Arnold AL, Bruining J, Girard C, Flanagan SE, Larkin B, Colclough K, Hattersley AT, Ashcroft FM, Ellard S: A heterozygous activating mutation in the sulphonylurea receptor SUR1 (ABCC8) causes neonatal diabetes. *Hum Mol Genet* 15:1793–800, 2006
- Babenko AP, Polak M, Cave H, Busiah K, Czernichow P, Scharfmann R, Bryan J, Aguilar-Bryan L, Vaxillaire M, Froguel P: Activating mutations in the ABCC8 gene in neonatal diabetes mellitus. *N Engl J Med* 355:456–466, 2006
- Hattersley AT, Ashcroft FM: Activating mutations in Kir6.2 and neonatal diabetes: new clinical syndromes, new scientific insights, and new therapy. *Diabetes* 54:2503–2513, 2005
- Pearson ER, Flechtner I, Njolstad PR, Malecki MT, Flanagan SE, Larkin B, Ashcroft FM, Klimes I, Codner E, Iotova V, Slingerland AS, Shield J, Robert J-J, Holst JJ, Clark PM, Ellard S, Sovik O, Polak M, Hattersley AT, the Neonatal Diabetes International Collaborative Group: Switching from insulin to oral sulfonylureas in patients with diabetes due to Kir6.2 mutations. *N Engl J Med* 355:467–477, 2006
- Sagen JV, Raeder H, Hathout E, Shehadeh N, Gudmundsson K, Baevre H, Abuelo D, Phornphutkul C, Molnes J, Bell GI, Gloyn AL, Hattersley AT, Molven A, Sovik O, Njolstad PR: Permanent neonatal diabetes due to mutations in KCNJ11 encoding Kir6.2: patient characteristics and initial response to sulfonylurea therapy. *Diabetes* 53:2713–2718, 2004
- Codner E, Flanagan S, Ellard S, Garcia H, Hattersley AT: High-dose glibenclamide can replace insulin therapy despite transitory diarrhea in early-onset diabetes caused by a novel R201L Kir6.2 mutation. *Diabetes Care* 28:758–759, 2005