

Prediction of Type 2 Diabetes Using Alternate Anthropometric Measures in a Multi-Ethnic Cohort

The Insulin Resistance Atherosclerosis Study

MEREDITH F. MACKEY, MSc¹
STEVEN M. HAFFNER, MD²
LYNNE E. WAGENKNECHT, DRPH³

RALPH B. D'AGOSTINO, JR., PHD³
ANTHONY J.G. HANLEY, PHD¹

OBJECTIVE — To compare different anthropometric measures in terms of their ability to predict type 2 diabetes and to determine whether predictive ability was modified by ethnicity.

RESEARCH DESIGN AND METHODS — Anthropometry was measured at baseline for 1,073 non-Hispanic white (nHW), African American (AA), and Hispanic (HA) subjects, of whom 146 developed type 2 diabetes after 5.2 years. Logistic regression models were used with areas under the receiver operator characteristic curve (AROCs) comparing the prediction of models.

RESULTS — Waist-to-height ratio (AROC 0.678) was the most predictive measure, followed by BMI (AROC 0.674). Results were similar in nHW and HA subjects, although in AA subjects, central adiposity measures appeared to best predict type 2 diabetes.

CONCLUSIONS — Measures of central and overall adiposity predicted type 2 diabetes to a similar degree, except in AA subjects, for whom results suggested that central measures were more predictive.

Diabetes Care 32:956–958, 2009

Various assessment approaches have been used to characterize the relationship between obesity and type 2 diabetes. Computed tomography and other imaging techniques allow for direct quantification of visceral adipose tissue (VAT) and subcutaneous adipose tissue (SAT). These methods are costly and invasive, however; thus, anthropometric measurements are more commonly utilized. The majority of previous studies have assessed only BMI, waist circumference (WC), and waist-to-hip ratio (WHR), and few data are available from multi-ethnic cohorts (1–6). Our objective, therefore, was to compare a wide range of anthropometric measures in

terms of their ability to predict incident type 2 diabetes in three ethnic groups.

RESEARCH DESIGN AND METHODS — The Insulin Resistance Atherosclerosis Study has been previously described (7). The current report was comprised of 1,073 nondiabetic subjects aged 40–69 years at baseline (1992–1994); 56% were women, 40% were non-Hispanic white (nHW), 34% were Hispanic (HA), and 26% were African American (AA).

Glucose tolerance was established using 1999 World Health Organization criteria for a 2-h oral glucose tolerance test. Height, weight, WC, hip circumference

(HC), and skinfold thicknesses (triceps and subscapular) were measured following a standardized protocol (7). The sum of the skinfold thicknesses (SumSF) was used to indicate overall obesity, and the ratio of subscapular to triceps (STratio) measures was used to determine the ratio of central to peripheral body fat, with a greater ratio indicating a larger proportion of centrally distributed adipose tissue. The RJL (RJL Systems, Clinton Township, MI) method of bioelectrical impedance (BIA) was used to determine percent body fat calculated according to the Segal formula (8). RJL BIA has previously been shown to have good reliability (interclass correlation ≥ 0.99) (9). Follow-up examinations occurred an average of 5.2 years later, with 146 cases of incident diabetes diagnosed by oral glucose tolerance test.

Logistic regression analysis was used to evaluate the association between baseline measures of obesity and incident diabetes. All anthropometric variables were log transformed, and odds ratios (ORs) were presented per one standard deviation change. Two models were run for each measure: model A controlled for age, sex, and ethnicity, whereas model B controlled for age, sex, and ethnicity as well as family history of diabetes and systolic blood pressure. These variables are common in current diabetes risk scores and are easily and noninvasively obtained in clinical settings. The area under the receiver operating characteristic (AROC) curve (c-statistic) for each model was calculated and used as the primary criterion upon which to judge a model's discriminative ability (10). Different AROCs were statistically compared using the method of DeLong (11). All analyses were performed using SAS, version 9.1 (SAS Institute, Cary, NC).

RESULTS — Associations between anthropometric measures and risk of diabetes for the overall cohort are presented in Table 1. Waist-to-height ratio (WHtR) was the most predictive of diabetes

From the ¹Department of Nutritional Sciences, University of Toronto, Toronto, Ontario, Canada; the ²Division of Clinical Epidemiology, University of Texas Health Science Center at San Antonio, San Antonio, Texas; and the ³Department of Public Health Sciences, Wake Forest University, Winston Salem, North Carolina.

Corresponding author: Anthony J.G. Hanley, anthony.hanley@utoronto.ca.

Received 12 September 2008 and accepted 22 January 2009.

Published ahead of print at <http://care.diabetesjournals.org> on 5 February 2009. DOI: 10.2337/dc08-1663.

© 2009 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Table 1—Association of baseline anthropometric measures with the 5-year incidence of type 2 diabetes: the Insulin Resistance Atherosclerosis Study, 1992–1994

	Full cohort (n = 1073)		Non-Hispanic white (n = 430)		African American (n = 282)		Hispanic (n = 361)	
	OR (95% CI)	AROC	OR (95% CI)	AROC	OR (95% CI)	AROC	OR (95% CI)	AROC
WHtR	1.79 (1.49–2.15)	0.678	2.25 (1.63–3.10)	0.730	1.62 (1.16–2.26)	0.645	1.55 (1.14–2.10)	0.650
BMI	1.76 (1.47–2.10)	0.674	2.22 (1.63–3.02)	0.734	1.46 (1.04–2.03)	0.616	1.62 (1.20–2.17)	0.658
WC	1.75 (1.45–2.12)	0.667	2.25 (1.59–3.17)	0.716	1.51 (1.08–2.11)	0.630	1.59 (1.15–2.19)	0.647
% BF	2.33 (1.74–3.12)	0.662	3.43 (2.06–5.71)	0.728	1.49 (0.88–2.54)	0.597*	2.12 (1.30–3.38)	0.657
STratio	1.85 (1.44–2.37)	0.659	2.34 (1.51–3.62)	0.729	2.78 (1.65–4.70)	0.714	1.12 (0.75–1.67)	0.564*
WHR	1.93 (1.49–2.50)	0.641	2.19 (1.39–3.45)	0.670	2.52 (1.57–4.05)	0.691	1.38 (0.90–2.13)	0.582
SumSF	1.65 (1.33–2.04)	0.630*	2.01 (1.40–2.90)	0.669*	1.28 (0.85–1.93)	0.585*	1.67 (1.18–2.38)	0.638
HC	1.42 (1.20–1.69)	0.625*	1.75 (1.31–2.23)	0.694*	1.05 (0.74–1.49)	0.567*	1.45 (1.09–1.93)	0.627

Model A adjusted for age, sex, and ethnicity. ORs are estimated by logistic regression analysis and refer to one standard deviation change in the natural log of the variable. Each row is an individual model with adjustments as indicated. *AROC $P < 0.05$ compared with Table's most predictive model. % BF, percent fat from BIA.

(AROC 0.678; OR 1.79 [95% CI 1.49–2.15]), followed closely by BMI (AROC 0.674; OR 1.76 [1.47–2.10]). The remainder of the indexes had AROCs ranging from 0.667 to 0.625 with no consistent patterning of central compared with overall measures in predicting diabetes. The only measures that were significantly less predictive than WHtR were SumSF and HC ($P < 0.05$) (Table 1). The results according to AROC ranking were similar after further adjustment for family history of diabetes and systolic blood pressure (see Table A, available in the online appendix at <http://care.diabetesjournals.org/cgi/content/full/dc08-1663/DC1>), or for fasting glucose concentration.

There were no significant interactions of sex with anthropometric measures in the prediction of diabetes. Given our a priori interest in ethnic differences in diabetes prediction by anthropometric measurements, logistic models were stratified by ethnicity.

In the nHW and HA populations, BMI was most predictive of diabetes, although there was no clear pattern in central versus overall measures in outcome prediction (Table 1). In contrast, for the AA subgroup, all measures of central adiposity ranked higher in predicting diabetes than overall obesity measures. STRatio had the greatest AROC (0.714) and OR (2.78 [95% CI 1.65–4.70]), followed by WHR, WHtR, and WC. HC, percent fat from BIA, and SumSF were significantly less predictive than STRatio in model A for this ethnic group (Table 1). Results of analyses in model B were generally similar for each ethnic subgroup (see Table A in the online appendix).

CONCLUSIONS— Our main finding was that measures of central adiposity were, in general, not superior to measures of overall obesity in predicting diabetes. In the full cohort, as well as in the nHW and HA subgroups, there was no clear distinction between measures of central versus overall obesity in prediction of diabetes. In contrast, among AA subjects, the results suggested a greater diabetes predictive ability for measures of central adiposity, although there was limited statistical power within this subgroup. Previous studies that have directly measured body fat distribution have shown that AA subjects have less VAT at levels of obesity similar to those of nHW subjects (12). Thus, it is possible that accumulation of greater amounts of VAT has a more detrimental effect on risk of diabetes in AA individuals compared with populations that have higher average levels of VAT.

This is the first paper to examine the predictive ability of a wide range of anthropometric measurements on incident diabetes in three ethnic groups. In previous studies of single or pooled ethnic groups, findings have been inconsistent (1–6), and a recent meta-analysis of 32 studies showed that there was no significant difference between the relative risks of BMI, WC, and WHR for incident diabetes (13).

There are two possible explanations for the nonsuperiority of central versus overall adiposity measures in the prediction of diabetes. First, both VAT and SAT are associated with inflammatory biomarkers and disease risk (14). Second, anthropometric measurements contain a misclassification error in characterizing

body fat depots. While WC has been shown to be the simple anthropometric measure that best correlates to VAT, WC actually captures both VAT and abdominal SAT, and it has been reported that WC is more highly correlated to SAT than to VAT (15). Limitations of the present study include a modest sample size for subgroup analysis, a relatively short follow-up period, and skinfold thickness measures from two rather than four sites.

In conclusion, we found no strong evidence that anthropometric measures of central adiposity were more predictive of diabetes than measures of overall obesity in the nHW and HA populations. However, our data suggest that central obesity measures may be useful in predicting diabetes among AA subjects, although additional studies in this population are needed.

Acknowledgments— This study was supported by National Heart, Lung and Blood Institute grants U01-HL47887, U01-HL47889, U01-HL47892, U01-HL47902, DK-29867, and R01 58329 and grant M01-RR-43 from the National Institutes of Health. A.J.G.H. is supported by the Canada Research Chairs Program and the Canadian Diabetes Association.

No potential conflicts of interest relevant to this article were reported.

References

- Meisinger C, Doring A, Thorn B, Heier M, Hannelore L. Body fat distribution and risk of type 2 diabetes in the general population: Are there differences between men and women? The MONICA/KORA Augsburg Cohort Study Am J Clin Nutr 2006;84:483–489

2. Wang Y, Rimm EB, Stampfer MJ, Willet WC, Hu FB. Comparison of abdominal adiposity and overall obesity in predicting risk of type 2 diabetes among men. *Am J Clin Nutr* 2005;81:555–563
3. Tulloch-Reid MK, Williams DE, Looker HC, Hanson RL, Knowler WC. Do measures of body fat distribution provide information on the risk of type 2 diabetes in addition to measures of general obesity? Comparison of anthropometric predictors of type 2 diabetes in Pima Indians. *Diabetes Care* 2003;26:2556–2561
4. Warne D, Charles M, Hanson R, Jacobson L, Mccance D, Knowler W, Pettitt D. Comparison of body size measurements as predictors of NIDDM in Pima Indians. *Diabetes Care* 1995;18:435–439
5. Sargeant LA, Bennett FI, Forrester TE, Cooper RS, Wilks RJ. Predicting incident diabetes in Jamaica: the role of anthropometry. *Obes Res* 2002;10:792–798
6. Stevens J, Couper D, Pankow J, Folsom AR, Duncan BB, Nietro FJ, Jones D, Tyroler HA. Sensitivity and specificity of anthropometrics for the prediction of diabetes in a biracial cohort. *Obes Res* 2001;9:696–705
7. Wagenknecht LE, Mayer EJ, Rewers M, Haffner S, Selby J, Borok GM, Henkin L, Howard G, Savage PJ, Saad MF. The Insulin Resistance Atherosclerosis Study (IRAS): Objectives, design and recruitment results. *Ann Epidemiol* 1995;5:464–472
8. Segal KR, Van Loan M, Fitzgerald PI, Hodgdon JA, Van Itallie TB. Lean body mass estimation by bioelectrical impedance analysis: a four-site cross validation study. *Am J Clin Nutr* 1988;47:1–14
9. Fornetti WC, Pivarnik JM, Foley JM, Fiechtner JJ. Reliability and validity of body composition measures in female athletes. *J Appl Physiol* 87(3):1114–22, 1999 Sep.
10. Pepe MS, Janes H, Longton G, Leisenring W, Newcomb P. Limitations of the Odds Ratio in gauging the performance of a diagnostic, prognostic, or screening marker. *Am J Epidemiol* 2004;159:882–890
11. DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated receiver operating characteristic curves: A nonparametric approach. *Biometrics* 1988;44:837–845
12. Hoffman DJ, Wang Z, Gallagher D, Heymsfield S. Comparison of visceral adipose tissue mass in adults African Americans and Whites. *Obes Res* 2005;13:66–74
13. Vazquez G, Duval S, Jacobs DR Jr, Silventoinen K. Comparison of body mass index, waist circumference, and waist/hip ratio in predicting incident diabetes: A meta-analysis. *Epidemiol Rev* 2007;29:115–128
14. Pou KM, Massaro JM, Hoffman U, Vasana RS, Maurovich-Horvat P, Larson MG, Keaney JF Jr, Meigs JB, Lipinska I, Kathiresan S, Murabito JM, O'Donnell CJ, Benjamin EJ, Fox CS. Visceral and subcutaneous adipose tissue volumes are cross-sectionally related to markers of inflammation and oxidative stress: The Framingham Heart Study. *Circulation* 2007;116:1234–1241
15. Fox CS, Massaro JM, Hoffman U, Pou KM, Maurovich-Horvat P, Liu CY, Vasana RS, Murabito JM, Meigs JB, Cupples LA, D'Agostino RB Sr, O'Donnell CJ. Abdominal visceral and subcutaneous adipose tissue compartments: Association with metabolic risk factors in the Framingham Heart Study. *Circulation* 2007;116:39–48