

An Association Between Subclinical Hypothyroidism and Sight-Threatening Diabetic Retinopathy in Type 2 Diabetic Patients

JIN-KUI YANG, MD, PHD¹
WEI LIU, MD¹

JING SHI, RN¹
YI-BING LI, MD, PHD²

OBJECTIVE — To determine the relationship between subclinical hypothyroidism (SCH) and the prevalence of diabetic retinopathy in type 2 diabetic patients.

RESEARCH DESIGN AND METHODS — A total of 1,170 type 2 diabetic patients were screened for thyroid function. There were 127 type 2 diabetic patients with SCH and 200 randomly selected euthyroid type 2 diabetic patients selected. Those with more severe than moderate nonproliferative diabetic retinopathy were classified as having sight-threatening diabetic retinopathy (STDR).

RESULTS — The trend for severe retinopathy was significantly higher in the SCH group than in the euthyroid group ($\chi^2 = 20.43$, $P = 0.000$). SCH was associated with greater prevalence of diabetic retinopathy, especially STDR [odds ratio (95% CI): 4.15 (2.17–7.96), $P = 0.000$] after an adjustment for age, sex, duration of diabetes, A1C, BMI, hypertension, and LDL cholesterol. Even euthyroid patients with thyroid-stimulating hormone levels between 2.0 and <4.0 $\mu\text{IU/ml}$ had a higher rate of STDR than those between 0.4 and <2.0 $\mu\text{IU/ml}$ ($P = 0.008$).

CONCLUSIONS — Type 2 diabetic patients with SCH are associated with an increased risk of STDR.

Diabetes Care 33:1018–1020, 2010

Diabetic retinopathy is one of the most common microvascular complications and the leading cause of blindness worldwide. Common risk factors for the development of microvascular complications include duration of diabetes, poor glycemic control, elevated blood pressure, and dyslipidemia (1,2).

Subclinical hypothyroidism (SCH) is defined as an asymptomatic state characterized by a normal serum thyroxine level and elevated serum concentration of thyrotropin (thyroid-stimulating hormone [TSH]). Patients with SCH sustain an obvious increase in cardiovascular event rates (3,4). Despite this, there is a distinct lack of relevant research into risk factors associated with microvascular complications in type 2 diabetes with SCH. In fact,

only a single study conducted by Chen et al. (5) has attempted to elucidate these issues. Yet this study focused predominantly on the issue of diabetic nephropathy, as defined solely by elevated microalbuminuria, rather than retinopathy. However, in most diabetic patients with elevated microalbuminuria, other chronic kidney diseases should be considered in the absence of diabetic retinopathy (6). Our investigation examined the relationship between SCH and diabetic retinopathy in large Chinese type 2 diabetic patient samples.

RESEARCH DESIGN AND METHODS

RESEARCH DESIGN AND METHODS — A total of 1,170 subjects comprising hospital-based patients with type 2 diabetes (aged 59.3 ± 14.0

years, with a mean duration of known diabetes for 8.8 ± 6.8 years) were investigated. Those with normal free triiodothyronine (FT3), free thyroxine (FT4), and an increased TSH (≥ 4 $\mu\text{IU/ml}$) level were diagnosed with SCH. We further divided euthyroid type 2 diabetic patients into two subgroups (TSH: 2.0 to <4.0 vs. 0.4 to <2.0 $\mu\text{IU/ml}$) and compared the two based on a more stringent “normal” reference interval of TSH suggested by the National Health and Nutrition Examination Survey (NHANES) III (7). All type 2 diabetic patients with SCH and 200 cases of euthyroid type 2 diabetic patients randomly selected from the 1,170 subjects were included in the further investigation. Digital retinal photographs (two eyes \times two fields), taken using a TRC-NW7SF (Topcon, Tokyo, Japan) non-mydiatic camera at 45° , were examined independently by two qualified retinal photography graders following quality assurance protocols. The severity of diabetic retinopathy was graded based on the international clinical diabetic retinopathy severity scale (8). Eyes with more severe than moderate nonproliferative diabetic retinopathy were classified as sight-threatening diabetic retinopathy (STDR). Others were considered as having non-sight-threatening diabetic retinopathy (NSTDR). Data were analyzed by an unpaired Student *t* test, Mann-Whitney *U* test, Fisher exact test, χ^2 test, and multiple logistic regression.

RESULTS — A total of 127 (10.9%) type 2 diabetic patients had SCH. No significant differences between the SCH and euthyroid groups in age, duration of known diabetes, BMI, systolic blood pressure, or biochemical parameters were found, except diastolic blood pressure, LDL cholesterol, and glomerular filtration rate (GFR). The trend to severe retinopathy in the SCH group was significantly higher than in the non-SCH group ($\chi^2 = 20.43$, $P = 0.000$) (Table 1).

After an adjustment for potential explanatory variables (age, duration of diabetes, A1C, BMI, blood pressure, and LDL

From the ¹Department of Endocrinology, Beijing Tongren Hospital, Capital Medical University, Beijing, China; and the ²Beijing Eye Institute, Beijing, China.

Corresponding author: Jin-kui Yang, jinkui.yang@gmail.com.

Received 23 September 2009 and accepted 22 January 2010. Published ahead of print at <http://care.diabetesjournals.org> on 11 February 2010. DOI: 10.2337/dc09-1784.

© 2010 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Table 1—Characteristics of type 2 diabetic patients with or without subclinical hypothyroidism

	Euthyroidism	Subclinical hypothyroidism	P
n	200	127	—
Age (years)	58.3 ± 14.16	61.0 ± 13.7	0.110
Sex (M/F)	106/94	43/84	0.001 †
Diabetes duration (years)	8.9 ± 6.9	8.3 ± 6.6	0.678
BMI (kg/m ²)	24.6 ± 3.4	24.8 ± 3.6	0.816
Systolic blood pressure (mmHg)	160.0 ± 20.1	168.7 ± 9.9	0.251
Diastolic blood pressure (mmHg)	82.1 ± 8.9	94.0 ± 3.5	0.037
Total cholesterol (mmol/l)	4.89 ± 1.24	5.18 ± 1.19	0.103
Triglycerides (mmol/l)	1.97 ± 1.92	1.90 ± 1.37	0.559
LDL cholesterol (mmol/l)	3.07 ± 0.95	3.31 ± 1.01	0.031
HDL cholesterol (mmol/l)	1.21 ± 0.33	1.23 ± 0.36	0.958
Blood urea nitrogen (mmol/l)	5.37 ± 2.52	5.20 ± 1.80	0.842
Creatinine (μmol/l)	74.5 ± 33.6	74.0 ± 25.3	0.586
Urinary albumin excretion rate (μg/min)	12.8 (1.18–4,631)	15.8 (1.32–1,875)	0.247*
Glomerular filtration rate	65.5 (24.3–113.8)	57.7 (24.8–106.3)	0.000 *
Alanine aminotransferase (IU/l)	16.0 ± 10.5	16.0 ± 9.0	0.638
Aspartate aminotransferase (IU/l)	17.0 ± 12.9	17.0 ± 11.0	0.322
Free triiodothyronine (pmol/l)	4.04 ± 2.46	3.84 ± 0.80	0.382
Free thyroid hormone (pmol/l)	17.0 ± 2.7	13.4 ± 2.4	0.000
TSH (μIU/l)	1.34 (0.16–3.77)	5.45 (4.12–28.3)	0.000 *
Diabetic retinopathy [n (%)]			
Proliferative diabetic retinopathy	16 (8.0)	21 (16.5)	0.000 ‡
Severe NPDR	12 (6.0)	22 (17.3)	
Mild and moderate NPDR	59 (29.5)	28 (22.0)	
Normal	100 (50.0)	44 (34.6)	
Unknown	13 (6.5)	12 (9.4)	

Data are means ± SD or median (range) unless otherwise indicated. MNPDR, mild and moderate NPDR; NPDR, nonproliferative diabetic retinopathy; SNPDR, severe NPDR. Unknown, with cataracts or could not check the ocular fundus. Student *t* test; *Mann-Whitney *U* test; †Fisher exact test; ‡Pearson χ^2 test, $\chi^2 = 20.43$. Data in bold are statistically significant.

cholesterol), SCH was also associated with diabetic retinopathy [odds ratio (95% CI): 2.02 (1.18–3.46), $P = 0.011$] and STDR [4.15 (2.17–7.96), $P = 0.000$]. Additionally, type 2 diabetic patients with SCH had a significantly lower glomerular filtration rate.

In 200 euthyroid type 2 diabetic patients, 187 patients without cataracts were analyzed. Of the 187 patients, 15 of 125 patients (12.0%) with a TSH level between 0.4 and 2.0 mIU/l had STDR, while 13 of 34 patients (38.2%) with a TSH level between 2.0 and <4.0 μIU/ml had STDR. A subgroup with a higher TSH level had a significantly higher rate of STDR (Fisher exact test, $P = 0.008$).

CONCLUSIONS— SCH is a common endocrine disorder and has been reported to range from 4 to 10% in large general population screening surveys (9) and has been found to be 4–17% in diabetic patients in previous studies (10).

SCH is an asymptomatic stage of hypothyroidism, but it is often complicated with endothelial dysfunction, including capillary and precapillary arterioles, manifested by thickening of the capillary basement membrane (11). Serum high-sensitive C-reactive protein levels in subjects with SCH were higher than control subjects (12). These changes lead to small vessel dysfunction (13), increasing the prevalence of retinopathy. The reference range for “normal” TSH has been the focus of considerable debate. Some clinicians have advocated reducing the upper limit of the normal reference interval for TSH to 2.5 or 3.0 mIU/l. Individuals in the 3.0–5.0 mIU/l TSH range are considered as possibly exhibiting the early signs of developing hypothyroidism, prompting continued monitoring (14). Our study supports this, since euthyroid patients with TSH levels between 2.0 and <4.0 μIU/ml demonstrated a higher rate of STDR than patients with levels between 0.4 and <2.0 μIU/ml.

Concerning limitations, this study is a cross-sectional study and cannot offer an explanation as to why type 2 diabetic patients with SCH had a high level of C-peptide and “better” glucose control, a finding which may warrant further investigation.

Routine screening for thyroid function in the general population and thyroxin replacement in patients with TSH within 4–10 μIU/ml is still controversial (15). Our findings suggest that type 2 diabetic patients with SCH demonstrated a higher prevalence of retinopathy than their non-SCH counterparts. A randomized controlled trial may provide evidence to support screening of thyroid function and the potential treatment of SCH in type 2 diabetic patients.

Acknowledgments— This work was supported by the National 863 Program of China (2006AA02A409).

No potential conflicts of interest relevant to this article were reported.

References

- Stratton IM, Kohner EM, Aldington SJ, Turner RC, Holman RR, Manley SE, Matthews DR. UKPDS 50: risk factors for incidence and progression of retinopathy in type II diabetes over 6 years from diagnosis. *Diabetologia* 2001;44:156–163
- Barnett AH, Dixon AN, Bellary S, Hanif MW, O'hare JP, Raymond NT, Kumar S. Type 2 diabetes and cardiovascular risk in the UK south Asian community. *Diabetologia* 2006;49:2234–2246
- Hak AE, Pols HA, Visser TJ, Drexhage HA, Hofman A, Witteman JC. Subclinical hypothyroidism is an independent risk factor for atherosclerosis and myocardial infarction in elderly women: the Rotterdam Study. *Ann Intern Med* 2000;132:270–278
- Imaizumi M, Akahoshi M, Ichimaru S, Nakashima E, Hida A, Soda M, Usa T, Ashizawa K, Yokoyama N, Maeda R, Nagataki S, Eguchi K. Risk for ischemic heart disease and all-cause mortality in subclinical hypothyroidism. *J Clin Endocrinol Metab* 2004;89:3365–3370
- Chen HS, Wu TE, Jap TS, Lu RA, Wang ML, Chen RL, Lin HD. Subclinical hypothyroidism is a risk factor for nephropathy and cardiovascular diseases in type 2 diabetic patients. *Diabet Med* 2007;24:1336–1344
- KDOQI. KDOQI Clinical Practice Guidelines and Clinical Practice Recommendations for Diabetes and Chronic Kidney Disease. *Am J Kidney Dis* 2007;49:S12–S154

7. Hollowell JG, Staehling NW, Flanders WD, Hannon WH, Gunter EW, Spencer CA, Braverman LE. Serum TSH, T(4), and thyroid antibodies in the United States population (1988 to 1994): National Health and Nutrition Examination Survey (NHANES III). *J Clin Endocrinol Metab* 2002;87:489–499
8. Wilkinson CP, Ferris FL 3rd, Klein RE, Lee PP, Agardh CD, Davis M, Dills D, Kampik A, Pararajasegaram R, Verdaguer JT, the Global Diabetic Retinopathy Project Group. Proposed international clinical diabetic retinopathy and diabetic macular edema disease severity scales. *Ophthalmology* 2003;110:1677–1682
9. McDermott MT, Ridgway EC. Subclinical hypothyroidism is mild thyroid failure and should be treated. *J Clin Endocrinol Metab* 2001;86:4585–4590
10. Chubb SA, Davis WA, Inman Z, Davis TM. Prevalence and progression of subclinical hypothyroidism in women with type 2 diabetes: the Fremantle Diabetes Study. *Clin Endocrinol (Oxf)* 2005;62:480–486
11. Cappola AR, Ladenson PW. Hypothyroidism and atherosclerosis. *J Clin Endocrinol Metab* 2003;88:2438–2444
12. Tuzcu A, Bahceci M, Gokalp D, Tuzun Y, Gunes K. Subclinical hypothyroidism may be associated with elevated high-sensitive c-reactive protein (low grade inflammation) and fasting hyperinsulinemia. *Endocr J* 2005;52:89–94
13. Baycan S, Erdogan D, Caliskan M, Pamuk BO, Ciftci O, Gullu H, Yildirim A, Guvener ND, Muderrisoglu H. Coronary flow reserve is impaired in subclinical hypothyroidism. *Clin Cardiol* 2007;30:562–566
14. Duggal J, Singh S, Barsano CP, Arora R. Cardiovascular risk with subclinical hyperthyroidism and hypothyroidism: pathophysiology and management. *J Cardiometab Syndr* 2007;2:198–206
15. Helfand M. Screening for subclinical thyroid dysfunction in nonpregnant adults: a summary of the evidence for the U.S. Preventive Services Task Force. *Ann Intern Med* 2004;140:128–141