

Physical Activity/Exercise and Type 2 Diabetes

A consensus statement from the American Diabetes Association

RONALD J. SIGAL, MD, MPH^{1,2,3}
GLEN P. KENNY, PHD^{2,3}
DAVID H. WASSERMAN, PHD⁴

CARMEN CASTANEDA-SCEPPA, MD, PHD⁵
RUSSELL D. WHITE, MD⁶

For decades, exercise has been considered a cornerstone of diabetes management, along with diet and medication. However, high-quality evidence on the importance of exercise and fitness in diabetes was lacking until recent years. The present document summarizes the most clinically relevant recent advances related to people with type 2 diabetes and the recommendations that follow from these. Our recently published technical review on physical activity/exercise and type 2 diabetes (1) includes greater detail on individual studies, on prevention of diabetes, and on the physiology of exercise.

The present statement focuses on type 2 diabetes. Issues primarily germane to type 1 diabetes will be covered in a subsequent technical review and ADA Statement. The levels of evidence used are defined by the ADA in ref. 2.

PHYSICAL ACTIVITY AND PREVENTION OF TYPE 2 DIABETES

Two randomized trials each found that lifestyle interventions including ~150 min/week of physical activity and diet-induced weight loss of 5–7% reduced the risk of progression from impaired glucose tolerance (IGT) to type 2 diabetes by 58% (3,4). A cluster-randomized trial found that diet alone, exercise alone, and combined diet and exercise were equally effective in reducing

the progression from IGT to diabetes (5). Therefore, there is firm and consistent evidence that programs of increased physical activity and modest weight loss reduce the incidence of type 2 diabetes in individuals with IGT.

EFFECTS OF STRUCTURED EXERCISE INTERVENTIONS ON GLYCEMIC CONTROL AND BODY WEIGHT IN TYPE 2 DIABETES

Boulé et al. (6) undertook a systematic review and meta-analysis on the effects of structured exercise interventions in clinical trials of ≥8 weeks duration on HbA_{1c} (A1C) and body mass in people with type 2 diabetes. Postintervention A1C was significantly lower in exercise than control groups (7.65 vs. 8.31%, weighted mean difference -0.66%; $P < 0.001$). In contrast, postintervention body weight did not differ between the exercise and control groups. Meta-regression confirmed that the beneficial effect of exercise on A1C was independent of any effect on body weight. Therefore, structured exercise programs had a statistically and clinically significant beneficial effect on glycemic control, and this effect was not primarily mediated by weight loss. A subsequent meta-analysis by the same authors (7) showed that exercise intensity predicted postintervention weighted mean difference in A1C ($r = -0.91$, $P = 0.002$) to a

larger extent than exercise volume ($r = -0.46$, $P = 0.26$). These results provide support for encouraging type 2 diabetic individuals who are already exercising at moderate intensity to consider increasing the intensity of their exercise in order to obtain additional benefits in both aerobic fitness and glycemic control.

PHYSICAL ACTIVITY, AEROBIC FITNESS, AND RISK OF CARDIOVASCULAR AND OVERALL MORTALITY

Large cohort studies have found that higher levels of habitual aerobic fitness and/or physical activity are associated with significantly lower subsequent cardiovascular and overall mortality (8–10), to a much greater extent than could be explained by glucose lowering alone. For example, Church et al. (8) found that men in the lowest, second, and third quartiles of cardiorespiratory fitness had 4.5-, 2.8-, and 1.6-fold greater risk for overall mortality than men in the highest quartile of cardiorespiratory fitness, even after adjustment for age, examination year, baseline cardiovascular disease (CVD), hypercholesterolemia, hypertriglyceridemia, BMI, hypertension, parental CVD, smoking, and baseline fasting glucose levels. Essentially all of the association between higher BMI and higher mortality was explained by confounding with cardiorespiratory fitness; there was no difference in mortality among normal-weight, overweight, and obese men after adjustment for cardiorespiratory fitness.

In the same cohort, it was shown that among moderately fit subjects (21st–60th percentile for age) whose only exercise was walking, the mean time spent per week on exercise was 130 min for men and 148 min for women (11). These times are consistent with recommendations from the U.S. Surgeon General (12) and other respected bodies (13–15) stating that 150 min/week of moderate-intensity exercise should be accumulated. Moderately fit subjects whose only exercise was jogging or running reported a mean of 90 min/week for men and 92 min/week for women. These times are consistent with

From the ¹Department of Medicine, University of Ottawa, Ottawa, Ontario, Canada; the ²School of Human Kinetics, University of Ottawa, Ottawa, Ontario, Canada; the ³Clinical Epidemiology Program, Ottawa Health Research Institute, Ottawa, Ontario, Canada; the ⁴Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, Tennessee; the ⁵Jean Mayer USDA Human Nutrition Research Center on Aging, Tufts University, Boston, Massachusetts; and the ⁶Department of Community and Family Medicine, Truman Medical Center-Lakehead, University of Missouri-Kansas City School of Medicine, Kansas City, Missouri.

Address correspondence and reprint requests to Ronald J. Sigal, Clinical Epidemiology Program, Ottawa Health Research Institute, 1053 Carling Ave., Ottawa, Ontario, Canada K1Y 4E9. E-mail: rsigal@ohri.ca.

Abbreviations: CAD, coronary artery disease; CVD, cardiovascular disease; ECG, electrocardiogram; IGT, impaired glucose tolerance.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

DOI: 10.2337/dc06-9910

© 2006 by the American Diabetes Association.

an alternative and equally valid recommendation for 30 min of vigorous activity three times a week.

To our knowledge, no meta-analysis of the effects of exercise training on lipids or blood pressure in people with diabetes has been published. In the general, predominantly nondiabetic population, the effects of exercise training on blood pressure (16) and lipids (17) are relatively modest. Greater increases in HDL cholesterol and decreases in plasma triglycerides have been seen with exercise programs that are more rigorous in terms of both volume and intensity than those that have been evaluated in diabetic subjects (18). Potential mechanisms through which exercise could improve cardiovascular health were reviewed by Stewart (19). These include decreased systemic inflammation, improved early diastolic filling (reduced diastolic dysfunction), improved endothelial vasodilator function, and decreased abdominal visceral fat accumulation.

FREQUENCY OF EXERCISE—The U.S. Surgeon General's report (12) recommended that most people accumulate ≥ 30 min of moderate-intensity activity on most, ideally all, days of the week. However, most clinical trials evaluating exercise interventions in people with type 2 diabetes have used a three-times-per-week frequency (6), and many people find it easier to schedule fewer longer sessions rather than five or more weekly shorter sessions. The effect of a single bout of aerobic exercise on insulin sensitivity lasts 24–72 h depending on the duration and intensity of the activity (20). Because the duration of increased insulin sensitivity is generally not > 72 h, we recommend that there should not be more than 2 consecutive days without aerobic physical activity. The effect of resistance exercise training on insulin sensitivity may last somewhat longer (21), perhaps because some of its effects are mediated by increases in muscle mass.

EXERCISE FOR WEIGHT LOSS AND WEIGHT MAINTENANCE—The most successful programs for long-term weight control have involved combinations of diet, exercise, and behavior modification (22). Exercise alone, without concomitant dietary caloric restriction and behavior modification, tends to produce only modest weight loss of ~ 2 kg. Weight loss is typically this small primarily because

obese people often have difficulty performing sufficient exercise to create a large energy deficit, and it is relatively easy to counterbalance increased energy expenditure through exercise by eating more or becoming less active outside of exercise sessions (22). However, in a randomized trial, high-volume aerobic exercise (700 kcal/day, ~ 1 h/day of moderate-intensity aerobic exercise) produced at least as much fat loss as the equivalent degree of caloric restriction and with greater resulting improvements in insulin sensitivity (23).

The optimal volume of exercise to achieve sustained major weight loss is probably much larger than needed to achieve improved glycemic control and cardiovascular health. In observational studies (24–27), individuals who successfully maintain large weight loss over at least a year have typically performed ~ 7 h/week of moderate- to vigorous-intensity exercise. Two randomized trials found that higher exercise volumes (2,000 and 2,500 kcal/week) produced greater and more sustained weight loss than lower exercise volumes (1,000 kcal/week) (28,29).

Because of the increased evidence for health benefits from resistance training during the past 10–15 years, the American College of Sports Medicine (ACSM) now recommends that resistance training be included in fitness programs for healthy young and middle-aged adults (13), older adults (30), and adults with type 2 diabetes (15). With increased age, there is a tendency for progressive declines in muscle mass, leading to “sarcopenia,” decreased functional capacity, decreased resting metabolic rate, increased adiposity, and increased insulin resistance, and resistance training can have a major positive impact on each of these (30). Resistance exercise improves insulin sensitivity to about the same extent as aerobic exercise (31).

RESISTANCE TRAINING

Studies of resistance training in type 2 diabetes

Two clinical trials provided the strongest evidence for the value of resistance training in type 2 diabetes (32,33). In both studies, the average age of participants was 66 years, and the resistance-training regimen involved multiple exercises at high intensity (three sets, three times per week), and absolute A1C declined 1.1–1.2% in resistance-training subjects ver-

sus no significant change in control subjects. One of these trials had a second phase (months 7–12), where training was home based rather than facility based (34). Body composition changes were maintained, but exercise intensity and adherence were lower than in the first 6 months and the A1C difference between groups became statistically nonsignificant. Other published studies of resistance exercise in type 2 diabetic participants have used less intense exercise regimens (35–40). All showed beneficial effects of the resistance exercise program but to a lesser extent than the Dunstan et al. (32) and Castaneda et al. (33) trials.

Safety of resistance training

Some medical practitioners have concerns about the safety of higher-intensity resistance exercise in middle-aged and older people who are at risk of CVD. Often, the main concern is that the acute rises in blood pressure associated with higher-intensity resistance exercise might be harmful, possibly provoking stroke, myocardial ischemia, or retinal hemorrhage. We have found no evidence that resistance training actually increases these risks. No serious adverse events have been reported in any research study of resistance training in patients with type 2 diabetes, although the total number of subjects enrolled in these studies was small (32,33,35–37,40). A review of 12 resistance exercise studies in a total of 246 male cardiac rehabilitation patients found no angina, ST depression, abnormal hemodynamics, ventricular dysrhythmias, or other cardiovascular complications (41). A study of 12 men with known coronary ischemia and electrocardiogram (ECG) changes inducible by moderate aerobic exercise found that even maximal-intensity resistance exercise did not induce ECG changes (42). Therefore, moderate- to high-intensity resistance training was found to be safe even in men at significant risk of cardiac events.

While it is well known that blood pressure rises while lifting a heavy weight, it is often not appreciated that blood pressure can also rise considerably in healthy older people performing aerobic exercise. Benn et al. (43) demonstrated that in healthy older men, the myocardial demands of high-intensity resistance exercise were comparable to those occasionally needed for activities of daily living, such as climbing stairs, walking up a hill, or carrying 20–30 lb of groceries.

FLEXIBILITY EXERCISE — Flexibility exercise (stretching) has frequently been recommended as a means of increasing range of motion and hopefully reducing risk of injury. However, two systematic reviews found that flexibility exercise did not reduce risk of exercise-induced injury (44,45). Most studies included in these systematic reviews evaluated younger subjects undertaking very vigorous activity programs, such as those in military basic training; these results may not be generalizable to older subjects. Flexibility exercise has been successfully used in clinical trials as a “placebo” exercise (32,46), since there is no evidence that flexibility exercise affects metabolic control or quality of life. One small randomized trial ($n = 19$) found that range-of-motion exercises modestly decreased peak plantar pressures (47). We found no studies that directly evaluated whether flexibility training reduced the risk of ulceration or injury in people with diabetes. Therefore, we feel that there is insufficient evidence to recommend for or against flexibility exercise as a routine part of the exercise prescription.

EVALUATION OF THE DIABETIC PATIENT BEFORE RECOMMENDING AN EXERCISE PROGRAM

— For a more detailed review on this subject, see ref. 48. Before beginning a program of physical activity more vigorous than brisk walking, people with diabetes should be assessed for conditions that might be associated with increased likelihood of CVD or that might contraindicate certain types of exercise or predispose to injury, such as severe autonomic neuropathy, severe peripheral neuropathy, and preproliferative or proliferative retinopathy. The patient's age and previous physical activity level should be considered.

The role of stress testing before beginning an exercise program is controversial. There is no evidence that such testing is routinely necessary for those planning moderate-intensity activity such as walking, but it should be considered for previously sedentary individuals at moderate to high risk of CVD who want to undertake vigorous aerobic exercise exceeding the demands of everyday living (49). The prevalence of both symptomatic and asymptomatic coronary artery disease (CAD) is greater in diabetic individuals compared with nondiabetic individuals of the same age-group. However, many younger diabetic patients have relatively low absolute risk for a coronary event.

Ten-year CVD risk can be estimated using the UKPDS (U.K. Prospective Diabetes Study) Risk Engine (50) (<http://www.dtu.ox.ac.uk/index.html?maindoc%20=%20/riskengine/>), while 30-year risk of a wide variety of outcomes, including CVD, can be estimated using the American Diabetes Association's Diabetes PHD (Personal Health Decisions) (51) (<http://diabetes.org/diabetesPHD>). The lower the absolute CAD risk, the higher the likelihood of a false-positive test. A recent systematic review for the U.S. Preventive Services Task Force came to the conclusion that stress tests should usually not be recommended to detect ischemia in asymptomatic individuals at low CAD risk (<10% risk of a cardiac event over 10 years) because the risks of subsequent invasive testing triggered by false-positive tests outweighed the expected benefits from detection of previously unsuspected ischemia (52,53).

There is little or no evidence to guide practitioners in terms of whether stress testing before undertaking resistance training is necessary. One might ask whether such testing should use resistance exercise rather than the usual aerobic exercise during a stress test in such circumstances. Very few test centers would currently be equipped for such testing, and such tests have not been standardized. In contrast, aerobic exercise stress testing is widely available, standardized, and of proven prognostic value.

EXERCISE IN THE PRESENCE OF NONOPTIMAL GLYCEMIC CONTROL

Hyperglycemia

When people with type 1 diabetes are deprived of insulin for 12–48 h and ketotic, exercise can worsen the hyperglycemia and ketosis (54). Previous ADA exercise position statements suggested that physical activity be avoided if fasting glucose levels are >13.9 mmol/l (>250 mg/dl) and ketosis is present and that it be performed with caution if glucose levels are >16.7 mmol/l (300 mg/dl), even if no ketosis is present (55). We agree that vigorous activity should probably be avoided in the presence of ketosis. However, the recommendation to avoid physical activity if plasma glucose is >300 mg/dl, even in the absence of ketosis, is probably more cautious than necessary for a person with type 2 diabetes, especially in a postprandial state. In the absence of very severe insulin deficiency, light- or moderate-intensity exercise would tend to decrease plasma glucose. Therefore, pro-

vided the patient feels well, the patient is adequately hydrated, and urine and/or blood ketones are negative, it is not necessary to postpone exercise based solely on hyperglycemia.

Hypoglycemia

In individuals taking insulin and/or insulin secretagogues, physical activity can cause hypoglycemia if medication dose or carbohydrate consumption is not altered. This is particularly so at times when exogenous insulin levels are at their peaks and if physical activity is prolonged. Hypoglycemia would be rare in diabetic individuals who are not treated with insulin or insulin secretagogues. Previous ADA guidelines suggested that added carbohydrate should be ingested if preexercise glucose levels are <5.6 mmol/l (100 mg/dl) (55). We agree with this recommendation for individuals on insulin and/or an insulin secretagogue. However, the revised guidelines clarify that supplementary carbohydrate is generally not necessary for individuals treated only with diet, metformin, α -glucosidase inhibitors, and/or thiazolidinediones without insulin or a secretagogue. We found no published studies examining responses to exercise in subjects taking pramlintide (synthetic amylin analog) or exenatide (incretin analog). Neither is likely to cause hypoglycemia when used as monotherapy or combined with only metformin or a thiazolidinedione. However, patients taking either of these drugs in combination with insulin or a secretagogue may need to take additional carbohydrate before physical activity and/or reduce doses of insulin or secretagogue to avoid hypoglycemia. For a detailed discussion of medication adjustments to reduce risk of hypoglycemia, see ref. 56.

Concomitant medications other than hypoglycemic agents. Diabetic patients frequently take diuretics, β -blockers, ACE inhibitors, aspirin, and lipid-lowering agents. In most type 2 diabetic individuals, medications will not interfere with the physical activities they choose to perform, but patients and health care providers should be aware of potential problems to minimize their impact. Diuretics, especially in higher doses, can interfere with fluid and electrolyte balance. β -Blockers can blunt the adrenergic symptoms of hypoglycemia, possibly increasing risk of hypoglycemia unawareness. They can reduce maximal exercise capacity to $\sim 87\%$ of what it would be without β blockade (57) through their

negative inotropic and chronotropic effects. However, most people with type 2 diabetes do not choose to exercise at very high intensity, so this reduction of maximum capacity is generally not problematic. In people with CAD, β blockade actually increases exercise capacity by reducing coronary ischemia (58). For additional discussion on the impact of concomitant medications on physical activity, see ref. 48.

EXERCISE IN THE PRESENCE OF SPECIFIC LONG-TERM COMPLICATIONS OF DIABETES

— There is a paucity of research on the risks and benefits of exercise in the presence of diabetes complications. Therefore, recommendations in this section are based largely on “expert opinion.”

Retinopathy

Exercise and physical activity are not known to have any adverse effects on vision or the progression of nonproliferative diabetic retinopathy or macular edema (59). This applies to both resistance and aerobic training. However, in the presence of proliferative or severe nonproliferative diabetic retinopathy, vigorous aerobic or resistance exercise may be contraindicated because of the potential risk of triggering vitreous hemorrhage or retinal detachment (59). We found no research studies providing guidance as to an appropriate time interval between successful laser photocoagulation and initiation or resumption of resistance exercise. Ophthalmologists, with whom one of us (C.C.-S.) consulted, suggested waiting 3–6 months after laser photocoagulation before initiating or resuming this type of exercise.

Peripheral neuropathy

We are unaware of research studies assessing the risk of exercise-induced injury in people with peripheral sensory neuropathy. Common sense, however, would indicate that decreased pain sensation in the extremities would result in increased risk of skin breakdown and infection and of Charcot joint destruction. Therefore, in the presence of severe peripheral neuropathy, it may be best to encourage non-weight-bearing activities such as swimming, bicycling, or arm exercises (60,61).

Autonomic neuropathy

Autonomic neuropathy can increase the risk of exercise-induced injury by decreasing cardiac responsiveness to exer-

cise, postural hypotension, impaired thermoregulation due to impaired skin blood flow and sweating, impaired night vision due to impaired papillary reaction, and impaired thirst, which increases the risk of dehydration and gastroparesis with unpredictable food delivery (60). Autonomic neuropathy is also strongly associated with CVD in people with diabetes (62,63). Individuals with diabetic autonomic neuropathy should undergo cardiac investigation before beginning physical activity more intense than that to which they are accustomed. Some experts advocate thallium scintigraphy as the preferred screening technique for CVD in this high-risk population (60).

Microalbuminuria and nephropathy

Physical activity can acutely increase urinary protein excretion. The magnitude of this increase is in proportion to the acute increase in blood pressure. This finding has led some experts to recommend that people with diabetic kidney disease perform only light or moderate exercise, such that blood pressure during exercise would not rise to more than 200 mmHg (64). However, there is no evidence from clinical trials or cohort studies demonstrating that vigorous exercise increases the rate of progression of diabetic kidney disease. Several randomized trials in animals with diabetes and proteinuria showed that aerobic exercise training decreased urine protein excretion (65,66), possibly in part due to improved glycemic control, blood pressure, and insulin sensitivity. Resistance training also may be of benefit in terms of muscle mass, nutritional status, functional capacity, and glomerular filtration rate (67). Because of these encouraging findings, we believe there may be no need for any specific exercise restrictions for people with diabetic kidney disease. However, because microalbuminuria and proteinuria are associated with increased risk for CVD, it is important to perform an exercise ECG stress test in previously sedentary individuals with these conditions before beginning exercise significantly more intense than the demands of everyday living.

RECOMMENDATIONS — Levels of evidence used are those defined by the ADA in ref. 2.

Lifestyle measures for prevention of type 2 diabetes

In people with IGT, a program of weight control is recommended, including at least 150 min/week of moderate to vigorous physical activity and a healthful diet

with modest energy restriction. *Level of evidence:* A (3,4,68,69).

Aerobic exercise

The amount and intensity recommended for aerobic exercise vary according to goals.

- To improve glycemic control (6,7), assist with weight maintenance, and reduce risk of CVD (9,10), we recommend at least 150 min/week of moderate-intensity aerobic physical activity (40–60% of $\dot{V}O_{2max}$ or 50–70% of maximum heart rate) and/or at least 90 min/week of vigorous aerobic exercise (>60% of $\dot{V}O_{2max}$ or >70% of maximum heart rate). The physical activity should be distributed over at least 3 days/week and with no more than 2 consecutive days without physical activity. *Level of evidence:* A (6,7).
- Performing ≥ 4 h/week of moderate to vigorous aerobic and/or resistance exercise physical activity is associated with greater CVD risk reduction compared with lower volumes of activity (10). *Level of evidence:* B (9,10).
- For long-term maintenance of major weight loss (≥ 13.6 kg/30 lb), larger volumes of exercise (7 h/week of moderate or vigorous aerobic physical activity) may be helpful (24–27,70). *Level of evidence:* B (24–27,70).

Resistance exercise

In the absence of contraindications, people with type 2 diabetes should be encouraged to perform resistance exercise three times a week, targeting all major muscle groups, progressing to three sets of 8–10 repetitions at a weight that cannot be lifted more than 8–10 times (8–10 RM). *Level of evidence:* A (32,33). To ensure resistance exercises are performed correctly, maximize health benefits, and minimize the risk of injury, we recommend initial supervision and periodic reassessments by a qualified exercise specialist, as was done in the clinical trials (32,33).

Prevention of hypoglycemia

Those who take insulin or secretagogues should check capillary blood glucose before, after, and several hours after completing a session of physical activity, at least until they know their usual glycemic responses to such activity. For those who show a tendency to hypoglycemia during or after exercise, several strategies can be used. Doses of insulin or secretagogues

can be reduced before sessions of physical activity, extra carbohydrate can be consumed before or during physical activity, or both strategies can be implemented. *Level of evidence:* E (consensus, clinical experience).

References

1. Sigal RJ, Kenny GP, Wasserman DH, Castaneda-Sceppa C: Physical activity/exercise and type 2 diabetes. *Diabetes Care* 27: 2518–2539, 2004
2. American Diabetes Association: Introduction. *Diabetes Care* 27 (Suppl. 1):S1–S2, 2004
3. Tuomilehto J, Lindstrom J, Eriksson JG, Valle TT, Hamalainen H, Ilanne-Parikka P, Keinanen-Kiukaanniemi S, Laakso M, Louheranta A, Rastas M, Salminen V, Uusitupa M: Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med* 344:1343–1350, 2001
4. Diabetes Prevention Program Research Group: Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 346:393–403, 2002
5. Pan XR, Li GW, Hu YH, Wang JX, Yang WY, An ZX, Hu ZX, Lin J, Xiao JZ, Cao HB, Liu PA, Jiang XG, Jiang YY, Wang JP, Zheng H, Zhang H, Bennett PH, Howard BV: Effects of diet and exercise in preventing NIDDM in people with impaired glucose tolerance: the Da Qing IGT and Diabetes Study. *Diabetes Care* 20:537–544, 1997
6. Boulé NG, Haddad E, Kenny GP, Wells GA, Sigal RJ: Effects of exercise on glycemic control and body mass in type 2 diabetes mellitus: a meta-analysis of controlled clinical trials. *JAMA* 286:1218–1227, 2001
7. Boulé NG, Kenny GP, Haddad E, Wells GA, Sigal RJ: Meta-analysis of the effect of structured exercise training on cardiorespiratory fitness in type 2 diabetes mellitus. *Diabetologia* 46:1071–1081, 2003
8. Church TS, Cheng YJ, Earnest CP, Barlow CE, Gibbons LW, Priest EL, Blair SN: Exercise capacity and body composition as predictors of mortality among men with diabetes. *Diabetes Care* 27:83–88, 2004
9. Wei M, Gibbons LW, Kampert JB, Nichaman MZ, Blair SN: Low cardiorespiratory fitness and physical inactivity as predictors of mortality in men with type 2 diabetes. *Ann Intern Med* 132:605–611, 2000
10. Hu FB, Stamper MJ, Solomon C, Liu S, Colditz GA, Speizer FE, Willett WC, Manson JE: Physical activity and risk for cardiovascular events in diabetic women. *Ann Intern Med* 134:96–105, 2001
11. Stofan JR, DiPietro L, Davis D, Kohl HW 3rd, Blair SN: Physical activity patterns associated with cardiorespiratory fitness and reduced mortality: the Aerobics Center Longitudinal Study. *Am J Public Health* 88:1807–1813, 1998
12. US Department of Health and Human Services: *Physical Activity and Health: A Report of the Surgeon General*. Atlanta, GA, U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, 1996
13. American College of Sports Medicine: American College of Sports Medicine Position Stand: the recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness, and flexibility in healthy adults. *Med Sci Sports Exerc* 30:975–991, 1998
14. Pate RR, Pratt M, Blair SN, Haskell WL, Macera CA, Bouchard C, Buchner D, Ettinger W, Heath GW, King AC, et al.: Physical activity and public health: a recommendation from the Centers for Disease Control and Prevention and the American College of Sports Medicine. *JAMA* 273:402–407, 1995
15. Albright A, Franz M, Hornsby G, Kriska A, Marrero D, Ullrich I, Verity LS: American College of Sports Medicine position stand: exercise and type 2 diabetes. *Med Sci Sports Exerc* 32:1345–1360, 2000
16. Whelton SP, Chin A, Xin X, He J: Effect of aerobic exercise on blood pressure: a meta-analysis of randomized, controlled trials. *Ann Intern Med* 136:493–503, 2002
17. Leon AS, Sanchez OA: Response of blood lipids to exercise training alone or combined with dietary intervention. *Med Sci Sports Exerc* 33:S502–S515, 2001
18. Kraus WE, Houmard JA, Duscha BD, Knetzer KJ, Wharton MB, McCartney JS, Bales CW, Henes S, Samsa GP, Otvos JD, Kulkarni KR, Slentz CA: Effects of the amount and intensity of exercise on plasma lipoproteins. *N Engl J Med* 347: 1483–1492, 2002
19. Stewart KJ: Exercise training and the cardiovascular consequences of type 2 diabetes and hypertension: plausible mechanisms for improving cardiovascular health. *JAMA* 288:1622–1631, 2002
20. Wallberg-Henriksson H, Rincon J, Zierath JR: Exercise in the management of non-insulin-dependent diabetes mellitus. *Sports Med* 25:25–35, 1998
21. Zachwieja JJ, Toffolo G, Cobelli C, Bier DM, Yarasheski KE: Resistance exercise and growth hormone administration in older men: effects on insulin sensitivity and secretion during a stable-label intravenous glucose tolerance test. *Metabolism* 45:254–260, 1996
22. Wing RR: Exercise and weight control. In *Handbook of Exercise in Diabetes*. 2nd ed. Ruderman N, Devlin JT, Schneider SH, Kriska A, Eds. Alexandria, VA, American Diabetes Association, 2002, p. 355–364
23. Ross R, Dagnone D, Jones PJ, Smith H, Paddags A, Hudson R, Janssen I: Reduction in obesity and related comorbid conditions after diet-induced weight loss or exercise-induced weight loss in men: a randomized, controlled trial. *Ann Intern Med* 133:92–103, 2000
24. Pavlou KN, Krey S, Steffee WP: Exercise as an adjunct to weight loss and maintenance in moderately obese subjects. *Am J Clin Nutr* 49:1115–1123, 1989
25. Schoeller DA, Shay K, Kushner RF: How much physical activity is needed to minimize weight gain in previously obese women? *Am J Clin Nutr* 66:551–556, 1997
26. Weinsier RL, Hunter GR, Desmond RA, Byrne NM, Zuckerman PA, Darnell BE: Free-living activity energy expenditure in women successful and unsuccessful at maintaining a normal body weight. *Am J Clin Nutr* 75:499–504, 2002
27. Saris WH, Blair SN, van Baak MA, Eaton SB, Davies PS, Di Pietro L, Fogelholm M, Rissanen A, Schoeller D, Swinburn B, Tremblay A, Westerterp KR, Wyatt H: How much physical activity is enough to prevent unhealthy weight gain? Outcome of the IASO 1st Stock Conference and consensus statement. *Obes Rev* 4:101–114, 2003
28. Jakicic JM, Marcus BH, Gallagher KI, Napolitano M, Lang W: Effect of exercise duration and intensity on weight loss in overweight, sedentary women: a randomized trial. *JAMA* 290:1323–1330, 2003
29. Jeffery RW, Wing RR, Sherwood NE, Tate DF: Physical activity and weight loss: does prescribing higher physical activity goals improve outcome? *Am J Clin Nutr* 78: 684–689, 2003
30. American College of Sports Medicine: American College of Sports Medicine Position Stand: exercise and physical activity for older adults. *Med Sci Sports Exerc* 30: 992–1008, 1998
31. Ivy JL: Role of exercise training in the prevention and treatment of insulin resistance and non-insulin-dependent diabetes mellitus. *Sports Med* 24:321–336, 1997
32. Dunstan DW, Daly RM, Owen N, Jolley D, de Courten M, Shaw J, Zimmet P: High-intensity resistance training improves glycemic control in older persons with type 2 diabetes. *Diabetes Care* 25:1729–1735, 2002
33. Castaneda C, Layne JE, Munoz-Orians L, Gordon PL, Walsmith J, Foldvari M, Roubenoff R, Tucker KL, Nelson ME: A randomized controlled trial of resistance exercise training to improve glycemic control in older adults with type 2 diabetes. *Diabetes Care* 25:2335–2341, 2002
34. Dunstan DW, Daly RM, Owen N, Jolley D, Vulikh E, Shaw J, Zimmet P: Home-based resistance training is not sufficient to maintain improved glycemic control following supervised training in older individuals with type 2 diabetes. *Diabetes*

- Care 28:3–9, 2005
35. Dunstan DW, Puddey IB, Beilin LJ, Burke V, Morton AR, Stanton KG: Effects of a short-term circuit weight training program on glycaemic control in NIDDM. *Diabetes Res Clin Pract* 40:53–61, 1998
 36. Honkola A, Forsen T, Eriksson J: Resistance training improves the metabolic profile in individuals with type 2 diabetes. *Acta Diabetol* 34:245–248, 1997
 37. Ishii T, Yamakita T, Sato T, Tanaka S, Fujii S: Resistance training improves insulin sensitivity in NIDDM subjects without altering maximal oxygen uptake. *Diabetes Care* 21:1353–1355, 1998
 38. Maiorana A, O'Driscoll G, Goodman C, Taylor R, Green D: Combined aerobic and resistance exercise improves glycemic control and fitness in type 2 diabetes. *Diabetes Res Clin Pract* 56:115–123, 2002
 39. Cuff DJ, Meneilly GS, Martin A, Ignaszewski A, Tildesley HD, Frohlich JJ: Effective exercise modality to reduce insulin resistance in women with type 2 diabetes. *Diabetes Care* 26:2977–2982, 2003
 40. Eriksson J, Taimela S, Eriksson K, Parvainen S, Peltonen J, Kujala U: Resistance training in the treatment of non-insulin-dependent diabetes mellitus. *Int J Sports Med* 18:242–246, 1997
 41. Wenger NK, Froelicher ES, Smith LK, Ades PA, Berra K, Blumenthal JA, Certo CM, Dattilo AM, Davis D, DeBusk RF, et al.: Cardiac rehabilitation as secondary prevention: Agency for Health Care Policy and Research and National Heart, Lung, and Blood Institute. *Clin Pract Guidel Quick Ref Guide Clin* 1–23, 1995
 42. Featherstone JF, Holly RG, Amsterdam EA: Physiologic responses to weight lifting in coronary artery disease. *Am J Cardiol* 71:287–292, 1993
 43. Benn SJ, McCartney N, McKelvie RS: Circulatory responses to weight lifting, walking, and stair climbing in older males. *J Am Geriatr Soc* 44:121–125, 1996
 44. Shrier I: Stretching before exercise does not reduce the risk of local muscle injury: a critical review of the clinical and basic science literature. *Clin J Sport Med* 9:221–227, 1999
 45. Yeung EW, Yeung SS: Interventions for preventing lower limb soft-tissue injuries in runners. *Cochrane Database Syst Rev* CD001256, 2001
 46. Beniamini Y, Rubenstein JJ, Zaichkowsky LD, Crim MC: Effects of high-intensity strength training on quality-of-life parameters in cardiac rehabilitation patients. *Am J Cardiol* 80:841–846, 1997
 47. Goldsmith JR, Lidtke RH, Shott S: The effects of range-of-motion therapy on the plantar pressures of patients with diabetes mellitus. *J Am Podiatr Med Assoc* 92:483–490, 2002
 48. Ganda OP: Patients on various drug therapies. In *Handbook of Exercise in Diabetes*. 2nd ed. Ruderman N, Devlin JT, Schneider SH, Kriska A, Eds. Alexandria, VA, American Diabetes Association, 2002, p. 587–599
 49. American Diabetes Association: Standards of Medical Care in Diabetes—2006. *Diabetes Care* 29:S4–S42, 2006
 50. Stevens RJ, Kothari V, Adler AI, Stratton IM: The UKPDS Risk Engine: a model for the risk of coronary heart disease in type II diabetes (UKPDS 56). *Clin Sci (Lond)* 101: 671–679, 2001
 51. Eddy DM, Schlessinger L: Archimedes: a trial-validated model of diabetes. *Diabetes Care* 26:3093–3101, 2003
 52. Fowler-Brown A, Pignone M, Pletcher M, Tice JA, Sutton SF, Lohr KN: Exercise tolerance testing to screen for coronary heart disease: a systematic review for the technical support for the U.S. Preventive Services Task Force. *Ann Intern Med* 140: W9–W24, 2004
 53. US Preventive Services Task Force: Screening for coronary heart disease: recommendation statement. *Ann Intern Med* 140:569–572, 2004
 54. Berger M, Berchtold P, Cuppers HJ, Drost H, Kley HK, Muller WA, Wiegelmann W, Zimmerman-Telschow H, Gries FA, Kruskemper HL, Zimmermann H: Metabolic and hormonal effects of muscular exercise in juvenile type diabetics. *Diabetologia* 13:355–365, 1977
 55. Zinman B, Ruderman N, Campaigne BN, Devlin JT, Schneider SH: Physical activity/exercise and diabetes mellitus. *Diabetes Care* 26 (Suppl. 1):S73–S77, 2003
 56. Berger M: Adjustment of insulin and oral agent therapy. In *Handbook of Exercise in Diabetes*. 2nd ed. Ruderman N, Devlin JT, Schneider SH, Kriska A, Eds. Alexandria, VA, American Diabetes Association, 2002, p. 365–376
 57. Sigal RJ, Purdon C, Bilinski D, Vranic M, Halter JB, Marliss EB: Glucoregulation during and after intense exercise: effects of beta-blockade. *J Clin Endocrinol Metab* 78:359–366, 1994
 58. de Muinck ED, Lie KI: Safety and efficacy of beta-blockers in the treatment of stable angina pectoris. *J Cardiovasc Pharmacol* 16 (Suppl. 5):S123–S128, 1990
 59. Aiello LP, Wong J, Cavallerano JD, Bursell S-E, Aiello LM: Retinopathy. In *Handbook of Exercise in Diabetes*. 2nd ed. Ruderman N, Devlin JT, Schneider SH, Kriska A, Eds. Alexandria, VA, American Diabetes Association, 2002, p. 401–413
 60. Vinik AI, Erbas T: Neuropathy. In *Handbook of Exercise in Diabetes*. 2nd ed. Ruderman N, Devlin JT, Schneider SH, Kriska A, Eds. Alexandria, VA, American Diabetes Association, 2002, p. 463–496
 61. Levin ME: The diabetic foot. In *Handbook of Exercise in Diabetes*. 2nd ed. Ruderman N, Devlin JT, Schneider SH, Kriska A, Eds. Alexandria, VA, American Diabetes Association, 2002, p. 385–399
 62. Wackers FJ, Young LH, Inzucchi SE, Chyun DA, Davey JA, Barrett EJ, Taillefer R, Wittlin SD, Heller GV, Filipchuk N, Engel S, Ratner RE, Iskandrian AE: Detection of silent myocardial ischemia in asymptomatic diabetic subjects: the DIAD study. *Diabetes Care* 27:1954–1961, 2004
 63. Valensi P, Sachs RN, Harfouche B, Lormeau B, Paries J, Cosson E, Paycha F, Leutenegger M, Attali JR: Predictive value of cardiac autonomic neuropathy in diabetic patients with or without silent myocardial ischemia. *Diabetes Care* 24:339–343, 2001
 64. Mogensen CE: Nephropathy: early. In *Handbook of Exercise in Diabetes*. 2nd ed. Ruderman N, Devlin JT, Schneider SH, Kriska A, Eds. Alexandria, VA, American Diabetes Association, 2002, p. 433–449
 65. Ward KM, Mahan JD, Sherman WM: Aerobic training and diabetic nephropathy in the obese Zucker rat. *Ann Clin Lab Sci* 24: 266–277, 1994
 66. Albright AL, Mahan JD, Ward KM, Sherman WM, Roehrig KL, Kirby TE: Diabetic nephropathy in an aerobically trained rat model of diabetes. *Med Sci Sports Exerc* 27:1270–1277, 1995
 67. Castaneda C, Gordon PL, Uhlin KL, Levey AS, Kehayias JJ, Dwyer JT, Fielding RA, Roubenoff R, Singh MF: Resistance training to counteract the catabolism of a low-protein diet in patients with chronic renal insufficiency: a randomized, controlled trial. *Ann Intern Med* 135:965–976, 2001
 68. Diabetes Prevention Program Research Group: The Diabetes Prevention Program (DPP): description of lifestyle intervention. *Diabetes Care* 25:2165–2171, 2002
 69. Lindstrom J, Louheranta A, Mannelin M, Rastas M, Salminen V, Eriksson J, Uusitupa M, Tuomilehto J: The Finnish Diabetes Prevention Study (DPS): lifestyle intervention and 3-year results on diet and physical activity. *Diabetes Care* 26:3230–3236, 2003
 70. Klem ML, Wing RR, McGuire MT, Seagle HM, Hill JO: A descriptive study of individuals successful at long-term maintenance of substantial weight loss. *Am J Clin Nutr* 66:239–246, 1997