

Type 2 Diabetes in the Young: The Evolving Epidemic

The International Diabetes Federation Consensus Workshop

GEORGE ALBERTI, FRCP, PHD
PAUL ZIMMET, FRACP, PHD
JONATHAN SHAW, FRACP, MD
ZACHARY BLOOMGARDEN, MD

FRANCINE KAUFMAN, MD
MARTIN SILINK, FRACP, MD
FOR THE CONSENSUS WORKSHOP GROUP

EXECUTIVE SUMMARY

1. The aims of the consensus meeting were to review the epidemiology, pathophysiology, management, and implications of the rising prevalence of type 2 diabetes in young people and to suggest means by which the continuing rise in incidence and prevalence might be prevented.
2. The overall global prevalence of type 2 diabetes is rising steadily. Previously, type 2 diabetes was predominantly a disease of middle-aged and older people. In recent decades, the age of onset has decreased and type 2 diabetes has been reported in adolescents and children worldwide, particularly in high-prevalence populations. Japan has seen an approximate fourfold rise in the incidence of type 2 diabetes in 6- to 15-year-olds, and between 8 and 45% of newly presenting children and adolescents in the U.S. have type 2 diabetes. The problem is particularly noticeable in indigenous peoples. Population-based data, however, are

sparse and indeed absent in most countries.

3. Additional cardiovascular risk factors are often associated with type 2 diabetes in the young, and microangiopathy is as common or commoner in those developing type 2 diabetes at a young age as in those with type 1 diabetes. This has profound societal implications.
4. Diagnostic separation of type 2 from other types of diabetes in young people can be difficult, and sophisticated testing may be necessary.
5. Data on the pathophysiology in the young are sparse, but there is no evidence to suggest differences from adults. The incidence of type 2 in the young is rising in parallel with the incidence of overweight and obesity, suggesting a possible causal relationship, particularly when the obesity is central and in relation to decreased physical activity. Other factors include family history, gestational diabetes in the mother, and low birth weight. All of these are associated with insulin re-

sistance, although decreased insulin secretion is also required.

6. Mass screening for type 2 diabetes in the young has been carried out in certain countries (Taiwan and Japan) but is probably appropriate only for individuals at very high risk. The best screening test for young people is not known.
7. Treatment should be aimed at physical and psychological well-being and avoidance of long-term complications. Lifestyle modification must accompany other forms of therapy. In certain countries, metformin is available for treatment of children, but efficacy remains unproven. Newer oral agents have not been tested systematically in children or adolescents. Insulin remains the most frequently used treatment. Hypertension and dyslipidemia are also common and require active intervention.
8. Prevention must be the main strategy for the future. School-based programs have been shown to be effective in the U.S. and Singapore. Major governmental actions that focus on lifestyle will be required.
9. Recommendations for action are made.

RECOMMENDATIONS

1. Population-based prevalence studies are urgently needed, particularly in "at-risk" populations. These should be followed by outcome studies.
2. Study methods need to be standardized with regard to classification, diagnostic methods, and diagnostic criteria.
3. Data on the natural history of type 2 diabetes in the young, particularly on complications, are required.
4. Further research is required to determine the role of the oral glucose tolerance test (OGTT) in screening asymptomatic young people, as well as the appropriate glucose load and tim-

The International Diabetes Federation Consensus Workshop, "Type 2 Diabetes in the Young: The Evolving Epidemic," took place in Santa Monica, California, on 7-9 February 2003. A complete list of workshop participants and their respective affiliations is listed in the APPENDIX.

Address correspondence and reprint requests to Dr. Jonathan Shaw, MD, MRCP(UK), FRACP, Director of Research, International Diabetes Institute, 250 Kooyong Rd., Caulfield, Victoria 3162, Australia. E-mail: jshaw@idi.org.au.

Received and accepted for publication on 13 April 2004.

F.K. has been a member of an advisory panel, a standing committee, or the board of directors for Eli Lilly, Novo Nordisk, Aventis, Medtronic MiniMed, Insulet, Clinical Products, LifeScan, and Amylin; has received grant and research support from Medtronic MiniMed, Novo Nordisk, BMS, and Merck; and holds stock in Clinical Products.

Abbreviations: ADA, American Diabetes Association; CVD, cardiovascular disease; DKA, diabetic ketoacidosis; IDF, International Diabetes Federation; HNF, hepatocyte nuclear factor; IFG, impaired fasting glucose; IGT, impaired glucose tolerance; MODY, maturity-onset diabetes of youth; OGTT, oral glucose tolerance test; PCOS, polycystic ovary syndrome.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

© 2004 by the American Diabetes Association.

- ing of blood sampling to be used in the OGTT.
5. The costs and psychosocial impact of screening in young people need to be examined.
 6. The safety and efficacy of oral hypoglycemic drugs in children need to be determined.
 7. The role of insulin therapy for children with type 2 diabetes needs to be determined in multicenter randomized controlled trials. These should examine the optimum type, timing, intensity, and duration of insulin therapy.
 8. Existing registers for type 1 diabetes in children should be expanded to include those with type 2 diabetes.
 9. Innovative ways of working with children with diabetes and their families to institute long-standing and sustainable lifestyle change need to be sought.
 10. Strategies for the prevention of type 2 diabetes in children need to involve government and societal, as well as individual, change. Programs should focus on the prevention of obesity through increasing physical activity and promoting healthy eating, the promotion of breast feeding, and achieving excellent pregnancy outcomes with regard to avoiding low birth weight and treating gestational diabetes.

AIM AND OBJECTIVES OF THE WORKSHOP

— The Consensus Workshop on “Type 2 Diabetes in the Young: The Evolving Epidemic” was convened by the International Diabetes Federation (IDF) in response to the many reports of the occurrence of type 2 diabetes in children and adolescents. The objective of this document is to review the current information on type 2 diabetes in youth and to reach a consensus on what actions need to be taken to slow or reverse this trend. The topic has become a clinical, research, and health economic priority, with important implications for the health status of future generations throughout the world.

The workshop sought to:

- Assess current information on type 2 diabetes in the pediatric population in terms of epidemiology and public health implications
- Review the classification and diagnosis of type 2 diabetes in children and adolescents

- Discuss the pathogenesis of type 2 diabetes in youth
- Evaluate screening recommendations
- Assess the burden of complications
- Make recommendations on therapeutic options
- Assess the opportunities for prevention in order to reach a consensus on what actions need to be taken to slow or reverse the epidemic.

INTRODUCTION AND BACKGROUND

— Until recently, type 2 diabetes was typically regarded as a disease of the middle-aged and elderly. While it still is true that this age-group maintains a higher risk than younger adults, evidence is accumulating that onset in those aged under 30 years is increasingly common. Even children and adolescents are now becoming caught up in the diabetes epidemic. Although type 1 diabetes remains the main form of the disease in children worldwide, it is likely that type 2 diabetes will be the predominant form within 10 years in many ethnic groups. Type 2 diabetes has already been reported in children in a number of countries, including Japan, the U.S., India, Australia, and the U.K. (1–5).

This new phenomenon brings a serious new aspect to the global diabetes epidemic and heralds an emerging public health problem of major proportions. Among children in Japan, type 2 diabetes is already more common than type 1 and accounts for 80% of childhood diabetes. The rising prevalence of obesity and type 2 diabetes in children is yet another symptom of the effects of sedentary lifestyles as part of globalization and industrialization affecting all societies (1).

This fall in the age of onset of type 2 diabetes is an important factor influencing the future burden of the disease and was part of the stimulus for the IDF to organize the workshop. Onset of diabetes in childhood or adolescence heralds many years of disease and an increased risk that the full range of both micro- and macrovascular complications will occur when affected individuals are still relatively young. Thus, future generations may be burdened with morbidity and mortality at the height of their productivity, potentially affecting the workforce and health care systems of countries across the world.

EPIDEMIOLOGY — The global burden of type 2 diabetes is both significant and rising, with most of the increase registered in the last two decades (6). From 2003 to 2025, the worldwide prevalence of diabetes in adults is expected to increase from 5.0 to 6.2%, or 328 million (7). The largest proportional and absolute increase will occur in developing countries, where the prevalence will rise from 4.2 to 5.6%. In India and China, the adult diabetic population is expected to double by 2025 to about 73 million and 46 million, respectively.

While most of the rise in the prevalence of type 2 diabetes has been seen in the middle-aged and elderly, there is now strong evidence of a rise among younger adults. In Australia, 1.7 and 1.4% of persons aged 35–44 and 45–54 years had diabetes in 1981 (8), and these prevalence rates increased to 2.4 and 6.2%, respectively, in 2000 (9). Chinese data show that the prevalence of diabetes among 35- to 44-year-olds rose from 1.7% in 1994 to 3.2% in 2000 (10). These data confirm a trend to an earlier age of onset of diabetes.

There are ever increasing reports of type 2 diabetes in children worldwide, with some as young as 8 years old being affected (11). These are mostly in ethnic groups with a high susceptibility to type 2 diabetes. However, there are now also reports of type 2 diabetes occurring among Europid (white Caucasoid) teenagers (4). In Japan, the prevalence of type 2 diabetes among junior high school children has doubled from 7.3 per 100,000 between 1976 and 1980 to 13.9 per 100,000 in 1991–1995, with type 2 diabetes now outnumbering type 1 diabetes in that country (1). Recent studies from the U.S. indicate that between 8 and 45% of recently diagnosed diabetes in the young is due to type 2 diabetes (12). Despite the large increase, the prevalence is still, fortunately, much lower than in the adult population.

Inadequate data at present

The available information on type 2 diabetes incidence and prevalence in childhood and adolescence is sparse compared with that for adults. Most surveys are clinic based or case series, with a paucity of population-based surveys, particularly outside North America (13) and Japan (1). Information on the natural history and etiology of type 2 diabetes in the pediatric age range is also sparse. There is

also a lack of uniformity in case definition (resulting in misclassification so that some children are being misdiagnosed and treated as type 1 diabetes), data collection, and follow-up (14).

The studies in the literature may be divided into population- and clinic-based studies.

Population-based studies

The largest study reported is from Japan (1), with ~7 million youth studied between 1976 and 1997. The initial screening step of the study was a urine glucose test, with blood testing reserved only for those with glycosuria; thus, the figures are probably an underestimate. Over the 20-year period, type 2 diabetes incidence increased 10-fold in children aged 6–12 years: 0.2 per 100,000 per year from 1976 to 1980 vs. 2.0 per 100,000 per year from 1991 to 1995. Similarly, over the same time period, type 2 diabetes incidence doubled among 13- to 15-year-olds: 7.3 vs. 13.9 per 100,000 per year.

A screening program in which fasting blood glucose was measured in those with persistent glycosuria carried out in 3 million students (aged 6–18 years) in Taiwan (15) found the prevalence of undiagnosed diabetes to be 9.0 and 15.3 per 100,000 boys and girls, respectively. The prevalence of undiagnosed diabetes was 62% higher in girls than boys, after adjustment for other factors, and the cases were most commonly identified between the ages of 12 and 14 years. A 3-year follow-up of these cases showed that 54% had type 2 diabetes, 10% had type 1 diabetes, 9% had secondary diabetes, 20% were nondiabetic, and 8% had no definite diagnosis. The cases identified as having type 2 diabetes had a higher mean BMI, cholesterol, and blood pressure than those with a normal fasting glucose, suggesting that even at this young age, cardiovascular risk was starting to rise.

In Pima Indians aged 15–19 years, the prevalence of type 2 diabetes markedly increased from 2.4% in males and 2.7% in females in 1967–1976 to 3.8% in males and 5.3% in females in 1987–1996 (14). In the U.S., among Navajo Indians, diabetes or impaired glucose tolerance (IGT) was found in 3 and 13%, respectively, of girls and boys aged 12–19 years (16). In Canada, among the Cree-Ojibway aboriginals, diabetes and impaired fasting glucose (IFG) were found in 1 and 3% of children aged 4–19 years (17), and IGT

was found in 10% of those aged 10–19 years (18). A 4% prevalence of diabetes among adolescent girls in native populations in Canada has been reported from several surveys (19). A cohort of indigenous Australian children aged 7–18 years was surveyed in 1989 and again in 1994. Over the 5 years, the prevalence of type 2 diabetes almost doubled to 1.3%, while that for IGT increased almost sevenfold to 8.1% (20). At the follow-up, 18% of the population was overweight or obese, and one-third had elevated cholesterol levels.

National U.S. data from 1988 to 1994 (21) show that of ~3,000 persons aged 12–19 years, the prevalence of IFG was 1.8% and the prevalence of elevated HbA_{1c} (>6%) was 0.4%, while diabetes of all types (9 of the 13 diagnosed with diabetes were on insulin) was diagnosed in 0.4%, suggesting some glycemic abnormality in ~600,000 adolescents in the U.S.

Clinic studies

Clinic and register-based studies make up the largest group of studies. They reveal type 2 diabetes occurring in children as young as 8 years old (11). In addition, they have demonstrated a female preponderance (22–24), a strong family history of diabetes (22,25,26), and associations with obesity (11,22,25–27), acanthosis nigricans (11,26,27), and polycystic ovary syndrome (PCOS) (28).

A number of clinic-based studies have estimated incidence rates. In Cincinnati, Ohio, the annual incidence of type 2 diabetes in 10- to 19-year-olds increased from 0.7 per 100,000 in 1982 to 7.2 per 100,000 in 1994 (26). Among African-American and Latino children (aged 0–17 years) in Chicago, Illinois, the incidence of type 2 diabetes rose by 9% per year from 1985, reaching 3.8 per 100,000 per year (29) by 1994. This study showed higher rates in girls and in African Americans.

The prevalence of diagnosed diabetes among American Indians aged 15–19 years, as reported right across the Indian Health Service, increased from 0.32 to 0.54% from 1990 to 1998 (30). Although the prevalence of type 2 diabetes was higher among females, the relative increase over this time period was greater among males (0.23 to 0.41% for males vs. 0.42 to 0.68% for females). Over the same time period, the prevalence among those

younger than 15 years was unchanged at 0.12%.

Some studies that analyzed referrals to diabetes clinics have reported the percentage of diabetes cases labeled as type 2. From 1994 to 1998, the proportion of new cases of pediatric diabetes in Florida that were labeled as having type 2 increased from 9.4 to 20% (31). In Cincinnati in 1994, type 2 diabetes accounted for 16% of all new cases of diabetes in children aged up to 19 years and accounted for 33% of new cases among the 10- to 19-year-old age-group (26). In Mexican-American youth aged 0–17 years, a California clinic reported that 31% of the diabetic children had type 2 diabetes (32). Among newly diagnosed diabetic children and adolescents in Bangkok, type 2 diabetes increased from 5% of all cases during 1986–1995 to 17.9% during 1996–1999 (27). In contrast, among European populations, which have lower overall prevalences of type 2 diabetes, two very large studies have reported that only 0.5% of children and adolescents with diabetes have been classified as having type 2 diabetes (33,34).

Complications—morbidity and mortality

As with adults, it can be expected that youth with type 2 diabetes will also develop diabetes-related micro- and macrovascular complications, and there is already evidence available of microvascular complications. In a study from Canada, subjects who developed type 2 diabetes as children were resurveyed as young adults aged between 18 and 33 years. Of the 51 subjects, 9% had died, 6% were on dialysis, 1 had a toe amputation, and 1 was blind (35). Another follow-up study from Japan compared those with type 1 and type 2 diabetes diagnosed at <30 years of age for development of nephropathy (36). After 30 years of diabetes, 44% of those with type 2 and 20.2% of those with type 1 had nephropathy. Krakoff et al. (37) looked at incidence of retinopathy and nephropathy among Pima Indians diagnosed with type 2 diabetes at <20 years of age (youth), 20–39 years of age (young adults), and 40–59 years of age (older). Nephropathy was equally common in all age-groups and was not related to age of diabetes onset (at <5 years' duration of type 2 diabetes, nephropathy incidence per 1,000

Table 1—Features to differentiate type 1 and 2 diabetes in young people

	Type 1 diabetes	Type 2 diabetes
Onset	Acute—symptomatic	Slow—often asymptomatic
Clinical picture	<ul style="list-style-type: none"> ● Weight loss ● Polyuria ● Polydipsia 	<ul style="list-style-type: none"> ● Obese ● Strong family history type 2 diabetes ● Ethnicity—high-prevalence populations ● Acanthosis nigricans ● PCOS
Ketosis	Almost always present	Usually absent
Insulin	<ul style="list-style-type: none"> ● C-peptide negative 	<ul style="list-style-type: none"> ● C-peptide positive
Antibodies	<ul style="list-style-type: none"> ● ICA positive ● Anti-GAD positive ● ICA 512 positive 	<ul style="list-style-type: none"> ● ICA negative ● Anti-GAD negative ● ICA 512 negative
Therapy	Insulin invariably	Oral hypoglycemic agents
Associated autoimmune diseases	Yes	No

person-years was 13/1,000 youth, 8/1,000 young adults, and 7/1,000 older). However, unlike those with adult-onset diabetes, in whom retinopathy was apparent at even this short duration, retinopathy did not appear among youth-onset cases until at least 5 years' duration. Retinopathy was less common in youth-onset than in adult-onset groups at all durations of diabetes.

Although it is not known if the complication rate is similar for other ethnic groups, these studies have important implications in that they highlight the risk of complications occurring at a young age and soon after diagnosis. This will place a significant burden on health budgets as well as on society as a whole, especially as these people would be entering their peak working and earning capacity at the time when complications begin to occur. Early detection and intervention are therefore essential to reduce the risk of future complications.

CLASSIFICATION AND DIAGNOSIS

— The appearance of type 2 diabetes in children and adolescents has exacerbated the existing issues in the classification of diabetes. There is now a problem in clinically distinguishing the etiology of diabetes in children and adolescents, often necessitating laboratory studies to differentiate type 1 from type 2 diabetes (Table 1). This requires tests that are not always available in the primary care environment or in many developing countries.

The presence of diabetic ketoacidosis (DKA) is a classic manifestation of type 1 diabetes. However, DKA may occur at

presentation in subjects who are eventually found to have type 2 diabetes (38); that is, they have elevated C-peptide and an absence of antibodies to islet cells (ICA) or glutamic acid decarboxylase (anti-GAD), and after the initial period of severe metabolic disturbance, they do not require insulin (13). Initial reports were among African-American adults, and an autosomal dominant inheritance was suggested (38,39). Subsequent studies have confirmed that type 2 diabetes often presents with ketosis or DKA and occurs in ~40% of African-American children (40,41) with type 2 diabetes, 4% of children with Canadian aboriginal backgrounds (42), and 30% of children with Mexican-American backgrounds (32). A strong family history of type 2 diabetes is common, but an autosomal dominant inheritance has not been confirmed. It has variously been called Flatbush diabetes, atypical diabetes (ADM), and phasic insulin dependence.

At the root of the classification problem is the incomplete understanding of the pathogenesis of the two major types of diabetes, as knowledge of causation is the optimal basis for classification. The association of autoimmune serologic reactions and particular HLA alleles in type 1 diabetes strongly implicates autoimmunity in the pathogenesis, and these tests can be used to help with classification. However, classification is complicated because autoimmunity (at least as determined by current antibody assays) does not contribute to type 1 diabetes as substantially in non-Europid as in Europid populations (43,44).

Further differential diagnoses are maturity-onset diabetes of youth (MODY), which is attributable to mutations of the glucokinase genes and hepatocyte nuclear factor (HNF)-1 α and -4 α genes (45), and diabetes due to mutations of mitochondrial DNA (46).

The increasing recognition of difficulties with diagnosis and the potentially major influence of diagnosis on therapy (particularly the decision to institute life-long insulin treatment) suggest the need to consider laboratory investigations, even in apparently typical cases. Therefore, in ideal circumstances it is recommended that apparently typical cases of type 1 diabetes are confirmed with a measurement of autoantibodies, and typical cases of type 2 diabetes are confirmed with an assessment of insulin resistance (e.g., fasting C-peptide). It is recognized that in many settings worldwide such an approach is not affordable and that precise cutoffs for some of the diagnostic tests (especially for insulin resistance) have not been established. When diagnostic tests do not confirm the clinical diagnostic label, further investigations are indicated. In such cases, particularly when there is a parental history of diabetes, genetic testing will often be required. Diagnostic molecular testing is now available (for details, visit www.diabetesgenes.org) and when used appropriately provides valuable clinical information on diagnosis, prognosis, and treatment since most patients with MODY respond best to sulfonylureas (47). It is relevant to note that among non-Hispanic Europid populations, the prevalence of monogenic diabetes in children is similar to or higher than that of type 2 diabetes and is not excluded by the presence of obesity (48).

PATHOGENESIS — There is sparse information on the pathophysiology of type 2 diabetes in the young; therefore, extrapolation from adults is necessary. Type 2 diabetes is characterized by disorders of insulin action and insulin secretion, either of which may be the predominant feature (49). Both are usually present at the time that type 2 diabetes clinically manifests. Persons with type 2 diabetes usually have plasma insulin concentrations that appear normal or elevated but insulin secretion, particularly first phase, is defective and insufficient to compensate for the insulin resistance (49). The specific reasons for the develop-




Figure 1—Annual incidence of type 2 diabetes and prevalence of obesity among Japanese school children. Adapted from Kitagawa et al. (1), with permission.

ment of these abnormalities are not yet known, but it is of heterogeneous etiology with behavioral and environmental risk factors unmasking the effects of genetic susceptibility (50).

The key factors involved in the development of type 2 diabetes are as follows:

Genetic

Due to complex inheritance patterns and interaction with the environment, identifying the genes involved in the common forms of type 2 diabetes has proved difficult. Only a small percentage of cases can be explained by the single gene defects such as those known to cause MODY. The genetic component of type 2 diabetes in children and adolescents has yet to be explored, but it is reasonable to assume that it is similar to that in adults, and indeed a strong family history of type 2 diabetes is often seen among adolescents with the disease (see “Other factors” below). A number of major predisposing genes have been found on chromosomes 1q, 12q, 20q, and 17q (51). Minor genes that have been defined by a candidate approach include the Pro12Ala polymorphisms in peroxisome proliferator-activated receptor (PPAR)- γ (52) and the Kir 6.2 E23K variant (53). In Ojibwa-Cree indigenous Canadians, a private mutation (G319S HNF1 α) present in 20% of the population predisposes to diabetes (phenotypically like type 2 diabetes), with a relative risk of 2 when a single copy is inherited

and a relative risk of 4 when two copies are inherited (54). This may yet prove to be another form of MODY. Sellers et al. showed an inverse relationship between number of copies of the mutation and markers of insulin resistance, suggesting that the gene is related to secretory defects (55). The data on the genetic component in adult type 2 diabetes are extensively reviewed elsewhere (51).

Obesity

On a global basis, the rise in type 2 diabetes rates mirrors the growth in urbanization and economic development (6,56). Obesity appears to be the key link, and this is exemplified by the situation in Japan, where a parallel rise in type 2 diabetes incidence in children and levels of obesity occurred from 1975 to 1995 (1) (Fig. 1). Overweight and obesity are be-

coming major problems in adolescents around the world (Table 2).

The prevalence of overweight (defined as at or above the 95th percentile of the sex-specific BMI for age growth charts) among children in the U.S. increased from 7% in 1988 to 10% in 1999 among those aged 2–5 years and from 5% in 1976 to 11% in 1988 and to 15% in 1999 among those aged 6–19 years (57). The rise in the prevalence has been much more pronounced among African-American and Hispanic children than among non-Hispanic whites (120% rise vs. 50% rise in 12 years) (58). Behavioral differences have been observed, which may explain some of these differences. While African-American and Europid youth may consume the same amount of calories, African-American children tend to get a greater percentage of their calories from fat, in addition to increased intake of sweetened drinks (59). Moreover, African-American adolescents do not perceive themselves to be heavy and actually express a desire to be heavier (60).

Secular increases in the prevalence of obesity in children have also been recorded in China (62), Hong Kong (63), the U.K. (64), and Australia (65). In Australia (using age- and sex-specific BMI cutoffs designed to be the equivalent percentiles to a BMI of 25 and 30 kg/m² in 18-year-olds), ~5% of children are currently obese and an additional 16% are overweight (66). These prevalences doubled over the past decade after being nearly stable at around 10% from 1969 to 1985 (65). In India, a recent study found that the age-adjusted prevalence of being overweight (using the same cutoffs as the Australian study) among 13- to 18-year-olds was 18% in boys and 16% in girls (67). Prevalence rates increased with age, decreasing physical activity, and higher

Table 2—Prevalence of overweight and obesity in six countries at age 18 years (61)

	Overweight and obesity (BMI \geq 25 kg/m ²)		Obesity (BMI \geq 30 kg/m ²)	
	Males	Females	Males	Females
Brazil	4.7	15.2	0.1	2.0
Great Britain	9.6	11.7	0.9	1.2
Hong Kong	11.7	9.8	3.1	1.8
Netherlands	5.5	6.5	0.3	0.3
Singapore	10.5	7.0	1.7	1.0
United States	18.1	16.5	3.3	4.0

Data are percent.

socioeconomic status (67). In the Canadian Ojibwa-Cree community, 48–51% of children aged 4–19 years were found to have a weight >90th percentile (17). Changes in traditional lifestyles among indigenous communities, such as a reduction in hunting and gathering and the adoption of a more sedentary life with a westernized diet, are thought to contribute to rising obesity levels (68).

Physical inactivity

Inactivity is one of the major contributors to overweight and obesity. A recent longitudinal study of girls in the U.S. showed a marked decline in physical activity over 10 years, such that by the age of 16–17 years, 56% of black and 31% of white girls reported no habitual leisure-time physical activity (69). Pregnancy, cigarette smoking, higher BMI, and lower parental education at baseline were all associated with a subsequent decline in physical activity. Participation in physical education in U.S. schools fell from 41.6% in 1991 to 24.5% in 1995 (70). Another study found that white students in the U.S. have higher physical activity levels than other ethnic groups, with boys usually more active than girls, regardless of race (71).

Television-viewing time has been linked with childhood obesity (72), which may be related to the associated consumption of high-energy foods (73). On average, children's programming in the U.S. includes 10 food advertisements hourly (74), more than twice that in adult viewing (75).

Insulin resistance

Published studies linking insulin resistance directly with type 2 diabetes in adolescents and children are very limited. However, observations about the typical age of onset of diabetes, evidence of insulin resistance in normoglycemic children from high-risk ethnic groups, and the vast literature on insulin resistance in adults make a compelling case. Most observations in youth with type 2 diabetes would suggest that insulin resistance is an early feature in these children. Initially, insulin resistance manifests itself in compensatory hyperinsulinemia. Over time there is β -cell failure, resulting in the clinical expression of type 2 diabetes (76) (77).

In a cross-sectional study of 14 adolescents with IGT matched with 14 control subjects of similar age, BMI, body fat, and leptin, the children with IGT were

found to have greater insulin resistance. Furthermore, they had higher visceral and lower subcutaneous abdominal fat and decreased first-phase insulin secretion and glucose disposition index (78). Intramyocellular fat content (as measured by nuclear magnetic resonance spectroscopy) showed a strong positive correlation with insulin resistance and with 2-h postload plasma glucose (78), suggesting that it may be important in pathogenesis. Further evidence of the importance of insulin resistance can be drawn from the observation that the onset of type 2 diabetes frequently occurs around the time of puberty, when insulin sensitivity declines (79–81). Even in healthy, prepubertal African-American children, a family history of diabetes is associated with an ~20% reduction in insulin sensitivity (82). The hyperinsulinemia and reduced insulin sensitivity observed in Hispanic and black children has been associated with both increased insulin secretion and reduced insulin clearance (83,84).

In a variety of ethnic groups, obesity is associated with evidence of insulin resistance and impaired insulin secretion among children, as in adults (85,86). Central obesity is of particular importance as a determinant of hyperinsulinemia (87). Compared with white children with obesity and similar insulin sensitivity levels, black children have lower hepatic glucose output, lower total and LDL cholesterol, and lower triglyceride levels, with considerably lower visceral fat levels. Visceral adiposity was associated with lower insulin sensitivity in both groups. This was compensated by higher insulin secretion in whites, but not in blacks (88). These findings suggest a greater diabetogenic risk of obesity among African Americans but greater atherogenic risk among whites. The dietary fat-to-carbohydrate ratio correlates significantly with insulin resistance and may partly explain the metabolic differences seen between black and white children (83). Indeed, a number of studies have shown that African-American children have higher total fat and cholesterol intake, prefer greater sweetness in liquids, are physically less active, and spend more time watching television (59,89).

It is possible to ameliorate insulin resistance by increasing the level of physical activity. This has been demonstrated in obese children and more recently in non-

diabetic, normal weight children (90), where more active subjects had lower fasting insulin values and greater insulin sensitivity, as measured by a glucose clamp.

Acanthosis nigricans and PCOS

Acanthosis nigricans is a well-established physical marker of insulin resistance and is reported to occur in up to 60–90% of young people with type 2 diabetes (91,92). This seems to be especially true for African Americans, Mexican Americans, and some Native Americans, but not in other populations, such as in Japan. PCOS is associated with a state of insulin resistance and compensatory hyperinsulinemia (83). Obese adolescents with PCOS have a 50% reduction in peripheral tissue insulin sensitivity and evidence of hepatic insulin resistance and compensatory hyperinsulinemia (93). Approximately 30% of adolescent girls with PCOS have been found to have IGT, with 4% having type 2 diabetes (28). In these adolescents, glucose intolerance is associated with decreased first-phase insulin secretion, decreased glucose disposition index, and increased hepatic glucose production (94).

Intrauterine environment

Two aspects relating to the intrauterine environment—birth weight and maternal hyperglycemia—may possibly affect the development of type 2 diabetes among the young. There is now abundant evidence that low birth weight predicts type 2 diabetes in middle age (95). Recent data from Taiwan confirm this in children (96). A study of 6- to 18-year-olds showed a U-shaped relationship, in which the risk of type 2 diabetes was lowest in those with a birth weight of 3–3.5 kg, and was significantly increased in those children who had either low (<2.5 kg) or high (>4.0 kg) birth weight, despite adjustment for other factors, including gestational diabetes, family history of diabetes, and socioeconomic status (96). Recent evidence seems to indicate that the greatest risk for obesity and glucose intolerance in adulthood is in those with low birth weight who gain weight rapidly in childhood (97,98). The risk in this group is higher than in those who are overweight from birth.

Gestational diabetes mellitus (GDM) also seems to increase the risk of diabetes developing in offspring (99). A prospec-

Table 3—ADA recommendations for testing for type 2 diabetes in children (12)

Criteria for considering screening for diabetes*

Overweight (BMI 85th percentile for age and sex, weight for height 85th percentile, or weight 120% of ideal for height)

Plus any two of the following risk factors:

- Family history of type 2 diabetes in first- or second-degree relative
- Race/ethnicity (American Indian, African American, Hispanic, Asian/Pacific Islander)
- Signs of insulin resistance or conditions associated with insulin resistance (acanthosis nigricans, hypertension, dyslipidemia, PCOS)

Age of initiation: age 10 years or at onset of puberty if puberty occurs at a younger age

Frequency: every 2 years

Test: fasting plasma glucose preferred

*Clinical judgment should be used to test for diabetes in high-risk subjects who do not meet these criteria.

tive study (100) found that the prevalence of IGT in the children of mothers with a diabetic pregnancy increased with time from 1.2% at <5 years of age to 19.3% at 10–16 years of age. This was compared with 2.5% at 10–16 years of age in control subjects. Higher levels of amniotic fluid insulin at 33–38 weeks of gestation was a strong predictor of later IGT (100). The specific environmental role of maternal hyperglycemia, as separate from maternal genetics, has been demonstrated in Pima Indians, in whom offspring born after the mother developed diabetes were more obese as children and more likely to have diabetes in their twenties than their siblings born before their mothers developed diabetes (14). No such differences were seen before and after fathers developed diabetes. Age at diagnosis in HNF-1 α mutation carriers was >10 years younger when the mother was diagnosed before pregnancy compared with when the mother was diagnosed after pregnancy (101). The father's age at diagnosis did not, however, influence the offspring's age at diagnosis.

Other factors

Family history. Many studies show a strong family history among affected youth, with 45–80% having at least one parent with diabetes and 74–100% having a first- or second-degree relative with type 2 diabetes (12,102). Children with diabetes are also more likely to have a family history of cardiovascular disease (CVD), with one study showing that up to 28% have a positive family history of CVD (103). In a study among Pima Indians, it was shown that the cumulative incidence of type 2 diabetes was highest in offspring if both parents had diabetes (104). Japanese children with type 2 diabetes show familial clustering, with siblings having a

175- to 250- fold increase in diabetes over the frequency in the general population, and parents have a 48–60% likelihood of having type 2 diabetes (105,106). A lifestyle predisposing to obesity and type 2 diabetes seems to characterize families with adolescents who have type 2 diabetes (26). Specifically, members of such families tend to be overweight and inactive and have a tendency to high-fat intake and even binge eating (107).

Sex. Girls are 1.7 times more likely than boys to develop type 2 diabetes in an analysis of a large set of studies (108). The reason for this is not clear, and it appears to be a special feature of type 2 diabetes in youth, as sex differences in diabetes prevalence are only a very minor feature in adults.

Socioeconomic status. Socioeconomic status has been associated with type 2 diabetes and obesity in adults. In developed countries, the risk of diabetes is highest in those with lower socioeconomic status (109,110); however, the opposite appears to apply in developing countries, where higher socioeconomic status is associated with an increased risk of diabetes (111). Physical inactivity probably explains these findings in both settings. In children, no data on diabetes are yet available, but a recent large study of 3-year-olds in the U.K. found that those in the most deprived group were 30% more likely to be obese than those in the least deprived group (112).

SCREENING

A major consequence of the appearance of type 2 diabetes in the younger age-group is that subjects with diabetes will have a longer duration of the disease that increases medical costs and increases the risk that both microvascular and macrovascular complications will develop at an

earlier age. This makes the issue of screening in the young a very relevant and, indeed, urgent issue. However, certain criteria need to be met before screening can be considered in the public health arena. These are summarized below.

- The disease is common among the general population or among easily identifiable high-risk subgroups
- The disease is serious in terms of morbidity and mortality
- The disease has a prolonged latency without symptoms
- The screening test is adequately sensitive and specific
- Intervention is available to prevent or delay disease onset or treat at an early stage.

Type 2 diabetes in children appears to meet a number of these criteria, suggesting that it may be an appropriate target for screening. However, even when considering the screening of adults for diabetes, there remains heated debate, as it has been suggested that evidence is not yet available to demonstrate that it fulfills the criteria for population-based mass screening (113). The IDF consensus group noted the recommendations of the 2000 American Diabetes Association (ADA) report on type 2 diabetes in children and adolescents (12), as outlined in Table 3. Consistent with those recommendations, only children at substantial risk for the presence or the development of type 2 diabetes should be considered for screening.

The Third National Health and Nutrition Examination Survey (NHANES III) data suggest that the ADA risk criteria would lead to testing of 10% of youths in the U.S., for a total of ~2.5 million adolescents between 12 and 19 years of age, of whom 5% might be expected to have IFG (fasting plasma glucose 6.1–6.9 mmol/l) or undiagnosed diabetes, while 1.8% of those not tested under such recommendations would be expected to have IFG (114). It should be noted that the ADA has recently recommended lowering the criteria for IFG to 5.7–6.9 mmol/l (115).

The question of what tests to use for the initial screening for type 2 diabetes is difficult. Evidence from adult populations clearly shows that ~30% of all those with undiagnosed diabetes have a nondiabetic fasting glucose (116) but are nev-

ertheless at high cardiovascular risk (117), suggesting that the OGTT should be part of the screening process. Others (118) have questioned this assertion, but the World Health Organization (WHO) and many national diabetes organizations continue to recommend its use for adults, particularly in those with a high but non-diabetic fasting glucose. It should be noted that it is currently unknown whether the diagnostic cut points used for adults are applicable to children. Furthermore, the evidence base for the standard 1.75-g/kg (to a maximum of 75 g) dose of glucose used in the OGTT is not strong, and the evidence needs to be further examined in order to review its validity.

Estimates from the Japan and Taiwan screening programs of the unselected general school population show a cost of approximately \$10,000 (U.S.) per case found, indicating the need to focus on high-risk groups. Cost-effectiveness analysis suggests that the optimal approach for screening adults is to use “opportunistic screening” for undiagnosed diabetes at routine medical system contacts (119). Such an approach may not work with children who generally have less frequent contact with health care systems.

The psychosocial impact of diagnosing an asymptomatic disease through screening warrants consideration. The potential for distress to occur when an asymptomatic individual is diagnosed with a lifelong disease is considerable, particularly when the diagnosis occurs during adolescence and when it carries with it implications about an individual's or family's lifestyle habits. Evidence would suggest that in adults there is not a significant detrimental effect of identifying diabetes through screening (120), but there are no such data available for children and adolescents, and it would be particularly dangerous to extrapolate such findings.

TREATMENT — The appearance of type 2 diabetes in a younger age-group raises new issues in management of diabetes apart from the difficulty in classification. What therapies are safe in this age-group? Most pharmacologic therapies for diabetes and its associated conditions (apart from insulin) are not approved for use in children, and the same applies for those for blood pressure and dyslipidemia.

The goals of treatment for type 2 dia-

betes in children and adolescents are to achieve:

- Physical well-being
- Long-term glycemic control
- Prevention of microvascular complications of diabetes
- Prevention of macrovascular disease
- Psychological well-being.

Physical well-being

The attainment of physical well-being has short-term and long-term goals. Short-term goals include treatment of acute metabolic decompensation if present (ketoacidosis), the elimination of symptoms associated with hyperglycemia (polyuria, polydipsia), treatment of acute and chronic infections (vaginal candidiasis is frequent at presentation), and treatment of medical problems associated with obesity (e.g., sleep apnea, steatosis).

The long-term goals for physical well-being include achieving normal growth for the younger adolescent with diabetes, achieving and maintaining a reasonable body weight, maintaining a reasonable level of fitness by a regular physical activity program, avoidance of smoking, and preventing the complications of diabetes.

Glycemic control

The theoretical glycemic goal is to achieve and maintain normoglycemia; however, a more pragmatic target is to maintain $HbA_{1c} < 7\%$. Lifestyle intervention (diet, exercise, and weight control) is the cornerstone of management and if successful is likely to benefit blood pressure and lipids as well as glycemia. If diabetes is diagnosed early, lifestyle intervention may suffice, although this only appears to be applicable to ~10% of patients in clinical practice (121).

In principle, there is no reason to suppose that pharmacological treatment of hyperglycemia should be any different in adults and children. What does pose a problem is our lack of knowledge about the long-term implications of drug treatment of diabetes in this age-group.

A proposed therapeutic algorithm for asymptomatic children with type 2 diabetes is to start with lifestyle intervention approaches; then add monotherapy, particularly emphasizing the use of metformin; and subsequently use combinations of two oral medications and, if adequate control is not achieved, the addition of insulin. For symptomatic

children, with blood glucose persistently exceeding 17 mmol/l or when DKA is present, insulin therapy is indicated, with subsequent efforts to taper this phenomenon and substitute metformin monotherapy once blood glucose levels are normal (108,121). Many youth maintain near-normal blood glucose levels for months to years after one course of insulin at diagnosis (122).

In clinical practice in the U.S., approximately one-half of young patients with type 2 diabetes receive insulin and one-half receive oral agents, most commonly metformin (108,121). In another analysis of treatment regimens for type 2 diabetes in young persons, 28% began therapy with metformin, with the remainder using insulin alone or the two in combination (123).

Pharmaceutical agents

Insulin. The apparent “overuse” of insulin therapy in adolescent type 2 diabetes is probably multifactorial. It is possible that significant insulin deficiency may be common among the symptomatic adolescents with type 2 diabetes presenting to diabetes centers. Insulin is very familiar to pediatricians and is necessary for treating acute metabolic decompensation. At presentation, there is often uncertainty about whether the patient has type 1 or type 2 diabetes. Experience with oral agents in children is limited and safety data are scant. Formal approval by the Food and Drug Administration (FDA) (and other national therapeutic goods agencies) for their use in childhood and adolescence is limited. Insulin may in addition convey a message of more serious illness, perhaps improving compliance.

The optimum schedule and type of insulin need to be determined in this age-group. It is a priority to study the optimum type of insulin, timing, intensity, duration, and follow-up in multicenter random controlled trials.

Metformin. This is approved in the U.S. for pediatric use. It is recommended as initial pharmacologic treatment in the absence of severe hyperglycemia. Data on metformin in children are limited, but a 16-week study (124,125) showed it to be safe and effective in 10- to 16-year-olds with type 2 diabetes. Long-term outcome studies are needed, as analysis of efficacy of metformin in a small group of Cree in northern Manitoba, Canada, showed little effect of metformin on HbA_{1c} or body

weight after 1 year among adolescents with type 2 diabetes (17). Gastrointestinal side effects were common with treatment. Poor adherence to oral therapy among relatively asymptomatic young persons with type 2 diabetes may be a major barrier to improvement in outcome.

Sulfonylureas. This class is well studied in adults; they are effective, safe, and inexpensive, although sulfonylureas have the potential to cause weight gain and hypoglycemia. They are the treatment of choice in patients with mutations in HNF1 α and HNF4 α , but only very low doses may be required (47).

Thiazolidinediones. This newer class of drugs is not yet approved for use in childhood and adolescence. Clinical studies for pediatric use are in progress, and although associated with weight gain, their reduction of visceral fat suggests overall benefit if long-term safety concerns can be addressed. Their safety profile with regard to edema and weight gain may be more favorable in children than adults, although this needs to be determined.

α -Glucosidase inhibitors. They have been used infrequently but are safe agents, although, like metformin, their gastrointestinal side effects may decrease acceptance among young persons.

Lipid-lowering therapy

The ADA has recently published consensus-based guidelines for the management of dyslipidemia in children (126). The recommendations for type 2 diabetes are that lipids should be measured every 2 years; optimal levels are as follows: LDL <2.6 mmol/l, HDL >0.9 mmol/l, and triglycerides <1.7 mmol/l. When optimal levels are not present, dietary advice should be provided and blood glucose should be treated aggressively. If after 6 months, LDL is >4.1 mmol/l, medication should be started (resins should be considered and if not accepted or tolerated, statins can be used). If LDL is 3.4–4.1 mmol/l after 6 months of diet and blood glucose management, medication should be considered in those with additional cardiovascular risk factors (including blood pressure, family history, and smoking status). Medication is not advised for the specific treatment of triglycerides (unless >11.2 mmol/l) or HDL.

Hypertension

Hypertension is a comorbidity of type 2 diabetes in youth and is a major risk factor

for nephropathy and atherosclerosis. Hypertension is defined in pediatric patients as an average systolic or diastolic blood pressure \geq 95th percentile for age, sex, and height, taken on three occasions. Blood pressure measurement should be obtained in all children with type 2 diabetes during diabetes health care visits. If hypertension does not respond to lifestyle changes, first-line pharmacological therapy for hypertension is an angiotensin-converting enzyme inhibitor, with angiotensin II receptor antagonists being reserved as second-line therapy due to little primary data in pediatric subjects.

Hypercoagulability

The risk of Reye's syndrome, which is extremely rare after puberty, has led the ADA to recommend against the use of aspirin in those <20 years of age.

Barriers to treatment

Attainment of treatment goals poses challenges, especially for the adolescent age-group when the developmental stage of moving toward independence is frequently marked by risk-taking behavior, lack of long-term planning, noncompliance, and resistance to lifestyle changes. Additional problems affecting the treatment of type 2 diabetes include family and/or psychosocial dysfunction and ethnic influences. Depression due to the high family burden of illness, poverty and isolation, and substance abuse are also not uncommon in lower socioeconomic groups and in many indigenous populations, leading to difficulties in the treatment of type 2 diabetes.

For young individuals, the important role of the family in diabetes management cannot be overestimated, as involvement of the family has been shown to lower HbA_{1c} (127).

PREVENTION — There is now very clear evidence supporting both lifestyle intervention and pharmaceutical agents in the prevention of type 2 diabetes in adults (128). However, the only evidence on diabetes prevention in children relates to breast-feeding, which may minimize excessive energy intake and perhaps improve insulin sensitivity by its higher polyunsaturated fat content. Studies of Pima Indians (129) and Native Canadian children (68) have shown a lower prevalence of type 2 diabetes among children and adolescents who were breast-fed dur-

ing infancy. Compared with formula feeding, breast-feeding has been associated with a lower weight for length and smaller skinfold thickness up to the age of 2 years (130) and with a lower plasma glucose in infants undergoing elective surgery (131).

Further relevant findings can be found in studies of overweight and obese children. In a 20-week study of 50 obese adolescents, those randomized to a weight loss program had, at the end of the study, lower serum insulin levels and blood pressure than those in the control group (132). In a similar study of 29 obese, hyperinsulinemic adolescents with a positive family history of type 2 diabetes randomized to metformin or placebo, BMI and fasting insulin improved modestly with metformin, but no change could be demonstrated in insulin sensitivity, HbA_{1c}, lipids, or glucose disposal (133).

In a trial of 192 children in two California schools, television and videotape viewing and video game use was reduced from 12 to 8 h/week in the intervention group, with no change in the control group. Those in the intervention group had a 0.45-kg/m² lesser increase in BMI and a 2.3-cm lesser increase in waist circumference during the 6-month study (134).

The "Trim and Fit" program in Singapore integrated nutrition education into the school curriculum, controlled the school canteens, encouraged water drinking by providing water coolers, targeted obese children for additional assistance, and rewarded schools achieving good health outcomes. Over 8 years of the program, from 1992 to 2000, the prevalence of obesity fell from 16.6 to 14.6% in 11- to 12-year-olds and from 15.5 to 13.1% in 15- to 16-year olds (135). An exercise intervention in Japan decreased the prevalence of overweight from 40 to 37% among boys and to 32% among girls between the ages of 10 and 13 years, with no change in a control group (136).

A school-based program among Mexican-American children has shown the importance of creating a network of social support in the classroom, the home, the school cafeteria, and among friends and classmates. In comparison to children in a control setting, those in the program have shown improved physical fitness, reduced numbers with fasting plasma glu-

cose >6.1 mmol/l, and reduced body fat (137,138).

Beyond individual and community-based interventions, to successfully prevent lifestyle diseases such as diabetes, changes in government policies and legislation are essential. Government intervention can include mandating a greater emphasis on more exercise and dietary education in schools, banning the advertising of unhealthy products, and subsidizing healthy food at the expense of less healthy food.

CONCLUSIONS— Until recently, type 2 diabetes has been viewed as a disease of older adults. With increasing rates of obesity, it is clear that the age of disease onset is falling in all ethnic groups and that type 2 diabetes is occurring in childhood (39). While much of the information is currently based on case reports and clinic-based series, and any generalizations should therefore be very guarded, the underlying problem of childhood obesity is, unfortunately, well documented. It would indeed be surprising if type 2 diabetes does not follow in its wake. The challenge to epidemiologists is to define its extent.

The pathophysiology of type 2 diabetes in children and adolescents appears to be very similar to that of adults. Insulin resistance (often exacerbated by puberty) is initially compensated by increased insulin secretion, but over time, β -cell function declines and hyperglycemia ensues. The pancreatic islet β -cell failure may occur very rapidly but the reasons for this are unclear.

The management of type 2 diabetes in the younger age-groups presents several major challenges. First, differentiation from type 1 diabetes can be difficult, notably when those presenting with DKA subsequently manifest many features typical of type 2 diabetes. Agreement on classification is required, and the framework needs to be useful in the clinical (both at the time of presentation and later on) and research settings. Successful treatment will require intensive efforts to alter lifestyle, which will need to focus on families as well as the individual. There do not appear to be any reasons why pharmacological treatment should differ from that used in adults, but more evidence is needed is about long-term safety and efficacy in relation to currently available oral hypoglycemic agents.

Prevention must remain a high priority and is only likely to be successful if governments and communities provide the environment within which individuals can make lifestyle changes that will prevent and, when necessary, reverse obesity. The school systems provide an ideal setting for prevention of obesity and diabetes, because the whole target population is available and both diet and physical activity can be influenced. In an ideal world, it would be possible to implement the proposed wide-reaching recommendations. In reality, this may be difficult given the socioeconomic constraints and the already tight health budgets of many governments. In addition, achieving good pregnancy outcomes with regard to avoiding low birth weight and treating gestational diabetes also provides an important prevention opportunity.

A consequence of urbanization is the parallel emergence of CVD, obesity, and type 2 diabetes, which until recently was mainly a problem of the developed world. Therefore, governments are going to be forced to deal with the problem of type 2 diabetes in children. As such, it would be better to address the problem as a public health issue under the heading of primary care and prevention instead of dealing with the consequences of an entrenched condition and its complications in a young population.

APPENDIX: WORKSHOP PARTICIPANTS

Organizing Group

- Co-chairperson: Sir George Alberti, FRCP (UK), Imperial College School of Medicine, Department of Diabetes, The Medical School. E-mail: george.alberti@ncl.ac.uk
- Co-chairperson: Paul Zimmet, AO (Australia), International Diabetes Institute. E-mail: pzimmet@idi.org.au
- Francine Kaufman, MD (USA), Children's Hospital Los Angeles. E-mail: fkaufman@chla.usc.edu
- Martin Silink, MD (Australia), Ray Williams Institute for Pediatric Endocrinology, Royal Alexandra Hospital for Children. E-mail: martins@chw.edu.au

Participants

- Stephanie Amiel, MD (UK), Guy's, King's and St. Thomas School of Medicine, Kings College. E-mail: Stephanie.amiel@kcl.ac.uk

- Silva Arslanian, MD (USA), Children's Hospital University of Pittsburgh. E-mail: silva.arslanian@chp.edu
- Peter Bennett, MD (USA), National Institute of Diabetes and Digestive Kidney Diseases. E-mail: pbennett@nih.gov
- Zachary Bloomgarden, MD (USA), Mt. Sinai Medical Center. E-mail: zblom@pol.net
- Sonia Caprio, MD (USA), Yale University School of Medicine. E-mail: sonia.caprio@yale.edu
- Juliana Chan, MD (China), The Chinese University of Hong Kong, Division of Clinical Pharmacology, Department of Medicine and Therapeutics, Prince of Wales Hospital. E-mail: jchan@cuhk.edu.hk
- Lee-Ming Chuang, MD (Taiwan), National Taiwan University Hospital. E-mail: leeming@ha.mc.ntu.edu.tw
- Dan Cooper, MD (USA), University of California, Irvine. E-mail: dcooper@uci.edu
- Paul Czernichow, MD (France), University of Paris, Hospital Robert Debre. E-mail: paul.czernichow@rdb.ap-hop-paris.fr
- Heather Dean, MD (Canada), University of Manitoba. E-mail: hdean@cc.umanitoba.ca
- Phillipe Halban, PhD (Switzerland), University of Geneva. E-mail: philippe.halban@medecine.unige.ch
- Andrew Hattersley, MD (UK), Diabetes & Vascular Medicine, Peninsula Medical School. E-mail: a.t.hattersley@ex.ac.uk
- Kenneth Lee Jones, MD (USA), University of California, San Diego, Department of Pediatrics. E-mail: kljones@ucsd.edu
- Richard Kahn, PhD (USA), American Diabetes Association. E-mail: rkhan@diabetes.org
- Kaichi Kida, MD, PhD (Japan), Department of Pediatrics, Ehime University, School of Medicine. E-mail: kaichi@m.ehime-u.ac.jp
- Pierre Lefebvre, MD, PhD (Belgium), Department of Medicine, University of Liege. E-mail: Pierre.lefebvre@ulg.ac.be
- Dinky Levitt, MD (South Africa), University of Cape Town. E-mail: dinky@uctgsh1.uct.ac.za
- Sir Michael Marmot, MD (UK), Department of Epidemiology and Public Health, University College London.

E-mail: m.marmot@ucl.ac.uk or michael@public-health.ucl.ac.uk

- Abdullah Nakhi, MD (Kuwait), Kuwait Diabetes Society. E-mail: ben_nakhi@hotmail.com
- Kerin O’Dea, PhD (Australia), Menzies School of Health Research, Royal Darwin Hospital. E-mail: kerin@menzies.edu.au
- Stefan Rössner, MD (Sweden), Obesity Unit, Huddinge University Hospital. E-mail: stephan.rossner@medhs.ki.se
- Jonathan Shaw, MD (Australia), International Diabetes Institute. E-mail: jshaw@idi.org.au
- Jay Skyler, MD (USA), Diabetes Research Institute. E-mail: jskyler@med.miami.edu
- Allen Spiegel, MD (USA), National Institute of Diabetes and Digestive and Kidney Diseases. E-mail: spiegelal@extra.niddk.nih.gov
- Robert Trevino, MD (USA), Social and Health Research Center. E-mail: rptrevino07@msn.com
- Frank Vinicor, MD (USA), Centers for Disease Control and Prevention. E-mail: fxv1@cdc.gov
- Rhys Williams, MD, PhD (UK), The Clinical School, University of Wales, Swansea. E-mail: d.r.r.williams@swansea.ac.uk
- Chittaragnan Yajnik, MD (India), King Edward Memorial Hospital. E-mail: csyajnik@hotmail.com

Acknowledgments—The Expert Workshop was supported by an educational grant from Johnson & Johnson.

References

1. Kitagawa T, Owada M, Urakami T, Yamachi K: Increased incidence of non-insulin dependent diabetes mellitus among Japanese school children correlates with an increased intake of animal protein and fat. *Clin Pediatr* 37:111–116, 1998
2. Dabelea D, Hanson RL, Bennett PH, Roumain J, Knowler WC, Pettitt DJ: Increasing prevalence of type II diabetes in American Indian children. *Diabetologia* 41:904–910, 1998
3. Braun B, Zimmerman MB, Kretchmer N, Spargo R, Smith R, Gracey M: Risk factors for diabetes and cardiovascular disease in young Australian Aborigines. *Diabetes Care* 19:472–479, 1996
4. Drake AJ, Smith A, Betts PR, Crowne EC, Shield JP: Type 2 diabetes in obese white children. *Arch Dis Child* 86:207–208,

- 2002
5. Ramachandran A, Snehalatha C, Satyavani K, Sivasankari S, Vijay V: Type 2 diabetes in Asian-Indian urban children. *Diabetes Care* 26:1022–1025, 2003
6. Zimmet P, Alberti KG, Shaw J: Global and societal implications of the diabetes epidemic. *Nature* 414:782–787, 2001
7. Sicree R, Shaw JE, Zimmet PZ: The global burden of diabetes. In *Diabetes Atlas*. 2nd ed. Gan D, Ed. Brussels, International Diabetes Federation, 2003, p. 15–71
8. Glatthaar C, Welborn TA, Stenhouse NS, Garcia-Webb P: Diabetes and impaired glucose tolerance: a prevalence estimate based on the Busselton 1981 survey. *Med J Australia* 143:436–440, 1985
9. Dunstan DW, Zimmet PZ, Welborn TA, De Courten MP, Cameron AJ, Sicree RA, Dwyer T, Colagiuri S, Jolley D, Knuiman M, Atkins R, Shaw JE: The rising prevalence of diabetes and impaired glucose tolerance: the Australian Diabetes, Obesity and Lifestyle Study. *Diabetes Care* 25: 829–834, 2002
10. Gu D, Reynolds K, Duan X, Xin X, Chen J, Wu X, Mo J, Whelton PK, He J: Prevalence of diabetes and impaired fasting glucose in the Chinese adult population: International Collaborative Study of Cardiovascular Disease in Asia (InterASIA). *Diabetologia* 46:1190–1198, 2003
11. Pihoker C, Scott CR, Lensing SY, Craddock MM, Smith J: Non-insulin dependent diabetes mellitus in African-American youths of Arkansas. *Clin Pediatr (Phila)* 37:97–102, 1998
12. American Diabetes Association: Type 2 diabetes in children and adolescents. *Diabetes Care* 23:381–389, 2000
13. Callahan ST, Mansfield MJ: Type 2 diabetes mellitus in adolescents. *Curr Opin Pediatr* 12:310–315, 2000
14. Dabelea D, Pettitt DJ, Jones KL, Arslanian SA: Type 2 diabetes mellitus in minority children and adolescents: an emerging problem. *Endocrinol Metab Clin North Am* 28:709–729, 1999
15. Wei JN, Sung FC, Lin CC, Lin RS, Chiang CC, Chuang LM: National surveillance for type 2 diabetes mellitus in Taiwanese children. *JAMA* 290:1345–1350, 2003
16. Freedman DS, Serdula MK, Percy CA, Ballew C, White L: Obesity, levels of lipids and glucose, and smoking among Navajo adolescents. *J Nutr* 127:2120S–2127S, 1997
17. Dean HJ, Young TK, Flett B, Wood-Steiman P: Screening for type-2 diabetes in aboriginal children in northern Canada (Letter). *Lancet* 352:1523–1524, 1998
18. Harris S, Gittelsohn J, Hanley A, Barnie A, Wolever T, Gao J, Logan A, Zinman B: The prevalence of NIDDM and associated risk factors in native Canadians. *Diabetes Care* 20:185–187, 1997
19. Fagot-Campagna A: Emergence of type 2 diabetes mellitus in children: epidemiological evidence. *J Pediatr Endocrinol Metab* 13 (Suppl. 6):1395–1402, 2000
20. Braun B, Zimmerman MB, Kretchmer N, Spargo RM, Smith RM, Gracey M: Risk factors for diabetes and cardiovascular disease in young Australian Aborigines: a 5-year follow-up study. *Diabetes Care* 19: 472–479, 1996
21. Fagot-Campagna A, Saaddine JB, Flegal KM, Beckles GL: Diabetes, impaired fasting glucose, and elevated HbA_{1c} in U. S. adolescents: the Third National Health and Nutrition Examination Survey. *Diabetes Care* 24:834–837, 2001
22. Harris SB, Perkins BA, Whalen-Brough E: Non-insulin-dependent diabetes mellitus among First Nations children: new entity among First Nations people of north western Ontario. *Can Fam Physician* 42: 869–876, 1996
23. Ehtisham S, Kirk J, McEvelly A, Shaw N, Rose Sea: Prevalence of type 2 diabetes in children in Birmingham (Letter). *Br Med J* 322:1428, 2001
24. Davis EA, Burrows MR, Russell MT, Jones TW: Increasing rates of type 2 diabetes in children and adolescents in Western Australia. Australian Diabetes Society and Australian Diabetes Educators Association Annual Scientific Meeting 2002. Adelaide, South Australia, 2002, p. 117
25. Glaser NS, Jones KL: Non-insulin dependent diabetes mellitus in Mexican-American children. *West J Med* 168:11–16, 1998
26. Pinhas-Hamiel O, Dolan L, Daniels S, Standiford D, Khoury P, Zeitler P: Increased incidence of non-insulin-dependent diabetes mellitus among adolescents. *J Pediatr* 128:608–615, 1996
27. Likitmaskul S, Kiattisathavee P, Chaichanwatanakul K, Punnakanta L, Angsusingha K, Tuchinda C: Increasing prevalence of type 2 diabetes mellitus in Thai children and adolescents associated with increasing prevalence of obesity. *J Pediatr Endocrinol Metab* 16:71–77, 2003
28. Palmert MR, Gordon CM, Kartashov AI, Legro RS, Emans SJ, Dunaif A: Screening for abnormal glucose tolerance in adolescents with polycystic ovary syndrome. *J Clin Endocrinol Metab* 87:1017–1023, 2002
29. Lipton R, Keenan H, Onyemere KU, Freels S: Incidence and onset features of diabetes in African-American and Latino children in Chicago, 1985–1994. *Diabetes Metab Res Rev* 18:135–142, 2002
30. Acton KJ, Burrows NR, Moore K, Querec L, Geiss LS, Engelgau MM: Trends in diabetes prevalence among American Indian and Alaska native children, adolescents, and young adults. *Am J Public Health* 92:1485–1490, 2002

31. Macaluso CJ, Bauer UE, Deeb LC, Malone JI, Chaudhari M, Silverstein J, Eidson M, Goldberg RB, Gaughan-Bailey B, Brooks RG, Rosenbloom AL: Type 2 diabetes mellitus among Florida children and adolescents, 1994 through 1998. *Public Health Rep* 117:373–379, 2002
32. Neufeld ND, Raffel LJ, Landon C, Chen Y-D, Vadheim CM: Early presentation of type 2 diabetes in Mexican-American youth. *Diabetes Care* 21:80–86, 1998
33. Holl RW, Grabert M, Krause U, Rami B, Schober E, Schweiggert F, Thon A: Prevalence and clinical characteristics of patients with non-type-1-diabetes in the pediatric age range: analysis of a multicenter database including 20,401 patients from 148 centers in Germany and Austria (Abstract). *Diabetologia* 46 (Suppl. 2):A26, 2003
34. Zachrisson I, Tibell C, Bang P, Ortqvist E: Prevalence of type 2 diabetes among known cases of diabetes aged 0–18 years in Sweden (Abstract). *Diabetologia* 46 (Suppl. 2):A25, 2003
35. Dean H, Flett B: Natural history of type 2 diabetes diagnosed in childhood: long term follow-up in young adult years (Abstract). *Diabetes* 51 (Suppl. 1):A24, 2002
36. Yokoyama H, Okudaira M, Otani T, Sato A, Miura J, Takaike H, Yamada H, Muto K, Uchigata Y, Ohashi Y, Iwamoto Y: Higher incidence of diabetic nephropathy in type 2 than in type 1 diabetes in early-onset diabetes in Japan. *Kidney Int* 58:302–311, 2000
37. Krakoff J, Lindsay RS, Looker HC, Nelson RG, Hanson RL, Knowler WC: Incidence of retinopathy and nephropathy in youth-onset compared with adult-onset type 2 diabetes. *Diabetes Care* 26:76–81, 2003
38. Banerji M, Chaiken R, Huey H, Tuomi T, Norin A, Mackay I, Rowley M, Zimmet P, Lebovitz H: GAD antibody negative NIDDM in adult black subjects with diabetic ketoacidosis and increased frequency of human leukocyte antigen DR3 and DR4: Flatbush diabetes. *Diabetes* 43:741–745, 1994
39. Winter WE, Maclaren NK, Riley WJ, Clarke DW, Kappy MS, Spillar RP: Maturity-onset diabetes of youth in black Americans. *N Engl J Med* 316:285–291, 1987
40. Pinhas-Hamiel O, Dolan LM, Zeitler PS: Diabetic ketoacidosis among obese African-American adolescents with NIDDM. *Diabetes Care* 20:484–486, 1997
41. Banerji MA: Impaired beta-cell and alpha-cell function in African-American children with type 2 diabetes mellitus—“Flatbush diabetes.” *J Pediatr Endocrinol Metab* 15 (Suppl 1):493–501, 2002
42. Sellers EA, Dean HJ: Diabetic ketoacidosis: a complication of type 2 diabetes in Canadian aboriginal youth. *Diabetes Care* 23:1202–1204, 2000
43. Lee YS, Ng WY, Thai AC, Lui KF, Loke KY: Prevalence of ICA and GAD antibodies at initial presentation of type 1 diabetes mellitus in Singapore children. *J Pediatr Endocrinol Metab* 14:767–772, 2001
44. Thai AC, Ng WY, Loke KY, Lee WR, Lui KF, Cheah JS: Anti-GAD antibodies in Chinese patients with youth and adult-onset IDDM and NIDDM. *Diabetologia* 40:1425–1430, 1997
45. Hattersley A: Maturity-onset diabetes of the young clinical heterogeneity explained by genetic heterogeneity. *Diabet Med* 15:15–24, 1998
46. Maassen JA: Mitochondrial diabetes: pathophysiology, clinical presentation, and genetic analysis. *Am J Med Genet* 115:66–70, 2002
47. Pearson ER, Starkey BJ, Powell RJ, Gribble FM, Clark PM, Hattersley AT: Genetic cause of hyperglycaemia and response to treatment in diabetes. *Lancet* 362:1275–1281, 2003
48. Ehtisham S, Barrett TG: The emergence of type 2 diabetes in childhood. *Ann Clin Biochem* 41:10–16, 2004
49. Porte D Jr: Banting Lecture 1990: β -cells in type II diabetes mellitus. *Diabetes* 40:166–180, 1991
50. Mooy J, Grootenhuys P, de Vries H, Valkenburg H, Bouter L, Kostense P, Heine R: Prevalence and determinants of glucose intolerance in a Dutch Caucasian population: the Hoorn study. *Diabetes Care* 18:1270–1273, 1995
51. Stern MP: The search for type 2 diabetes susceptibility genes using whole-genome scans: an epidemiologist's perspective. *Diabetes Metab Res Rev* 18:106–113, 2002
52. Altshuler D, Hirschhorn JN, Klannemark M, Lindgren CM, Vohl MC, Nemesh J, Lane CR, Schaffner SF, Bolk S, Brewer C, Tuomi T, Gaudet D, Hudson TJ, Daly M, Groop L, Lander ES: The common PPAR- γ Pro12Ala polymorphism is associated with decreased risk of type 2 diabetes. *Nat Genet* 26:76–80, 2000
53. Gloyn AL, Weedon MN, Owen KR, Turner MJ, Knight BA, Hitman G, Walker M, Levy JC, Sampson M, Halford S, McCarthy MI, Hattersley AT, Frayling TM: Large-scale association studies of variants in genes encoding the pancreatic β -cell KATP channel subunits Kir6.2 (KCNJ11) and SUR1 (ABCC8) confirm that the KCNJ11 E23K variant is associated with type 2 diabetes. *Diabetes* 52:568–572, 2003
54. Hegele RA, Zinman B, Hanley AJ, Harris SB, Barrett PH, Cao H: Genes, environment and Oji-Cree type 2 diabetes. *Clin Biochem* 36:163–170, 2003
55. Sellers EA, Triggs-Raine B, Rockman-Greenberg C, Dean HJ: The prevalence of the HNF-1 α G319S mutation in Canadian aboriginal youth with type 2 diabetes. *Diabetes Care* 25:2202–2206, 2002
56. Narayan KM, Fagot-Campagna A, Imperatore G: Type 2 diabetes in children: a problem lurking for India? *Indian Pediatr* 38:701–704, 2001
57. Ogden CL, Flegal KM, Carroll MD, Johnson CL: Prevalence and trends in overweight among US children and adolescents, 1999–2000. *JAMA* 288:1728–1732, 2002
58. Strauss RS, Pollack HA: Epidemic increase in childhood overweight, 1986–1998. *JAMA* 286:2845–2848, 2001
59. Kimm SY, Gergen PJ, Malloy M, Dresser C, Carroll M: Dietary patterns of U. S. children: implications for disease prevention. *Prev Med* 19:432–442, 1990
60. Desmond SM, Price JH, Hallinan C, Smith D: Black and white adolescents' perceptions of their weight. *J Sch Health* 59:353–358, 1989
61. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH: Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* 320:1240–1243, 2000
62. Luo J, Hu FB: Time trends of obesity in pre-school children in China from 1989 to 1997. *Int J Obes Relat Metab Disord* 26:553–558, 2002
63. Sung RY, Tong PC, Yu CW, Lau PW, Mok GT, Yam MC, Lam PK, Chan JC: High prevalence of insulin resistance and metabolic syndrome in overweight/obese pre-adolescent Hong Kong Chinese children aged 9–12 years (Letter). *Diabetes Care* 26:250–251, 2003
64. McCarthy HD, Ellis SM, Cole TJ: Central overweight and obesity in British youth aged 11–16 years: cross sectional surveys of waist circumference. *BMJ* 326:624, 2003
65. Booth ML, Chey T, Wake M, Norton K, Hesketh K, Dollman J, Robertson I: Change in the prevalence of overweight and obesity among young Australians, 1969–1997. *Am J Clin Nutr* 77:29–36, 2003
66. Booth ML, Wake M, Armstrong T, Chey T, Hesketh K, Mathur S: The epidemiology of overweight and obesity among Australian children and adolescents, 1995–97. *Aust N Z J Public Health* 25:162–169, 2001
67. Ramachandran A, Snehalatha C, Vinita R, Thayyil M, Kumar CK, Sheeba L, Joseph S, Vijay V: Prevalence of overweight in urban Indian adolescent school children. *Diabetes Res Clin Pract* 57:185–190, 2002
68. Young TK, Martens PJ, Taback SP, Sellers EA, Dean HJ, Cheang M, Flett B: Type 2 diabetes mellitus in children: prenatal and early infancy risk factors among native Canadians. *Arch Pediatr Adolesc Med* 156:

- 651–655, 2002
69. Kimm SY, Glynn NW, Kriska AM, Fitzgerald SL, Aaron DJ, Similo SL, McMahon RP, Barton BA: Longitudinal changes in physical activity in a biracial cohort during adolescence. *Med Sci Sports Exerc* 32:1445–1454, 2000
 70. Nesmith JD: Type 2 diabetes mellitus in children and adolescents. *Pediatr Rev* 22: 147–152, 2001
 71. Troiano RP: Physical inactivity among young people. *N Engl J Med* 347:706–707, 2002
 72. Lowry R, Wechsler H, Galuska DA, Fulton JE, Kann L: Television viewing and its associations with overweight, sedentary lifestyle, and insufficient consumption of fruits and vegetables among US high school students: differences by race, ethnicity, and gender. *J Sch Health* 72:413–421, 2002
 73. Campbell K, Crawford D, Jackson M, Cashel K, Worsley A, Gibbons K, Birch LL: Family food environments of 5–6-year-old-children: does socioeconomic status make a difference? *Asia Pac J Clin Nutr* 11 (Suppl. 3):S553–S561, 2002
 74. Hammond KM, Wyllie A, Casswell S: The extent and nature of televised food advertising to New Zealand children and adolescents. *Aust N Z J Public Health* 23:49–55, 1999
 75. Chestnutt IG, Ashraf FJ: Television advertising of foodstuffs potentially detrimental to oral health: a content analysis and comparison of children's and primetime broadcasts. *Community Dent Health* 19:86–89, 2002
 76. Gungor N, Arslanian S: Pathophysiology of type 2 diabetes mellitus in children and adolescents: treatment implications. *Treat Endocrinol* 1:359–371, 2002
 77. Arslanian S: Type 2 diabetes in children: clinical aspects and risk factors. *Horm Res* 57 (Suppl. 1):19–28, 2002
 78. Caprio S, Weiss R, Tamborlane W: Ethics of research involving vulnerable populations. *Lancet* 362:1857–1858, 2003
 79. Smith CP, Dunger DB, Williams AJ, Taylor AM, Perry LA, Gale EA, Preece MA, Savage MO: Relationship between insulin, insulin-like growth factor I, and dehydroepiandrosterone sulfate concentrations during childhood, puberty, and adult life. *J Clin Endocrinol Metab* 68:932–937, 1989
 80. Amiel SA, Sherwin RS, Simonson DC, Lauritano AA, Tamborlane WV: Impaired insulin action in puberty: a contributing factor to poor glycemic control in adolescents with diabetes. *N Engl J Med* 315:215–219, 1986
 81. Bloch CA, Clemons P, Sperling MA: Puberty decreases insulin sensitivity. *J Pediatr* 110:481–487, 1987
 82. Danadian K, Balasekaran G, Lewy V, Meza MP, Robertson R, Arslanian SA: Insulin sensitivity in African-American children with and without family history of type 2 diabetes. *Diabetes Care* 22:1325–1329, 1999
 83. Arslanian SA, Saad R, Lewy V, Danadian K, Janosky J: Hyperinsulinemia in African-American children: decreased insulin clearance and increased insulin secretion and its relationship to insulin sensitivity. *Diabetes* 51:3014–3019, 2002
 84. Goran MI, Bergman RN, Cruz ML, Watanabe R: Insulin resistance and associated compensatory responses in African-American and Hispanic children. *Diabetes Care* 25:2184–2190, 2002
 85. Whincup PH, Gilg JA, Papacosta O, Seymour C, Miller GJ, Alberti KG, Cook DG: Early evidence of ethnic differences in cardiovascular risk: cross sectional comparison of British South Asian and white children. *BMJ* 324:635, 2002
 86. Kobayashi K, Amemiya S, Higashida K, Ishihara T, Sawanobori E, Mochizuki M, Kikuchi N, Tokuyama K, Nakazawa S: Pathogenic factors of glucose intolerance in obese Japanese adolescents with type 2 diabetes. *Metabolism* 49:186–191, 2000
 87. Freedman DS, Srinivasan SR, Burke GL, Shear CL, Smoak CG, Harsha DW, Weber LS, Berenson GS: Relation of body fat distribution to hyperinsulinemia in children and adolescents: the Bogalusa Heart Study. *Am J Clin Nutr* 46:403–410, 1987
 88. Bacha F, Saad R, Gungor N, Janosky J, Arslanian SA: Obesity, regional fat distribution, and syndrome X in obese black versus white adolescents: race differential in diabetogenic and atherogenic risk factors. *J Clin Endocrinol Metab* 88: 2534–2540, 2003
 89. Aaron DJ, Kriska AM, Dearwater SR, Anderson RL, Olsen TL, Cauley JA, LaPorte RE: The epidemiology of leisure physical activity in an adolescent population. *Med Sci Sports Exerc* 25:847–853, 1993
 90. Schmitz KH, Jacobs DR, Jr, Hong CP, Steinberger J, Moran A, Sinaiko AR: Association of physical activity with insulin sensitivity in children. *Int J Obes Relat Metab Disord* 26:1310–1316, 2002
 91. Scott R, Brown L: Rates and causes of mortality in IDDM: a nine year follow-up study (Abstract). *Diabetes* 46 (Suppl. 1): 146A, 1997
 92. Fagot-Campagna A, Pettitt D, Engelgau M, Burrows N, Geiss L, Valdez R, Beckles G, Saaddine J, Gregg E, Williamson D, Narayan K: Type 2 diabetes among North American children and adolescents: an epidemiologic review and a public health perspective. *J Pediatr* 136:664–672, 2000
 93. Lewy VD, Danadian K, Witchel SF, Arslanian S: Early metabolic abnormalities in adolescent girls with polycystic ovarian syndrome. *J Pediatr* 138:38–44, 2001
 94. Arslanian SA, Lewy VD, Danadian K: Glucose intolerance in obese adolescents with polycystic ovary syndrome: roles of insulin resistance and beta-cell dysfunction and risk of cardiovascular disease. *J Clin Endocrinol Metab* 86:66–71, 2001
 95. Phillips DI: Birth weight and the future development of diabetes: a review of the evidence. *Diabetes Care* 21 (Suppl. 2): B150–B155, 1998
 96. Wei JN, Sung FC, Li CY, Chang CH, Lin RS, Lin CC, Chiang CC, Chuang LM: Low birth weight and high birth weight infants are both at an increased risk to have type 2 diabetes among schoolchildren in Taiwan. *Diabetes Care* 26:343–348, 2003
 97. Bhargava SK, Sachdev HS, Fall CH, Osmond C, Lakshmy R, Barker DJ, Biswas SK, Ramji S, Prabhakaran D, Reddy KS: Relation of serial changes in childhood body-mass index to impaired glucose tolerance in young adulthood. *N Engl J Med* 350:865–875, 2004
 98. Bavdekar A, Yajnik CS, Fall CH, Bapat S, Pandit AN, Deshpande V, Bhawe S, Kellingray SD, Joglekar C: Insulin resistance syndrome in 8-year-old Indian children: small at birth, big at 8 years, or both? *Diabetes* 48:2422–2429, 1999
 99. Pettitt DJ, Nelson RG, Saad MF, Bennett PH, Knowler WC: Diabetes and obesity in the offspring of Pima Indian women with diabetes during pregnancy. *Diabetes Care* 16:310–314, 1993
 100. Silverman BL, Metzger BE, Cho NH, Loeb CA: Impaired glucose tolerance in adolescent offspring of diabetic mothers: relationship to fetal hyperinsulinism. *Diabetes Care* 18:611–617, 1995
 101. Stride A, Shepherd M, Frayling TM, Bulman MP, Ellard S, Hattersley AT: Intrauterine hyperglycemia is associated with an earlier diagnosis of diabetes in HNF-1 α gene mutation carriers. *Diabetes Care* 25:2287–2291, 2002
 102. Sinha R, Fisch G, Teague B, Tamborlane WV, Banyas B, Allen K, Savoye M, Rieger V, Taksali S, Barbetta G, Sherwin RS, Caprio S: Prevalence of impaired glucose tolerance among children and adolescents with marked obesity. *N Engl J Med* 346:802–810, 2002
 103. Glowinska B, Urban M, Koput A: Cardiovascular risk factors in children with obesity, hypertension and diabetes: lipoprotein(a) levels and body mass index correlate with family history of cardiovascular disease. *Eur J Pediatr* 161:511–518, 2002
 104. McCance D, Pettitt D, Hanson R, Jacobsson L, Knowler W, Bennett P: Birth weight and non-insulin dependent diabetes: thrifty genotype, thrifty pheno-

- type, or surviving small baby genotype? *Br Med J* 308:942–945, 1994
105. Otani TO, Y. Shichiri, M.: [The present status of juvenile-onset NIDDM]. In *Progress in Diabetes Care and Education 1992*. Japan Diabetes Society, Ed. Tokyo, Shindan-to-Chiryosha, 1992, p. 155–159
 106. Owada M: [The present status and trend of lifestyle-related diseases in children: type II diabetes mellitus]. *Modern Physician* 19:829–834, 1999
 107. Pinhas-Hamiel O, Standiford D, Hamiel D, Dolan LM, Cohen R, Zeitler PS: The type 2 family: a setting for development and treatment of adolescent type 2 diabetes mellitus. *Arch Pediatr Adolesc Med* 153:1063–1067, 1999
 108. Rosenbloom A, Joe J, Young R, Winter W: Emerging epidemic of type 2 diabetes in youth. *Diabetes Care* 22:345–354, 1999
 109. Tang M, Chen Y, Krewski D: Gender-related differences in the association between socioeconomic status and self-reported diabetes. *Int J Epidemiol* 32: 381–385, 2003
 110. Brunner EJ, Marmot MG, Nanchahal K, Shipley MJ, Stansfeld SA, Juneja M, Alberti KG: Social inequality in coronary risk: central obesity and the metabolic syndrome: evidence from the Whitehall II Study. *Diabetologia* 40:1341–1349, 1997
 111. abu Sayeed M, Ali L, Hussain MZ, Rumi MA, Banu A, Azad Khan AK: Effect of socioeconomic risk factors on the difference in prevalence of diabetes between rural and urban populations in Bangladesh. *Diabetes Care* 20:551–555, 1997
 112. Armstrong J, Dorosty AR, Reilly JJ, Emmett PM: Coexistence of social inequalities in undernutrition and obesity in preschool children: population based cross sectional study. *Arch Dis Child* 88: 671–675, 2003
 113. Borch-Johnsen K, Beck-Nielsen H, Christiansen JS, Heickendorff L, Brandslund I, Faber JO, de Fine Olivarius N: [Guidelines on diagnosis of type 2 diabetes: investigation, diagnosis and glucose measurement]. *Ugeskr Laeger* 165: 1558–1561, 2003
 114. Fagot-Campagna A, Saaddine JB, Engलगau MM: Is testing children for type 2 diabetes a lost battle? *Diabetes Care* 23: 1442–1443, 2000
 115. Genuth S: Lowering the criterion for impaired fasting glucose is in order (Commentary). *Diabetes Care* 26:3331–3332, 2003
 116. Shaw J, Zimmet P, McCarty D, de Courten M: Type 2 diabetes worldwide according to the new classification and criteria. *Diabetes Care* 23 (Suppl. 2):B5–B10, 2000
 117. Shaw J, Hodge A, de Courten M, Chitson P, Zimmet P: Isolated post-challenge hyperglycaemia confirmed as a risk factor for mortality. *Diabetologia* 42:1050–1054, 1999
 118. Davidson MB: Counterpoint: the oral glucose tolerance test is superfluous. *Diabetes Care* 25:1883–1885, 2002
 119. The cost-effectiveness of screening for type 2 diabetes: CDC Diabetes Cost-Effectiveness Study Group, Centers for Disease Control and Prevention. *JAMA* 280:1757–1763, 1998
 120. Adriaanse MC, Snoek FJ, Dekker JM, Spijkerman AM, Nijpels G, van der Ploeg HM, Heine RJ: Perceived risk for type 2 diabetes in participants in a stepwise population-screening programme. *Diabet Med* 20:210–215, 2003
 121. Silverstein JH, Rosenbloom AL: Treatment of type 2 diabetes mellitus in children and adolescents. *J Pediatr Endocrinol Metab* 13 (Suppl. 6):1403–1409, 2000
 122. Sellers EHD: Short term insulin therapy in youth with type 2 diabetes (Abstract). *Diabetes* 51 (Suppl. 1):A613, 2002
 123. Zuhri-Yafi MI, Brosnan PG, Hardin DS: Treatment of type 2 diabetes mellitus in children and adolescents. *J Pediatr Endocrinol Metab* 15 (Suppl. 1):541–546, 2002
 124. Jones KL, Arslanian S, Peterokova VA, Park JS, Tomlinson MJ: Effect of metformin in pediatric patients with type 2 diabetes: a randomized controlled trial. *Diabetes Care* 25:89–94, 2002
 125. Arslanian SA, Lewy V, Danadian K, Saad R: Metformin therapy in obese adolescents with polycystic ovary syndrome and impaired glucose tolerance: amelioration of exaggerated adrenal response to adrenocorticotropin with reduction of insulinemia/insulin resistance. *J Clin Endocrinol Metab* 87:1555–1559, 2002
 126. American Diabetes Association: Management of dyslipidemia in children and adolescents with diabetes (Consensus Statement). *Diabetes Care* 26:2194–2197, 2003
 127. Bradshaw B: The role of the family in managing therapy in minority children with type 2 diabetes mellitus. *J Pediatr Endocrinol Metab* 15 (Suppl. 1):547–551, 2002
 128. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM: Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 346: 393–403, 2002
 129. Pettitt D, Forman M, Hanson R, Knowler W, Bennett P: Breast feeding in infancy is associated with lower rates of non-insulin-dependent diabetes mellitus. *Lancet* 350:166–168, 1997
 130. Dewey KG, Heinig MJ, Nommsen LA, Peerson JM, Lonnerdal B: Breast-fed infants are leaner than formula-fed infants at 1 y of age: the DARLING Study. *Am J Clin Nutr* 57:140–145, 1993
 131. Baur LA, O'Connor J, Pan DA, Kriketos AD, Storlien LH: The fatty acid composition of skeletal muscle membrane phospholipid: its relationship with the type of feeding and plasma glucose levels in young children. *Metabolism* 47:106–112, 1998
 132. Rocchini AP, Katch V, Schork A, Kelch RP: Insulin and blood pressure during weight loss in obese adolescents. *Hypertension* 10:267–273, 1987
 133. Freemark M, Bursey D: The effects of metformin on body mass index and glucose tolerance in obese adolescents with fasting hyperinsulinemia and a family history of type 2 diabetes. *Pediatrics* 107: E55, 2001
 134. Robinson TN: Reducing children's television viewing to prevent obesity: a randomized controlled trial. *JAMA* 282: 1561–1567, 1999
 135. Toh CM, Cutter J, Chew SK: School based intervention has reduced obesity in Singapore (Letter). *BMJ* 324:427, 2002
 136. Kida K, Ito T, Yang SW, Tamphaichitr V: Effects of Western diet on risk factors of chronic diseases in Asia. In *Preventive Nutrition: The Comprehensive Guide for Health Professionals*. Bendich A, Deckelbaum RJ, Eds. Totowa, NJ, Humana Press, 2001
 137. Trevino RP, Dempsey L, Garcoa O, Ramirez G, Yin Z: Reversing hyperglycemia in low-income Latino children participating in the Beinstar (Abstract). *Diabetes* 51 (Suppl. 1):A430, 2002
 138. Trevino RP, Pugh JA, Hernandez AE, Menchaca VD, Ramirez RR, Mendoza M: Bienestar: a diabetes risk-factor prevention program. *J Sch Health* 68:62–67, 1998