

Increased Adiposity at Diagnosis in Younger Children With Type 1 Diabetes Does Not Persist

SAMANTHA L. CLARKE, MND¹
 MARIA E. CRAIG, PHD, FRACP^{1,2,3}
 SARAH P. GARNETT, MND, PHD¹
 ALBERT K. CHAN, MAPPSTAT¹
 CHRISTOPHER T. COWELL, FRACP¹

JANINE M. CUSUMANO, RN¹
 OLGA KORDONOURI, MD⁴
 ABI SAMBASIVAN, MBBCHIR⁵
 KIM C. DONAGHUE, PHD, FRACP^{1,3}

Increased adiposity, as measured by BMI, has been described in younger children at diagnosis of type 1 diabetes compared with older children (1–3) in support of the “accelerator hypothesis” (1). Australia has one of the highest rates of childhood obesity in the Western world, but a disproportionate increase in type 1 diabetes incidence has not been observed in young children (4), which is contrary to recent reports of other populations (5,6). We therefore sought to investigate whether anthropometric data at diagnosis from 1976 to 2004 support the accelerator hypothesis in our population.

In addition to its potential contribution to type 1 diabetes etiology, the insulin resistance associated with increased adiposity places these children at greater risk of vascular disease (7–9). In a cross-sectional study of children with median diabetes duration of 3 years, we have previously found that those with the highest BMI SD score (SDS) were younger and had worse metabolic control (10). Hence, in this longitudinal study, we examined whether increased adiposity at diagnosis persisted in the youngest children after 5 years.

RESEARCH DESIGN AND METHODS

The study group consisted of children and adolescents attend-

ing the diabetes outpatient and outreach clinics of the Royal Alexandra Hospital for Children in Sydney and rural New South Wales, Australia, from 1976 to 2004. To exclude the initial weight loss due to acute destabilization, only height, weight, and BMI available 1–3 months postdiagnosis were analyzed ($n = 960$). In those diagnosed from 1976 to 2000 ($n = 661$), height, weight, and BMI were available in 443 children (67%) after a 5-year diabetes duration (range 4–6 years). Patients were stratified into three groups according to their age at diagnosis, 2–4.9 ($n = 68$), 5–9.9 ($n = 220$), and 10–14.9 years ($n = 155$).

Height was measured by Harpenden stadiometer, and weight was measured with digital scales. Height, weight, and BMI were expressed as SDSs using the Centers for Disease Control and Prevention 2000 growth data for the entire cohort to enable valid comparison over time (11). Data were analyzed using SAS version 8 (Cary, NC). The Kruskal-Wallis test was used to compare SDSs across the three age-groups. The signed-rank test was used to test whether SDSs changed 5 years after diagnosis. Factors associated with BMI SDSs (year, age at diagnosis, insulin dose, and their interaction terms) were examined using multiple regression.

RESULTS— In the entire cohort ($n = 960$), younger age at diagnosis was associated with higher BMI SDSs ($\beta = -0.05$, $P < 0.0001$); however, BMI SDSs did not change over the period 1976–2004 (regression coefficient for year: 0.011, $P = 0.11$). There was no significant interaction between age and year of diagnosis in the model, and no correlation between age at diagnosis and year of diagnosis ($r = 0.05$, $P = 0.12$).

In the longitudinal group of 443 subjects, height, weight, and BMI SDSs at diagnosis were higher in the youngest age-group compared with both the older groups ($P = 0.05$, $P < 0.001$, and $P = 0.004$, respectively) (Fig. 1).

Five years after diagnosis, differences in anthropometric measures were no longer significant across the three age-groups. Height and weight SDSs decreased significantly in the <5 years age-group ($P = 0.001$ and $P = 0.01$, respectively) (Fig. 1A), indicating regression toward the mean. In contrast, BMI SDSs increased significantly in the 10–14.9 years age-group ($P < 0.001$) (Fig. 1B).

At diagnosis, the median insulin dose was 0.44 units/kg (interquartile range 0.32–0.60) and was not significantly different across the three age-groups. After 5 years, insulin dose significantly increased in all groups (0.89, 1.19, and 1.12 units/kg in the 2–4.9, 5–9.9, and 10–14.9 years age-groups, respectively) ($P < 0.001$). The median HbA_{1c} (A1C) was 8.3% and was not significantly different between the three age-groups.

CONCLUSIONS— Younger children were taller and had higher BMI SDSs at diagnosis than older children, which is in line with previous studies (1,2,12). Our new finding is that after 5 years of diabetes duration, the <5 years age-group demonstrated a deceleration of growth with a decrease in height and weight SDSs. This is in contrast to the trend toward increasing BMI and weight SDSs in the general Australian population (13). The decrease in height SDSs could indicate that growth is slowing, following a period of rapid growth, but may also

From ¹The Institute of Endocrinology and Diabetes, The Children's Hospital at Westmead, Sydney, Australia; the ²School of Women's and Children's Health, University of New South Wales, Kensington, New South Wales, Australia; the ³Discipline of Paediatrics and Child Health, University of Sydney, New South Wales, Australia; ⁴Kinderkrankenhaus auf der Bult, Hannover, Germany; and ⁵The University of Cambridge, Cambridge, U.K.

Address correspondence and reprint requests to Dr. Kim C. Donaghue, Institute of Endocrinology and Diabetes, The Children's Hospital at Westmead, Locked Bag 4001, Westmead, NSW 2145, Australia. E-mail: kimd@chw.edu.au.

Received for publication 2 February 2006 and accepted in revised form 11 April 2006.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

DOI: 10.2337/dc06-0277

© 2006 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.




Figure 1— A: Comparison of height SDSs across three age-groups at diagnosis of type 1 diabetes (\square , $P = 0.05$) and after 5 years (\square , $P = 0.11$). B: Comparison of BMI SDSs across three age-groups at diagnosis ($P = 0.004$) and after 5 years ($P = 0.41$). Between the two time points, height SDSs decreased in the 2–4.9 years age-group ($P = 0.001$), and BMI SDSs increased in the 10–14.9 years group ($P = 0.0001$).

represent a loss of height associated with suboptimal glycemic control.

Recent studies have found a rise in BMI SDSs in children at diagnosis of type 1 diabetes, suggesting an increased role for adiposity as an environmental risk factor (1,14). Even though the Australian pediatric population as a whole has increased their BMI (13), there has not been an increase in BMI at diagnosis of type 1 diabetes; however, the youngest children have been overweight for almost 30 years. This temporarily supports the “accelerator hypothesis” as a consistent mechanism underlying diabetes over this time in Australia, particularly in young children. Combined with other recent studies (1–3,12,14), we speculate that there may be a threshold at which higher adiposity is associated with earlier onset of type 1 diabetes and that this threshold was reached some time ago in Australia (earlier than in Western Europe). Indeed, the BMI SDS at diagnosis was 0.93 in our 2–5 years age-group, which is higher than in the same age-group in the U.K. (0.21) and Germany (0.5) (12). Recent data from the U.S. also demonstrate an association between higher BMI and younger age at diagnosis of type 1 diabetes (3), but only in patients with low fasting C-peptide levels, suggesting that acceleration occurs late in the

evolution of type 1 diabetes, when β -cell function is already compromised.

While the younger group demonstrated a decrease in growth parameters after 5 years of diabetes, conversely, the oldest group increased their weight and BMI SDSs. The weight gain in the older children, who were less overweight at diagnosis, is likely to be related to changes of puberty and resulted in increased insulin requirements.

The deceleration in weight and BMI are encouraging for the future health of children with very-early-onset diabetes. While this may be a natural phenomenon after growth acceleration, such changes may also be influenced by frequent contact with health professionals discussing meal plans and demonstrating individual measurement on growth charts. Older children are at risk of sustained weight gain from diagnosis and may be an additional risk factor for development of microvascular complications.

References

1. Betts P, Mulligan J, Ward P, Smith B, Wilkin T: Increasing body weight predicts the earlier onset of insulin-dependent diabetes in childhood: testing the ‘accelerator hypothesis’ (2). *Diabet Med* 22:144–151, 2005
2. Kibirige M, Metcalf B, Renuka R, Wilkin TJ: Testing the accelerator hypothesis: the

relationship between body mass and age at diagnosis of type 1 diabetes. *Diabetes Care* 26:2865–2870, 2003

3. Dabelea D, D’Agostino RB Jr, Mayer-Davis EJ, Pettitt DJ, Imperatore G, Dolan LM, Pihoker C, Hillier TA, Marcovina SM, Linder B, Ruggiero AM, Hamman RF: Testing the accelerator hypothesis: body size, β -cell function, and age at onset of type 1 (autoimmune) diabetes. *Diabetes Care* 29:290–294, 2006
4. Taplin CE, Craig ME, Lloyd M, Talyor C, Crock P, Silink M, Howard NJ: The rising incidence of childhood type 1 diabetes in New South Wales, 1990–2002. *Med J Aust* 183:243–246, 2005
5. Gardner SG, Bingley PJ, Sawtell PA, Weeks S, Gale EA: Rising incidence of insulin dependent diabetes in children aged under 5 years in the Oxford region: time trend analysis: the Bart’s-Oxford Study Group. *BMJ* 315:713–717, 1997
6. Schoenle EJ, Lang-Muritano M, Gschwend S, Laimbacher J, Mullis PE, Torresani T, Bion-Laubert AML: Epidemiology of type 1 diabetes mellitus in Switzerland: steep rise in incidence in under 5 year old children in the past decade. *Diabetologia* 44:286–289, 2001
7. Yip J, Mattock MB, Morocutti A, Sethi M, Trevisan R, Viberti G: Insulin resistance in insulin-dependent diabetic patients with microalbuminuria. *Lancet* 342:883–887, 1993
8. Amin R, Williams RM, Frystyk J, Umpleby M, Matthews D, Orskov H, Dal-

- ton RN, Dunger DB: Increasing urine albumin excretion is associated with growth hormone hypersecretion and reduced clearance of insulin in adolescents and young adults with type 1 diabetes: the Oxford Regional Prospective Study. *Clin Endocrinol (Oxf)* 62:137–144, 2005
9. Amin R, Schultz C, Ong K, Frystyk J, Dalton RN, Perry L, Orskov H, Dunger DB: Low IGF-I and elevated testosterone during puberty in subjects with type 1 diabetes developing microalbuminuria in comparison to normoalbuminuric control subjects: the Oxford Regional Prospective Study. *Diabetes Care* 26:1456–1461, 2003
 10. Craig ME, Handelsman P, Donaghue KC, Chan A, Blades B, Laina R, Bradford D, Middlehurst A, Ambler G, Verge CF, Crock P, Moore P, Silink M: Predictors of glycaemic control and hypoglycaemia in children and adolescents with type 1 diabetes from NSW and the ACT. *Med J Aust* 177:235–238, 2002
 11. National Center for Health Statistics: CDC growth charts: United States [Internet], 2000. Available from <http://www.cdc.gov/nchs/about/major/nhanes/growthcharts/datafiles.htm>. Accessed 11 April 2006
 12. Kordonouri O, Hartmann R: Higher body weight is associated with earlier onset of type 1 diabetes in children: confirming the ‘accelerator hypothesis’ (Letter). *Diabetologia* 48:2501–2504, 2005
 13. Magarey AM, Daniels LA, Boulton JC: Prevalence of overweight and obesity in Australian children and adolescents: reassessment of 1985 and 1995 data against new standard international definitions. *Med J Aust* 174:561–564, 2001
 14. Knerr I, Wolf J, Reinehr T, Stachow R, Grabert M, Schober E, Rascher W, Holl RW, DPV Scientific Initiative of Germany and Austria: The ‘accelerator hypothesis’: relationship between weight, height, body mass index and age at diagnosis in a large cohort of 9,248 German and Austrian children with type 1 diabetes mellitus. *Diabetologia* 48:2501–2504, 2005