

# Is the Recent Rise in Type 2 Diabetes Incidence From 1984 to 2007 Explained by the Trend in Increasing BMI?

Evidence from a prospective study of British men

SARAH L. HARDOON, MSc<sup>1</sup>  
RICHARD W. MORRIS, PhD<sup>1</sup>  
MARY C. THOMAS, MSc<sup>1</sup>

S. GOYA WANNAMETHEE, PhD, FFPH<sup>1</sup>  
LUCY T. LENNON, MSc<sup>1</sup>  
PETER H. WHINCUP, PhD, FRCP<sup>2</sup>

**OBJECTIVE** — To estimate the extent to which increasing BMI may explain the rise in type 2 diabetes incidence in British men from 1984 to 2007.

**RESEARCH DESIGN AND METHODS** — A representative cohort ratio of 6,460 British men was followed-up for type 2 diabetes incidence between 1984 (aged 45–65 years) and 2007 (aged 67–89 years). BMI was ascertained at regular intervals before and during the follow-up.

**RESULTS** — Between 1984–1992 and 1999–2007, the age-adjusted hazard of type 2 diabetes more than doubled (hazard ratio 2.33 [95% CI 1.75–3.10]). Mean BMI rose by 1.42 kg/m<sup>2</sup> (95% CI 1.10–1.74) between 1984 and 1999; this could explain 26% (95% CI 17–38) of the type 2 diabetes increase.

**CONCLUSIONS** — An appreciable portion of the rise in type 2 diabetes can be attributed to BMI changes. A substantial portion remains unexplained, possibly associated with other determinants such as physical activity. This merits further research.

*Diabetes Care* 33:1494–1496, 2010

In the U.K., incidence of type 2 diabetes has increased by two-thirds in the last decade (1), and in the U.S. incidence has doubled over the last 30 years (2). Understanding the reasons for these unfavourable trends will help inform efforts to curb future type 2 diabetes increases. Marked rises in population adiposity have also occurred (3–5). However, few attempts have been made to quantify the contribution of changes in adiposity to the observed time trend in type 2 diabetes, partly reflecting the paucity of studies that have simultaneously monitored both type 2 diabetes and BMI levels in the same population over an extended period. We have estimated the proportion of the time-trend in type 2 diabetes incidence in British men over 24 years that may be ex-

plained by increasing population adiposity levels.

## RESEARCH DESIGN AND METHODS

Data came from the British Regional Heart Study (6), a socially and geographically representative cohort of middle-aged British men recruited between 1978 and 1980. The men were followed-up from baseline for all-cause mortality through National Health Service central registers and from 1984 for type 2 diabetes (the principal outcome) through a combination of biennial reviews of general practitioner (GP) records and self-report in questionnaires that were confirmed by GP records (7). Follow-up has been maintained for >99% of surviving men.

Type 2 diabetes incidence was compared in three consecutive follow-up periods, separated by intermittent questionnaires or examination sessions: period 1 from 1984 to 1992; period 2 from 1992 to 1999; and period 3 from 1999 to 2007 (Fig. 1). The principal exposure was BMI recorded in the questionnaire or examination at the start of each period (measurement techniques are detailed elsewhere [8]). BMI data were also available at recruitment in 1978–80. Analyses are based on 6,460 surviving men, aged 45–65 years at the start of the follow-up in 1984, with no prior diagnosis of diabetes. Men diagnosed within one year of the start of each period were excluded to limit the effect of reverse causality.

The proportion of the trend in type 2 diabetes hazard from period 1 to period 3 that may be statistically explained by increasing BMI was estimated by the expression  $(\beta_0 - \beta_1)/\beta_0$ , where  $\beta_0$  is the coefficient of an indicator for time period 3 in a Cox proportional hazard model for incident type 2 diabetes that only included time period, and  $\beta_1$  is the coefficient of the indicator for period 3 in a Cox proportional hazard model adjusting additionally for time-varying BMI (as a continuous variable with all significant powers) (9). A 95% CI for this estimate was obtained using bias-corrected bootstrap resampling (10). Age was used as the underlying time scale in the models, which were thereby automatically adjusted for age.

**RESULTS** — The hazard rate of type 2 diabetes increased more than twofold from period 1 (1984–1992) to period 3 (1999–2007) (hazard ratio [HR] 2.33 [95% CI 1.75–3.10], adjusting for age) (Fig. 1). An estimated 25.9% (95% CI 16.5–38.3) of this increase in type 2 diabetes hazard could be statistically explained by a population-averaged age-adjusted increase in BMI from 1984 to 1999 of 1.42 kg/m<sup>2</sup> (95% CI 1.10–1.74). A smaller percentage, 20.5% (95% CI 12.6–31.4), could be explained by “lagged” BMI changes, i.e., BMI ~5–8 years prior to each follow-up


From the <sup>1</sup>Division of Population Health, University College London, London, U.K.; and the <sup>2</sup>Division of Community Health Sciences, St George's, University of London, London, U.K.

Corresponding author: Sarah Hardoon, s.hardoon@ucl.ac.uk.

Received 16 December 2009 and accepted 29 March 2010. Published ahead of print at <http://care.diabetesjournals.org> on April 2010. DOI: 10.2337/dc09-2295.

© 2010 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.



**Figure 1**—Trend over time in incidence of type 2 diabetes and in mean BMI, by age-group. ○, age 45–49 years; ●, age 50–54 years; □, age 55–59 years; ■, aged 60–64 years; △, aged 65–69 years; ▲, aged 70–74 years.

period, corresponding to looking at the BMI increase over an earlier period from 1979 to 1992 of 1.13 kg/m<sup>2</sup> (95% CI 0.95–1.31). The combination of current and lagged measurements did not increase the contribution of BMI (percentage explained 25.8% [95% CI 16.8–38.1]).

The contribution of increasing BMI to the trend in type 2 diabetes incidence was examined separately in earlier and later portions of the follow-up. Between periods 1 (1984–1992) and 2 (1992–1999), incidence of type 2 diabetes increased by about a half (HR 1.59 [95% CI 1.23–

2.05]). Between periods 2 (1992–1999) and 3 (1999–2007), a similar increase was observed (HR 1.47 [1.17–1.84]). Of the increase in type 2 diabetes risk between periods 1 and 2, 21.7% (95% CI 11.9–47.9) could be statistically explained by an age-adjusted increase in BMI from 1984 to 1992 of 0.40 kg/m<sup>2</sup> (95% CI 0.29–0.52). During periods 2 and 3, 31.1% (95% CI 16.8–80.6) of the increase in type 2 diabetes could be statistically explained by an age-adjusted increase in BMI from 1992 to 1999 of 0.61 kg/m<sup>2</sup> (95% CI 0.44–0.78).

**CONCLUSIONS**— Increasing adiposity (assessed by BMI) appears to have accounted for about a quarter of the increase in type 2 diabetes incidence in periods 1 and 3 between 1984 and 2007 in this survivor cohort of British men. Control and reversal of the rise in adiposity levels is therefore an important priority in controlling the diabetes epidemic. A substantial proportion of the rise in type 2 diabetes remained unexplained. This may partly reflect study limitations; including the use of BMI as opposed to central adiposity measures and possible overestimation of the type 2 diabetes increase due to improved case ascertainment, changing diagnostic criteria, and the changing method of case identification (self-report in earlier years versus later use of GP records). However, trends in other determinants of type 2 diabetes, including declining physical activity levels (11) and dietary changes (12), could be playing an important role. Physical activity has a strong protective effect on type 2 diabetes risk, which operates at least partly independently of BMI (13). Dietary factors associated with reducing type 2 diabetes risk include a high fiber diet (14), and daily consumption of fiber declined between 1987 and 2000 (12).

Studies assessing population attributable risks of adiposity in type 2 diabetes have reported estimates similar to those in the present analysis (15). However in such studies, which examine overall type 2 diabetes incidence in the population, the contribution of modifiable risk factors, such as adiposity, are necessarily affected by contributions of static variables including genetic factors. In contrast our analysis examines time trends in type 2 diabetes, attributable only to modifiable factors that have changed over time in the cohort, thus carrying more immediate public health implications. Time-trend studies in other populations are needed to verify the findings and establish the roles of other risk factors. The presence of other contributing factors would suggest the need for a more multifactorial approach to combat rising type 2 diabetes in the population.

**Acknowledgments**— This work was supported by the British Heart Foundation (BHF Programme grant number RG/04/003). S.H. is supported by a Medical Research Council training fellowship grant (G0701739).

No potential conflicts of interest relevant to this article were reported.

References

1. Masso-González EL, Johansson S, Wal-lander MA, Garcia-Rodríguez LA. Trends in the prevalence and incidence of diabetes in the UK: 1996 to 2005. *J Epidemiol Community Health* 2009;63:332–336
2. Fox CS, Pencina MJ, Meigs JB, Vasan RS, Levitzky YS, D'Agostino RB Sr. Trends in the incidence of type 2 diabetes mellitus from the 1970s to the 1990s: the Framingham Heart Study. *Circulation* 2006;113:2914–2918
3. Silventoinen K, Sans S, Tolonen H, Monterde D, Kuulasmaa K, Kesteloot H, Tuomilehto J, WHO MONICA Project. Trends in obesity and energy supply in the WHO MONICA Project. *Int J Obes Relat Metab Disord* 2004;28:710–718
4. Flegal KM, Carroll MD, Ogden CL, Johnson CL. Prevalence and trends in obesity among US adults, 1999–2000. *JAMA* 2002;288:1723–1727
5. Health Survey for England 2004 [article online], 2005. Available from <http://www.ic.nhs.uk/statistics-and-data-collections/health-and-lifestyles-related-surveys/health-survey-for-england/health-survey-for-england-2004-updating-of-trend-tables-to-include-2004-data>. Accessed 9 June 2009
6. Walker M, Whincup PH, Shaper AG. The British Regional Heart Study 1975–2004. *Int J Epidemiol* 2004;33:1185–1192
7. Wannamethee SG, Shaper AG. Weight change and duration of overweight and obesity in the incidence of type 2 diabetes. *Diabetes Care* 1999;22:1266–1272
8. Hardoon SL, Whincup PH, Lennon LT, Wannamethee SG, Capewell S, Morris RW. How much of the recent decline in the incidence of myocardial infarction in British men can be explained by changes in cardiovascular risk factors? Evidence from a prospective population-based study. *Circulation* 2008;117:598–604
9. Freedman LS, Graubard BI, Schatzkin A. Statistical validation of intermediate endpoints for chronic diseases. *Stat Med* 1992;11:167–178
10. Efron B, Tibshirani RJ. *An Introduction to the Bootstrap*. London, Chapman and Hall, 1993
11. British Heart Foundation Statistics Website [article online], 2009. Available from <http://www.heartstats.org>. Accessed 20 June 2009
12. National Food Survey 2000 [article online], 2000. Available from <https://statistics.defra.gov.uk/esg/publications/nfs/2000>. Accessed 27 March 2009
13. Jeon CY, Lokken RP, Hu FB, van Dam RM. Physical activity of moderate intensity and risk of type 2 diabetes: a systematic review. *Diabetes Care* 2007;30:744–752
14. Hu FB, van Dam RM, Liu S. Diet and risk of Type II diabetes: the role of types of fat and carbohydrate. *Diabetologia* 2001;44:805–817
15. Wilson PW, D'Agostino RB, Sullivan L, Parise H, Kannel WB. Overweight and obesity as determinants of cardiovascular risk: the Framingham experience. *Arch Intern Med* 2002;162:1867–1872