

A Double-Blind, Randomized, Placebo-Controlled Clinical Trial on Benfotiamine Treatment in Patients With Diabetic Nephropathy

ALAA ALKHALAF, MD^{1,4}
 ASTRID KLOOSTER, BSC¹
 WILLEM VAN OEVEREN, PHD²
 ULRIKE ACHENBACH, PHD³
 NANNE KLEEFSTRA, MD^{4,5}
 ROBBERT J. SLINGERLAND, PHD⁶

G. SOPHIE MIJNHOUT, MD, PHD⁷
 HENK J.G. BILO, MD, PHD, FRCP^{1,4,7}
 REINOLD O.B. GANS, MD, PHD¹
 GERJAN J. NAVIS, MD, PHD¹
 STEPHAN J.L. BAKKER, MD, PHD¹

OBJECTIVE — To investigate the effect of benfotiamine on urinary albumin excretion (UAE) and the tubular damage marker kidney injury molecule-1 (KIM-1) in patients with type 2 diabetes and nephropathy.

RESEARCH DESIGN AND METHODS — Patients with type 2 diabetes and UAE equivalent to 15–300 mg/24 h, despite ACE inhibitors (ACE-Is) or angiotensin receptor blockers (ARBs), were randomly assigned to 12 weeks of benfotiamine (900 mg/day) ($n = 39$) or placebo ($n = 43$).

RESULTS — Compared with placebo, benfotiamine treatment resulted in significant improvement of thiamine status ($P < 0.001$). Benfotiamine treatment did not significantly decrease 24-h UAE or 24-h KIM-1 excretion.

CONCLUSIONS — In patients with type 2 diabetes and nephropathy, high-dose benfotiamine treatment for 12 weeks in addition to ACE-Is or ARBs did not reduce UAE or KIM-1 excretion, despite improvement of thiamine status.

Diabetes Care 33:1598–1601, 2010

The pathophysiology of diabetic nephropathy includes albuminuria as a consequence of glomerular endothelial damage and further progression due to tubulointerstitial inflammation and fibrosis (1,2). Despite protective treatment with ACE inhibitors (ACE-Is) and angiotensin receptor blockers (ARBs), many patients progress to end-stage renal disease (3).

Thiamine and benfotiamine have been proposed as protective agents for diabetes complications (4,5). Benfotiamine is a lipophilic thiamine derivative with

high bioavailability (6). In animal studies, both compounds had beneficial effects on microvascular complications, including diabetic nephropathy (5,7).

We investigated whether benfotiamine results in reduction in urinary albumin excretion (UAE) or tubulointerstitial damage markers in patients with type 2 diabetes and increased UAE despite ACE-Is or ARBs.

RESEARCH DESIGN AND METHODS

Participants were recruited at the Isala Clinics (Zwolle, the

Netherlands). Inclusion criteria were type 2 diabetes, age 40–75 years, active diabetic nephropathy (UAE 15–300 mg/24 h or equivalent albumin-to-creatinine ratio [males 1.25–25, females 1.75–35 mg/mmol] in two of three samples within 2–6 weeks) despite ACE-Is and/or ARBs in unchanged dose for at least 3 months, glycated hemoglobin (A1C) <8.5%, and an estimated glomerular filtration rate of >30 ml/min. Exclusion criteria were participation in another study, renal impairment by causes other than diabetes, elevated liver enzymes, hyper- or hypothyroidism, blood pressure >160/90 mmHg, neoplasm, severe general diseases, drug abuse, pregnancy, lactation, active menses, hypersensitivity to benfotiamine, use of vitamin B–containing supplements, changes in concomitant medication during the previous 3 months, and use of nonsteroidal anti-inflammatory drugs >3 times per week. In total, 2,711 patients were screened. Eligible patients were included after written informed consent was received. The trial was approved by the medical ethics committee.

Patients were randomized to oral benfotiamine 300 mg three times daily or placebo for 12 weeks. Study medication was prepared by Wörwag Pharma (Böblingen, Germany). Participants were evaluated at baseline and after 6 and 12 weeks. Each visit, 24-h urine, spot morning urine, and blood samples were collected. Noncompliance was considered if <80% of the study medication had been taken.

Thiamine concentration was measured in whole blood and plasma by high-performance liquid chromatography (8). Erythrocyte transketolase activity was measured in erythrocytes (9). Urinary albumin was measured by immunonephelometry (Behring Nephelometer; Mannheim, Germany), threshold 1.8–2.3 mg/l, intra- and interassay coefficients of variation (CV) 2.2 and 2.6%, respectively. Urinary kidney injury molecule-1 (KIM-1) was measured by ELISA, threshold 0.12 ng/ml, intra- and interassay CV 7.9 and 14.4%, respectively (10). Neuro-

From the ¹Department of Internal Medicine, University Medical Center Groningen, Groningen, the Netherlands; the ²Department of Biomedical Engineering, University Medical Center Groningen, Groningen, the Netherlands; ³Wörwag Pharma GmbH & Co. KG, Böblingen, Germany; the ⁴Diabetes Centre, Isala Clinics, Zwolle, the Netherlands; the ⁵Langerhans Medical Research Group, Zwolle, the Netherlands; the ⁶Department of Clinical Chemistry, Isala Clinics, Zwolle, the Netherlands; and the ⁷Department of Internal Medicine, Isala Clinics, Zwolle, the Netherlands.

Corresponding author: A. Alkhalaf, a.alkhalaf@int.umcg.nl.

Received 9 December 2009 and accepted 10 April 2010. Published ahead of print at <http://care.diabetesjournals.org> on 22 April 2010. DOI: 10.2337/dc09-2241. Clinical trial reg. no. NCT00565318; www.clinicaltrials.gov.

© 2010 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Table 1—Baseline characteristics and changes in thiamine status parameters, primary outcome measures, secondary outcome measures, and clinical characteristics over time

	Benfotiamine (n = 39)				Placebo (n = 43)				P
	Baseline	6 weeks	12 weeks	Baseline	6 weeks	12 weeks			
Males	30			33					
Age (years)	65.3 ± 5.9			64.6 ± 6.1				0.98	
BMI (kg/m ²)	32.1 ± 5.1			31.9 ± 5.9				0.63	
Duration of diabetes (years)	12 (9–18)			10 (7–18)				0.93	
Insulin treatment	31 (79)			29 (67)				0.41	
Oral hypoglycaemic agents	19 (49)			29 (67)				0.22	
Plasma thiamine (nmol/l)	31.8 ± 7.7			31.6 ± 9.8				0.05	
Thiamine status								0.92	
Whole blood thiamine (nmol/l)	126 ± 23	290 ± 31	300 ± 0	122 ± 23	124 ± 25	138 ± 30		<0.001	
TK activity (mU/mgHb)	0.41 ± 0.10	0.51 ± 0.12	0.53 ± 0.15	0.38 ± 0.11	0.39 ± 0.08	0.41 ± 0.10		<0.001	
Primary outcome parameters									
UAE (mg/24 h)	90 (38–267)	75 (49–280)	72 (38–199)	97 (48–177)	99 (43–200)	96 (45–200)		0.36	
U-KIM-1 (µg/24 h)	1.67 (0.95–2.47)	1.51 (0.86–2.59)	1.68 (1.06–2.40)	1.56 (1.06–1.83)	1.56 (1.06–1.83)	1.39 (1.02–2.01)		0.12	
Secondary outcome parameters									
24-h UACR (mg/mmol)	10.3 (3.7–23.4)	6.1 (3.0–17.7)	4.9 (2.5–18.4)	7.6 (4.3–13.3)	7.4 (2.8–11.0)	7.1 (4.0–12.5)		0.37	
Spot-urine UACR (mg/mmol)	9.3 (2.4–16.8)	5.8 (3.7–17.9)	7.1 (3.6–17.8)	6.2 (3.4–10.5)	8.2 (3.9–14.2)	8.1 (4.6–15.9)		0.58	
U-KIM-1/creatinine (ng/mmol)	103 (63–158)	95 (66–170)	96 (77–148)	99 (79–141)	89 (58–130)	81 (66–150)		0.37	
U-α1m (mg/24 h)	9.4 (4.3–24.4)	11.9 (4.4–20.2)	11.2 (4.1–18.8)	8.2 (4.3–20.3)	9.0 (5.7–21.1)	10.2 (2.5–19.7)		0.33	
U-α1m/creatinine (mg/mmol)	0.6 (0.3–1.4)	0.7 (0.3–1.3)	0.6 (0.3–1.2)	0.6 (0.3–1.3)	0.6 (0.3–1.4)	0.7 (0.2–1.1)		0.47	
U-Ngal (mg/24 h)	131 (67–227)	118 (77–229)	115 (73–284)	122 (53–224)	112 (52–218)	99 (52–222)		0.17	
U-Ngal/creatinine (mg/mmol)	6.7 (4.3–13.9)	6.2 (3.4–15.9)	5.1 (3.2–12.9)	7.7 (4.2–18.9)	6.4 (3.2–15.1)	8.5 (3.3–13.1)		0.18	
Clinical characteristics									
SBP (mmHg)	140 ± 16	139 ± 14	143 ± 17	137 ± 20	140 ± 20	140 ± 17		0.60	
DBP (mmHg)	76 ± 8	77 ± 10	76 ± 9	76 ± 10	76 ± 9	76 ± 10		0.68	
A1C (%)	7.3 ± 0.9	7.1 ± 0.9	7.3 ± 1.0	7.4 ± 0.9	7.2 ± 0.9	7.2 ± 0.9		0.33	
Plasma creatinine (µmol/l)	84 ± 19	89 ± 19	88 ± 20	87 ± 23	89 ± 25	87 ± 21		0.04	
Creatinine clearance (ml/min)	135 ± 51	129 ± 53	133 ± 45	130 ± 58	139 ± 58	131 ± 64		0.57	
Cystatin C (mg/l)	1.01 ± 0.21	1.06 ± 0.22	1.09 ± 0.23	1.03 ± 0.23	1.10 ± 0.26	1.11 ± 0.23		0.53	
LDL cholesterol (mmol/l)	1.9 ± 0.7	1.9 ± 0.8	2.1 ± 0.8	1.8 ± 0.9	1.8 ± 0.9	1.9 ± 0.9		0.55	
HDL cholesterol (mmol/l)	1.2 ± 0.3	1.1 ± 0.3	1.2 ± 0.3	1.1 ± 0.3	1.1 ± 0.3	1.1 ± 0.3		0.25	
Triglycerides (mmol/l)	1.8 (1.4–2.6)	1.9 (1.4–2.8)	1.7 (1.2–2.6)	2.1 (1.4–3.4)	2.2 (1.4–2.9)	2.0 (1.2–2.9)		0.06	

Data are means ± SD, n (%), or median (interquartile range). Comparison of baseline characteristics was performed by unpaired Student *t* test (for normally distributed variables) or Mann-Whitney *U* test (for non-normally distributed variables). χ^2 test was used to compare noncontinuous variables. Changes in thiamine status parameters, primary outcome measures, secondary outcome measures, and clinical characteristics over time were analyzed by ANOVA for repeated measures, with log-transformation of variables with skewed distribution prior to analysis. DBP, diastolic blood pressure; SBP, systolic blood pressure; TK, transketolase; U-α1m, urinary excretion of α1-microglobulin; U-KIM-1, urinary excretion of KIM-1; U-Ngal, urinary excretion of neutrophil gelatinase-associated lipocalin; UACR, urinary albumin-to-creatinine ratio.

phil gelatinase-associated lipocalin and α 1-microglobulin were measured by ELISA and cystatin C by immunoassay (Gentian; Moss, Norway). Other laboratory measurements were performed according to standard procedures.

Statistical analyses

Variables with normal distribution are presented as means \pm SD. Variables with skewed distribution were log-transformed before analysis and are presented as median (interquartile range). Changes were analyzed by ANOVA for repeated measurements. *P* values for change over time are presented. Results were considered statistically significant with *P* < 0.05.

To test whether benfotiamine reduces 24-h UAE or 24-h KIM-1 excretion, 38 evaluable patients per group were required to detect an effect of size 0.65 (power 80%, α = 0.05, one-sided test). To compensate for drop-out, we enrolled 43 patients per group. One-sided *P* values were calculated for primary outcome measures and two-sided *P* values for other outcomes. Statistics were done by SPSS, version 16.0 (Chicago, IL). Intention-to-treat analysis and per-protocol analyses were planned. In cases of drop-out, data were not replaced.

RESULTS — Baseline characteristics and results at 6 and 12 weeks are shown in Table 1. In the benfotiamine group, two patients did not complete the study because of newly diagnosed malignancy and two others withdrew informed consent (dizziness and urticaria). In patients receiving benfotiamine, parameters of thiamine status improved significantly. Benfotiamine treatment had no significant effect on primary or secondary outcome parameters. Change in UAE between baseline and 12 weeks was -18 mg/24 h in the benfotiamine group and -1 mg/24 h in the placebo group. For individual differences, respective changes were -9 (-53 to 34) mg/24 h and -7 mg/24 h (-56 to 65). With respect to clinical characteristics, benfotiamine resulted in a borderline significant increase in plasma creatinine, but this was not accompanied by changes in creatinine clearance or cystatin C.

During the study, two patients were noncompliant (one per group) and two protocol violations occurred in the placebo group (ACE-I stopped), resulting in 38 patients in the benfotiamine and 40 in the placebo group for per-protocol anal-

yses, of which results (data not shown) were not materially different from presented analyses.

CONCLUSIONS — We found that 12-week treatment with high-dose benfotiamine did not result in decreases in 24-h-UAE or 24-h-KIM-1 excretion despite significant improvement of thiamine status.

Our findings differ from those of a pilot study of 40 patients with type 2 diabetes in which 12 weeks of 300 mg/day of thiamine resulted in a significant decrease in UAE by 17.7 mg/24 h (11). In that study, baseline UAE was 44 mg/24 h (33–121) in the thiamine and 51 mg/24 h (32–122) in the placebo group, which is approximately 2 times lower than in our study, despite 100% of ACE-I and ARB treatment in our study versus <50% in the pilot study (12). Thus, thiamine derivatives may provide protective effects in earlier diabetic nephropathy stages, which is in line with an animal study in which development of albuminuria after induction of diabetes was inhibited by thiamine and benfotiamine (5). Furthermore, we investigated Caucasian patients, and the other study was of Pakistani patients. Thus, differences in diet, baseline prevalence of thiamine deficiency, or genetic susceptibility may also play a role. In our study, baseline plasma thiamine concentrations of 31.8 ± 7.7 nmol/l in the benfotiamine group and 31.6 ± 9.8 nmol/l in the placebo group were higher than the 16.3 ± 11.5 nmol/l reported for patients with type 2 diabetes in the U.K., but deficient compared to the 64.1 ± 12.0 nmol/l reported for healthy control subjects in that study (13).

It is important to realize that thiamine and benfotiamine are supposed to antagonize detrimental effects of hyperglycemia (7). Yet, in two large intervention studies, it took years of lowering A1C before a difference in UAE was found between strict metabolic control and standard therapy (14,15). Our study may therefore have been too short to demonstrate the effect of benfotiamine.

In conclusion, longer-term intervention studies and/or intervention studies in earlier stages of diabetic nephropathy are necessary to discern whether benfotiamine has an effect on the development of diabetic nephropathy.

Acknowledgments — This trial was partly funded by a grant from the European Union to

PREDICTIONS (Prevention of Diabetes Complication) network and by the medical research foundation in Zwolle.

U.A. is the head of medical affairs in Wörwag Pharma Co., and contributed in the preparation of the study protocol and other necessary documents needed for approval of the study by the ethics committee and authorities but was not directly or indirectly involved in the practical procedures, inclusion and evaluation of subjects, data collection, or data analysis.

No potential conflicts of interest relevant to this article were reported.

A.A. researched data and wrote the manuscript. A.K., W.V.O., and R.J.S. researched data. U.A. and G.S.M. contributed to the discussion. N.K., H.J.G.B., R.O.B.G., G.J.N., and S.J.L.B. contributed to the discussion and reviewed/edited the manuscript.

The authors acknowledge the efforts of Drs. J. Lambert and J.E. Heeg (Isala clinics, Zwolle, the Netherlands) and Dr. J. Jager (Diaconessenhuis Hospital, Meppel, the Netherlands) in preparing the study. The authors thank Dr. L.D. Dikkeschei, M. van der Saag, and M. Slingschröder (clinical chemical laboratories in the Isala Clinics, Zwolle, the Netherlands) and H. Breukelman (clinical chemical laboratories at the University Medical Center, Groningen, the Netherlands) for their collaboration. The authors also acknowledge the efforts of Dr. W. Gaus (University of Ulm, Germany) as independent statistician.

References

1. Fioretto P, Bruseghin M, Berto I, Gallina P, Manzato E, Mussap M. Renal protection in diabetes: role of glycemic control. *J Am Soc Nephrol* 2006;17:S86–S89
2. Nath KA. Tubulointerstitial changes as a major determinant in the progression of renal damage. *Am J Kidney Dis* 1992;20:1–17
3. Lewis EJ, Hunsicker LG, Clarke WR, Berl T, Pohl MA, Lewis JB, Ritz E, Atkins RC, Rohde R, Raz I, Collaborative Study Group. Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. *N Engl J Med* 2001;345:851–860
4. Bakker SJ, Heine RJ, Gans RO. Thiamine may indirectly act as an antioxidant. *Diabetologia* 1997;40:741–742
5. Babaei-Jadidi R, Karachalias N, Ahmed N, Battah S, Thornalley PJ. Prevention of incipient diabetic nephropathy by high-dose thiamine and benfotiamine. *Diabetes* 2003;52:2110–2120
6. Schreeb KH, Freudenthaler S, Vormfelde SV, Gundert-Remy U, Gleiter CH. Comparative bioavailability of two vitamin B1 preparations: benfotiamine and thiamine mononitrate. *Eur J Clin Pharmacol* 1997;52:319–320
7. Hammes HP, Du X, Edelstein D, Taguchi

- T, Matsumura T, Ju Q, Lin J, Bierhaus A, Nawroth P, Hannak D, Neumaier M, Bergfeld R, Giardino I, Brownlee M. Benfotiamine blocks three major pathways of hyperglycemic damage and prevents experimental diabetic retinopathy. *Nat Med* 2003;9:294–299
8. Schrijver J, Speek AJ, Klosse JA, van Rijn HJ, Schreurs WH. A reliable semiautomated method for the determination of total thiamine in whole blood by the thiochrome method with high-performance liquid chromatography. *Ann Clin Biochem* 1982;19:52–56
 9. Milner CR, Buttery JE, Chamberlain BR. The measurement of erythrocyte transketolase activity on a discrete analyser. *J Automat Chem* 1982;4:183–185
 10. van Timmeren MM, Vaidya VS, van Ree RM, Oterdoom LH, de Vries AP, Gans RO, van Goor H, Stegeman CA, Bonventre JV, Bakker SJ. High urinary excretion of kidney injury molecule-1 is an independent predictor of graft loss in renal transplant recipients. *Transplantation* 2007;84:1625–1630
 11. Rabbani N, Alam SS, Riaz S, Larkin JR, Akhtar MW, Shafi T, Thornalley PJ. High-dose thiamine therapy for patients with type 2 diabetes and microalbuminuria: a randomised, double-blind placebo-controlled pilot study. *Diabetologia* 2009;52:208–212
 12. Rabbani N, Alam SS, Riaz S, Larkin JR, Akhtar MW, Shafi T, Thornalley PJ. Thiamine in diabetic nephropathy: a novel treatment modality? Reply to Alkhalaf A, Kleefstra N, Groenier KH et al. (Letter). *Diabetologia* 2009;52:1214–1216
 13. Thornalley PJ, Babaei-Jadidi R, Al Ali H, Rabbani N, Antonysunil A, Larkin J, Ahmed A, Rayman G, Bodmer CW. High prevalence of low plasma thiamine concentration in diabetes linked to a marker of vascular disease. *Diabetologia* 2007;50:2164–2170
 14. Effect of intensive therapy on the development and progression of diabetic nephropathy in the Diabetes Control and Complications Trial. The Diabetes Control and Complications (DCCT) Research Group. *Kidney Int* 1995;47:1703–1720
 15. Bilous R. Microvascular disease: what does the UKPDS tell us about diabetic nephropathy? *Diabet Med* 2008;25(Suppl. 2):25–29