

COMMENTS AND RESPONSES

Pro12Ala Polymorphism in the PPARG Gene Contributes to the Development of Diabetic Nephropathy in Chinese Type 2 Diabetic Patients

Response to Lapice et al.

Lapice et al. (1) confirmed the findings in Chinese (2) or Caucasian patients (3) on the association between the peroxisome proliferator-activated receptor (PPAR)- γ 2 Pro12Ala and urinary albumin excretion rate (AER) in a large population with type 2 diabetic patients ($n = 750$) of Caucasian origin. Moreover, they observed that the Ala12 carrier showed significantly lower plasma urea and creatinine (Scr) and higher estimated glomerular filtration rate (eGFR) than the Pro12 carrier ($P < 0.05$ for each) in the study.

In our study, there were significantly elevated Scr and blood urea nitrogen [i.e., 84.0 (64.0–129.0) vs. 64.0 (55.0–78.0) $\mu\text{mol/l}$ (2) and 6.3 (5.1–9.1) vs. 5.6 (4.7–6.9) mmol/l, $P = 0.000$ for each], declined eGFR (77.6 ± 45.0 vs. 100.6 ± 29.7 ml/min/1.73 m^2 , $P = 0.000$), and Pro12Ala carriers (6.2% vs. 12.8%, $P = 0.003$) in diabetic nephropathy patients when compared with nondiabetic nephropathy patients ($P < 0.01$ for each) (2). In addition, except for the Pro12Ala patients who had lower AER levels than the Pro12Pro patients [9.22 (6.2–15.6)

vs. 173.0 (69.9–691.1) mg/24 h, $P = 0.000$] (2), Pro12Ala patients had lower Scr and blood urea nitrogen trends [i.e., 68.0 (57.0–104.0) vs. 76.0 (60.0–105.8) $\mu\text{mol/l}$ (2) and 6.0 (4.9–8.1) vs. 6.2 (4.8–8.7) mmol/l] and higher eGFR trend (87.2 ± 43.0 vs. 84.4 ± 42.3 ml/min/1.73 m^2), but the differences were not statistically significant ($P > 0.05$ for each). Similarly, the occurrence of subjects with mildly impaired renal function (i.e., GFR < 60 ml/min/1.73 m^2) has a lower tendency in patients with the Pro12Ala than that of the Pro12Pro genotype (24.1% vs. 28.0%, $P = 0.533$; odds ratio 0.82 [95% CI 0.44–1.53]); however, no statistical significance was observed.

Previous studies reported that the Ala12 allele was resistant to type 2 diabetes in Caucasians but not in Chinese, thus reflecting an ethnic genetic heterogeneity of type 2 diabetes at this locus (4). No difference in Scr between Ala12 carriers and noncarriers was reported in Caucasian type 2 diabetic patients (3). Jorsal et al. (5) reported that the Ala12 allele is associated with enhanced decline in GFR ($P = 0.04$) in Danish Caucasian type 1 diabetic patients.

In conclusion, except for AER, the Pro12Ala polymorphism of PPAR- γ 2 is not significantly associated with the overall better renal function in Chinese type 2 diabetic patients with and without diabetic nephropathy.

LIMEI LIU, MD, PHD¹
TAISHAN ZHENG, MD¹
FENG WANG, MD²
NIANSONG WANG, MD, PHD²
MING LI, MD¹

From the ¹Department of Endocrinology and Metabolism, Shanghai Diabetes Institute, Shanghai Jiaotong University Affiliated Sixth People's Hospital, Shanghai, China; and the ²Department of Nephrology, Shanghai Jiaotong University Affiliated Sixth People's Hospital, Shanghai, China.
Corresponding author: Limei Liu, lmliu@sjtu.edu.cn.

DOI: 10.2337/dc10-0825

© 2010 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

Acknowledgments—No potential conflicts of interest relevant to this article were reported.

L.L. researched the data, wrote the letter, and reviewed and edited the letter; T.Z. researched the data; F.W. reviewed the letter; N.W. reviewed the letter; and M.L. researched the data.

References

- Lapice E, Pinelli M, Riccardi G, Vaccaro O. Pro12Ala polymorphism in the PPARG gene contributes to the development of diabetic nephropathy in Chinese type 2 diabetic patients: comment on the study by Liu et al. (Letter). *Diabetes Care* 2010; 33:e114. DOI: 10.2337/dc10-0596
- Liu L, Zheng T, Wang F, Wang N, Song Y, Li M, Li L, Jiang J, Zhao W. Pro12Ala polymorphism in the PPARG gene contributes to the development of diabetic nephropathy in Chinese type 2 diabetic patients. *Diabetes Care* 2010;33:144–149
- Caramori ML, Canani LH, Costa LA, Gross JL. The human peroxisome proliferator-activated receptor γ 2 (PPAR γ 2) Pro12Ala polymorphism is associated with decreased risk of diabetic nephropathy in patients with type 2 diabetes. *Diabetes* 2003;52:3010–3013
- Radha V, Vimalaswaran KS, Babu HN, Abate N, Chandalia M, Satija P, Grundy SM, Ghosh S, Majumder PP, Deepa R, Rao SM, Mohan V. Role of genetic polymorphism peroxisome proliferator-activated receptor- γ 2 Pro12Ala on ethnic susceptibility to diabetes in South-Asian and Caucasian subjects: evidence for heterogeneity. *Diabetes Care* 2006;29:1046–1051
- Jorsal A, Tarnow L, Lajer M, Ek J, Hansen T, Pedersen O, Parving HH. The PPAR gamma 2 Pro12Ala variant predicts ESRD and mortality in patients with type 1 diabetes and diabetic nephropathy. *Mol Genet Metab* 2008;94:347–351