

Reduction of Visceral Fat Is Associated With Decrease in the Number of Metabolic Risk Factors in Japanese Men

YUKIYOSHI OKAUCHI, MD¹
HITOSHI NISHIZAWA, MD¹
TOHRU FUNAHASHI, MD¹
TOMOKO OGAWA, RN²
MIDORI NOGUCHI, RN²
MIWA RYO, MD¹

SHINJI KIHARA, MD¹
HIROMI IWAHASHI, MD¹
KAZUYA YAMAGATA, MD¹
TADASHI NAKAMURA, MD¹
IICHIRO SHIMOMURA, MD¹
YUJI MATSUZAWA, MD³

Visceral fat accumulation is associated with the development of metabolic disorders such as glucose intolerance, dyslipidemia, hypertension, and atherosclerotic cardiovascular diseases (1–8). However, the relationship between reduction of visceral fat and decrease in the number of metabolic risk factors has not been defined in the general population. Recently, we developed a new technique, the abdominal bioelectrical impedance analysis (BIA), to evaluate visceral fat area (VFA) (9). The aim of this study was to investigate whether reduction of visceral fat, estimated by the BIA, is associated with a decrease in the number of metabolic risk factors.

RESEARCH DESIGN AND METHODS

The study group comprised 2,336 Japanese men (aged mean \pm SD 48.0 ± 10.5 years, BMI 24.2 ± 2.9 kg/m²), who were employees of Amagasaki City Office, an urban area, and had undergone annual health check-ups in both 2004 and 2005. After the health check-up, the medical staff provided risk factor-oriented, rather than obesity-oriented, health promotion programs to select individuals with visceral fat accumulation and multiple risk factors, with the aim of encouraging a scientific understanding of the spectrum of metabolic syndrome from visceral fat accumulation

to atherosclerotic cardiovascular diseases. In this study, we used VFA estimated by the BIA, which was shown to correlate significantly with VFA determined by computed tomography (9). The measurement of VFA by BIA complied with the Guidelines of the Ethical Committees of Osaka University. Informed consent was obtained from all subjects.

Overall obesity was defined as BMI of ≥ 25 kg/m² (10). We investigated the presence of three metabolic risk factors: elevated blood pressure (systolic blood pressure ≥ 130 mmHg and/or diastolic blood pressure ≥ 85 mmHg), dyslipidemia, and dysglycemia/impaired glucose tolerance. Dyslipidemia represented hypertriglyceridemia (fasting or postprandial triglyceride of ≥ 1.69 or 2.27 mmol/l [11,12], respectively, and/or low HDL cholesterol [HDL cholesterol < 1.04 mmol/l]). Dysglycemia/impaired glucose tolerance represented hyperglycemia (fasting or postprandial serum glucose concentration of ≥ 6.1 or ≥ 7.77 mmol/l [13], respectively). Subjects who received specific treatment(s) for each of the metabolic risk factors were considered positive for that factor.

Statistical analysis

Fischer's protected least significant difference test and Kruskal-Wallis were used to analyze the relationship between the

number of metabolic risk factors and body fat distribution and between change in the number of metabolic risk factors and change in VFA, respectively. Significance level was set at $P < 0.05$.

RESULTS— BMI and VFA varied considerably among individuals. We divided subjects into two groups according to BMI and into two groups according to VFA (Fig. 1A). Visceral fat accumulation was defined as VFA of ≥ 100 cm² (10,14). Among 1,497 nonobese subjects (BMI < 25 kg/m²), 401 (26.8%) had visceral fat accumulation. The mean number of metabolic risk factors in subjects with VFA ≥ 100 cm² was significantly higher than in those with VFA < 100 cm², irrespective of BMI. Importantly, the mean number of metabolic risks was significantly higher in subjects with VFA ≥ 100 cm² plus BMI < 25 kg/m² than in those with VFA < 100 cm² plus BMI ≥ 25 kg/m² ($P < 0.0001$) (Fig. 1A). These results suggest that assessment of visceral fat accumulation is important in identifying subjects with multiple risk factors.

Next, we investigated the correlation between a 1-year change in VFA (Δ VFA) and change in the number of metabolic risk factors (Δn) within the same period in the 2,336 subjects. VFA decreased within 1 year in 53.1% (1,241 of 2,336) of participants, increased in 33.2% (776 of 2,336), and did not change in 13.7% (319 of 2,336).

We divided these subjects into six bins of Δ VFA (every 15 cm²). Δ VFA correlated significantly with Δn ($P < 0.0001$) (Fig. 1B). When the subjects who received new treatment after 2004 were excluded from the analysis, reduction of visceral fat was also associated with a significant decrease in the number of metabolic risk factors ($P < 0.0001$) (data not shown).

CONCLUSIONS— We demonstrated that 1) irrespective of BMI (< 25 kg/m²), subjects with visceral fat accumulation estimated by BIA had a cluster of metabolic risk factors and 2) falls in VFA within 1 year were associated with a significant decrease in the number of metabolic risk factors.

From the ¹Department of Metabolic Medicine, Graduate School of Medicine, Osaka University, Suita, Japan; the ²Amagasaki City Office, Hyogo, Japan; and the ³Sumitomo Hospital, Osaka, Japan.

Address correspondence and reprint requests to Hitoshi Nishizawa, Department of Metabolic Medicine, Graduate School of Medicine, Osaka University, 2-2 Yamadaoka, Suita, Osaka 565-0871, Japan. E-mail: hitoshin@imed2.med.osaka-u.ac.jp.

Received for publication 2 February 2007 and accepted in revised form 30 May 2007.

Published ahead of print at <http://care.diabetesjournals.org> on 11 June 2007. DOI: 10.2337/dc07-0218.

Abbreviations: BIA, bioelectrical impedance analysis; VFA, visceral fat area.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

© 2007 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.




Figure 1—A: Relationship between number of metabolic risk factors and body fat distribution. Subjects were divided according to their BMI (cutoff value 25 kg/m²) and VFA (cutoff value 100 cm²), measured in 2004. Data are means \pm SE. B: Correlation between changes in VFA and changes in the number of metabolic risk factors. Δ number of metabolic risk factors represents changes in the number of metabolic risk factors from 2004 to 2005. Δ VFA indicates change in VFA from 2004 to 2005. Subjects were divided into six 15-cm² bins of Δ VFA. Data are means \pm SE.

Importantly, our results also demonstrated that subjects with visceral fat accumulation but without overall obesity

(VFA \geq 100 cm² plus BMI <25 kg/m²) exhibited significantly more metabolic risk factors than overall obese subjects

without visceral fat accumulation (VFA <100 cm² plus BMI \geq 25 kg/m²). There is ample evidence for the role of visceral fat accumulation in the development of metabolic disorders (4–8,15). Collectively, the above results indicate that assessment of visceral fat accumulation using VFA estimated by BIA is useful for identifying high-risk groups for atherosclerotic cardiovascular diseases.

Our results also demonstrated in a large population sample that changes in VFA within 1 year correlated significantly with Δn . Several reports demonstrated in obese subjects that reduction of visceral fat correlated with improvement in glucose and lipid metabolism (16–19). However, there is little information on the effect of reduction of visceral fat on the number of metabolic risk factors in a large general population sample. Here, we showed in 2,336 subjects that changes in VFA within 1 year correlated significantly with changes in the number of metabolic risk factors. These results suggest that intervention strategies directed toward reduction of visceral fat could result in the reduction or disappearance of risks for atherosclerotic cardiovascular diseases. Since BIA is quite simple and noninvasive for evaluation of visceral fat amount, it could be used in routine clinical practice and large-scale studies for assessment of visceral fat accumulation.

In conclusion, we demonstrated that reduction of visceral fat was closely associated with a decrease in the number of metabolic risk factors in Japanese men.

References

- Eckel RH, Grundy SM, Zimmet PZ: Metabolic syndrome. *Lancet* 365:1415–1428, 2005
- Isomaa B, Almgren P, Tuomi T, Forsen B, Lahti K, Nissen M, Taskinen M, Groop L: Cardiovascular morbidity and mortality associated with the metabolic syndrome. *Diabetes Care* 24:683–689, 2001
- Alberti KG, Zimmet P, Shaw J: IDF Epidemiology Task Force Consensus Group: The metabolic syndrome: a new worldwide definition. *Lancet* 366:1059–1062, 2005
- Despres JP, Lemieux I: Abdominal obesity and metabolic syndrome. *Nature* 444:881–887, 2006
- Kissebah AH, Vydelingum N, Murray R, Evans DJ, Hartz AJ, Kalkhoff RK, Adams PW: Relation of body fat distribution to metabolic complications of obesity. *J Clin Endocrinol Metab* 54:254–260, 1982
- Lean MEJ, Han TS, Seidell JC: Impairment of health and quality of life in people with

- large waist circumference. *Lancet* 351: 853–856, 1998
7. Nakamura T, Tokunaga K, Shimomura I, Nishida M, Yoshida S, Kotani K, Islam AHMW, Keno Y, Kobatake T, Nagai Y, Fujioka S, Tarui S, Matsuzawa Y: Contribution of visceral fat accumulation to the development of coronary artery disease in non-obese men. *Atherosclerosis* 107:239–246, 1994
 8. Lemieux I, Pascot A, Couillard C, Lamarche B, Tchernof A, Almeras N, Berqeron J, Gaudet D, Tremblay G, Prud'homme D, Nadeau A, Despres JP: Hypertriglyceridemic waist: a marker of the atherogenic metabolic triad (hyperinsulinemia; hyperapolipoprotein B; small, dense LDL) in men. *Circulation* 102:179–184, 2000
 9. Ryo M, Maeda K, Onda T, Katashima M, Okumiya A, Nishida M, Yamaguchi T, Funahashi T, Matsuzawa Y, Nakamura T, Shimomura I: A new simple method for the measurement of visceral fat accumulation by bioelectrical impedance. *Diabetes Care* 28:451–453, 2005
 10. Examination Committee of Criteria for 'Obesity Disease' in Japan; Japan Society for the Study of Obesity: New criteria for 'obesity disease' in Japan. *Circ J* 66:987–992, 2002
 11. Eberly LE, Stamler J, Neaton JD, Multiple Risk Factor Intervention Trial Research Group: Relation of triglyceride levels, fasting and nonfasting, to fatal and nonfatal coronary heart disease. *Arch Intern Med* 163:1077–1083, 2003
 12. Ahmad J, Hameed B, Das G, Siddiqui M, Ahmad I: Postprandial hypertriglyceridemia and carotid intima-media thickness in north Indian type 2 diabetic subjects. *Diabetes Res Clin Pract* 69:142–150, 2005
 13. American Diabetes Association: Clinical practice recommendations 2000: screening for type 2 diabetes. *Diabetes Care* 23 (Suppl. 1):S20–S23, 2000
 14. Despres JP, Lamarche B: Effects of diet and physical activity on adiposity and body fat distribution: implications for the prevention of cardiovascular disease. *Nutr Res Rev* 6:137–159, 1993
 15. Janssen I, Katzmarzyk PT, Ross R: Waist circumference and not body mass index explains obesity-related health risk. *Am J Clin Nutr* 79:379–384, 2004
 16. Fujioka S, Matsuzawa Y, Tokunaga K, Kawamoto T, Kobatake T, Keno Y, Kotani K, Yoshida S, Tarui S: Improvement of glucose and lipid metabolism associated with selective reduction of intra-abdominal visceral fat in premenopausal women with visceral fat obesity. *Int J Obes* 15: 853–859, 1991
 17. Ross R, Dagnone D, Jones PJ, Smith H, Paddags A, Hudson R, Janssen I: Reduction in obesity and related comorbid conditions after diet-induced weight loss or exercise-induced weight loss in men: a randomized, controlled trial. *Ann Intern Med* 133:92–103, 2000
 18. Miyatake N, Nishikawa H, Morishita A, Kunitomi M, Wada J, Suzuki H, Takahashi K, Makino H, Kira S, Fujii M: Daily walking reduces visceral adipose tissue areas and improves insulin resistance in Japanese obese subjects. *Diabetes Res Clin Pract* 58:101–107, 2002
 19. Slentz CA, Aiken LB, Houmard JA, Bales CW, Johnson JL, Tanner CJ, Duscha BD, Kraus WE: Inactivity, exercise, and visceral fat: STRRIDE: a randomized, controlled study of exercise intensity and amount. *J Appl Physiol* 99:1613–1618, 2005