

# Possible Role of $\alpha$ -Cell Insulin Resistance in Exaggerated Glucagon Responses to Arginine in Type 2 Diabetes

NAOMI TSUCHIYAMA, MD  
TOSHINARI TAKAMURA, MD, PHD  
HITOSHI ANDO, MD, PHD  
MASARU SAKURAI, MD

AKIKO SHIMIZU, MD  
KEN-ICHIRO KATO, MD  
SEIICHIRO KURITA, MD, PHD  
SHUICHI KANEKO, MD, PHD

**OBJECTIVE** — Inappropriate excessive secretion of glucagon, which contributes to postprandial hyperglycemia, is a novel target for the treatment of diabetes. In this study, we sought to determine the factors associated with exaggerated glucagon secretion in response to an arginine challenge in patients with type 1 and type 2 diabetes.

**RESEARCH DESIGN AND METHODS** — Changes in circulating C-peptide immunoreactivity (CPR) and immunoreactive glucagon (IRG) after an arginine challenge were investigated in 35 patients with type 1 diabetes, 130 patients with type 2 diabetes, and 35 nondiabetic control subjects.

**RESULTS** — No significant differences were found in the basal level and the area under the concentration-time curve (AUC) of IRG ( $AUC_{IRG}$ ) among type 1 and type 2 diabetic patients and nondiabetic subjects. However, there was an inverse correlation between the  $AUC_{IRG}$  and the AUC of CPR ( $AUC_{CPR}$ ) for type 1 ( $r = -0.388$ ,  $P = 0.023$ ) and type 2 ( $r = 0.396$ ,  $P < 0.0001$ ) diabetic patients, whereas  $AUC_{IRG}$  was not correlated with  $AUC_{CPR}$  in nondiabetic subjects ( $r = -0.079$ ,  $P = 0.655$ ). In type 1 diabetic patients, the  $AUC_{CPR}$  decreased and the  $AUC_{IRG}$  increased with increasing disease duration. In type 2 diabetic patients, both  $AUC_{IRG}$  and  $AUC_{CPR}$  increased with increasing BMI, basal CPR level, and homeostasis model assessment of insulin resistance value.

**CONCLUSIONS** — Our findings suggest that the pathophysiology of the exaggerated glucagon response differs between type 1 and type 2 diabetes. Intra-islet insulin deficiency and  $\alpha$ -cell insulin resistance may be the primary contributors to this condition in type 1 and type 2 diabetes, respectively.

*Diabetes Care* 30:2583–2587, 2007

Diabetes is associated with increased hepatic glucose production, which is linked to fasting and postprandial hyperglycemia (1). This is caused by the reduced suppression of glucagon (2), along with the impairment of insulin secretion and insulin action. Arginine-stimulated

hyperglucagonemia in patients with various forms of diabetes was first discovered by Aronoff et al. (3,4). Their results demonstrated that excess glucagon or an elevated ratio of glucagon to insulin is etiologically important in the development of endogenous hyperglycemia in di-

abetes through the mediation of glucose overproduction from the liver (2,5–7). Thus, glucagon and its receptor have been examined extensively in recent years as being potential targets for the treatment of diabetes (8,9). In fact, glucagon-like peptide 1 analogs, which are forthcoming antidiabetes agents, lower postprandial hyperglycemia partly by inhibiting excessive secretion of glucagon in not only type 2 but also type 1 diabetes (10). An absolute deficiency in insulin secretion has been suggested to cause an exaggerated response of glucagon to arginine in patients with diabetes (11,12). Insulin replacement therapy can correct this deficiency in patients with type 1 diabetes (11,12) but not in those with type 2 diabetes (11). The pathophysiology associated with an exaggerated glucagon response to arginine remains unclear, particularly in patients with type 2 diabetes.

Here, we comprehensively analyze factors associated with the exaggerated glucagon secretion response to an arginine challenge in patients with diabetes and show that the glucagon response to arginine reflects a distinctly different pathophysiology in type 1 versus type 2 diabetes.

## RESEARCH DESIGN AND METHODS

A total of 223 Japanese patients with diabetes who were hospitalized in the Division of Endocrinology and Metabolism at Kanazawa University Hospital (Ishikawa, Japan) were studied for the management of diabetes and for patient education. The study was conducted from April 2003 to May 2006. Patients were diagnosed according to the criteria established by an expert committee on the diagnosis and classification of diabetes (13). Type 1 diabetes was determined based on the presence of islet autoantibodies (anti-GAD antibody). In addition, all type 2 diabetic patients met the requirements for insulin therapy, as defined by the American Diabetes Association. Among the 36 patients with type 1 diabetes, we excluded one patient with Wilson's disease (supplemental Fig. 1 [available in an online appendix at <http://dx.doi.org/10.2337/dc07-0066>]). Among the 187 patients with type 2 dia-

From the Department of Disease Control and Homeostasis, Kanazawa University Graduate School of Medical Science, Kanazawa, Japan.

Address correspondence and reprint requests to Toshinari Takamura, Department of Disease Control and Homeostasis, Kanazawa University Graduate School of Medical Science, 13-1 Takara-machi, Kanazawa, Ishikawa 920-8641, Japan. E-mail: [ttakamura@m-kanazawa.jp](mailto:ttakamura@m-kanazawa.jp)

Received for publication 11 January 2007 and accepted in revised form 13 July 2007.

Published ahead of print at <http://care.diabetesjournals.org> on 31 July 2007. DOI: 10.2337/dc07-0066. Additional information for this article can be found in an online appendix at <http://dx.doi.org/10.2337/dc07-0066>.

**Abbreviations:** AUC, area under the concentration-time curve; CPR, C-peptide immunoreactivity; HOMA-IR, homeostasis model assessment of insulin resistance; IRG, immunoreactive glucagon; QUICKI, quantitative insulin sensitivity index.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

© 2007 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Table 1—Characteristics of type 1 and type 2 diabetic patients and nondiabetic subjects

	Type 1 diabetic patients	Type 2 diabetic patients	Nondiabetic subjects
n	35	130	35
Acute/slowly progressive type 1 diabetes	26/9		
Sex (men/women)	14/21	81/49	15/20
Age (years)	49 ± 15	57 ± 13*	33 ± 6*†
BMI (kg/m <sup>2</sup> )	22.4 ± 4.3	23.8 ± 3.6	22.7 ± 4.0
Diabetes duration (months)	99.4 ± 92.5	123.9 ± 142.8	
A1C (%)	9.7 ± 2.8	8.9 ± 2.6	5.0 ± 0.4*†
Urinary C-peptide (mg/day)	9.9 ± 13.0	43.4 ± 42.1*	
Therapy			
Insulin	35	78	
Oral antidiabetes agent	0	34	
Diet alone	0	20	

Data are n or means ± SD. \*P < 0.01 vs. type 1 diabetic subjects. †P < 0.01 vs. type 2 diabetic subjects.

betes, we excluded 57 patients with secondary diabetes related to conditions including liver disease, pancreatic disease, and hormonal disorders (supplemental Fig. 1). Thus, 35 patients with type 1 diabetes and 130 with type 2 diabetes were enrolled in this study. Additionally, 35 nondiabetic healthy volunteers were enrolled as control subjects. The characteristics of the study subjects are shown in Table 1. Of those with type 1 diabetes, 26 patients were diagnosed with an acute-onset form and 9 with slowly progressive type 1 diabetes (14). Informed consent was obtained from all patients before study initiation, and the study was approved by the relevant ethics committee and conducted in accordance with the Declaration of Helsinki.

On admission, the patients received a standard diet of 30 kcal/kg. All patients with type 1 diabetes were treated with intensive insulin therapy. Among the patients with type 2 diabetes, 78 were treated with insulin (57 with premeal dosing of a rapid-acting insulin analog alone), 32 were treated with oral antidiabetes agents (14 with  $\alpha$ -glucosidase inhibitors, 16 with nateglinide, 4 with metformin, and 5 with sulfonylureas), and 20 received diet therapy alone.

### Arginine stimulation test

After an overnight fast, patients were kept at rest for  $\geq 30$  min, and endogenous insulin and glucagon levels, as measured by C-peptide immunoreactivity (CPR) in serum and immunoreactive glucagon (IRG) in plasma, respectively, were assessed at preloading baseline (0 min). Arginine (30 g) was administered by intravenous infu-

sion of a 10% L-arginine hydrochloride solution over 30 min. Blood was collected at seven time points: preloading (0 min) and 15, 30, 45, 60, 90, and 120 min after arginine loading. Circulating IRG and CPR were measured at each time point and were used to construct the arginine-stimulated time-response curves. The values of the area under the concentration-time curve for IRG ( $AUC_{IRG}$ ) and CPR ( $AUC_{CPR}$ ) between time 0 and 120 min were calculated by means of the trapezoidal rule and indicate the insulin- and glucagon-secreting responses to arginine. The homeostasis model assessment of insulin resistance (HOMA-IR) (15) and the quantitative insulin sensitivity index (QUICKI) (16) were used as conventional indexes for insulin resistance. The values for HOMA-IR and QUICKI were calculated using the following formulas:  $HOMA-IR = [\text{fasting insulin } (\mu\text{U/ml}) \times \text{fasting plasma glucose } (\text{mmol/l})] / 22.5$  and  $QUICKI = 1 / [\log[\text{fasting plasma glucose } (\mu\text{U/ml})] + \log[\text{fasting insulin } (\text{mmol/l})]]$ .

The immunoenzymometric assays used for quantifying C-peptide and insulin were performed using kits purchased from Tosoh (Shunan, Japan). Plasma glucagon levels were measured using a radioimmunoassay kit (Daiichi; Daiichi Radioisotope Labs, Tokyo, Japan). The lower limits of quantification for CPR and IRG were 0.2 ng/ml and 15.6 pg/ml, respectively. The intra- and interassay coefficients of variation were both <6%. Glucose and A1C were measured by standard methods.

### Statistical analysis

Data are expressed as means ± SD. Statistical analyses were performed with StatView software (SAS Institute, Cary, NC).

Differences among groups were tested by ANOVA with a post hoc test of Fisher's protected least significant differences. Pearson's product moment correlation coefficients were obtained to estimate linear correlation among the variables.  $P < 0.05$  was considered statistically significant.

## RESULTS

### Responses of insulin and glucagon to arginine challenge in patients with type 1 and type 2 diabetes and nondiabetic subjects


Responses of CPR and glucagon to arginine are shown in Fig. 1. Type 2 diabetic and nondiabetic subjects had relatively high levels of basal CPR and  $AUC_{CPR}$  compared with type 1 diabetic patients (Fig. 1). Interestingly, the mean glucagon response did not differ among the three treatment groups (i.e., type 1 and type 2 diabetes and control). However, some type 1 or type 2 diabetic patients exhibited an exaggerated glucagon response to arginine.

### Distinct relationship between $AUC_{IRG}$ and $AUC_{CPR}$ in patients with type 1 versus type 2 diabetes

To address the pathophysiology underlying the exaggerated glucagon response to an arginine challenge, we examined the relationship between  $AUC_{IRG}$  and  $AUC_{CPR}$  in type 1 and type 2 diabetic patients and in nondiabetic control subjects (Fig. 2). In patients with type 1 diabetes,  $AUC_{IRG}$  was negatively correlated with  $AUC_{CPR}$  ( $AUC_{IRG} = 32,462 - 17.8 AUC_{CPR}$ ;  $r = -0.388$ ,  $P = 0.023$ ) (Fig. 2A). In contrast,  $AUC_{IRG}$  was positively correlated with  $AUC_{CPR}$  in type 2 diabetic patients ( $AUC_{IRG} = 19,419 + 18.8 AUC_{CPR}$ ;  $r = 0.396$ ,  $P < 0.0001$ ) (Fig. 2B). No correlation was found between  $AUC_{IRG}$  and  $AUC_{CPR}$  in nondiabetic subjects ( $AUC_{IRG} = 30,803 - 4.7 AUC_{CPR}$ ;  $r = -0.079$ ,  $P = 0.655$ ) (Fig. 2C). These results suggest that the mechanisms involved in the glucagon response to arginine challenge are distinctly different between type 1 and type 2 diabetic patients.

### Relationship between $AUC_{IRG}$ and $AUC_{CPR}$ stratified by the class of diabetes duration in patients with type 1 diabetes

Intraislet insulin deficiency may determine the exaggerated glucagon response to arginine challenge (17). To test our hypothesis that type 1 diabetic patients with a longer diabetes duration are deficient in



**Figure 1**—IRG (A) and CPR (B) response curves to arginine. The glucagon responses were similar among type 1 and type 2 diabetic patients and nondiabetic control subjects, whereas the CPR responses of type 2 diabetic patients and control subjects were greater than those of type 1 diabetic patients. Data are shown as mean values; error bars denote SD. NS, not significant.


insulin secretory capacity, we analyzed the relationship between the  $AUC_{IRG}$  or  $AUC_{CPR}$  and diabetes duration in type 1 diabetic patients (Table 2 and supplemental Fig. 2A) and further investigated the correlation between  $AUC_{IRG}$  and  $AUC_{CPR}$  stratified by classes of diabetes duration, i.e., 0–72 months and 96–348 months (supplemental Fig. 2B). The patients with slowly progressive type 1 diabetes were excluded from this analysis because the onset of diabetes seemed to be unclear. As the diabetes duration increased,  $AUC_{CPR}$  decreased and  $AUC_{IRG}$  increased in type 1 diabetic patients (Table 2 and supplemental Fig. 2). This relationship was not evident in patients with type 2 diabetes (data not shown). These findings suggest that inislet insulin deficiency with long disease duration deter-

mines the exaggerated glucagon response to arginine challenge in patients with type 1 diabetes.

To rule out possible influences of insulin treatment on the glucagon response to arginine in type 1 diabetic patients, we analyzed the relationship of the  $AUC_{IRG}$  or  $AUC_{CPR}$  to insulin dose (supplemental Fig. 3) and glycemic control indexes such as A1C and fasting plasma glucose (Table 2 and supplemental Fig. 4). We found that neither the  $AUC_{IRG}$  nor the  $AUC_{CPR}$  was correlated with either insulin dose or glycemic control status; thus, exogenous insulin replacement therapy and glycemic control status were unlikely to have affected our results and conclusions in patients with type 1 diabetes.

#### Relationship between $AUC_{IRG}$ and $AUC_{CPR}$ stratified by class of BMI and insulin resistance indexes in patients with type 2 diabetes

Previous studies have suggested that the glucagon response to arginine is inversely related to insulin sensitivity (18–21). To test the hypothesis that obese type 2 diabetic patients have impaired insulin sensitivity, hyperinsulinemia, and hyperglucagonemia, we analyzed the relationship between the  $AUC_{IRG}$  or  $AUC_{CPR}$  and insulin resistance indexes such as BMI, basal CPR level, HOMA-IR, and QUICKI in patients with type 2 diabetes (Table 2 and supplemental Fig. 5A). Both the  $AUC_{IRG}$  and  $AUC_{CPR}$  were positively correlated with BMI, basal CPR level, and HOMA-IR, and both were negatively correlated with QUICKI. No significant cor-



**Figure 2**—Relationship between  $AUC_{IRG}$  and  $AUC_{CPR}$  in type 1 (A) and type 2 (B) diabetic patients and nondiabetic control subjects (C).  $AUC_{IRG}$  was negatively correlated with  $AUC_{CPR}$  in type 1 diabetic patients ( $r = -0.388$ ,  $P = 0.023$ ), whereas  $AUC_{IRG}$  was positively correlated with  $AUC_{CPR}$  in type 2 diabetic patients ( $r = 0.396$ ,  $P < 0.0001$ ). In the control group,  $AUC_{IRG}$  was not correlated with  $AUC_{CPR}$  ( $r = -0.079$ ,  $P = 0.655$ ).

Table 2—The relationship between  $AUC_{IRG}$  or  $AUC_{CPR}$  and clinical parameters in type 1 and type 2 diabetic patients and nondiabetic control subjects

	Diabetes duration	BMI	Basal CPR	HOMA-IR	QUICKI	A1C	FPG
$AUC_{IRG}$							
Type 1 diabetic patients							
<i>r</i>	0.445	−0.233	−0.391	ND	ND	0.022	0.009
<i>P</i>	0.033	0.208	0.024	ND	ND	0.910	0.961
Type 2 diabetic patients							
<i>r</i>	−0.174	0.467	0.269	0.361	−0.319	0.013	0.040
<i>P</i>	0.057	<0.001	0.003	0.001	0.005	0.889	0.721
Nondiabetic subjects							
<i>r</i>	ND	0.094	−0.042	0.036	−0.127	ND	ND
<i>P</i>	ND	0.592	0.813	0.851	0.502	ND	ND
$AUC_{CPR}$							
Type 1 diabetic patients							
<i>r</i>	−0.477	0.359	0.938	ND	ND	−0.194	−0.137
<i>P</i>	0.021	0.047	<0.001	ND	ND	0.323	0.447
Type 2 diabetic patients							
<i>r</i>	−0.260	0.455	0.844	0.343	−0.385	−0.204	−0.155
<i>P</i>	0.005	<0.001	<0.001	0.002	0.001	0.089	0.173
Nondiabetic subjects							
<i>r</i>	ND	0.348	0.695	0.511	−0.557	ND	ND
<i>P</i>	ND	0.043	<0.001	0.004	0.001	ND	ND

FPG, fasting plasma glucose; ND, not determined.

relation was found between the  $AUC_{IRG}$  and any of the insulin resistance indexes in either patients with type 1 diabetes (Table 2 and supplemental Fig. 6) or nondiabetic subjects (Table 2 and supplemental Fig. 7). We further analyzed the relationship between the  $AUC_{IRG}$  and  $AUC_{CPR}$  stratified by BMI class, basal CPR level, HOMA-IR, and QUICKI (supplemental Fig. 5B). Both the  $AUC_{IRG}$  and  $AUC_{CPR}$  significantly increased in relation to increases in BMI, basal CPR level, and HOMA-IR in patients with type 2 diabetes, whereas both significantly decreased with increases in QUICKI. In contrast, neither glycemic control status (Table 2 and supplemental Fig. 8) nor treatment (supplemental Fig. 9) affected the  $AUC_{IRG}$  in type 2 diabetic patients. These results suggest that insulin resistance contributes to the exaggerated glucagon response to arginine in patients with type 2 diabetes but not in those with type 1 diabetes or in nondiabetic subjects.

**CONCLUSIONS**— The arginine stimulation test has been demonstrated to be a valid method for evaluating residual  $\beta$ -cell function, even during periods of hyperglycemia (3,4,22–24). However, the relevance of the glucagon response to arginine remains uncertain and has not been comprehensively analyzed in pa-

tients with type 1 and type 2 diabetes. In the present study, we revealed the pathophysiology of the glucagon secretion response to an arginine challenge in patients with diabetes. Although no difference was observed in the glucagon response between patients with type 1 and type 2 diabetes, the glucagon response reflected a distinct pathophysiology in each type of diabetes.

In patients with type 1 diabetes,  $AUC_{IRG}$  was inversely correlated with  $AUC_{CPR}$  in the arginine challenge test and the insulin response to arginine inversely correlated with diabetes duration, suggesting that intraislet insulin deficiency determines the exaggerated glucagon response to arginine. Autoimmune type 1 diabetes is caused by a  $\beta$ -cell–targeted immune reaction and destruction with relatively conserved  $\alpha$ -cell mass (25). Therefore, the glucagon response to arginine cannot be suppressed by intrinsic insulin in hypoinsulinemic diabetic patients (12). Conversely, in patients with type 2 diabetes, the  $AUC_{CPR}$ , but not the  $AUC_{IRG}$ , was negatively correlated with diabetes duration (data not shown). In addition, patients with insulinopenic type 2 diabetes did not exhibit an exaggerated glucagon response to arginine (Fig. 2B). Based on these findings, we hypothesize that only absolute insulin deficiency, as

observed in type 1 diabetes, is associated with an exaggerated glucagon response. These results are in agreement with a previous study that suggested that absolute deficiencies in insulin secretion are the cause of an exaggerated glucagon response to arginine in patients with diabetes (11,12). In addition, insulin replacement therapy can correct these deficiencies in patients with type 1 diabetes (11,12). The molecular mechanism underlying this observation may involve the suppression of glucagon release by intraislet insulin via the GABA ( $\gamma$ -aminobutyric acid)-GABA<sub>A</sub> receptor system (26). Given that some patients displayed no C-peptide response and a relatively conserved glucagon response to arginine in the present study, it is possible that other currently unknown factors may also regulate glucagon and CPR responses to arginine.

In hyperinsulinemic patients with impaired glucose tolerance or type 2 diabetes, the glucagon response to arginine was reported to be inversely related to insulin sensitivity (19–21). In these studies, however, it remained unclear whether the exaggerated glucagon response to arginine was caused by intraislet insulin deficiency or  $\alpha$ -cell insulin resistance. In the present study, we demonstrated a positive correlation between  $AUC_{IRG}$  and  $AUC_{CPR}$  in type 2 diabetes, which was opposite the



relationship in type 1 diabetes. The exaggerated glucagon response to arginine was also related to insulin resistance. The mean AUC<sub>IRG</sub> and AUC<sub>CPR</sub> values increased with increases in BMI and insulin resistance indexes, suggesting that  $\beta$ -cell hypertrophy associated with obesity might determine the exaggerated glucagon secretion in response to arginine in patients with type 2 diabetes. This pathology was not evident in patients with type 1 diabetes. Thus, we speculate that  $\alpha$ -cell insulin resistance may be a key pathology causing the exaggerated glucagon response to arginine in patients with type 2 diabetes. In this regard, Hamaguchi et al. (18) previously reported that an exaggerated  $\alpha$ -cell response to arginine infusion in obese hyperinsulinemic patients with glucose intolerance was secondary to a reduction in insulin action on the pancreatic  $\alpha$ -cell. In addition, they observed that exogenous insulin replacement normalized these abnormalities (18), suggesting that compensatory hyperinsulinemia can overcome  $\alpha$ -cell insulin resistance. However, we did not observe such compensation in type 2 diabetic patients with severe hyperinsulinemia (Fig. 2B). Our results are in agreement with a previous report (11) that insulin replacement cannot correct an exaggerated glucagon response in patients with type 2 diabetes. These findings further support the hypothesis that  $\alpha$ -cell insulin resistance occurs with increasing systemic insulin resistance in patients with type 2 diabetes.

In summary, intraislet insulin deficiency and  $\alpha$ -cell insulin resistance may cause exaggerated glucagon secretion in response to arginine, which might in turn contribute to impaired suppression of hepatic glucose output in both type 1 and type 2 diabetes. Collectively, our data support the idea that exaggerated secretion of glucagon may be a therapeutic target in both insulinopenic diabetes and type 2 diabetes with insulin resistance. Therefore, the arginine challenge test could be useful for assessing the heterogeneous nature of diabetes and may be a valid method for identifying responders to therapy targeted at glucagon and its receptor, such as glucagon-like peptide 1 analogs. Large-scale clinical studies are needed to test this hypothesis.

## References

- Consoli A: Role of liver in pathophysiology of NIDDM. *Diabetes Care* 15:430–441, 1992
- Unger RH: Glucagon physiology and pathophysiology in the light of new advances. *Diabetologia* 28:574–578, 1985
- Aronoff SL, Bennett PH, Rushforth NB, Miller M, Unger RH: Arginine-stimulated hyperglucagonemia in diabetic Pima Indians. *Diabetes* 25:404–407, 1976
- Aronoff SL, Bennett PH, Rushforth NB, Miller M, Unger RH: Normal glucagon response to arginine infusion in “prediabetic” Pima Indians. *J Clin Endocrinol Metab* 43:279–286, 1976
- Unger RH: Glucagon physiology and pathophysiology. *N Engl J Med* 285:443–449, 1971
- Unger RH: Role of glucagon in the pathogenesis of diabetes: the status of the controversy. *Metabolism* 27:1691–1709, 1978
- Raskin P, Unger RH: Hyperglucagonemia and its suppression: importance in the metabolic control of diabetes. *N Engl J Med* 299:433–436, 1978
- Petersen KF, Sullivan JT: Effects of a novel glucagon receptor antagonist (Bay 27-9955) on glucagon-stimulated glucose production in humans. *Diabetologia* 44:2018–2024, 2001
- Jiang G, Zhang BB: Glucagon and regulation of glucose metabolism. *Am J Physiol Endocrinol Metab* 284:E671–E678, 2003
- Degn KB, Juhl CB, Sturis J, Jakobsen G, Brock B, Chandramouli V, Rungby J, Landau BR, Schmitz O: One week’s treatment with the long-acting glucagon-like peptide 1 derivative liraglutide (NN2211) markedly improves 24-h glycemia and  $\alpha$ - and  $\beta$ -cell function and reduces endogenous glucose release in patients with type 2 diabetes. *Diabetes* 53:1187–1194, 2004
- Raskin P, Aydin I, Unger RH: Effect of insulin on the exaggerated glucagon response to arginine stimulation in diabetes. *Diabetes* 25:227–229, 1976
- Gerich JE, Lorenzi M, Tsalikian E, Bohannon NV, Schneider V, Karam JH, Forsham PH: Effects of acute insulin withdrawal and administration on plasma glucagon responses to intravenous arginine in insulin-dependent subjects. *Diabetes* 25:955–960, 1976
- Expert Committee on the Diagnosis and Classification of Diabetes Mellitus: Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus (Committee Report). *Diabetes Care* 26 (Suppl. 1):S5–S20, 2003
- Kobayashi T, Tamemoto K, Nakanishi K, Kato N, Okubo M, Kajio H, Sugimoto T, Murase T, Kosaka K: Immunogenetic and clinical characterization of slowly progressive IDDM. *Diabetes Care* 16:780–788, 1993
- Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC: Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 28:412–419, 1985
- Katz A, Nambi SS, Mather K, Baron AD, Follmann DA, Sullivan G, Quon MJ: Quantitative insulin sensitivity check index: a simple, accurate method for assessing insulin sensitivity in humans. *J Clin Endocrinol Metab* 85:2402–2410, 2000
- Takamura T, Sakurai M, Nakamura M, Shimizu A, Ota T, Misu H, Takeshita Y, Tsuchiyama N, Kurita S, Ando H, Kaneko S: Factors associated with improvement of fasting plasma glucose level by meal-time dosing of a rapid-acting insulin analog in type 2 diabetes. *Diabetes Res Clin Pract* 75:278–284, 2007
- Hamaguchi T, Fukushima H, Uehara M, Wada S, Shirotani T, Kishikawa H, Ichinose K, Yamaguchi K, Shichiri M: Abnormal glucagon response to arginine and its normalization in obese hyperinsulinemic patients with glucose intolerance: importance of insulin action on pancreatic alpha cells. *Diabetologia* 34:801–806, 1991
- Kalkhoff RK, Gossain VV, Matute ML: Plasma glucagon in obesity: response to arginine, glucose and protein administration. *N Engl J Med* 289:465–467, 1973
- Gossain VV, Matute ML, Kalkhoff RK: Relative influence of obesity and diabetes on plasma alpha-cell glucagon. *J Clin Endocrinol Metab* 38:238–243, 1974
- Larsson H, Ahren B: Islet dysfunction in insulin resistance involves impaired insulin secretion and increased glucagon secretion in postmenopausal women with impaired glucose tolerance. *Diabetes Care* 23:650–657, 2000
- Palmer JP, Benson JW, Walter RM, Ensink JW: Arginine-stimulated acute phase of insulin and glucagon secretion in diabetic subjects. *J Clin Invest* 58:565–570, 1976
- Ahren B, Pacini G: Importance of quantifying insulin secretion in relation to insulin sensitivity to accurately assess beta cell function in clinical studies. *Eur J Endocrinol* 150:97–104, 2004
- Robertson RP: Consequences on  $\beta$ -cell function and reserve after long-term pancreas transplantation. *Diabetes* 53:633–644, 2004
- Sayama K, Imagawa A, Okita K, Uno S, Moriwaki M, Kozawa J, Iwashashi H, Yamagata K, Tamura S, Matsuzawa Y, Hanafusa T, Miyagawa J, Shimomura I: Pancreatic beta and alpha cells are both decreased in patients with fulminant type 1 diabetes: a morphometrical assessment. *Diabetologia* 48:1560–1564, 2005
- Xu E, Kumar M, Zhang Y, Ju W, Obata T, Zhang N, Liu S, Wendt A, Deng S, Ebina Y, Wheeler MB, Braun M, Wang Q: Intra-islet insulin suppresses glucagon release via GABA-GABAA receptor system. *Cell Metab* 3:47–58, 2006