

OBSERVATIONS

Changes in Serum Adiponectin Concentrations Correlate With Changes in BMI, Waist Circumference, and Estimated Visceral Fat Area in Middle-Aged General Population

Adiponectin was identified as an adipocytokine in the human adipose tissue cDNA library. It has antiatherosclerotic and antidiabetic properties in experimental studies, and its blood levels are low in obesity, diabetes, cardiovascular diseases, and metabolic syndrome. Several studies have reported that weight reduction in massively obese subjects is associated with a rise in serum adiponectin (APN) concentration (1–3). However, the relationship between changes in APN and BMI, waist circumference (WC), and visceral fat accumulation (VFA) in general population has not been reported. The present study investigated 1-year change in APN (Δ APN) in relation to changes in BMI (Δ BMI), WC (Δ WC), and estimated visceral fat area (Δ eVFA) in middle-aged general population.

The study subjects were 2,024 middle-aged Japanese (1,619 men [45.7 \pm 10.4 years] and 405 women [45.6 \pm 9.3 years], mean \pm SD) who were employees of Amagasaki city office and had undergone annual health checkup in both 2004 and 2005 and were not taking any medications for diabetes, hypertension, or dyslipidemia. The study was approved by the human ethics committee of Osaka University, and a signed informed consent was obtained from each participant. Height, weight, and WC were measured in standing position. BMI was calculated as weight in kilograms divided by the

square of height in meters. WC at umbilical level was measured with a nonstretchable tape in late expiration while standing (in cm). VFA was estimated noninvasively by bioelectrical impedance analysis (BIA) (4). Briefly, the voltage recorded at the flank to the flow of current between the umbilicus and the back correlates significantly with VFA and is not influenced by subcutaneous fat. We reported previously that VFA estimated by BIA correlates significantly with that determined by computed tomography (4). APN was measured using latex particle-enhanced turbidimetric assay (5). Δ APN correlated negatively with Δ BMI, both in men ($r = -0.256$, $P < 0.0001$) and women ($r = -0.223$, $P < 0.0001$) and with Δ WC in men only ($r = -0.191$, $P < 0.0001$), but no correlation was found in women ($P = 0.097$), and Δ APN correlated negatively with Δ eVFA in both men ($r = -0.189$, $P < 0.0001$) and women ($r = -0.121$, $P = 0.015$).

APN is likely influenced by genetic and environmental factors. It has been reported that APN is associated with single nucleotide polymorphisms in adiponectin gene. The present study demonstrated that changes in body fat, i.e., reduction in BMI, WC, and eVFA, correlated with a rise in APN in middle-aged general population, which to our knowledge is the first such report. We used BIA to evaluate VFA in the present study, and further research on both visceral and subcutaneous fat areas measured by computed tomography scan is needed to clarify the effects of visceral and subcutaneous adiposity on APN.

YUKIYOSHI OKAUCHI, MD¹
KEN KISHIDA, MD¹
TOHRU FUNAHASHI, MD¹
MIDORI NOGUCHI, RN²
TOMOKO OGAWA, RN²
MIWA RYO, MD¹
KOHEI OKITA, MD¹
HIROMI IWASHI, MD¹
AKIHISA IMAGAWA, MD¹
TADASHI NAKAMURA, MD³
YUJI MATSUZAWA, MD⁴
ICHIRO SHIMOMURA, MD¹

From the ¹Department of Metabolic Medicine, Graduate School of Medicine, Osaka University, Suita, Osaka, Japan; the ²Amagasaki City Office, General Affairs Bureau, Personal Department, Payroll Section, Employee Health Promotion Section, Amagasaki, Hyogo, Japan; ³Kawasaki Hospital, Kobe, Hyogo, Japan; and ⁴Sumitomo Hospital, Osaka, Japan.

Corresponding author: Ken Kishida, kkishida@imed2.med.osaka-u.ac.jp.

DOI: 10.2337/dc09-1130. Clinical trial reg. no. UMIN000002084, www.umin.ac.jp.

© 2009 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

Acknowledgments—No potential conflicts of interest relevant to this article were reported.

References

- Hotta K, Funahashi T, Arita Y, Takahashi M, Matsuda M, Okamoto Y, Iwahashi H, Kuriyama H, Ouchi N, Maeda K, Nishida M, Kihara S, Sakai N, Nakajima T, Hasegawa K, Muraguchi M, Ohmoto Y, Nakamura T, Yamashita S, Hanafusa T, Matsuzawa Y. Plasma concentrations of a novel, adipose-specific protein, adiponectin, in type 2 diabetic patients. *Arterioscler Thromb Vasc Biol* 2000;20:1595–1599
- Yang WS, Lee WJ, Funahashi T, Tanaka S, Matsuzawa Y, Chao CL, Chen CL, Tai TY, Chuang LM. Weight reduction increases plasma levels of an adipose-derived anti-inflammatory protein, adiponectin. *J Clin Endocrinol Metab* 2001;86:3815–3819
- Ng TW, Watts GF, Barrett PH, Rye KA, Chan DC. Effect of weight loss on LDL and HDL kinetics in the metabolic syndrome: associations with changes in plasma retinol-binding protein-4 and adiponectin levels. *Diabetes Care* 2007;30:2945–2950
- Ryo M, Maeda K, Onda T, Katashima M, Okumiyama A, Nishida M, Yamaguchi T, Funahashi T, Matsuzawa Y, Nakamura T, Shimomura I. A new simple method for the measurement of visceral fat accumulation by bioelectrical impedance. *Diabetes Care* 2005;28:451–453
- Nishimura A, Sawai T. Determination of adiponectin in serum using a latex particle-enhanced turbidimetric immunoassay with an automated analyzer. *Clin Chim Acta* 2006;371:163–168