

# Effectiveness of a Computerized Insulin Order Template in General Medical Inpatients With Type 2 Diabetes

A cluster randomized trial

DEBORAH J. WEXLER, MD, MSC<sup>1</sup>  
PETER SHRADER, MA<sup>2</sup>

SEAN M. BURNS, MD<sup>1</sup>  
ENRICO CAGLIERO, MD<sup>1</sup>

**OBJECTIVE**— To determine whether an electronic order template for basal-bolus insulin ordering improves mean blood glucose in hospitalized general medical patients with hyperglycemia and type 2 diabetes.

**RESEARCH DESIGN AND METHODS**— We randomly assigned internal medicine resident teams on acute general medical floors to the use of an electronic insulin order template or usual insulin ordering. We measured diabetes care parameters for 1 month on all patients with type 2 diabetes and blood glucose <60 mg/dl or >180 mg/dl treated by these physicians.

**RESULTS**— Intervention group patients ( $n = 65$ ) had mean glucose of  $195 \pm 66$  mg/dl. Control group patients ( $n = 63$ ) had mean glucose of  $224 \pm 57$  mg/dl ( $P = 0.004$ ). In the intervention group, there was no increase in hypoglycemia.

**CONCLUSIONS**— Access to a computer insulin order template was associated with improved mean glucose levels without increasing hypoglycemia in patients with type 2 diabetes.

*Diabetes Care* 33:2181–2183, 2010

Physiological, basal-bolus insulin prescribing is safe, effective (1), and the standard of care in hospitalized patients with type 2 diabetes and hyperglycemia (2). Yet only about half of such patients are prescribed basal insulin in the hospital (3). Order templates to support basal-bolus insulin prescribing (usually as part of a comprehensive inpatient diabetes quality improvement program) have been effective in improving glycemia in observational trials (4–8). Randomized trials have shown more modest effects (9,10). Knowledge of appropriate insulin ordering is a barrier to ordering basal-bolus insulin among inpatient providers (11–13).

We tested the hypothesis that giving internal medicine residents access to an electronic insulin order template would

be more effective than usual insulin ordering in lowering mean blood glucose in medical inpatients with type 2 diabetes.

## RESEARCH DESIGN AND METHODS

We piloted a simple electronic insulin order template based on previously studied protocols (1,4,14) and internal review by diabetologists and pharmacists at a tertiary care medical center with a proprietary computerized order entry system. The template presents sequential screens linking to a weight-based insulin dose calculator and facilitating prescription of a total daily dose of insulin of 0.5 units/kg, half in basal glargine and half in prandial aspart, with supplemental aspart (supplementary Appendix 1, available at <http://care.diabetesjournals.org/cgi/content/full/dc10-0964/DC1>).

Using a computerized coin toss, we randomly assigned seven teams of providers (42 internal medicine residents) working in general medical acute care units to have the option to use the order template (intervention group) or to use usual insulin ordering (control group). All residents received a 30-min educational session and a two-page pamphlet on how to prescribe and titrate basal-bolus insulin (supplementary Appendix 1).

We identified all patients prescribed insulin by these providers during the study period (30 April 2009 to 27 May 2009). We excluded patients with type 1 diabetes ( $n = 11$ ) and type 2 diabetes with blood glucose between 60 and 180 mg/dl because the order set was designed to apply to hyperglycemia in patients with type 2 diabetes without other risks for hypoglycemia (i.e., liver or renal failure). Diagnosis of type 2 diabetes was confirmed by chart review. We determined age, sex, race, length of stay, and primary diagnosis of the patients from the medical record. Glycemic parameters were calculated from point-of-care glucose values, using mean glucose as the main outcome for its easy interpretability and similarity to other metrics (15). Secondary outcomes included the rate of prolonged hyperglycemia (three consecutive glucose values >240 mg/dl); hospital-stay rate of hypoglycemia (any glucose value <60 mg/dl or <40 mg/dl); the rate of use of sliding scale insulin alone (having sliding scale insulin orders without any other insulin orders at any point during admission); rate of initial prescription of basal insulin (any order for long-acting insulin within 1 day of hospital admission); and basal insulin at any time (any order for long-acting insulin at any time during the hospital stay).

The study had 80% power to detect a difference in mean glucose of 30 mg/dl between groups, assuming SD of 60 mg/dl. We performed Student *t* tests and Wilcoxon rank sum tests to determine differences between means and medians of continuous and non-normally distributed variables and performed  $\chi^2$  or Fisher

From the <sup>1</sup>Massachusetts General Hospital Diabetes Center and Harvard Medical School, Boston, Massachusetts; and the <sup>2</sup>Division of General Internal Medicine, Massachusetts General Hospital, Boston, Massachusetts.

Corresponding author: Deborah J. Wexler, [dwexler@partners.org](mailto:dwexler@partners.org).

Received 25 May 2010 and accepted 17 July 2010. Published ahead of print at <http://care.diabetesjournals.org> on 27 July 2010. DOI: 10.2337/dc10-0964. Clinical trial reg. no. NCT01130558, [clinicaltrials.gov](http://clinicaltrials.gov).

© 2010 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

exact tests for categorical variables. We repeated mean glucose estimates accounting for repeated (glucose) and correlated (provider) measures and adjusting for baseline glucose (supplementary Appendix 2). All analyses were performed with SAS version 9.1. The protocol was approved by the Partners Healthcare Institutional Review Board.

**RESULTS**— During the 4-week study period, 144 insulin-treated patients with type 2 diabetes were admitted to the seven study teams. Mean glucose was  $186 \pm 56$  mg/dl in intervention patients and  $206 \pm 61$  mg/dl in control patients ( $P = 0.004$ ). Excluding 16 patients whose point-of-care glucose values were between 60 and 180 mg/dl left a study sample size of 128; 127 had at least one value  $>180$  mg/dl and 1 had a glucose value  $<60$  mg/dl. Sixty-three patients were treated by providers with access to the order template (intervention group), and 65 were treated by providers who received the pamphlet and brief teaching session alone (control group). There were no significant differences between groups in patient age, sex, race, length of stay, or primary diagnosis. Mean glucose was significantly better in the intervention group ( $194 \pm 66$  mg/dl vs.  $224 \pm 57$  mg/dl,  $P = 0.004$ ) (Table 1). The rates of basal insulin orders, any hypoglycemia, and severe hypoglycemia did not differ between groups.

**CONCLUSIONS**— In this randomized trial of a computer order template to support basal-bolus insulin prescribing for general acute medical inpatients with type 2 diabetes, we found an improved mean glucose level in patients of providers given access to the order template.

Mean glucose levels in both groups were higher than the goal for inpatient glucose levels in noncritically ill patients (2), and mean basal insulin doses were low, consistent with persistent inadequate treatment. The rate of basal insulin ordered at admission was low and increased equivalently in both groups to a rate over 60%; this compares favorably to a rate of 43% in a national sample of insulin-treated inpatients with type 2 diabetes (3). The rate of sliding scale insulin alone (35%) was equivalent to that of a comparable national sample (35%) (3). Given the efficacy of basal-bolus treatment, it seems even a low rate of appropriate insulin ordering may be associated with modest improvement in mean glucose levels. Use of the order template may

Table 1—Baseline characteristics and results

	Control	Intervention	P*
n	63	65	
Age (years)	$70 \pm 13.6$	$68 \pm 14.3$	0.4
Male subjects, n (%)	34 (54)	40 (61)	0.4
White subjects, n (%)	37 (84)	39 (80)	0.6
Length of stay (days), median (IQR)	5 (3–11)	6 (3–10)	0.6
Primary diagnosis, n (%)			
Cardiac	17 (27)	13 (20)	0.8
GI or liver disease	10 (16)	10 (15)	
Pulmonary (including pneumonia)	9 (14)	9 (14)	
Infection, nonpneumonia	7 (11)	11 (17)	
Diabetes	2 (3)	3 (5)	
Other	18 (29)	19 (29)	
Results			
Mean glucose (mg/dl)	$224 \pm 57$	$194 \pm 66$	0.004
Sliding scale alone, %	35	38	0.7
Basal insulin, day of admission, %	31	30	0.9
Basal insulin at any time, %	65	61	0.7
Prolonged hyperglycemia (3 consecutive glucose $>240$ mg/dl), %	38	26	0.2
Hypoglycemia ( $<60$ mg/dl at any time), %	14	12	0.7
Severe hypoglycemia ( $<40$ mg/dl), %	1	0	0.5
Basal insulin dose (units), median (IQR)	16 (10–34)	18 (10–28)	0.4

Data are means  $\pm$  SD, median (interquartile range [IQR]), percent, or n (%). \*Student *t* test for continuous variables, Wilcoxon rank sum test for length of stay and mean insulin dose, and  $\chi^2$  or Fisher exact test for categorical variables. GI, gastrointestinal.

have been limited by its optional use and the minimal support for the order set, in contrast to similar programs at other centers (9,10).

In conclusion, access to an electronic basal-bolus insulin order template was associated with a significant improvement in glycemic control among patients with type 2 diabetes without increasing the rate of hypoglycemia but did not substantially change insulin ordering behavior. A “smarter” template with alerts based on glucose levels and nutritional status and forced rather than optional use, coupled with more intensive implementation support, might further improve the care of hospitalized patients with type 2 diabetes.

**Acknowledgments**— This research study was funded by the Massachusetts General Hospital (MGH) Clinical Innovation Award program, an initiative of the MGH Quality and Safety Program. D.J.W. is supported by a National Institute of Diabetes and Digestive and Kidney Diseases Career Development Award (K23 DK 080 228). S.B. is supported in part by a National Institutes of Health institutional training grant (T32 DK007028-34).

No potential conflicts of interest relevant to this article were reported.

E.C. conceived of the project. E.C. and D.J.W. designed the study. E.C., D.J.W., and

S.M.B. developed the order template. E.C., D.J.W., and S.M.B. executed the study. P.S. managed the data and conducted the repeated-measures analysis. D.W. performed the remainder of the data analysis. D.J.W. drafted the manuscript. E.C., P.S., and S.M.B. reviewed/edited the manuscript. D.J.W. had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

The authors thank Linda Jones and Melissa Wright of Partners Healthcare Information Systems for extensive assistance with developing the computerized order template and Rhodes Berube and Mary Ann Walsh, RN, of Massachusetts General Hospital for assistance with electronic data acquisition.

## References

1. Umpierrez GE, Smiley D, Zisman A, Prieto LM, Palacio A, Ceron M, Puig A, Mejia R. Randomized study of basal-bolus insulin therapy in the inpatient management of patients with type 2 diabetes (RABBIT 2 trial). *Diabetes Care* 2007;30:2181–2186
2. Moghissi ES, Korytkowski MT, DiNardo M, Einhorn D, Hellman R, Hirsch IB, Inzucchi SE, Ismail-Beigi F, Kirkman MS, Umpierrez GE, American Association of Clinical Endocrinologists, American Diabetes Association. American Association of Clinical Endocrinologists and American Diabetes Association consensus state-

- ment on inpatient glycemic control. *Endocr Pract* 2009;15:353–369
3. Wexler DJ, Meigs JB, Cagliero E, Nathan DM, Grant RW. Prevalence of hyper- and hypoglycemia among inpatients with diabetes: a national survey of 44 U.S. hospitals. *Diabetes Care* 2007;30:367–369
  4. Maynard G, Lee J, Phillips G, Fink E, Renvall M. Improved inpatient use of basal insulin, reduced hypoglycemia, and improved glycemic control: effect of structured subcutaneous insulin orders and an insulin management algorithm. *J Hosp Med* 2009;4:3–15
  5. Donihi AC, DiNardo MM, DeVita MA, Korytkowski MT. Use of a standardized protocol to decrease medication errors and adverse events related to sliding scale insulin. *Qual Saf Health Care* 2006;15:89–91
  6. Murphy DM, Vercruyse RA, Bertucci TM, Wall MJ, Schriever AE, Nabhan FA, Barron WM, Emanuele MA. Reducing hyperglycemia hospitalwide: the basal-bolus concept. *Jt Comm J Qual Patient Saf* 2009;35:216–223
  7. Schnipper JL, Ndumele CD, Liang CL, Pendergrass ML. Effects of a subcutaneous insulin protocol, clinical education, and computerized order set on the quality of inpatient management of hyperglycemia: results of a clinical trial. *J Hosp Med* 2009;4:16–27
  8. Guerra YS, Das K, Antonopoulos P, Borkowsky S, Fogelfeld L, Gordon MJ, Palal BM, Witsil JC, Lacuesta EA. Computerized physician order entry- based hyperglycemia inpatient protocol and glycemic outcomes: The CPOE-HIP study. *Endocr Pract* 2010;16:389–397
  9. Schnipper JL, Liang CL, Ndumele CD, Pendergrass ML. Effects of a computerized order set on the inpatient management of hyperglycemia: a cluster-randomized controlled trial. *Endocr Pract* 2009;16:209–218
  10. Noschese M, Donihi AC, Koerbel G, Karslioglu E, Dinardo M, Curll M, Korytkowski MT. Effect of a diabetes order set on glycaemic management and control in the hospital. *Qual Saf Health Care* 2008;17:464–468
  11. Cheekati V, Osburne RC, Jameson KA, Cook CB. Perceptions of resident physicians about management of inpatient hyperglycemia in an urban hospital. *J Hosp Med* 2009;4:E1–8
  12. Cook CB, Jameson KA, Hartsell ZC, Boyle ME, Leonhardi BJ, Farquhar-Snow M, Beer KA. Beliefs about hospital diabetes and perceived barriers to glucose management among inpatient midlevel practitioners. *Diabetes Educ* 2008;34:75–83
  13. Rubin DJ, Moshang J, Jabbour SA. Diabetes knowledge: are resident physicians and nurses adequately prepared to manage diabetes? *Endocr Pract* 2007;13:17–21
  14. Inzucchi SE. Clinical practice. Management of hyperglycemia in the hospital setting. *N Engl J Med* 2006;355:1903–1911
  15. Goldberg PA, Bozzo JE, Thomas PG, Mesmer MM, Sakharova OV, Radford MJ, Inzucchi SE. “Glucometrics”—assessing the quality of inpatient glucose management. *Diabetes Technol Ther* 2006;8:560–569