

The Natural History of LDL Control in Type 2 Diabetes

A prospective study of adherence to lipid guidelines

DIANA RUCKER, MD, MSC¹
JEFFREY A. JOHNSON, PHD^{1,2,3}
T.K. LEE, MD¹
DEAN T. EURICH, MSC^{2,3}

RICHARD Z. LEWANCZUK, MD, PHD¹
SCOT H. SIMPSON, PHARM, D, MSC^{3,4}
ELLEN L. TOTH, MD^{1,3}
SUMIT R. MAJUMDAR, MD, MPH^{1,2,3}

Despite randomized trials repeatedly showing the benefits of lowering LDL cholesterol with hydroxymethylglutaryl (HMG)-CoA reductase inhibitors (statins) (1–3), these medications are suboptimally used in type 2 diabetes (4–10). Although this care gap in type 2 diabetes has been frequently described in cross-sectional studies (6–11), it may be as informative to understand LDL control over time. In particular, there is growing recognition of care gaps in diabetes when comparing urban and academic settings with rural settings (11,12). Therefore, we examined changes over an 18-month period for adherence to guideline-recommended LDL cholesterol targets in a rural cohort with type 2 diabetes and determined the rates and correlates of 1) losing control of LDL cholesterol in those who were initially at target and 2) achieving control of LDL cholesterol in those who were not initially at target.

RESEARCH DESIGN AND METHODS

The Diabetes Outreach Van Enhancement (DOVE) study was a controlled trial of a multifaceted intervention directed at health care providers to improve the quality of care for rural patients with type 2 diabetes in northern Alberta, Canada. The intervention consisted of an educational outreach (“academic detailing”) service, whereby specialist phy-

sicians promoted aggressive cardiovascular risk reduction for diabetes to primary care physicians. The study rationale, design, and outcomes have been previously published (12–16). All subjects provided written consent, and the study was approved by the University of Alberta.

All patients had universal health care coverage and fee-for-service primary care physicians, with the nearest specialists being ~6 h away by vehicle. Patients were eligible if they were aged ≥ 20 years, had type 2 diabetes, and understood English. They were excluded for shortened life expectancy or inability to consent. Data were collected from 2000 through 2001, and patients were assessed at baseline and 18 months. Clinical data were collected by in-person interviews and physical assessments, while laboratory measurements were drawn locally and analyzed in one central laboratory.

Primary outcomes were defined according to LDL cholesterol levels < 2.5 mmol/l, which was the target recommended during the study (12–14,17). To explore independent correlates of guideline adherence, three multivariable logistic regression analyses were built using Stata (version 8.2; StataCorp, College Station, TX). The first model represented LDL guideline adherence at baseline; the second, loss of LDL control among those well controlled at baseline; and the third,

new achievement of LDL guideline adherence over 18 months among those not controlled at baseline. The educational intervention had no effect on lipid levels or lipid-lowering medications (14–16), but we adjusted for it in all analyses. Otherwise, candidate variables had to have a univariate significance of at least $P < 0.1$ and a multivariate significance of $P < 0.05$.

RESULTS — Overall, 393 patients with type 2 diabetes were enrolled, and 346 (88%) had complete data for these analyses. Patient-level characteristics, according to LDL target status over time, are displayed in Table 1. At baseline, 216 patients (62%) were not at LDL targets. In a cross-sectional multivariable model, independent correlates of LDL > 2.5 mmol/l were older age ($P = 0.014$) and not using lipid-lowering medications (adjusted odds ratio 2.1 [95% CI 1.2–3.6], $P = 0.013$).

After 18 months, 80% ($n = 278$) of the study population was not at recommended LDL targets. Of the 130 patients who had met targets at baseline, 83 (64%) lost control of LDL cholesterol over 18 months (Table 1). The variables independently associated with loss of LDL control were younger age ($P = 0.026$) and better education ($P = 0.041$) (Table 1).

Of the 216 patients not at LDL targets at study entry, 195 (90%) never achieved recommended LDL levels over 18 months (Table 1). In multivariable analyses, the only independent correlate of achieving LDL < 2.5 mmol/l was starting lipid-lowering medication (adjusted odds ratio 9.2 [95% CI 3.5–24.2], $P < 0.0001$).

CONCLUSIONS — Nearly two-thirds of rural patients with a median duration of type 2 diabetes of 5 years were not at guideline-recommended targets for LDL cholesterol at the start of our study, and 18 months later, nonadherence with LDL targets had actually risen to 80%. This occurred in the context of their health care providers being exposed to an educational intervention during this period that was intended to help improve cardiovascular risk management.

From the ¹Department of Medicine, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, Alberta, Canada; the ²Department of Public Health Sciences, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, Alberta, Canada; the ³Institute of Health Economics, Edmonton, Alberta, Canada; and the ⁴Faculty of Pharmacy and Pharmaceutical Sciences, University of Alberta, Edmonton, Alberta, Canada.

Address correspondence and reprint requests to Dr. Sumit R. Majumdar, University of Alberta, 2E3.07 Walter Mackenzie Health Sciences Centre, University of Alberta Hospital, 8440-112th St., Edmonton, Alberta, Canada, T6G 2B7. E-mail: me2.majumdar@ualberta.ca.

Received for publication 9 June 2006 and accepted in revised form 21 July 2006.

Abbreviations: HMG, hydroxymethylglutaryl.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

DOI: 10.2337/dc06-1205

© 2006 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Table 1—Characteristics of 346 patients with type 2 diabetes according to achievement of LDL cholesterol target <2.5 mmol/L, at baseline and over 18 months of follow-up

Characteristics	Baseline (cross-sectional data)			Over 18 months of follow-up					
	Not at target (n = 216)	Already at target (n = 130)	Adjusted odds ratio (95% CI)*	Never at target (n = 195)	Achieved target (n = 21)	Adjusted odds ratio (95% CI)*	Maintains target (n = 47)	Lost target (n = 83)	Adjusted odds ratio (95% CI)*
Age >60 years	62	50	1.8 (1.1–2.8)	62	62	—	66	41	0.6 (0.4–0.9)
Female	60	52	—	62	38	—	43	59	—
High school or more	28	39	—	22	40	—	23	48	2.4 (1.0–5.5)
Income >\$40,000/year	24	27	—	22	40	—	17	33	—
Aboriginal	26	24	—	24	19	—	28	26	—
Current smoker	16	21	—	30	10	—	20	22	—
Median duration of diabetes (years)	5	5	—	5	7	—	5	5	—
HbA _{1c} (%)	7.1 ± 1.4	7.4 ± 1.7	—	7.1 ± 1.4	7.4 ± 1.2	—	7.3 ± 1.7	7.4 ± 1.7	—
BMI >27 kg/m ²	77	84	—	79	57	—	77	88	—
Systolic blood pressure (mmHg)	132 ± 19	129 ± 17	—	131 ± 19	134 ± 20	—	129 ± 16	130 ± 17	—
Total cholesterol (mmol/L)	5.4 ± 0.8	4.1 ± 1.0	—	5.4 ± 0.8	3.6 ± 0.3	—	3.5 ± 0.4	4.8 ± 0.5	—
LDL cholesterol (mmol/L)	3.3 ± 0.6	2.0 ± 1.4	—	3.4 ± 0.5	2.3 ± 0.2	—	2.2 ± 0.3	3.0 ± 0.3	—
Lipid-lowering therapy									
Taking	14	24	2.1 (1.2–3.6)	23	71	—	30	18	—
Started	NA	NA	NA	13	57	9.2 (3.5–24.0)	9	5	—
Stopped	NA	NA	NA	2	0	—	4	4	—

Data are means ± SD or percent unless otherwise indicated. *All odds ratios adjusted for intervention status (P > 0.5, all analyses, see text) and those variables presented in the table that were significant at P < 0.05 in the relevant multivariable model. NA, not applicable; —, not included in multivariable analyses.

To our knowledge, the only comparable study examining temporal trends in hyperlipidemia management in type 2 diabetes was conducted by Mehler et al. (18), who demonstrated a stable nonadherence rate to lipid guidelines of ~50% over 5 years; specific losses or gains of adherence were not reported. Similar to our baseline findings, other cross-sectional studies have shown nonadherence to LDL guidelines ranging from 50 to 80% in Canadian (6), Australian (7), and U.S. (8–11) populations. Our results suggest that cross-sectional studies likely overestimate quality of care.

Recently, it has been suggested that it may be more useful to promote absolute reductions in LDL cholesterol rather than unattainable LDL targets, given that there is a 20% reduction in major coronary events for every 1-mmol reduction in LDL irrespective of initial LDL levels (2). Using this approach, only 12% of our population achieved a 1-mmol LDL reduction over 18 months, suggesting even modest improvements are not being achieved. This could be explained by physician chart audits, which suggest that 4% of dyslipidemic patients are treated with maximal doses of statins and even fewer are treated with combination regimens (19). Conversely, physicians themselves cite patient adherence as the most common barrier to attaining LDL targets (20). Clearly, this example of clinical inertia is a systems problem, with barriers (and potential solutions) at multiple levels (21).

This study had several limitations. First, we did not have information regarding doses of, or adherence to, lipid-lowering medications. Second, we did not collect data about changes in diet and lifestyle. Third, since patients were volunteers, our results may overestimate rates of achieving LDL targets and underestimate losses of LDL control. Fourth, given how few patients' LDL levels changed (for better or worse) over time, we were underpowered to determine all factors potentially associated with guideline adherence. Last, we examined rural patients in one Canadian province, and our results may not be generalizable to urban populations or to other nations.

In summary, we found that target LDL cholesterol levels are rarely attained and, even when achieved, are all too easily lost in patients with type 2 diabetes. Much remains to be done to learn how to translate evidence into everyday clinical practice (21).

Acknowledgments—This work was supported by the Canadian Diabetes Association, the Institute of Health Economics, and the Alliance of Canadian Health Outcomes Researchers in Diabetes (funded by the Canadian Institutes of Health Research [CIHR] and its partners). D.R. is supported by the SCOLAR Training Program funded by CIHR and its partners. J.A.J. holds a Canada Research Chair and receives salary support from the Alberta Heritage Foundation for Medical Research (AHFMR). D.T.E. receives salary support from AHFMR. S.H.S. receives salary support from CIHR. S.R.M. receives salary support from AHFMR and from CIHR.

References

1. Pyorala K, Pedersen TR, Kjekshus J, Faergeman O, Olsson AG, Thorgeirsson G: Cholesterol lowering with simvastatin improves prognosis of diabetic patients with coronary heart disease: a subgroup analysis of the Scandinavian Simvastatin Survival Study (4S). *Diabetes Care* 20: 614–620, 1997
2. Baigent C, Keech A, Kearney PM, Blackwell L, Buck G, Pollicino C, Kirby A, Sourjina T, Peto R, Collins R, Simes R: Efficacy and safety of cholesterol-lowering treatment: prospective meta-analysis of data from 90,056 participants in 14 randomised trials of statins. *Lancet* 366:1267–1278, 2005
3. Vijan S, Hayward RA: Pharmacologic lipid-lowering therapy in type 2 diabetes mellitus: background paper for the American College of Physicians. *Ann Intern Med* 140:650–658, 2004
4. Hill-Briggs F, Gary TL, Bone LR, Hill MN, Levine DM, Brancati FL: Medication adherence and diabetes control in urban African Americans with type 2 diabetes. *Health Psychol* 24:349–357, 2005
5. Majumdar SR, Gurwitz JH, Soumerai SB: Undertreatment of hyperlipidemia in the secondary prevention of coronary artery disease. *J Gen Intern Med* 14:711–717, 1999
6. Harris SB, Ekoe JM, Zdanowicz Y, Web-

- ster-Bogaert S: Glycemic control and morbidity in the Canadian primary care setting (results of the diabetes in Canada evaluation study). *Diabetes Res Clin Pract* 70:90–97, 2005
7. Kemp TM, Barr EL, Zimmet PZ, Cameron AJ, Welborn TA, Colagiuri S, Phillips P, Shaw JE: Glucose, lipid, and blood pressure control in Australian adults with type 2 diabetes: the 1999–2000 AusDiab. *Diabetes Care* 28:1490–1492, 2005
8. Grant RW, Buse JB, Meigs JB: Quality of diabetes care in U.S. academic medical centers: low rates of medical regimen change. *Diabetes Care* 28:337–442, 2005
9. Resnick HE, Foster GL, Bardsley J, Ratner RE: Achievement of American Diabetes Association clinical practice recommendations among U.S. adults with diabetes, 1999–2002. *Diabetes Care* 29:531–537, 2006
10. Pearson TA, Laurora I, Chu H, Kafonek S: The lipid treatment assessment project (L-TAP): a multicenter survey to evaluate the percentages of dyslipidemic patients receiving lipid-lowering therapy and achieving low-density lipoprotein cholesterol goals. *Arch Intern Med* 160:459–467, 2000
11. Saadine JB, Engelgau MM, Beckles GL, Gregg EW, Thompson TJ, Venkat Narayan KM: A diabetes report card for the United States: quality of care for the 1990s. *Ann Intern Med* 136:565–574, 2002
12. Majumdar SR, Johnson JA, Guirguis LM, Lewanczuk RZ, Lee TK, Toth EL: Rationale and design for the DOVE study: a prospective, controlled trial of an intervention to improve care for patients with diabetes in rural communities. *Canadian Journal of Diabetes* 25:173–179, 2001
13. Toth EL, Majumdar SR, Guirguis LM, Lewanczuk RZ, Lee TK, Johnson JA: Compliance with clinical practice guidelines for type 2 diabetes in rural patients: treatment gaps and opportunities for improvement. *Pharmacotherapy* 23:659–665, 2003
14. Majumdar SR, Guirguis LM, Toth EL, Lewanczuk RZ, Lee TK, Johnson JA: Controlled trial of a multifaceted intervention for improving quality of care for rural patients with type 2 diabetes. *Diabetes Care* 26:3061–3066, 2003
15. Johnson JA, Eurich DT, Toth EL, Lewanczuk RZ, Lee TK, Majumdar SR: Generalizability and persistence of a multifaceted intervention for improving quality of care for rural patients with type 2 diabetes. *Diabetes Care* 28:783–788, 2005
16. Maddigan SL, Majumdar SR, Guirguis LM, Lewanczuk RZ, Lee TK, Toth EL, Johnson JA: Improvements in patient-reported outcomes associated with an intervention to enhance quality of care for rural patients with type 2 diabetes: results of a controlled trial. *Diabetes Care* 27: 1306–1312, 2004
17. Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III): Executive Summary of the Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA* 285:2486–2497, 2001
18. Mehler PS, Esler A, Estacio RO, MacKenzie TD, Hiatt WR, Schrier RW: Lack of improvement in the treatment of hyperlipidemia among patients with type 2 diabetes. *Am J Med* 114:377–382, 2003
19. Rodondi N, Peng T, Karter AJ, Bauer DC, Vittinghoff E, Tang S, Pettitt D, Kerr EA, Selby JV: Therapy modifications in response to poorly controlled hypertension, dyslipidemia, and diabetes mellitus. *Ann Intern Med* 144:475–484, 2006
20. Leiter LA, Betteridge DJ, Charca AR, Chait A, Ferrannini E, Haffner SM, Kadowaki T, Tuomilehto J, Zimmet PZ, Newman CB, Hey-Hadavi J, Walkinshaw C: AUDIT study: evidence of global undertreatment of dyslipidaemia in patients with type 2 diabetes mellitus. *British Journal of Diabetes and Vascular Disease* 6:31–40, 2006
21. Majumdar SR, McAlister FA, Furberg CD: From knowledge to practice in chronic cardiovascular disease: a long and winding road. *J Am Coll Cardiol* 43:1738–1742, 2004

Downloaded from http://ada.silverchair.com/care/article-pdf/29/11/2506/668255/zdc01106002506.pdf by guest on 02 March 2024