



References

1. Leese GP, Morris A: Prediction of diabetic foot ulcer occurrence using commonly available clinical information (Letter). *Diabetes Care* 29:2562–2563, 2006
2. Leese GP, Reid F, Green V, McAlpine R, Cunningham S, Emslie-Smith AM, Morris AD, McMurray B, Connacher AC: Stratification of foot ulcer risk in patients with diabetes: a population-based study. *Int J Clin Pract* 60:541–545, 2006
3. Boyko EJ, Ahroni JH, Cohen V, Nelson KM, Heagerty PJ: Prediction of diabetic foot ulcer occurrence using commonly available clinical information: the Seattle Diabetic Foot Study. *Diabetes Care* 29:1202–1207, 2006
4. Boyko EJ, Ahroni JH, Stensel V, Forsberg RC, Davignon DR, Smith DG: A prospective study of risk factors for diabetic foot ulcer: the Seattle Diabetic Foot Study. *Diabetes Care* 22:1036–1042, 1999
5. Boyko EJ, Ahroni JH, Smith DG, Davignon D: Increased mortality associated with diabetic foot ulcer. *Diabet Med* 13: 967–972, 1996

Standards of Medical Care in Diabetes—2006

Response to the American Diabetes Association

I recently encountered a discrepancy in the American Diabetes Association's (ADA's) recommendations regarding conventional versus SI units for HDL cholesterol (1). The article states "raise HDL cholesterol to >40 mg/dl (1.15 mmol/l)." Simple calculation shows that 40 mg/dl = 40 × 0.02586 mmol/l = 1.03 mmol/l, rather than 1.15 mmol/l. The same error is also noted in the 2005 version.

Furthermore, in regard to the ADA's recommendation to use statin therapy for diabetic patients without overt cardiovascular disease (CVD), the recommendation to treat "regardless of baseline LDL" might have extended beyond the evidence quoted. I find the recommendation's evidence rather weak, despite a similar recommendation elsewhere (2). For evidence on diabetic patients without overt cardiovascular problems, two studies are listed (3,4). Here is an abbreviated synopsis:

The Heart Protection Study (3) showed cardiovascular benefit of statin therapy for LDL >3.0 mmol/l similar to

Downloaded from http://ada.silverchair.com/article-pdf/29/11/2563/666190/zdc011062563-a.pdf by guest on 01 March 2024

available clinical information. *Diabetes Care* 29:1202–1207, 2006

2. Leese GP, Reid F, Green V, McAlpine R, Cunningham S, Emslie-Smith A, Morris AD, McMurray B, Connacher A: Predicting foot ulceration in diabetes: validation of a clinical tool in a population-based study. *Int J Clin Pract* 60:541–545, 2006
3. Scottish Intercollegiate Guideline Network (SIGN): *SIGN Guideline Publication No.55: The Management of Diabetes*. Edinburgh, U.K., Royal College of Physicians, Edinburgh, 2001. Available from <http://http://www.sign.ac.uk/guidelines/fulltext/55/index.html>. Accessed 19 September 2006
4. International Diabetes Federation: *International Concensus on the Diabetic Foot: Five Cornerstones, Identification of the Foot at Risk* (Consensus Meeting). May 2003. Available from http://www.iwgdf.org/index.php?option=com_content&task=view&id=39&Itemid=60. Accessed 19 September 2006

Prediction of Diabetic Foot Ulcer Occurrence Using Commonly Available Clinical Information

Response to Leese and Morris

We appreciate the interest of Leese and Morris (1) in our article and enjoyed reading their publication (2), which was published 1 month before our own and hence was not cited by us (3). We wish to point out that their characterization of our population as having been "recruited from a hospital diabetes clinic" is not correct, as we clearly describe that our subjects were recruited from a general internal medicine clinic (3). With regard to the generalizability of our findings, they should be applicable to the ~25 million U.S. veterans and other males with similar clinical and demographic characteristics enrolled in a primary care setting. Although Leese and Morris write that they have "already addressed" foot ulcer prediction in their publication, they do not cite our publication from 1999, which presented a prediction model for diabetic foot ulcer and anticipated several of the findings of our recent article (4).

Leese and Morris describe their findings as useful for "all-comers" in a general community setting and in specialized

clinics; they also suggest that their findings are valid for male and female subjects and for both types of diabetes (1). However, their data and analysis do not provide strong support for these statements for several reasons. First, although they used a population-sampling strategy that targeted 8,923 subjects with diabetes, only 3,526 (40%) underwent a clinical foot risk assessment. Thus, most subjects in the sampling frame were not included in the analysis, which certainly raises concerns regarding the validity of findings and limits the degree of confidence with which one can recommend these results for "all-comers." Second, although both sexes and diabetes types were included in their study, the appropriate analysis of interaction to determine whether the results apply similarly in these subgroups of interest was not performed.

Leese and Morris bring up the issue of competing risks, in that individuals at higher risk for diabetic foot ulcer are also at higher risk for death. They refer to our "suspicion" that the death rate in people at high risk for ulcer is increased, but this would not be our preferred wording, as we feel certain about this association, having published data reporting this finding in 1996 (5). Comparison of cumulative risks, as in their report, may thus lead to biased results due to differential follow-up times by degree of risk. Fortunately, analytic methods that compare failure times and employ censoring, which we used in our recent publication, can address this problem.

Hopefully these recent efforts to develop foot ulcer prediction models and others will lead to more accurate identification of individuals at higher risk of foot ulcer and stimulate the development of better preventive strategies to reduce morbidity and mortality from these complications.

EDWARD J. BOYKO, MD, MPH^{1,2}
 JESSIE H. AHRONI, ARNP, PHD¹
 KARIN M. NELSON, MD^{1,2}

From the ¹Seattle Epidemiologic Research and Information Center, VA Puget Sound Health Care System, Seattle, Washington; and the ²Department of Medicine, University of Seattle, Seattle, Washington.

Address correspondence to Edward J. Boyko, MD, MPH, VA Epidemiologic Research and Information Center, 1100 Olive Way, Suite 1400, Seattle, WA 98101. E-mail: eboyko@u.washington.edu.

DOI: 10.2337/dc-06-1661

© 2006 by the American Diabetes Association.

