

# Two-Year Statin Therapy Does Not Alter the Progression of Intima-Media Thickness in Patients With Type 2 Diabetes Without Manifest Cardiovascular Disease

EDITH D. BEISHUIZEN, MD<sup>1</sup>  
MARCEL A. VAN DE REE, MD, PHD<sup>2</sup>  
J. WOUTER JUKEMA, MD, PHD<sup>3</sup>  
JOUKE T. TAMMSMA, MD, PHD<sup>1</sup>

J. CAREL M. VAN DER VIJVER, MD, PHD<sup>4</sup>  
A. EDO MEINDERS, MD, PHD<sup>1</sup>  
HEIN PUTTER, PHD<sup>5</sup>  
MENNO V. HUISMAN, MD, PHD<sup>1</sup>

**OBJECTIVE** — Cardiovascular disease (CVD) is the most important cause of mortality in patients with type 2 diabetes. We aimed to determine the effect of statin therapy versus placebo on the progression of carotid intima-media thickness (IMT) in type 2 diabetic patients without manifest CVD.

**RESEARCH DESIGN AND METHODS** — A randomized, placebo-controlled, double-blind clinical trial was performed in 250 patients with type 2 diabetes. Patients were given either 0.4 mg cerivastatin or placebo daily. In August 2001, when cerivastatin was withdrawn from the market, 0.4 mg cerivastatin was replaced by 20 mg simvastatin without deblinding the study. The primary end point was the change of mean common carotid IMT, as measured by B-mode ultrasound, over 2 years.

**RESULTS** — Common carotid IMT at baseline was 0.780 mm in the placebo group and 0.763 mm in the statin group and did not change significantly after 2 years. There was no significant difference in IMT change in any carotid segment between the groups. LDL cholesterol was reduced by 25% in the statin group and increased by 8% in the placebo group ( $P < 0.001$ ). Cardiovascular events occurred in 12 patients in the placebo group and two patients in the statin group ( $P = 0.006$ ).

**CONCLUSIONS** — There was no effect of 2 years' statin therapy on carotid IMT in type 2 diabetic subjects. The natural history of IMT in our patients was milder than anticipated. In contrast, we observed a significantly lower cardiovascular event rate on statin therapy. Prognostic tools other than IMT should be explored in this patient group.

*Diabetes Care* 27:2887–2892, 2004

From the <sup>1</sup>Department of General Internal Medicine, Leiden University Medical Center, Leiden, the Netherlands; the <sup>2</sup>Department of Internal Medicine, Diaconessenhuis, Utrecht, the Netherlands; the <sup>3</sup>Department of Cardiology, Leiden University Medical Center, Leiden, the Netherlands; the <sup>4</sup>Department of Internal Medicine, Leyenburg Hospital, the Hague, the Netherlands; and the <sup>5</sup>Department of Biostatistics, Leiden University Medical Center, Leiden, the Netherlands.

Address correspondence and reprint requests to E.D. Beishuizen, Department of General Internal Medicine, C1-R41, Leiden University Medical Center, P.O. Box 9600, 2300 RC Leiden, Netherlands. E-mail: e.d.beishuizen@lumc.nl or m.v.huisman@lumc.nl.

Received for publication 25 June 2004 and accepted in revised form 11 September 2004.

**Abbreviations:** ALT, alanine aminotransferase; CAD, coronary artery disease; CVD, cardiovascular disease; IMT, intima-media thickness.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

© 2004 by the American Diabetes Association.

Cardiovascular disease (CVD), including cerebrovascular disease, coronary artery disease (CAD), and peripheral vascular disease, is the most important cause of mortality in patients with type 2 diabetes (1). The severity and progression of atherosclerosis can be assessed noninvasively by ultrasonographic measurements of the intima-media thickness (IMT) in the carotid and femoral arteries. Ultrasonographic IMT measurements of the far wall relate to histological IMT measurements (2). Carotid IMT correlates with prevalent CVD (3), angiographically proven coronary atherosclerosis (4), and risk factors for CVD, including LDL cholesterol (5,6). In prospective studies, carotid IMT has proven to be predictive of CVD (7–9), and as a consequence, IMT is increasingly used as an intermediate end point in clinical trials. Mean common carotid IMT in middle-aged subjects without CAD is reported to range from 0.71 to 0.91 mm in diabetic patients vs. 0.66 to 0.74 mm in control subjects (10,11). In diabetes, IMT is less consistently correlated to classical risk factors such as LDL cholesterol. Importantly, at the time our study was designed, data on the progression and predictive value of IMT in type 2 diabetes were lacking.

During the last 10 years, large clinical trials have demonstrated that HMG-CoA (hydroxy-methylglutaryl coenzyme A) reductase inhibitors (statins) reduce the risk of cardiovascular events in the setting of secondary and primary prevention (12–15). Subgroup analyses in diabetic patients in several of these studies showed conflicting results (14–16). However, these trials were not specifically designed for type 2 diabetic patients in the setting of primary prevention. We therefore set out to evaluate the effect of statin therapy versus placebo on the progression of ca-

rotid and femoral IMT in type 2 diabetic patients without established CVD.

## RESEARCH DESIGN AND METHODS

Patients were recruited from the departments of internal medicine at two nonacademic teaching hospitals, the Leyenburg Hospital and the Red Cross Hospital, the Hague, the Netherlands. Subjects were eligible for the study if they had been diagnosed with type 2 diabetes for at least 1 year, were aged 30–80 years, and were without a history of CVD (defined as CAD, electrocardiographic criteria for a past myocardial infarction, ischemic stroke, peripheral artery bypass surgery, percutaneous transluminal angioplasty, or amputation because of atherosclerotic disease).

At a screening visit, fasting blood samples were drawn and a resting electrocardiogram performed. Patients with fasting total cholesterol >6.9 mmol/l or <4.0 mmol/l, triglycerides >6.0 mmol/l, creatinine kinase values more than three times and alanine aminotransferase (ALT) more than two times the upper limit of normal, or creatinine clearance <30 ml/min were excluded. Any lipid-lowering therapy had to be discontinued 8 weeks before the screening visit. The study was approved by the medical ethics committees of both hospitals and performed in accordance with the Declaration of Helsinki.

The primary end point of the study was the change in mean IMT of the common carotid artery after 24 months. Secondary end points were the changes in mean and maximum IMT of the carotid bifurcation, internal carotid artery, common femoral artery, and superficial femoral artery and the changes in aggregate carotid IMT (defined as the average of the mean IMT of the three carotid segments), all after 24 months. The change in mean IMT after 12 months was also considered a secondary end point. The following predefined cardiovascular events were evaluated during the study: cardiovascular death, nonfatal myocardial infarction, percutaneous transluminal coronary angioplasty, coronary artery bypass graft surgery, nonfatal stroke, peripheral artery bypass graft, percutaneous transluminal angioplasty, or amputation because of atherosclerotic disease.

After giving written informed consent, 250 patients participated at least 1 week after the screening visit. Patients were randomly assigned to receive 0.4 mg

cerivastatin (Bayer, Mijdrecht, the Netherlands) or placebo daily for a period of 2 years. Double-blind study medication was assigned using a predetermined computer-generated randomization scheme with a block size of 10. On 18 August 2001, cerivastatin was withdrawn from the market due to reports of serious morbidity and mortality possibly related to the drug (17). At that moment, all 250 patients had been included in the study with a mean follow-up of 15.4 months (range 6–23).

All patients were instructed to discontinue the study drug. The study was not unblinded at any time point. No patient had developed myopathy, and creatinine kinase values above five times the upper limit of normal had not been observed. After consultation with independent experts in lipid-lowering treatment studies, it was decided to continue the study. Cerivastatin (0.4 mg) was replaced by simvastatin (20 mg) daily, on the basis of a comparable LDL reduction (18,19). Both simvastatin and matching placebo tablets (Merck Sharp & Dohme, Haarlem, the Netherlands) were given according to the original allocation. The study was continued 1 month after the discontinuation of cerivastatin. The total use of study medication was kept at 24 months, resulting in the study being prolonged for 1 month.

### Follow-up

Patients returned to the study site after a 12-h fast at 3, 6, 12, 18, and 24 months, when blinded lipid and safety measurements (creatinine kinase and ALT) were performed. Carotid IMT was measured at baseline, 12 months, and 24 months. Femoral IMT was performed at baseline and 24 months. Two-year follow-up for clinical events was performed for all 250 patients.

### Ultrasound measurements

Ultrasound imaging was performed with an Acuson Aspen scanner with a linear array 7.5-MHz probe. All images were recorded digitally and on a S-VHS videotape for off-line, blinded analysis by an independent core laboratory (Heartcore, Leiden, the Netherlands). During the study, all measurements were performed by the same two certified ultrasonographers.

In the supine position, the left and right carotid arteries, near and far walls, were examined longitudinally at the angle that resulted in an optimal and maximal

IMT (while avoiding plaques) for each segment. The segments scanned were the distal 1.0 cm of the common carotid artery, the carotid bifurcation, and the proximal 1.0 cm of the internal carotid artery. The optimal angle was used for follow-up. The same procedure was done for the common femoral artery and superficial femoral artery.

For each segment, three R wave-triggered images were stored. Mean and maximal IMT were measured, when possible, over the entire 1 cm of the vessel segment. The three IMT measurements were averaged. To obtain mean and maximal IMT per vessel segment, far and near wall, left and right values were averaged. During the first year of the study, a reproducibility investigation was performed for the two ultrasonographers in 16 subjects. For the common carotid artery, interobserver variability (expressed as mean difference  $\pm$  SD) was  $0.0082 \pm 0.050$  mm and intraobserver variability was  $0.0067 \pm 0.049$  and  $0.00036 \pm 0.058$  mm for the two observers. For the common femoral artery, interobserver variability was  $0.039 \pm 0.11$  mm and intraobserver variability was  $0.0085 \pm 0.10$  and  $0.060 \pm 0.078$  mm for the two observers.

### Laboratory investigations

All laboratory measurements were performed at the Department of Clinical Chemistry and Hematology of the Leyenburg Hospital, according to ISO 15189 standard procedures.

### Statistical analysis

When this study was designed, no data were available on IMT progression in type 2 diabetes. From clinical studies in patients with CAD, we assumed a progression rate of 0.03 mm per 2 years for the common carotid artery IMT. The number of patients needed to detect a difference in mean common carotid artery IMT of 0.04 mm after 2 years (expected SD 0.10) with a power of 80% ( $\alpha = 0.05$ ) was 100 patients in each group. To allow for a 20% drop-out rate, the total number of patients randomized would be 250.

The primary treatment comparison is between placebo and statin therapy in patients completing the study (on-treatment analysis). Changes from baseline within each treatment group were analyzed using Student's paired *t* test. Comparisons of the effects between the treatment

Table 1—Baseline characteristics of 250 randomized patients

	Placebo	Statin
n	125	125
Male sex	57 (46)	61 (49)
Age (years)	58.2 ± 11.4	58.8 ± 11.3
Ethnicity		
Caucasian	86 (69)	83 (66)
Indo-Asian	20 (16)	28 (22)
Other	19 (15)	14 (11)
BMI (kg/m <sup>2</sup> )	31.0 ± 6.0	31.0 ± 6.3
Waist-to-hip ratio	0.99 ± 0.09	0.98 ± 0.08
Current smoker	33 (26)	28 (22)
Hypertension	66 (53)	60 (48)
Diabetes duration (years)*	7 ± 8	6 ± 7
Insulin use	69 (55)	62 (50)
HbA <sub>1c</sub> (%)	7.60 ± 1.48	7.53 ± 1.10
Microalbuminuria†	19 (15)	24 (19)

Data are means ± SD or n (%). \*Median ± SD. †Men, >2.5 g/mol creatinine; women, >3.5 g/mol creatinine.

groups were performed using Student's independent samples *t* test. Mixed-model analysis was used as a sensitivity analysis to assess the influence of missing values on the results, under the assumption of "missing at random" (20), and to investigate systematic differences between replications, positions, and between far and near wall.

Stepwise regression techniques were used to investigate the effect on baseline IMT and on changes in IMT of sex, age, smoking habits, ethnicity, blood pressure, anthropometric parameters, and duration of cerivastatin versus simvastatin use. To test the equivalence of 0.4 mg cerivastatin and 20 mg simvastatin, LDL levels before and after the switch to simvastatin were compared using Student's paired *t* test. Correlation between changes in IMT and changes in lipid levels were evaluated by calculating Pearson's correlation coefficients. The occurrence of clinical events was expressed as a proportion and evaluated using  $\chi^2$  test or Fisher's ex-

act test as appropriate. All analyses were two sided with a level of significance of  $\alpha = 0.05$ .

**RESULTS**— Of a total of 302 patients screened, 52 did not fulfill the entry criteria. The baseline characteristics of the 250 randomized patients did not differ between the groups and are reported in Table 1. None of the patients had recently been on lipid-lowering therapy.

Of the 250 patients randomized, 68 did not complete the study: 46 in the placebo group and 22 in the statin group. In 16 patients in the placebo group and in 8 in the statin group, the only reason for discontinuation was the withdrawal of cerivastatin from the market. Drop-out rates were slightly lower in the Caucasian group than in the Indo-Asian and other ethnic groups (22 vs. 35%, respectively, 42%,  $P = 0.02$ ). The other baseline characteristics of the 182 patients who completed the study did not differ from the 68 drop outs (data not shown). Overall com-

pliance, as assessed by pill counting, was 97% and was equal in the statin and placebo groups. Compliance was not reduced after the switch to simvastatin.

### Cholesterol reductions

LDL cholesterol was reduced by 25% in the statin group and increased by 8% in the placebo group ( $P < 0.001$ ), and changes in HDL cholesterol and triglycerides were not significantly different between the groups (Table 2). Average LDL cholesterol levels were higher after the switch to simvastatin (2.34 before vs. 2.56 mmol/l after the switch,  $P < 0.001$ ).

### IMT

Baseline mean IMTs were not significantly different between the groups. Common carotid artery IMT in the placebo group was  $0.780 \pm 0.129$  mm at baseline and  $0.774 \pm 0.124$  mm at 2 years ( $P = 0.50$ ), and in the statin group, it was  $0.763 \pm 0.124$  mm at baseline and  $0.765 \pm 0.116$  mm at 2 years ( $P = 0.78$ ) (Table 3). There was no significant difference between the change in IMT in the placebo and statin groups (mean difference  $-0.0075$  mm [95% CI  $-0.0281$  to  $0.0132$  mm],  $P = 0.48$ ). After 2 years, the mean changes in IMT of the other segments were also not significantly different between the groups and compared with baseline. This was also observed for the changes in maximal IMT (data not shown). Finally, the changes in mean common carotid artery IMT after 1 year were equal in both groups ( $-0.0155$  mm in the placebo group vs.  $-0.0166$  mm in the statin group,  $P$  for difference in change = 0.90). Mixed-model analysis confirmed these results.

Determinants for baseline IMT were age ( $r = 0.358$ ,  $P < 0.001$ ) and systolic blood pressure ( $r = 0.26$ ,  $P < 0.001$ ). Baseline IMT and changes in IMT were

Table 2—Plasma lipid and lipoprotein concentrations

	Placebo (n = 79)			Statin (n = 103)			
	Baseline	2 years	P	Baseline	2 years	P	P*
Total cholesterol (mmol/l)	5.60 ± 0.77	5.74 ± 0.93	0.058	5.49 ± 0.72	4.49 ± 1.01	<0.001	<0.001
LDL cholesterol (mmol/l)	3.55 ± 0.71	3.78 ± 0.81	0.003	3.44 ± 0.71	2.58 ± 0.95	<0.001	<0.001
HDL cholesterol (mmol/l)	1.21 ± 0.37	1.22 ± 0.38	0.963	1.23 ± 0.39	1.20 ± 0.36	0.144	0.284
Triglycerides (mmol/l)	1.88 ± 0.79	1.72 ± 1.22	0.206	1.82 ± 0.97	1.60 ± 1.38	0.043	0.371
ApoB100 (mg/l)	1.15 ± 0.23	1.11 ± 0.24	0.094	1.10 ± 0.21	0.84 ± 0.26	<0.001	<0.001

Data are means ± SD. \*P value for difference in percent change between placebo and statin group. To convert to mg/dl: cholesterol, multiply by 38.6; triglycerides, multiply by 88.5. Apo, apolipoprotein.

Table 3—Mean IMT of 182 patients who completed the study

	Baseline		2 years		Mean change		P
	IMT	SD	IMT	SD	IMT	95% CI	
Placebo (n = 79)							
Primary end point							
Common carotid artery	0.780	0.129	0.774	0.124	-0.006	-0.0223 to 0.0109	0.50
Secondary end point							
Carotid bifurcation	0.815	0.148	0.805	0.143	-0.010	-0.0267 to 0.0072	0.25
Internal carotid artery	0.640	0.136	0.670	0.130	0.031	-0.0104 to 0.0718	0.14
Aggregate carotid IMT	0.757	0.137	0.763	0.120	0.006	-0.0117 to 0.0229	0.52
Common femoral artery	0.663	0.149	0.652	0.141	-0.011	-0.0390 to 0.0165	0.42
Superficial femoral artery	0.551	0.111	0.549	0.099	-0.002	-0.0236 to 0.0197	0.86
Statin (n = 103)							
Primary end point							
Common carotid artery	0.763	0.124	0.765	0.116	0.002	-0.0112 to 0.0149	0.78
Secondary end point							
Carotid bifurcation	0.823	0.140	0.806	0.122	-0.017	-0.0358 to 0.0017	0.07
Internal carotid artery	0.684	0.183	0.689	0.181	0.005	-0.0211 to 0.0314	0.69
Aggregate carotid IMT	0.759	0.116	0.762	0.105	0.003	-0.0116 to 0.0175	0.69
Common femoral artery	0.630	0.151	0.635	0.148	0.005	-0.0152 to 0.0258	0.61
Superficial femoral artery	0.543	0.092	0.538	0.107	-0.005	-0.0187 to 0.0078	0.42

Values are in millimeters. Mean change: mean change from baseline to 2 years.

not correlated with LDL cholesterol or any other lipid parameter. Baseline IMT and changes in IMT were also not related to sex, ethnicity, diabetes duration, insulin use, HbA<sub>1c</sub>, anthropometric parameters, or smoking habits. The effect of the two statins used was analyzed by correcting the change in IMT for duration of cerivastatin treatment (range 6–23 months). This did not change the results.

**Clinical events**

Cardiovascular events occurred in 12 patients in the placebo group and 2 in the statin group (P = 0.006). Coronary events occurred in four patients in the placebo group and none in the statin group (P = 0.122). Four patients in the placebo group and three in the statin group died. The causes of death were cancer (n = 4), sepsis (n = 1), and hemorrhagic stroke (n = 2). Malignancies occurred in eight patients: four in the placebo group and four in the statin group. Myalgia was reported 18 times in the statin group and 26 times in the placebo group and was never accompanied by an increase in creatinine kinase. In one patient in the statin group, at his 24-month visit ALT was raised more than three times above the upper limit of normal. This was attributed to steatosis hepatis.

**CONCLUSIONS**— This is the first prospective study in patients with type 2 diabetes but without overt CVD that investigated the effect of statins versus placebo on carotid and femoral IMT. Despite a mean LDL cholesterol reduction of 25%, we did not find any effect of 2 years' statin therapy on mean common carotid IMT.

In patients with familial hypercholesterolemia and in patients with established CAD, statin therapy has resulted in significantly less progression or even regression of carotid IMT (21–23). In the only other randomized controlled lipid intervention IMT study in type 2 diabetic patients, 3 years' therapy with bezafibrate did not have any effect on carotid and femoral IMT (24). Our findings warrant several remarks. First, the mean LDL cholesterol reduction of 25% is fully comparable to statin-induced LDL cholesterol reductions ranging between 22 and 29% in studies of nondiabetic patients, showing a significant effect after 18–24 months on carotid IMT (21,23,25,26). Second, contrary to the postulated progression of mean common carotid artery IMT of 0.03 mm per 2 years, in the present study there was a nonsignificant regression of 0.006 mm per 2 years in the placebo group. It could be argued that our patient population had been low risk. However, we included diabetic patients with a broad

range in age and diabetes duration, while their baseline common carotid artery IMT was quite comparable to that of patients in other studies (10,27). Moreover, the observed rate of first major vascular events (myocardial infarction, strokes, and revascularizations) in our placebo group, which translates to 14% per 5 years, is similar to the incidence rate of 13.5% in the diabetic placebo subgroup without prior CVD in the Heart Protection Study (16). Thus, it seems unlikely that our results have been influenced by any healthy volunteer effect. Third, in our study, we observed no association between LDL cholesterol reduction and IMT reduction. This is at variance with the effect of statin therapy on IMT in nondiabetic patients (22) but in agreement with the effect of 3 years' bezafibrate treatment (LDL cholesterol reduction 9.6%) on carotid and femoral IMT in type 2 diabetic patients (24). Equally, baseline IMT in the present study was not correlated to baseline LDL cholesterol or any other lipid level, similar to the results of several previous cross-sectional studies in type 2 diabetes (10,28). Finally, given the low inter- and intra-observer variability in our IMT measurements compared with other studies (29), we strongly feel that the quality of the assessments has not biased the results.

As our study was designed with the assumption of similar IMT progression rates in diabetic and coronary patients, we conclude from the lack of IMT progression in our placebo group that IMT progression rates in diabetic patients are lower than those of patients with CAD. This is supported by recently published cohort studies in which diabetes was not predictive of IMT progression (30,31). Moreover, the ARIC (Atherosclerosis Risk in Communities) study (32) and the IRAS (Insulin Resistance Atherosclerosis Study) (33) have recently shown progression rates for diabetic subjects of 0.011 and 0.0072 mm/year, respectively, both of which are lower than those observed in patients with CAD (34).

We observed a statistically significant effect on the incidence of predefined cardiovascular events. This is in agreement with the results of the recently published CARDS (Collaborative Atorvastatin Diabetes Study) (35) and with the results of a meta-analysis on the effects of lipid management for type 2 diabetic patients in primary prevention (36). As we did not find any IMT regression, we postulate that statin-induced cardiovascular event reduction in diabetic patients is not related to IMT regression. From a pathophysiological point of view, the intimal and medial layers of the vessel wall in type 2 diabetes are most likely irreversibly changed by processes such as extracellular matrix glycosylation and media calcification (37,38). These changes may resist global regression based on interference with local intravascular cholesterol metabolism. We hypothesize that although statins do not influence the irreversibly changed glycosylated extracellular matrix, it may well have an effect on outcome in type 2 diabetic patients by its beneficial influence on plaque vulnerability. To accurately measure IMT in our study, we avoided eccentric plaques; therefore, we cannot address this hypothesis with the available data.

Our study has its limitations. First, cerivastatin was withdrawn from the market, resulting in a change from cerivastatin to simvastatin. After correcting the change in IMT for duration of cerivastatin treatment, however, the results remained unchanged. Second, as a result of the withdrawal of cerivastatin, we had a higher withdrawal rate than anticipated in our sample size estimation. However, except for ethnicity, which was not a deter-

minant of IMT or IMT progression, baseline characteristics did not differ between the drop outs and the 182 patients who fulfilled the study. Moreover, given the narrow CI of the mean difference in common carotid artery IMT change between placebo and statin (95% CI  $-0.0281$  to  $0.0132$  mm), we can exclude a type II error.

In conclusion, 2 years of statin therapy in a broad range of type 2 diabetic patients without prior manifest atherosclerotic disease did not have any effect on carotid and femoral IMT. The natural history of atherosclerosis progression, as measured by IMT in type 2 diabetic patients, was milder than previously postulated.

We observed a lower cardiovascular event rate in patients on statin therapy, which is in line with other clinical trials. As this benefit has not been related to IMT regression, other mechanistic explanations, like a beneficial effect on plaque vulnerability, might be of importance. Vessel wall biology in type 2 diabetes is distinct from other high-risk patients, and this implies that prognostic tools other than IMT should be evaluated in this patient group.

**Acknowledgments**—Bayer (Mijdrecht, the Netherlands) supplied the study medication and supported the study until cerivastatin was withdrawn from the market (8 August 2001). Merck Sharp & Dohme (Haarlem, the Netherlands) supplied the simvastatin/placebo medication for the remaining study period. J.W.J. is an established clinical investigator of the Netherlands Heart Foundation 2001 (D032). The study concept, design, and data analysis and writing of the manuscript were the responsibilities of the authors.

We thank M. Spaans for his invaluable role as research nurse; J. Ruiterman, R. Pijpers, and N. de Jonge for laboratory investigations; A. van Weert and analysts from Heartcore for their skilled ultrasound analysis; and all the patients for their participation.

## References

1. Stamler J, Vaccaro O, Neaton JD, Wentworth D: Diabetes, other risk factors, and 12-yr cardiovascular mortality for men screened in the Multiple Risk Factor Intervention Trial. *Diabetes Care* 16:434–444, 1993
2. Pignoli P, Tremoli E, Poli A, Oreste P, Paoletti R: Intimal plus medial thickness of the arterial wall: a direct measurement with ultrasound imaging. *Circulation* 74: 1399–1406, 1986

3. Burke GL, Evans GW, Riley WA, Sharrett AR, Howard G, Barnes RW, Rosamond W, Crow RS, Rautaharju PM, Heiss G: Arterial wall thickness is associated with prevalent cardiovascular disease in middle-aged adults: the Atherosclerosis Risk in Communities (ARIC) study. *Stroke* 26: 386–391, 1995
4. Lekakis JP, Papamichael CM, Cimponeriu AT, Stamatelopoulos KS, Papaioannou TG, Kanakakis J, Alevizaki MK, Papanagiotou A, Kalofoutis AT, Stamatelopoulos SF: Atherosclerotic changes of extracoronary arteries are associated with the extent of coronary atherosclerosis. *Am J Cardiol* 85:949–952, 2000
5. Salonen JT, Salonen R: Risk factors for carotid and femoral atherosclerosis in hypercholesterolaemic men. *J Intern Med* 236:561–566, 1994
6. O'Leary DH, Polak JF, Kronmal RA, Savage PJ, Borhani NO, Kittner SJ, Tracy R, Gardin JM, Price TR, Furberg CD: Thickening of the carotid wall: a marker for atherosclerosis in the elderly? Cardiovascular Health Study Collaborative Research Group. *Stroke* 27:224–231, 1996
7. O'Leary DH, Polak JF, Kronmal RA, Manolio TA, Burke GL, Wolfson SK Jr: Carotid-artery intima and media thickness as a risk factor for myocardial infarction and stroke in older adults: Cardiovascular Health Study Collaborative Research Group. *N Engl J Med* 340: 14–22, 1999
8. Bots ML, Hoes AW, Koudstaal PJ, Hofman A, Grobbee DE: Common carotid intima-media thickness and risk of stroke and myocardial infarction: the Rotterdam Study. *Circulation* 96:1432–1437, 1997
9. Chambless LE, Heiss G, Folsom AR, Rosamond W, Szklo M, Sharrett AR, Clegg LX: Association of coronary heart disease incidence with carotid arterial wall thickness and major risk factors: the Atherosclerosis Risk in Communities (ARIC) study, 1987–1993. *Am J Epidemiol* 146: 483–494, 1997
10. Niskanen L, Rauramaa R, Miettinen H, Haffner SM, Mercuri M, Uusitupa M: Carotid artery intima-media thickness in elderly patients with NIDDM and in nondiabetic subjects. *Stroke* 27:1986–1992, 1996
11. Goff DC Jr, D'Agostino RB Jr, Haffner SM, Saad MF, Wagenknecht LE: Lipoprotein concentrations and carotid atherosclerosis by diabetes status: results from the Insulin Resistance Atherosclerosis Study. *Diabetes Care* 23:1006–1011, 2000
12. Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet* 344:1383–1389, 1994
13. MRC/BHF Heart Protection Study of cho-

- lesterol lowering with simvastatin in 20,536 high-risk individuals: a randomised placebo-controlled trial. *Lancet* 360: 7–22, 2002
14. Major outcomes in moderately hypercholesterolemic, hypertensive patients randomized to pravastatin vs usual care: the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT-LLT). *JAMA* 288:2998–3007, 2002
  15. Sever PS, Dahlof B, Poulter NR, Wedel H, Beevers G, Caulfield M, Collins R, Kjeldsen SE, Kristinsson A, McInnes GT, Mehlsen J, Nieminen M, O'Brien E, Ostergren J: Prevention of coronary and stroke events with atorvastatin in hypertensive patients who have average or lower-than-average cholesterol concentrations, in the Anglo-Scandinavian Cardiac Outcomes Trial–Lipid Lowering Arm (ASCOT-LLA): a multicentre randomised controlled trial. *Lancet* 361:1149–1158, 2003
  16. Collins R, Armitage J, Parish S, Sleight P, Peto R: MRC/BHF Heart Protection Study of cholesterol-lowering with simvastatin in 5963 people with diabetes: a randomised placebo-controlled trial. *Lancet* 361: 2005–2016, 2003
  17. Staffa JA, Chang J, Green L: Cerivastatin and reports of fatal rhabdomyolysis. *N Engl J Med* 346:539–540, 2002
  18. Jones P, Kafonek S, Laurora I, Hunninghake D: Comparative dose efficacy study of atorvastatin versus simvastatin, pravastatin, lovastatin, and fluvastatin in patients with hypercholesterolemia (the CURVES study). *Am J Cardiol* 81:582–587, 1998
  19. Stein E: Cerivastatin in primary hyperlipidemia: a multicenter analysis of efficacy and safety. *Atherosclerosis* 139 (Suppl. 1): S15–S22, 1998
  20. Laird NM, Ware JH: Random-effects models for longitudinal data. *Biometrics* 38:963–974, 1982
  21. MacMahon S, Sharpe N, Gamble G, Hart H, Scott J, Simes J, White H: Effects of lowering average of below-average cholesterol levels on the progression of carotid atherosclerosis: results of the LIPID Atherosclerosis Substudy: LIPID Trial Research Group. *Circulation* 97:1784–1790, 1998
  22. Smilde TJ, van Wissen S, Wollersheim H, Trip MD, Kastelein JJ, Stalenhoef AF: Effect of aggressive versus conventional lipid lowering on atherosclerosis progression in familial hypercholesterolaemia (ASAP): a prospective, randomised, double-blind trial. *Lancet* 357:577–581, 2001
  23. de Groot E, Jukema JW, van Boven AJ, Reiber JH, Zwinderman AH, Lie KI, Ackersstaff RA, Bruschke AV: Effect of pravastatin on progression and regression of coronary atherosclerosis and vessel wall changes in carotid and femoral arteries: a report from the Regression Growth Evaluation Statin Study. *Am J Cardiol* 76:40C–46C, 1995
  24. Elkeles RS, Diamond JR, Poulter C, Dhanjil S, Nicolaides AN, Mahmood S, Richmond W, Mather H, Sharp P, Feher MD: Cardiovascular outcomes in type 2 diabetes: a double-blind placebo-controlled study of bezafibrate: the St. Mary's, Ealing, Northwick Park Diabetes Cardiovascular Disease Prevention (SENDCAP) study. *Diabetes Care* 21:641–648, 1998
  25. Furberg CD, Adams HP Jr, Applegate WB, Byington RP, Espeland MA, Hartwell T, Hunninghake DB, Lefkowitz DS, Probstfield J, Riley WA: Effect of lovastatin on early carotid atherosclerosis and cardiovascular events: Asymptomatic Carotid Artery Progression Study (ACAPS) Research Group. *Circulation* 90:1679–1687, 1994
  26. Mercuri M, Bond MG, Sirtori CR, Veglia F, Crepaldi G, Feruglio FS, Descovich G, Ricci G, Rubba P, Mancini M, Gallus G, Bianchi G, D'Alo G, Ventura A: Pravastatin reduces carotid intima-media thickness progression in an asymptomatic hypercholesterolemic mediterranean population: the Carotid Atherosclerosis Italian Ultrasound Study. *Am J Med* 101:627–634, 1996
  27. Keven K, Gullu S, Cesur V, Gurses MA, Aytac S, Kamel N, Erdogan G: Carotid intima media thickness in patients with newly diagnosed T2DM and impaired glucose tolerance. *Diabetes Nutr Metab* 12: 49–51, 1999
  28. Kanters SD, Algra A, Banga JD: Carotid intima-media thickness in hyperlipidemic type I and type II diabetic patients. *Diabetes Care* 20:276–280, 1997
  29. Kanters SD, Algra A, van Leeuwen MS, Banga JD: Reproducibility of in vivo carotid intima-media thickness measurements: a review. *Stroke* 28:665–671, 1997
  30. Lakka TA, Laukkanen JA, Rauramaa R, Salonen R, Lakka HM, Kaplan GA, Salonen JT: Cardiorespiratory fitness and the progression of carotid atherosclerosis in middle-aged men. *Ann Intern Med* 134: 12–20, 2001
  31. van der Meer IM, Iglesias del Sol A, Hak AE, Bots ML, Hofman A, Witteman JC: Risk factors for progression of atherosclerosis measured at multiple sites in the arterial tree: the Rotterdam Study. *Stroke* 34:2374–2379, 2003
  32. Chambless LE, Folsom AR, Davis V, Sharrett R, Heiss G, Sorlie P, Szklo M, Howard G, Evans GW: Risk factors for progression of common carotid atherosclerosis: the Atherosclerosis Risk in Communities Study, 1987–1998. *Am J Epidemiol* 155: 38–47, 2002
  33. Wagenknecht LE, Zaccaro D, Espeland MA, Karter AJ, O'Leary DH, Haffner SM: Diabetes and progression of carotid atherosclerosis: the Insulin Resistance Atherosclerosis Study. *Arterioscler Thromb Vasc Biol* 23:1035–1041, 2003
  34. Bots ML, Evans GW, Riley WA, Grobbee DE: Carotid intima-media thickness measurements in intervention studies: design options, progression rates, and sample size considerations: a point of view. *Stroke* 34:2985–2994, 2003
  35. Colhoun HM, Betteridge DJ, Durrington PN, Hitman GA, Neil HA, Livingstone SJ, Thomason MJ, Mackness MI, Charlton-Menys V, Fuller JH: Primary prevention of cardiovascular disease with atorvastatin in type 2 diabetes in the Collaborative Atorvastatin Diabetes Study (CARDS): multicentre randomised placebo-controlled trial. *Lancet* 364:685–696, 2004
  36. Vijan S, Hayward RA: Pharmacologic lipid-lowering therapy in type 2 diabetes mellitus: background paper for the American College of Physicians. *Ann Intern Med* 140:650–658, 2004
  37. Schmidt AM, Yan SD, Wautier JL, Stern D: Activation of receptor for advanced glycation end products: a mechanism for chronic vascular dysfunction in diabetic vasculopathy and atherosclerosis. *Circ Res* 84:489–497, 1999
  38. Sakata N, Takeuchi K, Noda K, Saku K, Tachikawa Y, Tashiro T, Nagai R, Horiuchi S: Calcification of the medial layer of the internal thoracic artery in diabetic patients: relevance of glycoxidation. *J Vasc Res* 40:567–574, 2003