

# Age, Sex, and Ethnic Variations in Serum Insulin Concentrations Among U.S. Youth

Findings from the National Health and Nutrition Examination Survey 1999–2002

EARL S. FORD, MD, MPH<sup>1</sup>  
CHAOYANG LI, MD, PHD<sup>1</sup>

GIUSEPPINA IMPERATORE, MD, PHD<sup>2</sup>  
STEPHEN COOK, MD<sup>3</sup>

**OBJECTIVE**— Distributions of serum concentrations of insulin among adolescents and young adults are poorly understood in the U.S. The objective of this study was to describe the distribution of serum insulin across demographic characteristics of U.S. adolescents and young adults.

**RESEARCH DESIGN AND METHODS**— A total of 1,791 male and female subjects aged 12–19 years who participated in the National Health and Nutrition Examination Surveys for 1999–2002 were included in the analyses.

**RESULTS**— Among male participants, serum concentrations of insulin increased from age 12 to 14 years before decreasing. Among female participants, concentrations were highest at age 13 years before decreasing steadily through age 19 years. Among participants aged 12–17 years but not those aged 18–19 years, females had higher mean log-transformed concentrations than males ( $P$ , Wald,  $F = 0.038$  and  $0.125$ , respectively) after adjusting for age and ethnicity. After adjusting for age and BMI percentile, mean log-transformed concentrations were higher in African-American females aged 12–17 years than in white or Mexican-American participants. No significant ethnic differences were found among female participants aged 18–19 years or male participants aged 12–19 years. Concentrations of insulin increased strongly with increasing levels of BMI.

**CONCLUSIONS**— These results provide detailed information about serum concentrations of insulin in a representative sample of U.S. adolescents and young adults and may be useful to monitor future trends of this risk factor for diabetes and cardiovascular disease.

*Diabetes Care* 29:2605–2611, 2006

The increase in obesity among children in the U.S. has generated great concern about their future health because of the increased risk of cardiovascular disease and diabetes associated with obesity (1,2). Obesity is associated with insulin resistance, a state characterized by the body's resistance to the actions of insulin, which results in increased insulin production by the pancreas and hyperin-

sulinemia. Measures of adiposity are generally strongly related to concentrations of insulin (3). In children and adolescents, elevated concentrations of circulating insulin are associated with risk factors for cardiovascular disease such as high blood pressure, dyslipidemia, inflammatory markers, endothelial dysfunction, and cardiovascular reactivity (4–12). Thus, early increases in insulin concen-

trations may promote early vascular changes that lead to early onset of cardiovascular disease. Furthermore, because high concentrations of insulin may serve as a surrogate marker for insulin resistance (13), a step on the road to diabetes, changes in circulating concentrations of insulin may serve as early indications of future trends in diabetes incidence and prevalence.

These considerations suggest that surveillance of circulating concentrations of insulin may be of critical importance during a time in world history when increases in the prevalence of obesity are occurring in many countries and are expected to continue (14). Such information may be useful in anticipating which particular population groups may be at increased risk for developing adverse events and may help target resources for intervention programs. In general, however, such data are not available in most countries, and even more rarely are such data available from nationally representative samples of the population of country, including the U.S. Our objective was to characterize serum concentrations of insulin for adolescents and young adults in the U.S.

## RESEARCH DESIGN AND METHODS

We examined data from National Health and Nutrition Examination Surveys (NHANES) for 1999–2002 for participants aged 12–19 years. Detailed information about the methods and procedures of this survey is available elsewhere (15,16). In brief, representative samples of the noninstitutionalized civilian U.S. population were selected through a stratified multistage design. Trained interviewers, using a computer-assisted personal interview system, interviewed participants at home. Participants were asked to attend the mobile examination center, where they completed additional questionnaires, underwent various examinations, and provided a blood sample. The study received human subjects approval from the Centers for Disease Control and Prevention.

From the <sup>1</sup>Division of Adult and Community Health, National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention, Atlanta, Georgia; the <sup>2</sup>Division of Diabetes Translation, National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention, Atlanta, Georgia; and the <sup>3</sup>Strong Children's Research Center, Department of Pediatrics, University of Rochester School of Medicine and Dentistry, Rochester, New York.

Address correspondence and reprint requests to Earl Ford, MD, MPH, Centers for Disease Control and Prevention, 4770 Buford Hwy., MS K66, Atlanta, GA 30341. E-mail: eford@cdc.gov.

Received for publication 25 May 2006 and accepted in revised form 17 August 2006.

**Abbreviations:** NHANES, National Health and Nutrition Examination Survey.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

DOI: 10.2337/dc06-1083


© 2006 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Table 1—Unadjusted mean and geometric mean concentrations of serum insulin\* in 1,791 participants aged 12–19 years, National Health and Nutrition Examination Survey 1999–2002

	n	Serum insulin (pmol/l)		
		Mean $\pm$ SE	Geometric mean $\pm$ SE	Range
Aged 12–19 years				
Males				
Whites	254	70.63 $\pm$ 3.11	60.75 $\pm$ 2.12	10.62–358.32
African Americans	282	75.20 $\pm$ 2.97	65.22 $\pm$ 2.43	15.60–283.38
Mexican Americans	331	83.42 $\pm$ 3.63	70.06 $\pm$ 2.39	17.76–478.32
BMI percentile				
<85th	602	55.40 $\pm$ 1.15	51.34 $\pm$ 1.04	10.62–277.02
85th–<95th	142	77.66 $\pm$ 4.01	71.06 $\pm$ 4.31	4.26–341.58
$\geq$ 95th	187	133.16 $\pm$ 7.13	115.45 $\pm$ 5.79	30.90–478.32
Females				
Whites	234	70.78 $\pm$ 3.03	63.92 $\pm$ 2.64	22.38–345.84
African Americans	225	98.47 $\pm$ 4.23	83.21 $\pm$ 2.71	23.04–473.16
Mexican Americans	335	85.21 $\pm$ 2.82	74.96 $\pm$ 2.27	22.80–450.78
BMI percentile				
<85th	564	64.50 $\pm$ 2.19	59.95 $\pm$ 2.00	22.38–286.14
85th–<95th	136	85.50 $\pm$ 3.96	80.44 $\pm$ 3.32	31.92–313.14
$\geq$ 95th	160	135.00 $\pm$ 6.58	118.98 $\pm$ 4.61	43.32–473.16
Aged 12–17 years				
Males				
Whites	707	72.93 $\pm$ 2.53	62.41 $\pm$ 1.63	4.26–478.32
African Americans	184	70.97 $\pm$ 3.56	60.92 $\pm$ 2.08	10.62–358.32
Mexican Americans	218	76.27 $\pm$ 3.44	66.58 $\pm$ 2.72	15.60–283.38
Mexican Americans	255	83.99 $\pm$ 4.14	70.71 $\pm$ 2.94	17.76–478.32
BMI percentile				
<85th	443	55.47 $\pm$ 1.41	51.40 $\pm$ 1.31	10.62–277.02
85th–<95th	116	76.14 $\pm$ 4.16	69.74 $\pm$ 4.56	4.26–341.58
$\geq$ 95th	148	135.91 $\pm$ 9.78	117.27 $\pm$ 7.85	30.90–478.32
Females				
Whites	676	78.83 $\pm$ 2.74	69.23 $\pm$ 2.41	22.38–473.16
African Americans	190	71.34 $\pm$ 3.57	63.76 $\pm$ 3.10	22.38–345.84
African Americans	180	100.73 $\pm$ 4.54	85.51 $\pm$ 2.75	23.04–473.16
Mexican Americans	257	85.63 $\pm$ 3.07	75.24 $\pm$ 2.27	22.80–450.78
BMI percentile				
<85th	447	65.30 $\pm$ 2.69	60.35 $\pm$ 2.49	22.38–286.14
85th–<95th	101	84.20 $\pm$ 4.63	78.64 $\pm$ 3.80	31.92–313.14
$\geq$ 95th	128	142.12 $\pm$ 8.00	125.16 $\pm$ 5.84	43.32–473.16
Aged 18–19 years				
Males				
Whites	224	70.92 $\pm$ 3.66	61.40 $\pm$ 2.93	21.24–395.40
Whites	70	69.73 $\pm$ 5.56	60.29 $\pm$ 4.21	21.60–243.54
African Americans	64	71.04 $\pm$ 5.31	60.22 $\pm$ 4.40	21.24–259.32
Mexican Americans	76	81.41 $\pm$ 6.07	67.78 $\pm$ 4.00	28.14–395.40
BMI percentile				
<85th	159	55.19 $\pm$ 3.01	51.15 $\pm$ 2.48	21.24–151.14
85th–<95th	26	84.42 $\pm$ 8.72	77.25 $\pm$ 8.14	22.14–294.18
$\geq$ 95th	39	125.16 $\pm$ 11.45	110.30 $\pm$ 11.27	35.76–395.40
Females				
Whites	184	74.38 $\pm$ 2.53	68.19 $\pm$ 2.20	26.46–443.34
Whites	44	68.34 $\pm$ 3.70	64.61 $\pm$ 3.38	32.10–129.00
African Americans	45	90.27 $\pm$ 9.87	75.40 $\pm$ 8.17	26.46–260.46
Mexican Americans	78	83.96 $\pm$ 6.99	74.16 $\pm$ 5.16	34.38–443.34
BMI percentile				
<85th	117	61.01 $\pm$ 2.60	58.25 $\pm$ 2.43	26.46–158.88
85th–<95th	35	88.74 $\pm$ 5.22	85.11 $\pm$ 4.91	46.62–190.32
$\geq$ 95th	32	111.42 $\pm$ 9.11	100.60 $\pm$ 8.57	52.56–443.34

\*To convert from picomoles per liter to units per milliliter, divide by 6.945.



**Figure 1**—Unadjusted geometric mean concentrations (95% CI) of serum insulin among 1,791 youth aged 12–19 years by sex and age, NHANES 1999–2002.

All insulin assays for NHANES 1999–2002 were performed in the same laboratory at the University of Missouri, Columbia, using the same method. Serum specimens were frozen at  $< -70^{\circ}\text{C}$ , shipped on dry ice, and stored at  $< -70^{\circ}\text{C}$  until analysis. The serum insulin concentration was measured using a radioimmunoassay kit (Pharmacia Diag-

nostics, Uppsala, Sweden). Quality control procedures following modified Westgard rules were used; these included both within- and between-assay quality control measures as well as monitoring of trend. The coefficient of variation of the method was 5.0–6.9% during the 4 years of the study.


We included participants aged 12–19

years who attended the morning examination and who had fasted at least 8 h. We excluded females who were pregnant, and participants with diabetes. We included age, sex, ethnicity, and BMI percentiles as covariates. We categorized age into two groups: 12–17 years and 18–19 years. This categorization was done for several reasons. First, the gap in terms of maturation between a 12 year-old and a 19 year-old is substantial. Second, the division was based on sociocultural reasons. Third, using a cut point of 18 years may facilitate comparisons of the results from the present analyses with those of other studies. Finally, having adequate sample size for analyses also played a role in determining how many age categories were feasible. For analyses stratified by ethnicity, we only included whites, African Americans, and Mexican Americans because the sample sizes for other participants were relatively small. We present means and geometric means with SEM or 95% CIs. For regression analyses, concentrations of insulin were log-transformed to improve the distribution of this variable. To account for the complex sampling design, SUDAAN version 9.0 software was used to calculate means and to perform linear regression analyses. Percentiles were calculated with SAS version

**Table 2**—Selected percentiles of serum concentrations of insulin\* among 1,791 participants aged 12–19 years, National Health and Nutrition Examination Survey 1999–2002

	n	Percentiles of serum insulin (pmol/L)								
		15	20	25	40	50	60	75	80	85
<b>Males</b>										
Age (years)										
12–13	249	33.90	40.86	42.30	50.82	56.40	64.86	81.72	89.76	98.94
14–15	199	43.74	45.60	48.66	59.40	63.00	71.94	92.88	96.66	120.30
16–17	259	38.22	40.86	47.16	54.12	59.58	65.46	78.54	84.60	88.62
18–19	224	37.08	38.82	41.34	51.36	56.58	62.40	84.72	99.30	108.90
<b>Females</b>										
Age (years)										
12–13	253	47.22	51.24	52.92	61.50	69.96	78.84	102.48	108.90	118.20
14–15	214	43.02	46.38	49.80	60.54	65.34	74.28	90.00	101.88	114.42
16–17	209	43.20	47.16	49.26	58.50	64.56	73.02	82.56	88.74	98.46
18–19	184	46.74	50.34	51.42	58.86	63.00	70.62	86.46	92.70	105.30
<b>Race or ethnicity</b>										
<b>Males</b>										
White	254	37.62	40.86	44.76	53.28	57.78	63.00	78.66	88.20	98.22
African American	282	40.02	42.90	46.32	55.38	62.58	71.64	88.74	98.28	108.60
Mexican American	331	40.74	43.14	46.92	57.78	65.64	74.22	95.28	108.48	126.24
<b>Females</b>										
White	234	42.54	46.44	49.02	57.96	62.40	69.54	84.00	87.72	101.64
African American	225	47.04	52.02	55.14	67.74	79.44	90.48	120.72	132.36	156.66
Mexican American	335	48.30	51.06	54.42	63.06	70.98	79.68	100.02	109.08	120.84

\*To convert from picomoles per liter to units per milliliter, divide by 6.945.



**Figure 2**—Geometric mean (95% CI) concentrations of insulin among 1,791 participants aged 12–19 years, by age, sex, and ethnicity, NHANES 1999–2002. A: Unadjusted results. B: Adjusted for age and BMI percentile. W, whites; AA, African Americans; MA, Mexican Americans.

8.2. Sampling weights were used to calculate measures of central tendency, percentiles, and regression estimates.

**RESULTS**— A total of 1,791 participants aged 12–19 years who attended the morning examination had a value for insulin. These included 931 male and 860 female subjects, 488 whites, 507 African Americans, 666 Mexican Americans, and 130 other participants.

Unadjusted mean and geometric mean concentrations of serum insulin are shown in Table 1. The pattern of geometric mean concentrations of insulin by age differed somewhat according to sex (Fig. 1). Among male participants, concentrations increased from age 12 to 14 years

before decreasing somewhat. Among female participants, concentrations were highest at age 13 years before decreasing steadily through age 19 years. Selected percentiles of concentrations of insulin are shown in Table 2. Among participants aged 12–17 years but not among those aged 18–19 years, females had higher log-transformed concentrations of insulin than males ( $P$ , Wald,  $F = 0.038$  and  $0.125$ , respectively) after adjusting for age and ethnicity.


In unadjusted analyses of participants aged 12–17 years, ethnic variation was present among both males ( $P$ , Wald,  $F = 0.006$ ) and females ( $P$ , Wald,  $F < 0.001$ ). No significant ethnic variation occurred among male or female participants aged

18–19 years. Among participants aged 12–17 years, mean log-transformed concentrations did not vary significantly by ethnicity among males ( $P$ , Wald,  $F = 0.199$ ) but did so among females ( $P$ , Wald,  $F = 0.021$ ) after adjusting for age and BMI percentile in linear regression analysis (Fig. 2). African-American females had higher concentrations of insulin than white or Mexican-American females. Among participants aged 18–19 years, no significant ethnic differences were present among male and female participants. BMI was strongly associated with concentrations of insulin among both male and female participants (Fig. 3).

**CONCLUSIONS**— These data represent the first nationally representative information about the distributions of fasting insulin concentrations among youth in the U.S., and they show significant demographic variation. Concentrations were higher among females than males. Significant ethnic variation was present among female participants aged 12–17 years but not among those aged 18–19 years.

In many studies conducted in the U.S., African-American children and adolescents were found to have higher concentrations of insulin than white children (4,6,17–26). Among 2,996 Bogalusa Study participants aged 5–17 years, the difference in median concentrations of insulin between whites and African Americans was larger among females than males (3). However, in one study, the ethnic difference was eliminated after adjustment for fat mass quantified using dual energy X-ray absorptiometry (26). Factors that may account for the difference between African-American and white children and adolescents include lower insulin clearance, lower insulin sensitivity, and higher insulin secretion (24–28). Given the high prevalence of diabetes among African Americans and Mexican Americans compared with whites in the U.S. and the high mortality rates from coronary heart disease among African Americans compared with whites, the high concentrations of insulin among these groups, especially females, underscore the fact that the roots for these diseases can often be traced back to childhood. Our findings also underscore the wisdom of trying to prevent the development of disease and risk factors at an early age.

Fewer studies have compared concentrations of insulin among Hispanic



**Figure 3**—Unadjusted geometric mean (95% CI) concentrations of insulin among 1,791 participants aged 12–19 years by percentiles of BMI, NHANES 1999–2002.

and white children and adolescents. In a study of 403 third graders in Corpus Christi, Texas, median concentrations were higher in Mexican Americans than in whites (29). Among children from the Los Angeles area, concentrations of insulin were higher among 28 Hispanic children than 14 among Caucasian children (25). In a study of 491 Hispanic children aged 2–3 years from New York City, concentrations of insulin were higher in girls than in boys (11). In the San Diego Study of Children's Activity and Nutrition, concentrations of insulin were higher in 54 Mexican American children than in 29 white children at age 11 years (30). Although the median concentration of insulin was still higher in Mexican American children than in white children, 6 years later the difference was not significant.

Most studies conducted in U.S. youth have also shown that concentrations of insulin are higher in females than in males (31). Among 90 Pima Indian children aged 5 and 10 years, concentrations of insulin were similar in boys and girls aged 5 years but were higher in girls than in boys at the age of 10 years (32). In addition, many studies conducted in other countries have also reported higher circulating concentrations of insulin in females than in males (33–38). However, in some studies no sex differences were found (38,39).

Our results showing a large increase in fasting concentrations of insulin in those who were overweight compared with those who were not reiterate some of the negative physiological consequences of excess weight. In addition, the significant increase in concentrations of insulin among those who were at risk for being overweight shows that even this degree of excess weight produces unfavorable physiological changes.

A limitation of our study is the lack of data on pubertal status and serum androgens, estrogens, or growth hormone. Such information would be helpful to describe the difference in distribution of insulin concentrations among teens at different pubertal stages, when elevated values may reflect either a normal physiological state or a pathological one. For now, our results suggest that fasting serum insulin concentrations peak earlier in girls than in boys. The lack of insulin values for subjects <12 years of age, when some youth, mostly girls, may enter early pubertal development, precludes our ability to comment on the distribution of insulin among preadolescent children. Previous NHANES surveys did not collect serum insulin on adolescents, so comparisons to a group of U.S. teens that were less overweight than this sample are not possible.

Although the relative merit of surrogate

markers for insulin resistance continues to be researched, fasting concentrations of insulin correlate about as well with measures of insulin resistance determined by gold standard methods as other surrogate markers. In four studies, correlation coefficients between fasting concentrations of insulin and measures of insulin sensitivity/resistance determined by gold standard methods ranged from 0.4 to 0.91 (28,40–42). In three of these studies, the correlation coefficients ranged from 0.8 to 0.91 (28,41,42).

In our analysis, we examined fasting serum concentrations of insulin, one facet of a complex physiological picture. Glucose loading or meals elicit a greater insulin response in African Americans than in whites (17,43). Unfortunately, measures of stimulated insulin response were not available in the NHANES data set.

In summary, fasting concentrations of insulin showed significant demographic differences. Geometric mean concentrations were higher in females than in males and peaked at age 14 years in males and 13 years in females. The only ethnic or racial differences occurred in females aged 12–17 years among whom geometric mean concentrations were higher in African Americans than in whites or Mexican Americans. The increases in obesity among U.S. youth, the strong links between obesity and hyperinsulinemia and insulin resistance, and the effects of hyperinsulinemia and insulin resistance on cardiovascular disease and diabetes suggest that surveillance of concentrations of insulin in the population, particularly in children and adolescents, may be useful to assess early effects of obesity on health and to allow international comparisons. For example, a report on the distribution of serum insulin in Canadian youth collected in 1999 showed median values of plasma insulin to be lower than those in comparable groups of U.S. youth (35). The values reported for 13-year-old boys were lower than those for the 12- to 13-year-old boys in this study (41.9 vs. 56.4 pmol/l), and values for the 16-year-old boys were lower than those for the 16- to 17-year-old boys (38.7 vs. 59.58 pmol/l). These differences may be attributable to differences in race or ethnicity or distribution of obesity. Assay variability may be a significant reason for these differences, highlighting the need for achieving standardization in insulin measurements among laboratories (44) to facilitate surveillance.

## References

- Hedley AA, Ogden CL, Johnson CL, Carroll MD, Curtin LR, Flegal KM: Prevalence of overweight and obesity among US children, adolescents, and adults, 1999–2002. *JAMA* 291:2847–2850, 2004
- Ford ES, Mokdad AH, Ajani UA: Trends in risk factors for cardiovascular disease among children and adolescents in the United States. *Pediatrics* 114:1534–1544, 2004
- Freedman DS, Dietz WH, Srinivasan SR, Berenson GS: The relation of overweight to cardiovascular risk factors among children and adolescents: the Bogalusa Heart Study. *Pediatrics* 103:1175–1182, 1999
- Burke GL, Webber LS, Srinivasan SR, Radhakrishnamurthy B, Freedman DS, Berenson GS: Fasting plasma glucose and insulin levels and their relationship to cardiovascular risk factors in children: Bogalusa Heart Study. *Metabolism* 35:441–446, 1986
- Ronnemaa T, Knip M, Lautala P, Viikari J, Uhari M, Leino A, Kaprio EA, Salo MK, Dahl M, Nuutinen EM, et al: Serum insulin and other cardiovascular risk indicators in children, adolescents and young adults. *Ann Med* 23:67–72, 1991
- Gutin B, Islam S, Manos T, Cucuzzo N, Smith C, Stachura ME: Relation of percentage of body fat and maximal aerobic capacity to risk factors for atherosclerosis and diabetes in black and white seven- to eleven-year-old children. *J Pediatr* 125:847–852, 1994
- Bao W, Srinivasan SR, Berenson GS: Persistent elevation of plasma insulin levels is associated with increased cardiovascular risk in children and young adults: the Bogalusa Heart Study. *Circulation* 93:54–59, 1996
- Moussa MA, Shaltout AA, Nkansa-Dwamena D, Mourad M, Al-Sheikh N, Agha N, Galal DO: Association of fasting insulin with serum lipids and blood pressure in Kuwaiti children. *Metabolism* 47:420–424, 1998
- Reaven P, Nader PR, Berry C, Hoy T: Cardiovascular disease insulin risk in Mexican-American and Anglo-American children and mothers. *Pediatrics* 101:E12, 1998
- Ford ES, Giles WH, Myers GL, Rifai N, Ridker PM, Mannino DM: C-reactive protein concentration distribution among US children and young adults: findings from the National Health and Nutrition Examination Survey, 1999–2000. *Clin Chem* 49:1353–1357, 2003
- Shea S, Aymong E, Zybert P, Berglund L, Shamoon H, Deckelbaum RJ, Basch CE: Fasting plasma insulin modulates lipid levels and particle sizes in 2- to 3-year-old children. *Obes Res* 11:709–721, 2003
- Steinberger J, Daniels SR, American Heart Association Atherosclerosis, Hypertension, and Obesity in the Young Committee (Council on Cardiovascular Disease in the Young), American Heart Association Diabetes Committee (Council on Nutrition, Physical Activity, and Metabolism): Obesity, insulin resistance, diabetes, and cardiovascular risk in children: an American Heart Association scientific statement from the Atherosclerosis, Hypertension, and Obesity in the Young Committee (Council on Cardiovascular Disease in the Young) and the Diabetes Committee (Council on Nutrition, Physical Activity, and Metabolism). *Circulation* 107:1448–1453, 2003
- Monzillo LU, Hamdy O: Evaluation of insulin sensitivity in clinical practice and in research settings. *Nutr Rev* 61:397–412, 2003
- Wang Y, Lobstein T: Worldwide trends in childhood overweight and obesity. *Int J Pediatr Obes* 1:111–125, 2006
- Centers for Disease Control and Prevention: NHANES 1999–2000 [article online], 2006. Available from [http://www.cdc.gov/nchs/about/major/nhanes/nhanes99\\_00.htm](http://www.cdc.gov/nchs/about/major/nhanes/nhanes99_00.htm). Accessed 15 March 2006
- Centers for Disease Control and Prevention: NHANES 2001–2002 [article online], 2006. Available from <http://www.cdc.gov/nchs/about/major/nhanes/nhanes01-02.htm>. Accessed 15 March 2006
- Freedman DS, Srinivasan SR, Burke GL, Shear CL, Smoak CG, Harsha DW, Webber LS, Berenson GS: Relation of body fat distribution to hyperinsulinemia in children and adolescents: the Bogalusa Heart Study. *Am J Clin Nutr* 46:403–410, 1987
- Svec F, Nastasi K, Hilton C, Bao W, Srinivasan SR, Berenson GS: Black-white contrasts in insulin levels during pubertal development: the Bogalusa Heart Study. *Diabetes* 41:313–317, 1992
- Arslanian S, Suprasongsin C: Differences in the in vivo insulin secretion and sensitivity of healthy black versus white adolescents. *J Pediatr* 129:440–443, 1996
- Arslanian S, Suprasongsin C, Janosky JE: Insulin secretion and sensitivity in black versus white prepubertal healthy children. *J Clin Endocrinol Metab* 82:1923–1927, 1997
- Arslanian S: Insulin secretion and sensitivity in healthy African-American vs American white children. *Clin Pediatr (Phila)* 37:81–88, 1998
- Schuster DP, Kien CL, Osei K: Differential impact of obesity on glucose metabolism in black and white American adolescents. *Am J Med Sci* 316:361–367, 1998
- Gower BA, Nagy TR, Goran MI: Visceral fat, insulin sensitivity, and lipids in prepubertal children. *Diabetes* 48:1515–1521, 1999
- Arslanian SA, Saad R, Lewy V, Danadian K, Janosky J: Hyperinsulinemia in African-American children: decreased insulin clearance and increased insulin secretion and its relationship to insulin sensitivity. *Diabetes* 51:3014–3019, 2002
- Goran MI, Bergman RN, Cruz ML, Watanabe R: Insulin resistance and associated compensatory responses in African-American and Hispanic children. *Diabetes Care* 25:2184–2190, 2002
- Uwaifo GI, Nguyen TT, Keil MF, Russell DL, Nicholson JC, Bonat SH, McDuffie JR, Yanovski JA: Differences in insulin secretion and sensitivity of Caucasian and African American prepubertal children. *J Pediatr* 140:673–680, 2002
- Gower BA, Granger WM, Franklin F, Shewchuk RM, Goran MI: Contribution of insulin secretion and clearance to glucose-induced insulin concentration in African-American and Caucasian children. *J Clin Endocrinol Metab* 87:2218–2224, 2002
- Huang TT, Johnson MS, Goran MI: Development of a prediction equation for insulin sensitivity from anthropometry and fasting insulin in prepubertal and early pubertal children. *Diabetes Care* 25:1203–1210, 2002
- Batey LS, Goff DC Jr, Tortolero SR, Nichaman MZ, Chan W, Chan FA, Grunbaum J, Hanis CL, Labarthe DR: Summary measures of the insulin resistance syndrome are adverse among Mexican-American versus non-Hispanic white children: the Corpus Christi Child Heart Study. *Circulation* 96:4319–4325, 1997
- Reaven PD, Traustadottir T, Brennan J, Nader PR: Cardiovascular risk factors associated with insulin resistance in children persist into late adolescence. *Diabetes Care* 28:148–150, 2005
- Young-Hyman D, Schlundt DG, Herman L, De Luca F, Counts D: Evaluation of the insulin resistance syndrome in 5- to 10-year-old overweight/obese African-American children. *Diabetes Care* 24:1359–1364, 2001
- Bunt JC, Salbe AD, Harper IT, Hanson RL, Tataranni PA: Weight, adiposity, and physical activity as determinants of an insulin sensitivity index in Pima Indian children. *Diabetes Care* 26:2524–2530, 2003
- Lautala P, Akerblom HK, Viikari J, Louhivuori K, Uhari M, Dahlstrom S, Dahl M, Lahde PL, Pesonen E, Pietikainen M, et al: Atherosclerosis precursors in Finnish children and adolescents. VII. Serum immunoreactive insulin. *Acta Paediatr Scand Suppl* 318:127–133, 1985
- Matsui I, Nambu S, Baba S: Evaluation of fasting serum insulin levels among Japanese school-age children. *J Nutr Sci Vitaminol (Tokyo)* 44:819–828, 1998
- Allard P, Delvin EE, Paradis G, Hanley JA, O'Loughlin J, Lavallee C, Levy E, Lambert M: Distribution of fasting plasma insulin, free fatty acids, and glucose concentrations

- and of homeostasis model assessment of insulin resistance in a representative sample of Quebec children and adolescents. *Clin Chem* 49:644–649, 2003
36. Misra A, Vikram NK, Arya S, Pandey RM, Dhingra V, Chatterjee A, Dwivedi M, Sharma R, Luthra K, Guleria R, Talwar KK: High prevalence of insulin resistance in postpubertal Asian Indian children is associated with adverse truncal body fat patterning, abdominal adiposity and excess body fat. *Int J Obes Relat Metab Disord* 28:1217–1226, 2004
  37. Garces C, Cano B, Granizo JJ, Benavente M, Viturro E, Gutierrez-Guisado J, de Oya I, Lasuncion MA, de Oya M: Insulin and HOMA in Spanish prepubertal children: relationship with lipid profile. *Clin Biochem* 38:920–924, 2005
  38. Hsieh AT, Chu NF, Shen MH, Wu DM, Wang DJ, Hsieh SM: Insulin, proinsulin and insulin resistance status in relation to lipid profiles among school children in Taiwan-The Taipei Children Heart Study. *Clin Biochem* 36:367–372, 2003
  39. Ehtisham S, Crabtree N, Clark P, Shaw N, Barrett T: Ethnic differences in insulin resistance and body composition in United Kingdom adolescents. *J Clin Endocrinol Metab* 90:3963–3969, 2005
  40. Cutfield WS, Jefferies CA, Jackson WE, Robinson EM, Hofman PL: Evaluation of HOMA and QUICKI as measures of insulin sensitivity in prepubertal children. *Pediatr Diabetes* 4:119–125, 2003
  41. Gungor N, Saad R, Janosky J, Arslanian S: Validation of surrogate estimates of insulin sensitivity and insulin secretion in children and adolescents. *J Pediatr* 144:47–55, 2004
  42. Conwell LS, Trost SG, Brown WJ, Batch JA: Indexes of insulin resistance and secretion in obese children and adolescents: a validation study. *Diabetes Care*; 27:314–319, 2004
  43. Osei K, Schuster DP: Ethnic differences in secretion, sensitivity, and hepatic extraction of insulin in black and white Americans. *Diabet Med* 11:755–762, 1994
  44. Robbins DC, Andersen L, Bowsher R, Chance R, Dinesen B, Frank B, Gingerich R, Goldstein D, Widemeyer HM, Haffner S, Hales CN, Jarett L, Polonsky K, Porte D, Skyler J, Webb G, Gallagher K: Report of the American Diabetes Association's Task Force on standardization of the insulin assay. *Diabetes* 45:242–256, 1996