

# Targeting Glucose in Acute Myocardial Infarction

## Has glucose, insulin, and potassium infusion missed the target?

AJAY CHAUDHURI, MD<sup>1</sup>  
MICHAEL MILLER, PHD<sup>2</sup>  
RICHARD NESTO, MD<sup>3</sup>

NOAH ROSENBERG, MD<sup>2</sup>  
PARESH DANDONA, MD, PHD<sup>1</sup>

**P**roinflammatory mechanisms may contribute to hyperglycemia-associated adverse outcomes in acute myocardial infarction (AMI) (1–3). Insulin exerts anti-inflammatory effects in ST elevation myocardial infarction and coronary artery bypass graft patients (4–6). Inflammation plays an important role in the pathogenesis of atherosclerosis and thrombosis (7,8). Thus, insulin infusion in AMI should be beneficial. However, glucose, insulin, and potassium (GIK) infusion was neutral in its benefit in the Clinical Trial of Metabolic Modulation in Acute Myocardial Infarction Treatment Evaluation—Estudios Cardiológicos Latinoamericana (CREATE-ECLA) study (9).

The GIK regimen (fixed combination of 1 l of 25% dextrose with 50 units regular insulin and 80 meq/l potassium) used in CREATE-ECLA is known to lower serum free fatty acid concentrations (10) and to reduce mortality by 18% in AMI, as previously reviewed (11). However, the infusion of 30 g/h glucose without titration of glucose or insulin in CREATE-ECLA led to a significant increase in blood glucose concentrations. Mehta et al. suggested that the adverse effect of GIK-induced hyperglycemia may have neutralized any potential benefit of insulin in the GIK regimen. However, Mehta

et al. did not comment on the magnitude of the potential contribution of hyperglycemia to neutralize benefits of GIK. Since admission hyperglycemia was predictive of mortality in AMI in CREATE-ECLA, we have estimated the potential effects of GIK-induced hyperglycemia on mortality in this study.

### RESEARCH DESIGN AND METHODS

Based on the admission blood glucose–related mortality rates in control subjects in CREATE-ECLA, we have constructed a model: 30-day % mortality =  $100\% \times [1 - c \times \exp(-d \times BG)]$  predicting 30-day mortality as a function of admission blood glucose, where  $\exp$  = exponential function and BG = blood glucose. The constants  $c$  and  $d$  were estimated using a regression involving 3 points from this constructed curve. This model was then applied to the blood glucose at admission and at 6 and 24 h for both the GIK and control groups, assuming that the relationship between glycemia and mortality is maintained even after admission in AMI. This assumption is probably valid because for every 0.6-mmol/l reduction in glucose postadmission, there is an 8% reduction in mortality in patients with AMI (12) and because

glucose levels after admission predict mortality in AMI (13,14).

The projected mortality at 0, 6, and 24 h (Table 1) yielded trapezoids of which the areas were calculated. After dividing by 24 h, the following weighted average formula for mortality was obtained: % mortality (average) =  $(0.125) \times \% \text{ mortality at 0 h} + (0.5) \times \% \text{ mortality at 6 h} + (0.375) \times \% \text{ mortality at 24 h}$ .

**RESULTS**— In our model, the estimated 30-day mortality for control subjects based on blood glucose achieved during 24 h is 9.9%, which is similar to the observed mortality of 9.7% for the control subjects in CREATE-ECLA. However, the estimated mortality rate for the GIK group on the basis of the GIK-induced hyperglycemia during 24 h was 12.2%, which was 2.2% higher than the observed mortality of 10% for the GIK group.

**CONCLUSIONS**— We suggest that the insulin in the GIK infusion used in CREATE-ECLA might have neutralized the 2.2% (12.2% [expected] – 10% [observed]) increase in mortality that should have been observed in the GIK group because of the effect of hyperglycemia induced by this infusion. Thus, if hyperglycemia was not induced by the GIK infusion used in CREATE-ECLA, the administration of insulin in this trial could have resulted in a 2.2% absolute and a 22% relative reduction in mortality in the GIK group. Indeed, CREATE-ECLA investigators have recently reported an excess mortality and congestive cardiac failure in the GIK group in the first 3 days, when GIK-induced hyperglycemia was present and probably had its maximal effect. In contrast, there was a reduction in mortality and congestive cardiac failure between 4 and 30 days (15), when glucose levels had probably approximated the levels in the control group. In a canine model of AMI, low-dose insulin alone reduced the infarct size, while glucose and potassium (16) caused hyperglycemia and increased infarct size. These find-

From the <sup>1</sup>Department of Endocrinology, State University at New York at Buffalo and Kaleida Health, Buffalo, New York; <sup>2</sup>Sanofi-Aventis Pharmaceuticals, Bridgewater, New Jersey; and the <sup>3</sup>Department of Cardiovascular Medicine, Lahey Clinic and Harvard Medical School, Boston, Massachusetts.

Address correspondence and reprint requests to Paresh Dandona, MD, PHD, Diabetes-Endocrinology Center of WNY, 3 Gates Cir., Buffalo, NY 14209. E-mail: pdandona@kaleidahealth.org.

Received for publication 28 May 2007 and accepted in revised form 4 September 2007.

Published ahead of print at <http://care.diabetesjournals.org> on 11 September 2007. DOI: 10.2337/dc07-1220.

P.D. is currently affiliated with the Diabetes-Endocrinology Center of Western New York, Buffalo, New York.

**Abbreviations:** AMI, acute myocardial infarction; CREATE-ECLA, Clinical Trial of Metabolic Modulation in Acute Myocardial Infarction Treatment Evaluation—Estudios Cardiológicos Latinoamericana; GIK, glucose, insulin, and potassium.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

© 2007 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

**Table 1—Hyperglycemia-related mortality in CREATE-ECLA based on relationship of admission glucose to mortality in control subjects**

Time	Control group		GIK group	
	Blood glucose (mmol/l)	Mortality	Blood glucose (mmol/l)	Mortality
0	9	11.4	9	11.4
6 h	8.2	10.2	10.4	13.4
24 h	7.5	9.1	8.6	10.8

Data are percentages unless otherwise indicated.

ings and the proinflammatory, prothrombotic effects of hyperglycemia may explain how GIK-induced hyperglycemia can neutralize the potential benefits of insulin (2).

A limitation of our study is that our analysis is based on the published data in CREATE-ECLA. Because of the absence of detailed data and the derivational nature of our methodology, which arrives at 12.2% expected mortality, we are not in a position to provide a *P* value or SEs. However, based on the level of the expected precision in a large study like CREATE-ECLA, a potential absolute reduction in mortality of 2.2% or relative reduction of 22% would be statistically significant and well outside the confidence bounds (95% CI 9.4–10.6) of the observed mortality of 10% for the GIK. Although our model accurately predicted the death rate for the control subjects, it is possible that we could have overestimated the death rate for the GIK patients. We have also speculated that the adverse effects of the reactive hyperglycemia observed following AMI are equivalent to the iatrogenic hyperglycemia induced by the GIK infusion. Iatrogenic hyperglycemia is known to induce proinflammatory cytokines and endothelial dysfunction (17,18). These mechanisms may be responsible for the adverse cardiovascular outcomes associated with hyperglycemia. The important point for discussion is not whether the model is fundamentally wrong but whether, if these assumptions are “essentially correct,” is there then a factor (insulin) that protected the GIK patients from the toxic effects of hyperglycemia? Using the model based on the observations in CREATE-ECLA, the answer is probably in the affirmative. Thus, hyperglycemia needs to be avoided when designing studies investigating whether insulin administration is beneficial in AMI.

We have now designed a trial to test the hypothesis that insulin is cardioprotective in AMI because of its anti-

inflammatory, profibrinolytic, antioxidant, antiapoptotic, vasodilatory, and antiaggregatory actions and that these effects are enhanced by lowering glucose into the normoglycemic range. We are using intravenous insulin infusion to lower glucose to 90–130 mg/dl in ST elevation myocardial infarction patients. To allow us to infuse a minimal dose of 2.5 units/h (the anti-inflammatory dose of insulin) in this trial, ~7 g/h dextrose appropriately titrated will be infused simultaneously to maintain euglycemia (R.N., P.D., personal communication). With such an insulin infusion regimen, the anti-inflammatory and potentially cardioprotective effect of insulin is likely to be observed.

## References

1. The ACE/ADA Task Force on Inpatient Diabetes: American College of Endocrinology and American Diabetes Association consensus statement on inpatient diabetes and glycemic control: a call to action. *Diabetes Care* 29:1955–1962, 2006
2. Dandona P, Mohanty P, Chaudhuri A, Garg R, Aljada A: Insulin infusion in acute illness. *J Clin Invest* 115:2069–2072, 2005
3. Vanhorebeek I, Langouche L, Van den Berghe G: Glycemic and nonglycemic effects of insulin: how do they contribute to a better outcome of critical illness? *Curr Opin Crit Care* 11:304–311, 2005
4. Chaudhuri A, Janicke D, Wilson MF, Tripathy D, Garg R, Bandyopadhyay A, Caleri J, Hoffmeyer D, Tufail S, Ghanim H, Aljada A, Dandona P: Anti-inflammatory and pro-fibrinolytic effect of insulin in acute ST-elevation myocardial infarction. *Circulation* 109:849–854, 2004
5. Visser L, Zuurbier CJ, Hoek FJ, Opmeer BC, de Jonge E, de Mol BA, van Wezel HB: Glucose, insulin and potassium applied as perioperative hyperinsulinaemic normoglycaemic clamp: effects on inflammatory response during coronary artery surgery. *Br J Anaesth* 95:448–457, 2005
6. Dandona P, Aljada A, Mohanty P, Ghanim H, Hamouda W, Assian E, Ahmad S: Insulin inhibits intranuclear nuclear factor

kappaB and stimulates IkappaB in mononuclear cells in obese subjects: evidence for an anti-inflammatory effect? *J Clin Endocrinol Metab* 86:3257–3265, 2001

7. Libby P, Simon DI: Inflammation and thrombosis: the clot thickens. *Circulation* 103:1718–1720, 2001
8. Ross R: Atherosclerosis: an inflammatory disease. *N Engl J Med* 340:115–126, 1999
9. Mehta SR, Yusuf S, Diaz R, Zhu J, Pais P, Xavier D, Paolasso E, Ahmed R, Xie C, Kazmi K, Tai J, Orlandini A, Pogue J, Liu L: Effect of glucose-insulin-potassium infusion on mortality in patients with acute ST-segment elevation myocardial infarction: CREATE-ECLA randomized controlled trial. *JAMA* 293:437–446, 2005
10. Rackley CE, Russell RO Jr, Rogers WJ, Mantle JA, McDaniel HG, Papapietro SE: Clinical experience with glucose-insulin-potassium therapy in acute myocardial infarction. *Am Heart J* 102:1038–1049, 1981
11. Yusuf S, Mehta SR, Diaz R, Paolasso E, Pais P, Xavier D, Xie C, Ahmed RJ, Khazmi K, Zhu J, Liu L: Challenges in the conduct of large simple trials of important generic questions in resource-poor settings: CREATE and ECLA trial program evaluating GIK (glucose, insulin and potassium) and low-molecular-weight heparin in acute myocardial infarction. *Am Heart J* 148:1068–1078, 2004
12. Goyal A, Mahaffey KW, Garg J, Nicolau JC, Hochman JS, Weaver WD, Theroux P, Oliveira GB, Todaro TG, Mojcik CF, Armstrong PW, Granger CB: Prognostic significance of the change in glucose level in the first 24 h after acute myocardial infarction: results from the CARDINAL study. *Eur Heart J* 27:1289–1297, 2006
13. Cheung NW, Wong VW, McLean M: The Hyperglycemia: Intensive Insulin Infusion in Infarction (HI-5) Study: a randomized controlled trial of insulin infusion therapy for myocardial infarction. *Diabetes Care* 29:765–770, 2006
14. van der Horst IC, Nijsten MW, Vogelzang M, Zijlstra F: Persistent hyperglycemia is an independent predictor of outcome in acute myocardial infarction (Letter). *Cardiovasc Diabetol* 6:2, 2007
15. Diaz R: Glucose-insulin-potassium in STEMI patients: results of OASIS-6 study and a combined analysis with CREATE ECLA trial. Paper presented at the European Society of Cardiology Scientific Congress, 3 September 2006, Fira Gran Via M2, Barcelona, Spain
16. Zhang HX, Zang YM, Huo JH, Liang SJ, Zhang HF, Wang YM, Fan Q, Guo WY, Wang HC, Gao F: Physiologically tolerable insulin reduces myocardial injury and improves cardiac functional recovery in myocardial ischemic/reperfused dogs.

- J Cardiovasc Pharmacol* 48:306–313, 2006
17. Esposito K, Nappo F, Marfella R, Giugliano G, Giugliano F, Ciotola M, Quagliaro L, Ceriello A, Giugliano D: Inflammatory cytokine concentrations are acutely increased by hyperglycemia in humans: role of oxidative stress. *Circulation* 106:2067–2072, 2002
  18. Srinivasan M, Herrero P, McGill JB, Ben-  
nik J, Heere B, Lesniak D, Davila-Roman  
VG, Gropler RJ: The effects of plasma  
insulin and glucose on myocardial  
blood flow in patients with type 1 diabe-  
tes mellitus. *J Am Coll Cardiol* 46:42–48,  
2005