

Efficacy and Safety of the Dipeptidyl Peptidase-4 Inhibitor Alogliptin in Patients With Type 2 Diabetes and Inadequate Glycemic Control

A randomized, double-blind, placebo-controlled study

RALPH A. DEFONZO, MD¹
PENNY R. FLECK, MT²
CRAIG A. WILSON, PHD²

QAIS MEKKI, MD, PHD²
ON BEHALF OF THE ALOGLIPTIN STUDY 010
GROUP*

OBJECTIVE — To evaluate the dipeptidyl peptidase-4 (DPP-4) inhibitor alogliptin in drug-naïve patients with inadequately controlled type 2 diabetes.

RESEARCH DESIGN AND METHODS — This double-blind, placebo-controlled, multicenter study included 329 patients with poorly controlled diabetes randomized to once-daily treatment with 12.5 mg alogliptin ($n = 133$), 25 mg alogliptin ($n = 131$), or placebo ($n = 65$) for 26 weeks. Primary efficacy end point was mean change from baseline in A1C at the final visit.

RESULTS — At week 26, mean change in A1C was significantly greater ($P < 0.001$) for 12.5 mg (-0.56%) and 25 mg (-0.59%) alogliptin than placebo (-0.02%). Reductions in fasting plasma glucose were also greater ($P < 0.001$) in alogliptin-treated patients than in those receiving placebo. Overall, incidences of adverse events (67.4–70.3%) and hypoglycemia (1.5–3.0%) were similar across treatment groups.

CONCLUSIONS — Alogliptin monotherapy was well tolerated and significantly improved glycemic control in patients with type 2 diabetes, without raising the incidence of hypoglycemia.

Diabetes Care 31:2315–2317, 2008

Inhibition of dipeptidyl peptidase-4 (DPP-4) increases the concentration of glucagon-like peptide-1, an incretin hormone that stimulates glucose-dependent insulin release, suppresses glucagon production, slows gastric emptying, reduces appetite, and may promote preservation of β -cell function in patients with type 2 diabetes (1). Alogliptin is a novel, high-affinity, high-specificity DPP-4 inhibitor that produces rapid and sustained DPP-4 inhibition and significantly reduces postprandial plasma glucose concentrations in patients with type

2 diabetes (2,3). A phase-three study was conducted to evaluate the efficacy and safety of alogliptin in adults with type 2 diabetes that was inadequately controlled with diet and exercise.

RESEARCH DESIGN AND METHODS — Eligible patients were treatment-naïve (i.e., no current antidiabetes therapy and <7 days of therapy in the past 3 months) men and women, aged 18–80 years, with type 2 diabetes. Key inclusion criteria included A1C 7–10%, BMI 23–45 kg/m², treatment with diet

and exercise for ≥ 1 month, and systolic/diastolic blood pressure $\leq 180/\leq 110$ mmHg. Patients received counseling on diet and exercise. Patients who completed a 4-week, single-blind run-in period with fasting plasma glucose (FPG) < 275 mg/ml (15.27 mmol/l) and $\geq 75\%$ compliance (by tablet count) were randomized (2:2:1) to 26 weeks of double-blind treatment with 12.5 mg alogliptin, 25 mg alogliptin, or placebo taken once daily before the first meal. Additional antidiabetes agents were prohibited.

Efficacy assessments included all randomized patients who received the double-blind study drug. The primary end point was mean change from baseline in A1C at week 26. Other efficacy measures included changes in FPG, clinical response rates, incidences of marked hyperglycemia (FPG ≥ 200 mg/dl [11.10 mmol/l]) and hyperglycemic rescue, and changes in body weight. Exploratory end points included changes in measures of pancreatic function (fasting insulin, fasting proinsulin, and homeostasis model assessment of β -cell function) and lipid profiles. Treatment group differences for the primary end point were analyzed through ANCOVA, with treatment and geographic region as variables and baseline A1C and diabetes duration as covariates. The last observation carried forward method was used for imputing missing data; testing was two-sided at a significance level of 0.05. Continuous secondary efficacy analyses were performed as for the primary analysis, except that the baseline covariate corresponds with the tested end point. Incidence variables were compared using nonparametric, extended Mantel-Haenszel X^2 tests; covariates were the same as for the primary end point.

The safety population included patients who took at least one dose of the study drug (double blind). Safety assessments included adverse events, clinical laboratory findings, 12-lead electrocar-

From the ¹University of Texas Health Science Center at San Antonio, San Antonio, Texas; and the ²Takeda Global Research & Development Center, Deerfield, Illinois.

Corresponding author: Ralph A. DeFronzo, albarado@uthscsa.edu.

Received 5 June 2008 and accepted 13 September 2008.

Published ahead of print at <http://care.diabetesjournals.org> on 22 September 2008. DOI: 10.2337/dc08-1035. Clinical trial reg. no. NCT00286455, clinicaltrials.gov.

*A full list of investigators of the Alogliptin Study 010 Group is available in an online appendix at <http://dx.doi.org/10.2337/dc08-1035>.

© 2008 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Table 1—Results of secondary efficacy end points and exploratory assessments

Parameter	Placebo	12.5 mg Alogliptin	P (vs. placebo)	25 mg Alogliptin	P (vs. placebo)
n	64	133	—	131	—
A1C (%)					
≤6.5%	7 (10.9)	23 (17.3)	0.818	27 (20.6)	0.294
≤7.0%	15 (23.4)	63 (47.4)	0.001	58 (44.3)	0.008
Reduction ≥0.5%	19 (29.7)	67 (50.4)	0.005	72 (55.0)	<0.001
Reduction ≥1.0%	7 (10.9)	38 (28.6)	0.003	39 (29.8)	<0.001
FPG	11.3 ± 5.24	−10.3 ± 3.6	<0.001	−16.4 ± 3.7	<0.001
Hyperglycemic rescue	19 (29.7)	13 (9.8)	0.001	10 (7.6)	<0.001
Marked hyperglycemia	30 (46.9)	44 (33.1)	0.110	33 (25.4)	0.005
Body weight (kg)	0.18 ± 0.37	−0.09 ± 0.26	0.539	−0.22 ± 0.26	0.379
Proinsulin-to-insulin ratio	0.046 ± 0.022	−0.040 ± 0.015	0.001	−0.038 ± 0.015	0.002
HOMA-B	−0.26 ± 5.98	7.53 ± 3.99	0.279	9.70 ± 4.08	0.172
Total cholesterol (mg/dl)	10.1 ± 3.3	−1.2 ± 2.3	0.006	−3.9 ± 2.3	<0.001
HDL cholesterol (mg/dl)	1.3 ± 0.9	0.9 ± 0.6	0.724	0.4 ± 0.6	0.417
LDL cholesterol (mg/dl)	4.6 ± 3.0	−0.5 ± 2.1	0.169	−0.4 ± 2.1	0.178
Triglycerides (mg/dl)	26.5 ± 14.8	−5.8 ± 10.2	0.074	−17.8 ± 10.4	0.015

Data are n (%) or change in least-squares means ± SE unless otherwise indicated. *HOMA-B, homeostasis model of assessment of β -cell function.

diagrams, physical examination findings, vital signs, and hypoglycemic events. Skin and digits were specifically examined because of lesions previously observed in monkeys given DPP-4 inhibitors other than alogliptin (4–6). Safety findings were summarized with descriptive statistics.

RESULTS — Of 420 patients enrolled (see online appendix, available at <http://dx.doi.org/10.2337/dc08-1035>), 329 were randomized to double-blind treatment. Baseline characteristics were similar among treatment groups. Participants had a mean ± SD age of 53.4 ± 11.1 years and baseline A1C 7.9 ± 0.08% and were predominantly male (53.2%) and white (66.9%).

Mean A1C decreased significantly more with 12.5 mg (−0.56%; $P < 0.001$) or 25 mg (−0.59%; $P < 0.001$) alogliptin than with placebo (−0.02%) by week 26. Significant A1C reductions were observed as early as week 4 ($P < 0.001$). FPG reductions were significantly greater with alogliptin than with placebo at week 26 ($P < 0.001$) and as early as week 1 ($P \leq 0.002$). The percentage of patients who required hyperglycemic rescue was significantly less with alogliptin (12.5 mg, 9.8%; 25 mg, 7.6%; $P \leq 0.001$) than with placebo (29.7%). Minor reductions in weight with alogliptin were neither clinically nor statistically significant relative to placebo. Results for secondary efficacy measures and exploratory assessments are summarized in Table 1.

Overall incidences of adverse events (67.4–70.3%) and proportions of patients who discontinued because of adverse events (1.5–2.3%) were similar across treatment groups. Most adverse events were mild or moderate in intensity. Serious adverse events occurred without relation to dose (12.5 mg, 3.8%; 25 mg, 0.8%; and placebo, 3.1%) and were considered unrelated to treatment. No patient died during the study. Adverse events from the most commonly observed categories occurred with similar or lower frequency in those given alogliptin versus placebo (infection, 28.0–37.6%; gastrointestinal, 12.1–14.3%). Headache occurred more frequently with alogliptin (6.8–7.5%) than with placebo (4.7%). Despite increased surveillance for skin-related adverse events, their overall incidence remained low (12.5%), albeit higher with alogliptin (12.8–15.2%) than with placebo (6.3%), mostly because of pruritic events. Two patients discontinued because of skin-related adverse events: one adverse event was considered possibly related to the study drug (25 mg, moderate subcorneal pustular dermatosis); the other was judged unrelated to the study drug (12.5 mg, moderate exacerbation of contact dermatitis). No skin lesions resembling those noted in nonclinical studies of other DPP-4 inhibitors were observed. Hypoglycemia was rare (1.5–3.0%), and no hypoglycemic event was considered an adverse event or was severe enough to require assistance. No clinically meaningful changes in labo-

ratory test results, vital sign measurements, or electrocardiogram recordings were observed.

CONCLUSIONS — Alogliptin monotherapy administered for 26 weeks to treatment-naïve patients with type 2 diabetes produced significant and clinically meaningful improvements in A1C. Glycemic improvements with alogliptin were rapid, sustained, and independent of age, race, and sex. Alogliptin treatment was well tolerated and was not associated with treatment-related serious adverse events. A low incidence of hypoglycemia occurred with alogliptin and with placebo.

Weight gain is common among patients taking sulfonylureas and thiazolidinediones and may reduce treatment adherence (7,8); thus, the weight neutrality of alogliptin and other DPP-4 inhibitors may offer a therapeutic advantage (9–15). Increases in the proinsulin-to-insulin ratio with alogliptin versus placebo and a trend toward increased homeostasis model assessment of β -cell function suggest that alogliptin, similarly to other DPP-4 inhibitors (9–11,14,15), may modestly improve pancreatic function.

In summary, the efficacy and safety of alogliptin monotherapy were comparable with those of other DPP-4 inhibitors (9–15). Alogliptin represents an effective treatment option whether given alone or in combination with antihyperglycemic agents from other classes.

Acknowledgments—Financial support for completion of this study and for data analysis was provided by Takeda Global Research & Development Center, Deerfield, Illinois.

This study was presented at the 68th Annual Scientific Sessions of the American Diabetes Association, San Francisco, CA, 6–10 June 2008, and at the 44th European Association for the Study of Diabetes Annual Meeting, Rome, Italy, 7–11 September 2008.

Editorial assistance with manuscript preparation was provided by Scientific Connections, Newtown, PA.

References

- Drucker DJ, Nauck MA: The incretin system: glucagon-like peptide-1 receptor agonists and dipeptidyl peptidase-4 inhibitors in type 2 diabetes. *Lancet* 368: 1696–1705, 2006
- Feng J, Zhang Z, Wallace MB, Stafford JA, Kaldor SW, Kassel DB, Navre M, Shi L, Skene RJ, Asakawa T, Takeuchi K, Xu R, Webb DR, Gwaltney SL 2nd: Discovery of alogliptin: a potent, selective, bioavailable, and efficacious inhibitor of dipeptidyl peptidase IV. *J Med Chem* 50:2297–2300, 2007
- Covington P, Christopher R, Davenport M, Fleck P, Mekki QA, Wann ER, Karim A: Pharmacokinetic, pharmacodynamic, and tolerability profiles of the dipeptidyl peptidase 4 inhibitor alogliptin: a randomized, double-blind, placebo-controlled multiple-dose study in adult patients with type 2 diabetes. *Clin Ther* 30:499–512, 2008
- European Medicines Agency: European public assessment report for Galvus—scientific discussion, 2007. Available from <http://www.emea.europa.eu/humandocs/PDFs/EPAR/galvus/H-771-en6.pdf>. Accessed 29 July 2008
- Lee B, Shi L, Kassel DB, Asakawa T, Takeuchi K, Christopher RJ: Pharmacokinetic, pharmacodynamic, and efficacy profiles of alogliptin, a novel inhibitor of dipeptidyl peptidase-4, in rats, dogs, and monkeys. *Eur J Pharmacol* 589:306–314, 2008
- Sato K, Ozaki H, Salamon CM, Christopher RJ, Yamamoto M: Lack of dermal toxicity with the highly selective dipeptidyl peptidase-4 inhibitor alogliptin in monkeys (Abstract). *Diabetes* 57 (Suppl. 1): A580, 2008
- Bolen S, Feldman L, Vassy J, Wilson L, Yeh HC, Marinopoulos S, Wiley C, Selvin E, Wilson R, Bass EB, Brancati FL: Systematic review: comparative effectiveness and safety of oral medications for type 2 diabetes mellitus. *Ann Intern Med* 147: 386–399, 2007
- Barnett A, Allsworth J, Jameson K, Mann R: A review of the effects of antihyperglycaemic agents on body weight: the potential of incretin targeted therapies. *Curr Med Res Opin* 23:1493–1507, 2007
- Aschner P, Kipnes MS, Lunceford JK, Sanchez M, Mickel C, Williams-Herman DE, the Sitagliptin Study 021 Group: Effect of the dipeptidyl peptidase-4 inhibitor sitagliptin as monotherapy on glycemic control in patients with type 2 diabetes. *Diabetes Care* 29:2632–2637, 2006
- Pratley RE, Jauffret-Kamel S, Galbreath E, Holmes D: Twelve-week monotherapy with the DPP-4 inhibitor vildagliptin improves glycemic control in subjects with type 2 diabetes. *Horm Metab Res* 38:423–428, 2006
- Scott R, Wu M, Sanchez M, Stein P: Efficacy and tolerability of the dipeptidyl peptidase-4 inhibitor sitagliptin as monotherapy over 12 weeks in patients with type 2 diabetes. *Int J Clin Pract* 61:171–180, 2007
- Pi-Sunyer FX, Schweizer A, Mills D, Dejager S: Efficacy and tolerability of vildagliptin monotherapy in drug-naïve patients with type 2 diabetes. *Diabetes Res Clin Pract* 76: 132–138, 2007
- Dejager S, Razac S, Foley JE, Schweizer A: Vildagliptin in drug-naïve patients with type 2 diabetes: a 24-week, double-blind, randomized, placebo-controlled, multiple-dose study. *Horm Metab Res* 39:218–223, 2007
- Bosi E, Camisasca RP, Collober C, Rochotte E, Garber AJ: Effects of vildagliptin on glucose control over 24 weeks in patients with type 2 diabetes inadequately controlled with metformin. *Diabetes Care* 30: 890–895, 2007
- DeFronzo RA, Hissa M, Blauwet MB, Chen RS: Saxagliptin added to metformin improves glycemic control in patients with type 2 diabetes (Abstract). *Diabetes* 56 (Suppl. 1):A74, 2007