

Skeletal Muscle Insulin Resistance Is the Primary Defect in Type 2 Diabetes

RALPH A. DEFONZO, MD
DEVJIT TRIPATHY, MD

Insulin resistance is a characteristic feature of type 2 diabetes and plays a major role in the pathogenesis of the disease (1,2). Although β -cell failure is the sine qua non for development of type 2 diabetes, skeletal muscle insulin resistance is considered to be the initiating or primary defect that is evident decades before β -cell failure and overt hyperglycemia develops (3,4). Insulin resistance is defined as a reduced response of target tissues (compared with subjects with normal glucose tolerance [NGT] without a family history of diabetes), such as the skeletal muscle, liver, and adipocytes, to insulin. Because skeletal muscle is the predominant site of insulin-mediated glucose uptake in the postprandial state, here we will focus on recent advances about the time of onset, as well as the mechanism, of the skeletal muscle insulin resistance.

RESEARCH DESIGN AND METHODS

The euglycemic insulin clamp technique (5) is considered to be the gold standard for measuring insulin action in vivo. With this technique, whole-body insulin action is quantified as the rate of exogenous glucose infusion (plus any residual hepatic glucose production) required to maintain the plasma glucose concentration at euglycemic levels in response to a fixed increment in the plasma insulin concentration. Because 80–90% of the infused glucose is taken up by skeletal muscle under conditions of euglycemic hyperinsulinemia, insulin sensitivity measured with the insulin clamp technique primarily reflects skeletal muscle (6). Another advantage of this technique is that it can be combined with indirect calorimetry to measure different substrate oxidation rates and with muscle

biopsy to examine the biochemical/molecular etiology of the insulin resistance. Measurement of insulin sensitivity by the frequently sampled intravenous glucose tolerance test reflects both hepatic and peripheral insulin resistance and correlates well with the insulin clamp technique (7).

Because insulin clamp studies are not feasible in large epidemiological studies, other surrogate markers of insulin sensitivity from glucose and insulin values in the fasting state or after an oral glucose tolerance test (OGTT) have been developed (8–10). The homeostatic model assessment correlates reasonably well with the insulin clamp (10), but it primarily reflects hepatic insulin sensitivity, since the fasting plasma glucose is determined mainly by the rate of hepatic glucose production (HGP) and insulin is the primary regulator of HGP. The correlation between homeostatic model assessment and the insulin clamp also is less robust when analyzed in subgroups of glucose tolerance (11). During an OGTT, significant (~30–40%) amounts of glucose are taken up by the splanchnic bed, and HGP is less completely suppressed than during the insulin clamp technique (12). As a result, the plasma glucose concentration during OGTT is affected by both hepatic and peripheral (primarily muscle) insulin resistance. Therefore, indexes of insulin resistance from the OGTT, e.g., the Matsuda index, reflect both hepatic and peripheral insulin resistance and correlate well (R value ~ 0.70) with insulin sensitivity measured with the euglycemic insulin clamp (9).

Normal glucose homeostasis

Skeletal muscle is the major site of glucose uptake in the postprandial state in hu-

mans. Under euglycemic hyperinsulinemic conditions, ~80% of glucose uptake occurs in skeletal muscle (13). Studies using the euglycemic hyperinsulinemic clamp and femoral artery/vein catheterization to quantitate glucose uptake have allowed investigators to quantify leg muscle glucose uptake. Because adipose tissue uses <5% of an infused glucose load and bone is metabolically inert, the great majority of leg glucose uptake can be accounted for by skeletal muscle. During physiological hyperinsulinemia (80–100 μ U/ml), leg muscle glucose uptake increases linearly with time, reaching a plateau value of ~10 mg/kg leg weight per minute after 60 min (13,14). In contrast, in lean type 2 diabetic subjects, the onset of insulin action is delayed and the ability of insulin to maximally stimulate glucose uptake is markedly blunted. During the last hour of the insulin clamp, insulin-stimulated leg muscle glucose uptake is reduced by ~50% in type 2 diabetes (14). These studies support the notion that the primary defect in insulin action in patients with type 2 diabetes resides in the skeletal muscle. Similarly, using the forearm catheterization technique, a number of investigators have demonstrated reduced insulin-mediated glucose uptake by the peripheral tissues, primarily muscle (15). Quantitation of leg muscle glucose uptake in type 2 diabetes with positron emission tomography has provided additional evidence for the presence of severe muscle insulin resistance in type 2 diabetes (16). In the postabsorptive state, the majority (~70–75%) of glucose uptake (~2 mg/kg per min) occurs in insulin-insensitive tissues (brain, erythrocytes, and splanchnic tissues) (17,18), with only ~25% of glucose uptake occurring in insulin-sensitive tissues. In the postabsorptive state, total body glucose uptake is precisely matched by the rate of endogenous glucose production, primarily by the liver and to a smaller extent by the kidney (19). Thus, hepatic glucose production is the main determinant of the fasting plasma glucose concentration and is regulated primarily by the plasma insulin and, to a lesser extent, glucagon concentrations (20).

From the University of Texas Health Science Center, San Antonio, Texas.

Corresponding author: Ralph A. DeFronzo, albarado@uthscsa.edu.

The publication of this supplement was made possible in part by unrestricted educational grants from Eli Lilly, Ethicon Endo-Surgery, Genex Biotechnology, Hoffmann-La Roche, Johnson & Johnson, LifeScan, Medtronic, MSD, Novo Nordisk, Pfizer, sanofi-aventis, and WorldWIDE.

DOI: 10.2337/dc09-S302

© 2009 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

Glucose disposal during OGTT

After a mixed-meal or oral glucose load, the ensuing hyperglycemia and hyperinsulinemia work in concert to suppress HGP and stimulate glucose uptake by the splanchnic (liver) and peripheral (muscle) tissues (12,21). After glucose is transported into the cell, it is phosphorylated and subsequently oxidized to carbon dioxide and water or converted to glycogen, which can be stored in the liver or skeletal muscle (22). Glycogen synthesis is regulated by the enzyme glycogen synthase, and glycolysis primarily is controlled by the enzyme complex of pyruvate dehydrogenase. In humans, the rate of glucose oxidation and the nonoxidative glucose metabolism (primarily reflects glycogen synthesis) in the insulin-stimulated state can be estimated by indirect calorimetry or magnetic response spectroscopy (23,24). Approximately 75% of glucose is metabolized nonoxidatively, and impaired glycogen synthesis is one of the earliest metabolic defects seen in the pathogenesis of type 2 diabetes (1,2).

Intracellular pathways of glucose disposal

Under physiologic conditions, approximately two-thirds of all glucose-6-phosphate is converted to glycogen, and one-third enters glycolysis. Of the glucose that enters the glycolytic pathway, the majority (80–90%) is converted to carbon dioxide and water, whereas the remaining 10–20% is converted to lactate. Studies using indirect calorimetry and euglycemic clamp technique have shown that the glucose oxidation is more sensitive (lower half-maximum) but saturates earlier (lower maximum) than glycogen synthesis, which has low sensitivity but high capacity. Skeletal muscle is the predominant site of glycogen synthesis.

Molecular basis of insulin action

In the early stages of development of type 2 diabetes, impaired glycogen synthesis in muscle is the primary defect responsible for the insulin resistance. The initial step in muscle glucose metabolism involves activation of the glucose transport system, leading to influx of glucose into insulin target tissues. The free glucose that has entered the cell is then metabolized by a series of enzymatic steps that are under the control of insulin. In skeletal muscle and adipose tissue, insulin promotes glucose uptake into the cells by activating a complex cascade of phosphorylation-dephosphorylation reactions. In skeletal

muscle, insulin binds to the insulin receptor leading to phosphorylation of three key tyrosine molecules on the insulin receptor. Once the insulin receptor has been phosphorylated, insulin receptor substrate (IRS)-1 moves to the cell membrane and becomes phosphorylated on contiguous tyrosine molecules. Tyrosine phosphorylation of IRS-1 results in activation of the p85 regulatory subunit of phosphatidylinositol (PI)-3 kinase and activates the p110 catalytic subunit, leading to an increase in phosphatidylinositol-3,4,5 triphosphate. This results in activation of downstream protein kinase B (also called Akt) and phosphorylation of Akt substrate 160 (AS160), which facilitates the translocation of GLUT4 to the sarcolemma and subsequent entry of glucose into the cell (25). Intracellular glucose is rapidly phosphorylated by hexokinase II and directed to oxidative or nonoxidative pathways. Maintaining the integrity of the IRS-1/PI-3 kinase/Akt pathway is essential for normal insulin-mediated glucose uptake in skeletal muscle (26).

What is the initial metabolic defect in the pathogenesis of type 2 diabetes?

By the time that hyperglycemia is manifest, multiple metabolic abnormalities are present in individuals with type 2 diabetes. Because the majority of type 2 diabetic subjects are obese, they also have daylong elevation of the plasma free fatty acid (FFA) concentration and increased circulating levels of inflammatory cytokines (1,2). Because elevated plasma glucose, FFA, and cytokine concentrations all can induce insulin resistance, it is extremely difficult to separate the contribution of each of these metabolic defects in the pathogenesis of type 2 diabetes. To examine what is the earliest defect(s) in the development of type 2 diabetes, investigators have used two approaches. First, one can study the lean, normal glucose tolerant, first-degree relatives of two parents with type 2 diabetes. These individuals have a very high lifetime risk (~40%) of developing type 2 diabetes (4). In certain high-risk populations, such as Mexican Americans, the prevalence of diabetes in the offspring of two type 2 diabetic parents can reach 70–80%. The advantage of studying this genetically predisposed group is that they do not have other confounding factors that contribute to insulin resistance, such as obesity and hyperglycemia. Thus, they represent an ideal

model to study the early metabolic defects in the pathogenesis of type 2 diabetes. A second approach uses the long-term follow-up of normal glucose tolerant subjects as they progress to impaired glucose tolerance and subsequently to type 2 diabetes. This prospective approach has been used in Pima Indians (27).

What is the evidence that muscle insulin resistance is the initial metabolic defect in type 2 diabetes?

Multiple investigators unequivocally have demonstrated that lean NGT offspring of two parents with type 2 diabetes exhibit moderate to severe skeletal muscle insulin resistance (28–33). The natural history of type 2 diabetes is depicted in Fig. A1 (all figures can be found in an online appendix available at <http://care.diabetesjournals.org/cgi/content/full/dc09-S302/DC1>) (1,34). As Europoid individuals progress from NGT to IGT, insulin sensitivity declines markedly but glucose tolerance deteriorates minimally because of a marked increase in insulin secretion. Similar observations have been made in Pima Indians (Fig. A2) (35) and Mexican Americans and Caucasians residing in San Antonio (Fig. A3) (30,32,36). These results, spanning a wide range of ethnic groups, clearly demonstrate that insulin resistance, and not insulin deficiency, initiates the sequence of events leading to the development of type 2 diabetes. However, progressive β -cell failure is required and ultimately responsible for type 2 diabetes to become fully manifest (1,34,35,37,38).

In humans, ~75–80% of insulin-stimulated muscle glucose disposal during a euglycemic insulin clamp is converted to glycogen, whereas the remaining 20–25% is oxidized to CO₂ and H₂O (39). In type 2 diabetes, impaired glycogen synthesis secondary to reduced glycogen synthase activity is the earliest detectable metabolic defect (39–41). Gulli et al. (32) studied the lean NGT offspring of two diabetic Mexican American parents using a two-step euglycemic insulin clamp (20 and 40 mU/m² per min) that produced steady-state elevations in the plasma insulin concentration of ~40 and ~80 μ U/ml, respectively, which are close to the half-maximal suppression of HGP and stimulation of muscle glucose uptake and glycogen synthesis. At both plasma insulin concentrations, glucose uptake was decreased by 33 and 43%, respectively (32) (Fig. A4). The impairment in glucose uptake was accounted for en-

tirely by reduced nonoxidative glucose metabolism (Fig. A4), which represents glycogen synthesis. No defect was noted in the suppression of hepatic glucose production by insulin (Fig. A5). These results indicate that the defect in insulin action primarily is localized to skeletal muscle and involves the glycogen synthetic pathway. During both the OGTT and hyperglycemic clamp (Fig. A6), insulin secretion (total, early, and late responses) was significantly increased, excluding a primary defect in β -cell function in the pathogenesis of type 2 diabetes in this ethnic population.

A similar defect in insulin-stimulated glycogen synthesis has been reported by Groop et al. in the Botnia study (42,43). Using nuclear magnetic resonance (NMR) spectroscopy, Perseghin et al. (44) directly quantified the defect in muscle glycogen synthesis in NGT offspring of two diabetic parents. This technique allows one to monitor glycogen synthesis noninvasively over real time. These investigators demonstrated that reduced glycogen synthesis could account for almost all of the decrease in insulin-stimulated glucose disposal in skeletal muscle. Of note, the severity of skeletal muscle insulin resistance in the offspring of two diabetic parents is of similar magnitude to that seen in type 2 diabetic individuals.

Before the incorporation of glucose into glycogen, it first must be transported into the cell and phosphorylated by hexokinase II to glucose-6-phosphate. Using a novel triple isotope (^{12}C -mannitol, ^{13}C -0-methylglucose, $3\text{-}^3\text{H}$ -glucose) technique in combination with brachial arterial/deep vein forearm catheterization and the euglycemic clamp, Pendergrass et al. (45) quantitated muscle glucose uptake and phosphorylation in the insulin-resistant NGT offspring of two type 2 diabetic parents (Fig. A7). The NGT offspring demonstrated a severity of whole-body and forearm muscle insulin resistance similar to that observed in lean and obese type 2 diabetic subjects and in NGT obese insulin-resistant individuals. Similarly, the NGT offspring manifested defects in muscle glucose transport and phosphorylation that were similar to those in lean and obese type 2 diabetic subjects (Fig. A7). Using ^{14}C -NMR to measure muscle glucose-6-phosphate levels, Rothman et al. (46) demonstrated a similar defect in combined glucose transport/phosphorylation in the NGT offspring of two type 2 diabetic parents.

Another strategy to examine what is

the earliest defect responsible for the development of type 2 diabetes is to prospectively follow individuals who are at high risk for the development of type 2 diabetes. This has been done by Weyer et al. (47) in Pima Indians. At-risk individuals received a euglycemic insulin clamp to measure insulin sensitivity (primarily reflects muscle) and an intravenous glucose tolerance test to quantitate insulin secretion and were followed sequentially until they developed diabetes. At-risk individuals were markedly resistant to insulin but, at the stage of NGT, their β -cells were able to secrete sufficient amounts of insulin to offset the insulin resistance (Fig. A8). With time, both progressors (to type 2 diabetes) and nonprogressors (remained as NGT) experienced a further modest reduction (11–14%) in insulin sensitivity. However, nonprogressors were able to offset the worsening insulin resistance in muscle by augmenting insulin secretion (30%), whereas progression to type 2 diabetes was associated with a 78% decline in the acute insulin response to an intravenous glucose challenge (Fig. A8). Using a similar approach, Warram et al. (4) also demonstrated that insulin resistance was a strong predictor for the future diabetes. Because the great majority of glucose disposal after intravenous glucose administration occurs in muscle, these results provide strong evidence that insulin resistance in muscle is the earliest demonstrable defect in the natural history of type 2 diabetes, but that the development of overt type 2 diabetes occurs only in those individuals whose β -cells are unable to compensate for the defect in insulin action (37,38,48,49).

Insulin resistance begets insulin resistance

The normal β -cell response to insulin resistance, irrespective of the etiology of the insulin resistance, is to increase its secretion of insulin (1,2,39). However, a chronic physiologic increase in the plasma insulin concentration has a detrimental effect on skeletal muscle insulin sensitivity. Del Prato et al. (50) demonstrated that a 72 pmol/l (11 $\mu\text{U/ml}$) increase in the plasma insulin concentration in healthy NGT insulin-sensitive individuals for as little as 72–96 h reduced insulin-stimulated glucose disposal (insulin clamp technique) by 30–40%. The defect in insulin action was accounted for entirely by impaired nonoxidative glucose disposal (Fig. A9). On the other hand, chronic euglycemic hyperinsulinemia did

not alter insulin-mediated suppression of hepatic glucose production (50). Koopmans et al. (51,52) showed that chronic hyperinsulinemia (threefold increase above baseline) in conscious rats for 7 days resulted in a reduction in insulin-mediated total-body glucose uptake, glucose storage, and glycolysis by 39, 62, and 26%, respectively. Hepatic glucose production was normally suppressed after 7 days of hyperinsulinemia. Because the majority (>80–90%) of glucose disposal during the euglycemic insulin clamp occurs in muscle, these results demonstrate that a physiologic elevation in the plasma insulin concentration will exacerbate the underlying muscle insulin resistance. Izzo et al. (53) performed a 240-min euglycemic insulin clamp study with muscle biopsies in healthy volunteers. Subjects then received a low-dose insulin infusion for 72 h (plasma insulin concentration 143 ± 25 pmol/l [21 ± 2 $\mu\text{U/ml}$]), followed by a repeat insulin clamp with muscle biopsies. After 72 h of sustained physiologic hyperinsulinemia, insulin-stimulated muscle glycogen synthase activity, total body glucose uptake, and nonoxidative glucose disposal (primarily reflects glycogen synthesis in muscle) were significantly reduced. Taken together, these findings indicate that hyperinsulinemia is not only a compensatory response to insulin resistance, but also a self-perpetuating cause of the defect in muscle insulin action.

Molecular etiology of the skeletal muscle insulin resistance in genetically predisposed individuals

Using the euglycemic insulin clamp with skeletal muscle biopsy, a number of investigators have examined the insulin signal transduction system in human skeletal muscle of type 2 diabetic subjects and consistently demonstrated defects in IRS-1 tyrosine phosphorylation and PI-3 kinase and Akt activation (26,54,55). To examine whether similar defects are present in genetically predisposed individuals, Pratipanawat et al. (36) examined insulin signaling in NGT subjects with a strong family history of type 2 diabetes and demonstrated that both the basal and insulin-stimulated IRS-1 tyrosine phosphorylation and PI 3-kinase activity associated with IRS-1 were significantly decreased (Fig. A10). Insulin stimulation of PI 3-kinase activity is a requisite for activation of glucose transport and glycogen synthesis. Increased serine phosphorylation of IRS-1 has been shown

to impair insulin signaling (tyrosine phosphorylation of both insulin resistance and IRS-1) in type 2 diabetes (56). In lean insulin-resistant NGT offspring of type 2 diabetic parents, increased serine phosphorylation of IRS-1 in skeletal muscle has been documented in association with impaired activation of Akt (57) (Fig. A11). Thus, at the earliest stage in the natural history of type 2 diabetes, i.e., the NGT insulin-resistant offspring of two type 2 diabetic parents, the molecular etiology of the muscle insulin resistance already is well established and is virtually identical to that in their diabetic parents.

Relationship between muscle insulin resistance and altered FFA/muscle lipid metabolism

Gulli et al. (32) were the first to demonstrate that the NGT offspring of two type 2 diabetic parents demonstrated marked muscle insulin resistance but normal sensitivity to the suppressive effect of insulin on hepatic glucose production. However, a normal basal rate of HGP in the face of fasting hyperinsulinemia could be construed to indicate the presence of hepatic insulin resistance. More impressive was the elevated fasting plasma FFA concentration in the presence of fasting hyperinsulinemia and the impaired suppression of plasma FFA during the euglycemic insulin clamp (Fig. A12). These findings indicate the presence of marked adipocyte resistance to the antilipolytic effect of insulin. Impaired insulin-mediated suppression of whole-body lipid oxidation also was present in the NGT offspring (Fig. A12). Petersen et al. (58) documented an increase in intramyocellular lipid content in the offspring of two type 2 diabetic parents. This observation is of important clinical significance, since diacylglycerol, long-chain fatty acyl CoAs, and ceramides all have been shown to cause serine phosphorylation of insulin resistance and IRS-1 and lead to the development of insulin resistance in skeletal muscle (59,60). Collectively, these results suggest that intramyocellular accumulation of toxic lipid metabolites plays an important role in the pathogenesis of muscle insulin resistance.

To further address this question, Kashyap et al. (30) infused a lipid emulsion for 4 days to cause a physiologic elevation in the plasma FFA concentration in NGT insulin-resistant offspring of two type 2 diabetes parents and in NGT insulin-sensitive subjects without any family history of diabetes. Four days of physio-

logical elevation in the plasma FFA concentration in the offspring did not cause any further worsening of insulin-stimulated whole-body glucose disposal, nonoxidative glucose disposal, glucose oxidation, or preexisting defects in insulin-stimulated insulin receptor tyrosine phosphorylation (30). In contrast, in healthy control subjects, chronic lipid infusion was associated with a marked decline in insulin-stimulated glucose uptake and insulin receptor tyrosine phosphorylation (30). When the insulin-resistant offspring were treated with acipimox for 7 days to reduce the plasma FFA concentration and intramyocellular FFA concentration, a marked improvement in insulin sensitivity was observed (61). These data lend further support to the observation that insulin resistance in skeletal muscle is an early metabolic defect in the pathogenesis of type 2 diabetes and that muscle lipid accumulation plays a central role in the etiology of the muscle insulin resistance.

Mitochondria are the main organelles where fatty acids are oxidized and investigators have focused on their structure and function in patients with type 2 diabetes. Studies using the leg balance technique have documented that fat oxidation is reduced in both type 2 diabetic and obese insulin-resistant nondiabetic individuals (62), suggesting that muscle mitochondrial oxidative capacity is impaired. Recently, two groups independently showed that NGT offspring of two type 2 diabetic parents had a reduced expression of key mitochondrial genes involved in the regulation of oxidative metabolism in skeletal muscle (63,64). The most commonly underexpressed functional genes were those coding for energy generation, including multiple glycolytic, tricarboxylic acid cycle, and oxidative phosphorylation genes. Evidence in support of a role for mitochondrial dysfunction as a cause of muscle insulin resistance in the NGT offspring of two type 2 diabetic parents has been provided by Shulman and colleagues. Using ³¹P-NMR, these investigators demonstrated impaired mitochondrial activity in NGT insulin-resistant offspring of type 2 diabetic parents (57,58,65). Whereas mitochondria from NGT subjects without any family history of diabetes responded to insulin by increasing ATP production by 90%, mitochondria from insulin-resistant offspring increased ATP production by only 5% (Fig. A13). The authors postulated that muscle mitochondrial

dysfunction was the primary defect, leading to elevated intramyocellular fatty acid metabolites (as a consequence of reduced fat oxidation) and subsequent insulin resistance (58,66). However, recent studies by Abdul-Ghani et al. (67) have shown that even small increases in palmitoyl carnitine (5–10 μmol/l) can markedly impair ATP synthesis in mitochondria isolated from human muscle. Thus, it is unclear which is the cart and which is the horse: mitochondrial dysfunction leading to increased intramyocellular lipid content and insulin resistance or increased muscle lipid content (i.e., secondary to elevated plasma FFA levels and/or excessive lipid ingestion) leading to mitochondrial dysfunction and insulin resistance.

SUMMARY — The maintenance of normal glucose homeostasis depends on a finely balanced dynamic interaction between tissue (muscle, liver, and fat) sensitivity to insulin and insulin secretion. Even in the presence of severe insulin resistance, a perfectly normal β-cell is capable of secreting sufficient amounts of insulin to offset the defect in insulin action. Thus, the evolution of type 2 diabetes requires the presence of defects in both insulin secretion and insulin action, and both of these defects can have a genetic as well as an acquired component. When type 2 diabetic patients initially present to the physician, they will have had their diabetes for many years, and defects in insulin action (in muscle, liver, and adipocytes) and insulin secretion will be well established (1,2,39). At this stage, it is not possible to define which defect came first in the natural history of the disease and which tissue is the primary defect responsible for the insulin resistance. Although insulin resistance represents the earliest detectable abnormality in the great majority of type 2 diabetic people, in a minority of individuals (i.e., glucokinase deficiency), it is clear that a β-cell defect initiates the disturbance in glucose homeostasis. Nevertheless, it is now clear that in any given diabetic patient, whatever defect (insulin resistance or impaired insulin secretion) initiates the disturbance in glucose metabolism, it will eventually be followed by the emergence of its counterpart (Fig. A14).

Insulin resistance is a nearly universal finding in patients with established type 2 diabetes. In normal-weight and obese individuals with IGT and in type 2 diabetic subjects with mild fasting hyperglycemia (110–140 mg/dl, 6.1–7.8 mmol/l), both

the basal and glucose-stimulated plasma insulin levels are increased. Although the first-phase insulin response may be decreased in some, but not all, of these subjects, the first phase consistently is increased in the NGT offspring of two type 2 diabetic parents and the total insulin response is increased in NGT offspring and in IGT subjects. In each of these groups, tissue sensitivity to insulin, measured with the insulin clamp technique, has been shown to be diminished. Prospective studies conclusively have demonstrated that hyperinsulinemia and insulin resistance precede the development of IGT and that IGT represents the forerunner of type 2 diabetes. This scenario has been well documented in Pima Indians, Mexican Americans, and Pacific Islanders. It is noteworthy that all of these populations are characterized by obesity and a younger age at onset of diabetes. Such results provide conclusive evidence that insulin resistance is the inherited defect that initiates the diabetic condition in the majority of type 2 diabetic patients. Studies in NGT first-degree relatives of diabetic individuals and in the offspring of two diabetic parents indicate that the inherited defect in insulin action results from an abnormality in the glycogen synthetic pathway in muscle and more proximal defects in glucose transport/phosphorylation and insulin signal transduction. As the insulin resistance progresses and muscle glucose uptake becomes further impaired, the postprandial rise in plasma glucose concentration becomes excessive, but the increase in basal hyperinsulinemia is sufficient to maintain the fasting plasma glucose concentration and HGP within the normal range. Nonetheless, there is an excessive postprandial rise in plasma glucose concentration, and a longer time is required to restore normoglycemia after each meal. Eventually, however, the insulin resistance becomes so severe that the compensatory hyperinsulinemia is no longer sufficient to maintain the fasting glucose concentration at the basal level. The development of hyperglycemia further stimulates β -cell secretion of insulin, and the resultant hyperinsulinemia causes a downregulation of insulin receptor number and of the intracellular events involved in insulin action, thus exacerbating the insulin resistance. Initially, the hyperglycemia-induced increase in insulin secretion serves a compensatory function to maintain near-NGT. In some individuals, the persistent stimulus to the β -cell to over-

secrete insulin leads to a progressive loss of β -cell function. Chronic hyperglycemia (glucose toxicity) and/or disturbances in lipid metabolism (lipotoxicity) may contribute to the defect in insulin secretion. The resultant insulinopenia leads to the emergence/exacerbation of postreceptor defects in insulin action. Many of the intracellular events involved in glucose metabolism depend on the surge of insulin that occurs three to four times per day in response to nutrient ingestion. When the insulin response becomes deficient, the activity of the glucose transport system becomes severely impaired and a number of key intracellular enzymatic steps involved in glucose metabolism become depressed. Additionally, when severe insulinopenia ensues, plasma FFA levels rise, further contributing to the defects in intracellular glucose disposal. There is also compelling evidence that hyperglycemia per se can downregulate the glucose transport system, as well as a number of other intracellular events involved in insulin action (glucose toxicity), and a similar argument can be made concerning the intracellular derangement in lipid metabolism. This pathogenetic sequence can explain all of the clinical and laboratory features observed in type 2 diabetic patients. Insofar as the cellular defect is generalized, both hepatic and peripheral tissues (skeletal muscle and adipocytes), and possibly the β -cells themselves, would manifest insulin resistance, and the numerous metabolic alterations characteristic of the diabetic state could be related to one and the same primary defect.

The NGT offspring of two type 2 diabetic parents also manifest marked adipocyte resistance to the suppressive effects of insulin on lipolysis. One could argue, therefore, that the adipocyte represents the primary tissue responsible for the insulin resistance. According to this scenario, the elevated plasma FFA levels produce insulin resistance in muscle and liver and impair β -cell function. Adipocytes in the NGT offspring of two type 2 diabetic parents also secrete excessive amounts of inflammatory and insulin resistance producing adipocytokines that could initiate/exacerbate the insulin resistance in skeletal muscle. As reviewed by Iozzo in this symposium, the adipocyte insulin resistance could be genetic in origin or induced in utero during the third trimester by nutritional deprivation or overfeeding.

There is less evidence to support a role for the liver as the organ responsible for the insulin resistance. However, the NGT offspring of two type 2 diabetic parents have a normal rate of HGP in the presence of fasting hyperinsulinemia, suggesting the presence of hepatic resistance to the suppressive effect of insulin on glucose production. Therefore, one could argue that the resultant fasting hyperinsulinemia leads to the development of insulin resistance in skeletal muscle.

Acknowledgments—No potential conflicts of interest relevant to this article were reported.

References

1. DeFronzo RA. Lilly lecture 1987. The triumvirate: beta-cell, muscle, liver: a collusion responsible for NIDDM. *Diabetes* 1988;37:667–687
2. DeFronzo RA. Pathogenesis of type 2 diabetes mellitus. *Med Clin North Am* 2004;88:787–835
3. Lillioja S, Mott DM, Howard BV, Bennett PH, Yki-Jarvinen H, Freymond D, Nyomba BL, Zurlo F, Swinburn B, Bogardus C. Impaired glucose tolerance as a disorder of insulin action: longitudinal and cross-sectional studies in Pima Indians. *N Engl J Med* 1988;318:1217–1225
4. Warram JH, Martin BC, Krolewski AS, Soeldner JS, Kahn CR. Slow glucose removal rate and hyperinsulinemia precede the development of type II diabetes in the offspring of diabetic parents. *Ann Intern Med* 1990;113:909–915
5. DeFronzo RA, Tobin JD, Andres R. Glucose clamp technique: a method for quantifying insulin secretion and resistance. *Am J Physiol* 1979;237:E214–E223
6. Ferrannini E, Simonson DC, Katz LD, Reichard G Jr, Bevilacqua S, Barrett EJ, Olsson M, DeFronzo RA. The disposal of an oral glucose load in patients with non-insulin-dependent diabetes. *Metabolism* 1988;37:79–85
7. Pacini G, Finegood DT, Bergman RN. A minimal-model-based glucose clamp yielding insulin sensitivity independent of glycemia. *Diabetes* 1982;31:432–441
8. Katz A, Nambi SS, Mather K, Baron AD, Follmann DA, Sullivan G, Quon MJ. Quantitative insulin sensitivity check index: a simple, accurate method for assessing insulin sensitivity in humans. *J Clin Endocrinol Metab* 2000;85:2402–2410
9. Matsuda M, DeFronzo RA. Insulin sensitivity indices obtained from oral glucose tolerance testing: comparison with the euglycemic insulin clamp. *Diabetes Care* 1999;22:1462–1470
10. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Ho-

- meostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 1985;28:412–419
11. Tripathy D, Almgren P, Tuomi T, Groop L. Contribution of insulin-stimulated glucose uptake and basal hepatic insulin sensitivity to surrogate measures of insulin sensitivity. *Diabetes Care* 2004;27:2204–2210
 12. Ferrannini E, Bjorkman O, Reichard GA Jr, Pilo A, Olsson M, Wahren J, DeFronzo RA. The disposal of an oral glucose load in healthy subjects: a quantitative study. *Diabetes* 1985;34:580–588
 13. Thiebaud D, Jacot E, DeFronzo RA, Maeder E, Jequier E, Felber JP. The effect of graded doses of insulin on total glucose uptake, glucose oxidation, and glucose storage in man. *Diabetes* 1982;31:957–963
 14. DeFronzo RA, Jacot E, Jequier E, Maeder E, Wahren J, Felber JP. The effect of insulin on the disposal of intravenous glucose: results from indirect calorimetry and hepatic and femoral venous catheterization. *Diabetes* 1981;30:1000–1007
 15. Pendergrass M, Bertoldo A, Bonadonna R, Nucci G, Mandarino L, Cobelli C, DeFronzo RA. Muscle glucose transport and phosphorylation in type 2 diabetic, obese nondiabetic, and genetically predisposed individuals. *Am J Physiol Endocrinol Metab* 2007;292:E92–E100
 16. Utriainen T, Takala T, Luotolahti M, Ronnema T, Laine H, Ruotsalainen U, Haaparanta M, Nuutila P, Yki-Jarvinen H. Insulin resistance characterizes glucose uptake in skeletal muscle but not in the heart in NIDDM. *Diabetologia* 1998;41:555–559
 17. Landau BR, Wahren J, Chandramouli V, Schumann WC, Ekberg K, Kalhan SC. Contributions of gluconeogenesis to glucose production in the fasted state. *J Clin Invest* 1996;98:378–385
 18. DeFronzo RA, Ferrannini E. Regulation of hepatic glucose metabolism in humans. *Diabetes Metab Rev* 1987;3:415–459
 19. Gerich JE, Meyer C, Woerle HJ, Stumvoll M. Renal gluconeogenesis: its importance in human glucose homeostasis. *Diabetes Care* 2001;24:382–391
 20. DeFronzo RA, Ferrannini E, Simonson DC. Fasting hyperglycemia in non-insulin-dependent diabetes mellitus: contributions of excessive hepatic glucose production and impaired tissue glucose uptake. *Metabolism* 1989;38:387–395
 21. Katz LD, Glickman MG, Rapoport S, Ferrannini E, DeFronzo RA. Splanchnic and peripheral disposal of oral glucose in man. *Diabetes* 1983;32:675–679
 22. Del Prato S, Bonadonna RC, Bonora E, Gulli G, Solini A, Shank M, DeFronzo RA. Characterization of cellular defects of insulin action in type 2 (non-insulin-dependent) diabetes mellitus. *J Clin Invest* 1993;91:484–494
 23. Groop LC, Bonadonna RC, Simonson DC, Petrides AS, Shank M, DeFronzo RA. Effect of insulin on oxidative and nonoxidative pathways of free fatty acid metabolism in human obesity. *Am J Physiol* 1992;263:E79–E84
 24. Simonson DC, DeFronzo RA. Indirect calorimetry: methodological and interpretative problems. *Am J Physiol* 1990;258:E399–E412
 25. Taniguchi CM, Emanuelli B, Kahn CR. Critical nodes in signalling pathways: insights into insulin action. *Nat Rev Mol Cell Biol* 2006;7:85–96
 26. Krook A, Bjornholm M, Galuska D, Jiang XJ, Fahlman R, Myers MG Jr, Wallberg-Henriksson H, Zierath JR. Characterization of signal transduction and glucose transport in skeletal muscle from type 2 diabetic patients. *Diabetes* 2000;49:284–292
 27. McCance DR, Pettitt DJ, Hanson RL, Jacobsson LT, Bennett PH, Knowler WC. Glucose, insulin concentrations and obesity in childhood and adolescence as predictors of NIDDM. *Diabetologia* 1994;37:617–23
 28. Perseghin G, Ghosh S, Gerow K, Shulman GI. Metabolic defects in lean nondiabetic offspring of NIDDM parents: a cross-sectional study. *Diabetes* 1997;46:1001–1009
 29. Ferrannini E, Gastaldelli A, Matsuda M, Miyazaki Y, Pettiti M, Glass L, DeFronzo RA. Influence of ethnicity and familial diabetes on glucose tolerance and insulin action: a physiological analysis. *J Clin Endocrinol Metab* 2003;88:3251–3257
 30. Kashyap SR, Belfort R, Berria R, Suraamornkul S, Pratipanawatr T, Finlayson J, Barrentine A, Bajaj M, Mandarino L, DeFronzo R, Cusi K. Discordant effects of a chronic physiological increase in plasma FFA on insulin signaling in healthy subjects with or without a family history of type 2 diabetes. *Am J Physiol Endocrinol Metab* 2004;287:E537–E546
 31. Vaag A, Henriksen JE, Beck-Nielsen H. Decreased insulin activation of glycogen synthase in skeletal muscles in young nonobese Caucasian first-degree relatives of patients with non-insulin-dependent diabetes mellitus. *J Clin Invest* 1992;89:782–788
 32. Gulli G, Ferrannini E, Stern M, Haffner S, DeFronzo RA. The metabolic profile of NIDDM is fully established in glucose-tolerant offspring of two Mexican-American NIDDM parents. *Diabetes* 1992;41:1575–1586
 33. Tripathy D, Lindholm E, Isomaa B, Saloranta C, Tuomi T, Groop L. Familiarity of metabolic abnormalities is dependent on age at onset and phenotype of the type 2 diabetic proband. *Am J Physiol* 2003;285:E1297–E303
 34. Jallut D, Golay A, Munger R, Frascarolo P, Schutz Y, Jequier E, Felber JP. Impaired glucose tolerance and diabetes in obesity: a 6-year follow-up study of glucose metabolism. *Metabolism* 1990;39:1068–1075
 35. Lillioja S, Mott DM, Spraul M, Ferraro R, Foley JE, Ravussin E, Knowler WC, Bennett PH, Bogardus C. Insulin resistance and insulin secretory dysfunction as precursors of non-insulin-dependent diabetes mellitus: prospective studies of Pima Indians. *N Engl J Med* 1993;329:1988–1992
 36. Pratipanawatr W, Pratipanawatr T, Cusi K, Berria R, Adams JM, Jenkinson CP, Mazzone K, DeFronzo RA, Mandarino LJ. Skeletal muscle insulin resistance in normoglycemic subjects with a strong family history of type 2 diabetes is associated with decreased insulin-stimulated insulin receptor substrate-1 tyrosine phosphorylation. *Diabetes* 2001;50:2572–2578
 37. Abdul-Ghani MA, Tripathy D, DeFronzo RA. Contributions of beta-cell dysfunction and insulin resistance to the pathogenesis of impaired glucose tolerance and impaired fasting glucose. *Diabetes Care* 2006;29:1130–1139
 38. Gastaldelli A, Ferrannini E, Miyazaki Y, Matsuda M, DeFronzo RA. Beta-cell dysfunction and glucose intolerance: results from the San Antonio metabolism (SAM) study. *Diabetologia* 2004;47:31–39
 39. DeFronzo RA. Pathogenesis of type 2 diabetes: metabolic and molecular implications for identifying genes. *Diabetes Rev* 1997;5:177–269
 40. Bogardus C, Lillioja S, Stone K, Mott D. Correlation between muscle glycogen synthase activity and in vivo insulin action in man. *J Clin Invest* 1984;73:1185–1190
 41. Shulman GI, Rothman DL, Jue T, Stein P, DeFronzo RA, Shulman RG. Quantitation of muscle glycogen synthesis in normal subjects and subjects with non-insulin-dependent diabetes by ¹³C nuclear magnetic resonance spectroscopy. *N Engl J Med* 1990;322:223–228
 42. Eriksson J, Franssila-Kallunki A, Ekstrand A, Saloranta C, Widen E, Schalin C, Groop L. Early metabolic defects in persons at increased risk for non-insulin-dependent diabetes mellitus. *N Engl J Med* 1989;321:337–343
 43. Groop L, Forsblom C, Lehtovirta M, Tuomi T, Karanko S, Nissen M, Ehrnstrom BO, Forsen B, Isomaa B, Snickars B, Taskinen MR. Metabolic consequences of a family history of NIDDM (the Botnia study): evidence for sex-specific parental effects. *Diabetes* 1996;45:1585–1593
 44. Perseghin G, Price TB, Petersen KF, Roden M, Cline GW, Gerow K, Rothman DL, Shulman GI. Increased glucose transport-phosphorylation and muscle glycogen synthesis after exercise training in insulin-resistant subjects. *N Engl J Med* 1996;335:1357–1362

45. Pendergrass M, Bertoldo A, Bonadonna R, Nucci G, Mandarino L, Cobelli C, DeFronzo RA. Muscle glucose transport and phosphorylation in type 2 diabetic, obese nondiabetic, and genetically predisposed individuals. *Am J Physiol Endocrinol Metab* 2007;292:E92–E100
46. Rothman DL, Magnusson I, Cline G, Gerard D, Kahn CR, Shulman RG, Shulman GI. Decreased muscle glucose transport/phosphorylation is an early defect in the pathogenesis of non-insulin-dependent diabetes mellitus. *Proc Natl Acad Sci U S A* 1995;92:983–987
47. Weyer C, Bogardus C, Mott DM, Pratley RE. The natural history of insulin secretory dysfunction and insulin resistance in the pathogenesis of type 2 diabetes mellitus. *J Clin Invest* 1999;104:787–794
48. Ferrannini E, Natali A, Bell P, Cavallo-Perin P, Lalic N, Mingrone G. Insulin resistance and hypersecretion in obesity: European Group for the Study of Insulin Resistance (EGIR). *J Clin Invest* 1997;100:1166–1173
49. Tripathy D, Eriksson KF, Orho-Melander M, Fredriksson J, Ahlqvist G, Groop L. Parallel manifestation of insulin resistance and beta cell decompensation is compatible with a common defect in type 2 diabetes. *Diabetologia* 2004;47:782–793
50. Del Prato S, Leonetti F, Simonson DC, Sheehan P, Matsuda M, DeFronzo RA. Effect of sustained physiologic hyperinsulinemia and hyperglycemia on insulin secretion and insulin sensitivity in man. *Diabetologia* 1994;37:1025–1035
51. Koopmans SJ, Ohman L, Haywood JR, Mandarino LJ, DeFronzo RA. Seven days of euglycemic hyperinsulinemia induces insulin resistance for glucose metabolism but not hypertension, elevated catecholamine levels, or increased sodium retention in conscious normal rats. *Diabetes* 1997;46:1572–1578
52. Koopmans SJ, Kushwaha RS, DeFronzo RA. Chronic physiologic hyperinsulinemia impairs suppression of plasma free fatty acids and increases de novo lipogenesis but does not cause dyslipidemia in conscious normal rats. *Metabolism* 1999;48:330–337
53. Iozzo P, Pratipanawatr T, Pijl H, Vogt C, Kumar V, Pipek R, Matsuda M, Mandarino LJ, Cusi KJ, DeFronzo RA. Physiological hyperinsulinemia impairs insulin-stimulated glycogen synthase activity and glycogen synthesis. *Am J Physiol Endocrinol Metab* 2001;280:E712–E719
54. Cusi K, Maezono K, Osman A, Pendergrass M, Patti ME, Pratipanawatr T, DeFronzo RA, Kahn CR, Mandarino LJ. Insulin resistance differentially affects the PI 3-kinase- and MAP kinase-mediated signaling in human muscle. *J Clin Invest* 2000;105:311–320
55. Bouzakri K, Roques M, Gual P, Espinosa S, Guebre-Egziabher F, Riou JP, Laville M, Le Marchand-Brustel Y, Tanti JF, Vidal H. Reduced activation of phosphatidylinositol-3 kinase and increased serine 636 phosphorylation of insulin receptor substrate-1 in primary culture of skeletal muscle cells from patients with type 2 diabetes. *Diabetes* 2003;52:1319–1325
56. Morino K, Petersen KF, Dufour S, Befroy D, Frattini J, Shatzkes N, Neschen S, White MF, Bilz S, Sono S, Pypaert M, Shulman GI. Reduced mitochondrial density and increased IRS-1 serine phosphorylation in muscle of insulin-resistant offspring of type 2 diabetic parents. *J Clin Invest* 2005;115:3587–3593
57. Morino K, Petersen KF, Dufour S, Befroy D, Frattini J, Shatzkes N, Neschen S, White MF, Bilz S, Sono S, Pypaert M, Shulman GI. Reduced mitochondrial density and increased IRS-1 serine phosphorylation in muscle of insulin-resistant offspring of type 2 diabetic parents. *J Clin Invest* 2005;115:3587–3593
58. Petersen KF, Dufour S, Shulman GI. Decreased insulin-stimulated ATP synthesis and phosphate transport in muscle of insulin-resistant offspring of type 2 diabetic parents. *PLoS Med* 2005;2:e233
59. Hotamisligil GS, Peraldi P, Budavari A, Ellis R, White MF, Spiegelman BM. IRS-1-mediated inhibition of insulin receptor tyrosine kinase activity in TNF- α - and obesity-induced insulin resistance. *Science* 1996;271:665–668
60. Jiang ZY, Lin YW, Clemont A, Feener EP, Hein KD, Igarashi M, Yamauchi T, White MF, King GL. Characterization of selective resistance to insulin signaling in the vasculature of obese Zucker (fa/fa) rats. *J Clin Invest* 1999;104:447–457
61. Bajaj M, Suraamornkul S, Kashyap S, Cusi K, Mandarino L, DeFronzo RA. Sustained reduction in plasma free fatty acid concentration improves insulin action without altering plasma adipocytokine levels in subjects with strong family history of type 2 diabetes. *J Clin Endocrinol Metab* 2004;89:4649–4655
62. Kelley DE. Skeletal muscle fat oxidation: timing and flexibility are everything. *J Clin Invest* 2005;115:1699–702
63. Mootha VK, Lindgren CM, Eriksson KF, Subramanian A, Sihag S, Lehar J, Puigserver P, Carlsson E, Ridderstrale M, Laurila E, Houstis N, Daly MJ, Patterson N, Mesirov JP, Golub TR, Tamayo P, Spiegelman B, Lander ES, Hirschhorn JN, Altshuler D, Groop LC. PGC- α -responsive genes involved in oxidative phosphorylation are coordinately down-regulated in human diabetes. *Nat Genet* 2003;34:267–273
64. Patti ME, Butte AJ, Crunkhorn S, Cusi K, Berria R, Kashyap S, Miyazaki Y, Kohane I, Costello M, Saccone R, Landaker EJ, Goldfine AB, Mun E, DeFronzo R, Finlayson J, Kahn CR, Mandarino LJ. Coordinated reduction of genes of oxidative metabolism in humans with insulin resistance and diabetes: potential role of PGC1 and NRF1. *Proc Natl Acad Sci U S A* 2003;100:8466–8471
65. Short KR, Nair KS, Stump CS. Impaired mitochondrial activity and insulin-resistant offspring of patients with type 2 diabetes. *N Engl J Med* 2004;350:2419–2421
66. Morino K, Neschen S, Bilz S, Sono S, Tsigotis D, Reznick RM, Moore I, Nagai Y, Samuel V, Sebastian D, White M, Philbrick W, Shulman GI. Muscle-specific IRS-1 Ser \rightarrow Ala transgenic mice are protected from fat-induced insulin resistance in skeletal muscle. *Diabetes* 2008;57:2644–2651
67. Abdul-Ghani MA, Muller FL, Liu Y, Chavez AO, Balas B, Zuo P, Chang Z, Tripathy D, Jani R, Molina-Carrion M, Monroy A, Folli F, Van Remmen H, DeFronzo RA. Deleterious action of FA metabolites on ATP synthesis: possible link between lipotoxicity, mitochondrial dysfunction, and insulin resistance. *Am J Physiol Endocrinol Metab* 2008;295:E678–E685