

Brief report

Mutation of the β 1-tubulin gene associated with congenital macrothrombocytopenia affecting microtubule assemblyShinji Kunishima,¹ Ryoji Kobayashi,² Tomohiko J. Itoh,³ Motohiro Hamaguchi,¹ and Hidehiko Saito⁴¹Department of Hemostasis and Thrombosis, Clinical Research Center, National Hospital Organization Nagoya Medical Center, Nagoya, Japan; ²Department of Pediatrics, Hokkaido University Graduate School of Medicine, Sapporo, Japan; ³Division of Biological Sciences, Graduate School of Science, Nagoya University, Nagoya, Japan; and ⁴Nagoya Central Hospital, Nagoya, Japan

Congenital macrothrombocytopenia is a genetically heterogeneous group of rare disorders. We identified the first *TUBB1* mutation, R318W, in a patient with congenital macrothrombocytopenia. The patient was heterozygous for Q43P, but this single-nucleotide polymorphism (SNP) did not relate to macrothrombocytopenia. Although no abnormal platelet β 1-tubulin localization/marginal band organization was observed, the level of β 1-tubulin was

decreased by approximately 50% compared with healthy controls. Large and irregular bleb protrusions observed in megakaryocytes derived from the patient's peripheral blood CD34⁺ cells suggested impaired megakaryocyte fragmentation and release of large platelets. In vitro transfection experiments in Chinese hamster ovary (CHO) cells demonstrated no incorporation of mutant β 1-tubulin into microtubules, but the formation of punc-

tuated insoluble aggregates. These results suggested that mutant protein is prone to aggregation but is unstable within megakaryocytes/platelets. Alternatively, mutant β 1-tubulin may not be transported from the megakaryocytes into platelets. W318 β 1-tubulin may interfere with normal platelet production, resulting in macrothrombocytopenia. (Blood. 2009; 113:458-461)

Introduction

Congenital macrothrombocytopenia is a genetically heterogeneous group of rare disorders.¹⁻³ The most frequent forms include *MYH9* disorders, such as May-Hegglin anomaly, and Bernard-Soulier syndrome. In approximately half of the cases the pathogenesis remains unknown; thus, a definite diagnosis is not possible. The linkage between the membrane skeleton and cytoskeletal actin filaments as well as the marginal microtubule band maintains normal platelet morphology.^{4,5} Defects in these systems may result in macrothrombocytopenia. The microtubules are assembled from α - and β -tubulin heterodimers. β 1-Tubulin expression is restricted in the megakaryocyte/platelet lineage.⁶ *Tubb1* knockout mice show thrombocytopenia and spherical platelets.⁷ *TUBB1* Q43P functional polymorphism has been reported. However, it may not be relevant to macrothrombocytopenia.⁸ We identified the first *TUBB1* mutation affecting microtubule assembly in congenital macrothrombocytopenia.

GPIb/IX. An initial bone marrow examination revealed normal megakaryocyte number and morphology. The mother of the patient also had macrothrombocytopenia. Peripheral blood samples were obtained after the mother gave informed consent in accordance with the Declaration of Helsinki for the study, which was approved by the institutional review boards (IRBs) of Nagoya Medical Center and Hokkaido University. Platelet size and *TUBB1* gene frequencies were determined in 108 healthy controls and 16 consecutive, unrelated patients with macrothrombocytopenia who were enrolled in an institutional review board–approved collaborative study on congenital thrombocytopenia.

Antibodies

We raised an anti- β 1-tubulin antibody (NB2301) in rabbits against a synthetic peptide corresponding to C-terminal 425-451aa of human β 1-tubulin (KAVLEEDDEEVTEEAEMEPEDKGH). Antiserum was collected and affinity-purified. NB2301 specifically reacted with recombinant β 1-tubulin among 7 known recombinant human β -tubulin isoforms on immunoblots, and only stained platelets on blood smears (Figure S1, available on the *Blood* website; see the Supplemental Materials link at the top of the online article). Other antibodies used were anti-GPIIb (SZ22, Immunotech, Marseille, France), anti- α -tubulin (DM1A and RB9281, LabVision, Fremont, CA), anti- β -tubulin (TUB2.1), and anti- β 5-tubulin (SAP.4G5, Abcam, Cambridge, United Kingdom).

Mutational analysis

The entire coding sequence of exons and exon-intron boundaries of *TUBB1* was amplified by PCR, and the products were subjected to DNA sequence analysis (Table S1).

Methods

Patient

The patient was a 7-year-old boy who was incidentally found to have thrombocytopenia (platelets, $40\text{-}60 \times 10^9/\text{L}$). He was diagnosed with immune thrombocytopenic purpura. Peripheral blood smears showed the prominent appearance of giant platelets. Electron microscopy showed no other abnormalities (Figure 1A,B). There were no leukocyte inclusion bodies, confirmed by myosin IIA localization.⁹ The platelets aggregated normally with adenosine diphosphate (ADP), collagen, and ristocetin. Flow cytometry showed normal expression of platelet


Submitted June 11, 2008; accepted September 17, 2008. Prepublished online as *Blood* First Edition paper, October 10, 2008; DOI 10.1182/blood-2008-06-162610.

The online version of this article contains a data supplement.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2009 by The American Society of Hematology

Figure 1. Platelet morphology and the *TUBB1* R318W mutation. (A) Peripheral blood smears were stained with May-Grünwald Giemsa for a normal control (*TUBB1* 43QQ/307RR) and the patient (original magnification $\times 1000$). The patient showed giant platelets with morphologically normal leukocytes. The number in each panel shows the mean platelet size in microns ($n = 200$). (B) Ultrastructure of platelets. Platelet-rich plasma prepared from acid-citrate-dextrose citrated whole blood was fixed in 2% glutaraldehyde and postfixed in 1% osmium tetroxide. The platelet samples were embedded in epoxy resin (Quetol 651; Nissin EM, Tokyo, Japan). Ultra-thin sections were cut, stained with uranyl acetate and lead citrate, and observed under a transmission electron microscope (JEM 1011, JEOL, Tokyo, Japan). Original magnification $\times 8000$. (C) The entire coding regions of the patient's *TUBB1* gene were amplified from genomic DNA by polymerase chain reaction (PCR) and amplified DNA fragments were subjected to direct cycle sequence analysis. A C-to-T transition at nucleotide 952 (952C>T), changing Arg318 to Trp (R318W), was detected. The arrow shows the position of the substitution. (D) β -Tubulin sequence alignment. Amino acid sequence alignment is shown for the 7 human β -tubulin isoforms and the known $\beta 1$ -tubulin from other species. R318 is conserved in other species and other human β -tubulin isoforms. The substituted amino acid is indicated in bold. (E) Structural analysis of R318 using a tubulin 3D model. (i) Position of R318 (red) in the whole tubulin molecule. The residue is almost buried beneath the taxol pocket of β -tubulin but exposed slightly. Blue indicates β -tubulin; gray, α -tubulin. (ii) R318 (orange) is located near the α and β intradimer interface. D249 (yellow) is located at the interface. GTP (magenta) bound on the N-site of α -tubulin is also shown. (iii), Residues surrounding the side chain of R318. All the figures are depicted from 1JFF with the aid of MacPyMol (DeLano Scientific, Palo Alto, CA; <http://www.delanoscientific.com/>).



TUBB1 cloning, mutagenesis, and transfection

Full-length *TUBB1* sequences were amplified from the patient's platelet cDNA (Table S1). *TUBB1* cDNA in-frame with the C-terminal myc epitope tag was cloned into pcDNA3.1 (Invitrogen, Carlsbad, CA). We prepared 3 mutant constructs (P43, W318, and P43/W318), and used them to transfect Chinese hamster ovary (CHO) cells. At 24 hours after transfection, the cells were replated on fibronectin-coated (10 $\mu\text{g}/\text{mL}$) chamber slides, and cultured for 48 hours. For microtubule-depolymerization experiments, cells were incubated at 4°C for 4 hours or 33 mM nocodazole at 37°C for 2 hours.

Analysis of cultured megakaryocytes

CD34⁺ cells isolated from 10 mL of peripheral blood were cultured for 14 days in serum-free medium supplemented with thrombopoietin (TPO), stem cell factor (SCF), and interleukin-3 (IL-3).¹⁰ Cytospin preparations of differentiated megakaryocytes were analyzed by immunofluorescence staining.¹⁰

Results and discussion

We searched for *TUBB1* mutations in the patient, and found a novel conserved R318W mutation (Figure 1C,D). Restriction analysis confirmed heterozygosity in the patient and mother. The mutation was not found in 108 healthy controls or in the SNP database (<http://www.ncbi.nlm.nih.gov/SNP/>). In a tubulin 3D model, it is

buried inside the molecule and is near the α and β intradimer interface (Figure 1E).¹¹ The side chain is oriented toward the inside of the molecule and faces that of E27. An R318W mutation replacing the charged arginine with bulky aromatic tryptophan will disrupt the side chain interactions. Because D249N, located at the interface, has recently been reported to cause canine macrothrombocytopenia,^{12,13} mutations disrupting the structure of the intradimer interface may affect platelet morphology.

Two other nonsynonymous substitutions, Q43P and R307H, were identified. The former was recently described as a functional polymorphism.^{8,14} The prevalence of the P43 allele was higher in patients with congenital macrothrombocytopenia (0.12) than controls (0.06).⁸ We also found a similar gene frequency of 0.16 and 0.06 in our patients and controls, respectively (Table S2). Although R307 is conserved, this position contains basic histidine in other human β -tubulins, and the gene frequency was the same in patients and controls. Control individuals with Q43P or R307H even in their homozygous state had normal-sized platelets (Table S3). We therefore suggested that both substitutions are not related to macrothrombocytopenia.

The expression and localization of platelet $\beta 1$ -tubulin were investigated using the newly produced NB2301 antibody. $\beta 1$ -Tubulin was normally localized in the marginal microtubule band in resting patient platelets (Figure 2A). Immunoblotting showed the normal electrophoretic mobility. However, the expression level was decreased: the




Figure 2. Biochemical analysis of β 1-tubulin in platelets and transfected cells. (A) Immunofluorescence analysis of platelets. After being fixed in methanol and permeabilized in acetone, the peripheral blood smears were hydrated with PBS and blocked with normal goat serum. Slides were concomitantly incubated with DM1A and NB2301, and then reacted with Alexa 555-labeled anti-mouse IgG and Alexa 488-labeled anti-rabbit IgG (Invitrogen). The stained cells were examined under a BX50 fluorescence microscope with a $100\times/1.35$ numeric aperture oil objective (Olympus, Tokyo, Japan). Image slides were taken with a DP70 digital camera using DP Manager software (Olympus). (B) Immunoblot analysis of platelets. Whole platelet proteins were isolated using a NucleoSpin RNA/Protein kit (Macherey-Nagel, Düren, Germany), separated by sodium dodecyl sulfate–polyacrylamide gel electrophoresis on 4% to 12% gradient acrylamide slab gels (Invitrogen), and electroblotted onto polyvinylidene difluoride membranes. The blots were incubated with RB9281 and NB2301, and reacted with horseradish peroxidase-conjugated secondary antibody. The bound antibodies were visualized using an enhanced chemiluminescent substrate. Densitometric analysis of blots performed using ImageQuant software (Molecular Dynamics, Sunnyvale, CA) showed that the β 1-tubulin/ α -tubulin ratio was reduced by 50% in the patient compared with controls (0.75 vs 1.57 ± 0.11 ; $n = 7$). (C) Megakaryocytes derived from the peripheral blood CD34⁺ cells of a normal control and the patient were doubly stained with anti-GPIIb mouse monoclonal antibody SZ22 (not shown) and NB2301 (green). Although the distribution of β 1-tubulin in the patient's megakaryocytes was normal, irregular and large bleb protrusions were observed in the patient. (D) Immunofluorescence analysis of transfected CHO cells. *TUBB1*-myc transfected CHO cells grown on chamber slides were doubly stained with anti-myc tag antibody and DM1A, and reacted with Alexa 488-labeled anti-rabbit IgG and Alexa 555-labeled anti-mouse IgG. Note that W318 mutants were only localized as small punctuate structures. (E) *TUBB1*-myc transfected CHO cells grown on chamber slides were exposed to 4°C for 4 hours to facilitate the depolymerization of microtubules, and then fixed and stained. Aggregates of β 1-tubulin W318 did not dissociate under conditions of microtubule depolymerization. (F) Low temperature-exposed *TUBB1*-myc transfected CHO cells were lysed in 10 mM Pipes, pH 6.8, 1 mM MgCl₂, 1 mM EGTA, 10% dimethyl sulfoxide, 0.5% Nonidet P-40, and Complete Protease Inhibitor Cocktail (Roche Diagnostics, Mannheim, Germany), and incubated for 30 minutes at 37°C. Cell lysates were centrifuged at 1000g for 10 minutes at 37°C before supernatants and pellets were collected for immunoblot analysis with anti-myc tag antibody. Densitometric analysis of the blots showed that a large proportion of β 1-tubulin W318 was recovered in the pellets. The arrowhead indicates recombinant myc-tagged β 1-tubulin.

β 1-tubulin/ α -tubulin ratio was decreased by approximately 50% compared with controls (Figure 2B). Q43P is reportedly associated with the reduced platelet β 1-tubulin.⁸ We found in a control P43 homozygote that the β 1-tubulin/ α -tubulin ratio was not decreased (Figure 2B). Thus, the decreased expression of β 1-tubulin in the patient is likely to be due to R318W and not Q43P. We examined megakaryocytes cultured from CD34⁺ peripheral blood cells. Cultured mature megakaryocytes frequently extend blebs, similar to demarcation membrane system-like structure.^{15,16} We observed large and irregular bleb protrusions in the

patient (Figure 2C). This suggested impaired megakaryocyte fragmentation and release of large platelets. The plasma glycosialin concentration, a proteolytic fragment of GPIIb α (molecular marker of platelet production/destruction),¹⁷ was below the normal limit (1.02 μ g/mL; normal values, 1.40 ± 0.25 μ g/mL), indicating that thrombocytopenia is not due to underproduction of platelets but peripheral destruction.

To determine the functional and structural consequences of the R318W mutation on microtubule assembly, we monitored mutant β 1-tubulins after their transfection into CHO cells. The wild-type

and P43 mutant myc-tagged β 1-tubulin were localized as fine filamentous cytoplasmic networks, indicating normal incorporation of recombinant β 1-tubulin into microtubules with endogenous α -tubulin. In contrast, W318 and P43/W318 mutants accumulated in punctuate structures in the cytoplasm (Figure 2D). Under conditions of microtubule depolymerization by low temperature (Figure 2E,F) or nocodazole treatment, aggregates of W318 mutants did not dissociate.

Our results suggested that the net effect of the *TUBB1* R318W mutation in platelets is the instability of mutant protein. These data seem to contradict those obtained by in vitro expression experiments, whereby β 1-tubulin W318 formed insoluble aggregates. One important difference is that the latter employs an expression vector with strong cytomegalovirus (CMV) promoter. Thus, the unstable but aggregate-prone W318 mutant was detected as insoluble aggregates. The coordinated production of α - and β -tubulins necessary for proper heterodimer assembly are regulated both transcriptionally and translationally and, when tubulin is overexpressed in cells, synthesis of the endogenous form is strongly inhibited.¹⁸⁻²⁰ In *Tubb1* knockout mice, up-regulation of β 2- and β 5-tubulins partially compensates for the null expression of β 1-tubulin.⁷ Because heterozygous *Tubb1* knockout mice do not exhibit giant platelets, simple haploinsufficiency alone cannot explain the pathogenesis of macrothrombocytopenia. Mutant β 1-tubulin may dominantly affect microtubule assembly in some manner. Normal expression of mutant *TUBB1* mRNA (not shown) and decreased β 1-tubulin in platelets lead to an alternative hypothesis that mutant β 1-tubulin is not transported from the megakaryocyte into platelets.

Elucidating the precise molecular mechanisms will not only advance our classification and diagnosis of congenital macrothrombocytopenias, but also our comprehensive understanding of megakaryopoiesis and platelet genesis.

Acknowledgments

We thank Ms Yoshimi Ito for her skillful technical assistance.

This work was supported by grants to S.K. from the Japan Society for the Promotion of Science (18591094 and 20591161), the Ministry of Health, Labor and Welfare (Grant for Child Health and Development 19C-2), Charitable Trust Laboratory Medicine Foundation of Japan, and National Hospital Organization (network research grant for congenital thrombocytopenia).

Authorship

Contributions: S.K. designed and performed research, analyzed data, and wrote the paper; R.K. contributed patient samples and clinical data; T.J.I. designed the tubulin experiments and wrote the paper; and M.H. and H.S. supervised the research.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Shinji Kunishima, Department of Advanced Diagnosis, Clinical Research Center, National Hospital Organization Nagoya Medical Center 4-1-1 Sannomaru, Naka-ku, Nagoya 4600001, Japan. E-mail: kunishis@nnh.hosp.go.jp.

References

- Balduini CL, Cattaneo M, Fabris F, et al. Inherited thrombocytopenias: a proposed diagnostic algorithm from the Italian Gruppo di Studio delle Piastri. *Haematologica*. 2003;88:582-592.
- Kunishima S, Saito H. Congenital macrothrombocytopenias. *Blood Rev*. 2006;20:111-121.
- Nurden P, Nurden AT. Congenital disorders associated with platelet dysfunctions. *Thromb Haemost*. 2008;99:253-263.
- Italiano JE Jr, Bergmeier W, Tiwari S, et al. Mechanisms and implications of platelet discoid shape. *Blood*. 2003;101:4789-4796.
- Schulze H, Shivdasani RA. Mechanisms of thrombopoiesis. *J Thromb Haemost*. 2005;3:1717-1724.
- Lecine P, Italiano JE Jr, Kim SW, Villeval JL, Shivdasani RA. Hematopoietic-specific β 1 tubulin participates in a pathway of platelet biogenesis dependent on the transcription factor NF-E2. *Blood*. 2000;96:1366-1373.
- Schwer HD, Lecine P, Tiwari S, Italiano JE Jr, Hartwig JH, Shivdasani RA. A lineage-restricted and divergent β -tubulin isoform is essential for the biogenesis, structure and function of blood platelets. *Curr Biol*. 2001;11:579-586.
- Freson K, De Vos R, Wittevrongel C, et al. The *TUBB1* Q43P functional polymorphism reduces the risk of cardiovascular disease in men by modulating platelet function and structure. *Blood*. 2005;106:2356-2362.
- Kunishima S, Matsushita T, Kojima T, et al. Immunofluorescence analysis of neutrophil nonmuscle myosin heavy chain-A in *MYH9* disorders: association of subcellular localization with *MYH9* mutations. *Lab Invest*. 2003;83:115-122.
- Kunishima S, Hamaguchi M, Saito H. Differential expression of wild-type and mutant NMMHC-IIA polypeptides in blood cells suggests cell-specific regulation mechanisms in *MYH9* disorders. *Blood*. 2008;111:3015-3023.
- Lowe J, Li H, Downing KH, Nogales E. Refined structure of α beta-tubulin at 3.5 angstrom resolution. *J Mol Biol*. 2001;313:1045-1057.
- Cowan SM, Bartges JW, Gompf RE, et al. Giant platelet disorder in the Cavalier King Charles spaniel. *Exp Hematol*. 2004;32:344-350.
- Davis B, Toivio-Kinnucan M, Schuller S, Boudreaux MK. Mutation in beta1-tubulin correlates with macrothrombocytopenia in Cavalier King Charles spaniels. *J Vet Intern Med*. 2008;22:540-545.
- Navarro-Nunez L, Lozano ML, Rivera J, et al. The association of the beta1-tubulin Q43P polymorphism with intracerebral hemorrhage in men. *Haematologica*. 2007;92:513-518.
- De Botton S, Sabri S, Daugas E, et al. Platelet formation is the consequence of caspase activation within megakaryocytes. *Blood*. 2002;100:1310-1317.
- Lepage A, Leboeuf M, Cazenave JP, de la Salle C, Lanza F, Uzan G. The α IIb β 3 integrin and GPIb-V-IX complex identify distinct stages in the maturation of CD34+ cord blood cells to megakaryocytes. *Blood*. 2000;96:4169-4177.
- Kunishima S, Kobayashi S, Takagi A, Naoe T, Ohno R. Rapid detection of plasma glycofocalin by a latex agglutination test: a useful adjunct in the differential diagnosis of thrombocytopenia. *Am J Clin Pathol*. 1993;100:579-584.
- Cleveland DW. Autoregulated instability of tubulin mRNAs: a novel eukaryotic regulatory mechanism. *Trends Biochem Sci*. 1988;13:339-343.
- Gonzalez-Garay ML, Cabral F. Overexpression of an epitope-tagged beta-tubulin in Chinese hamster ovary cells causes an increase in endogenous α -tubulin synthesis. *Cell Motil Cytoskeleton*. 1995;31:259-272.
- Gonzalez-Garay ML, Cabral F. alpha-Tubulin limits its own synthesis: evidence for a mechanism involving translational repression. *J Cell Biol*. 1996;135:1525-1534.