

Clinical and molecular predictors of thrombocytopenia and risk of bleeding in patients with von Willebrand disease type 2B: a cohort study of 67 patients

Augusto B. Federici,¹ Pier M. Mannucci,¹ Giancarlo Castaman,² Luciano Baronciani,¹ Paolo Bucciarelli,¹ Maria T. Canciani,¹ Alessandro Pecci,³ Peter J. Lenting,⁴ and Philip G. De Groot⁴

¹Angelo Bianchi Bonomi Hemophilia Thrombosis Center, Department of Medicine and Medical Specialties, Istituti di Ricovero e Cura a Carattere Scientifico (IRCCS) Maggiore Hospital, Mangiagalli and Regina Elena Foundation, and University of Milan, Milan, Italy; ²Department of Hematology, Hemophilia Thrombosis Center, San Bortolo Hospital, Vicenza, Italy; ³Department of Internal Medicine, IRCCS Policlinico S Matteo Foundation and University of Pavia, Pavia, Italy; and ⁴Department of Clinical Chemistry and Hematology, University of Utrecht, Utrecht, The Netherlands

Type 2B von Willebrand disease (VWD2B) is caused by an abnormal von Willebrand factor (VWF) with increased affinity for the platelet receptor glycoprotein Ib- α (GPIb- α) that may result in moderate to severe thrombocytopenia. We evaluated the prevalence and clinical and molecular predictors of thrombocytopenia in a cohort of 67 VWD2B patients from 38 unrelated families characterized by VWF mutations. Platelet count, mean platelet volume, and morphologic evaluations of blood smear were obtained at baseline

and during physiologic (pregnancy) or pathologic (infections, surgeries) stress conditions. Thrombocytopenia was found in 20 patients (30%) at baseline and in 38 (57%) after stress conditions, whereas platelet counts were always normal in 16 patients (24%) from 5 families carrying the P1266L/Q or R1308L mutations. VWF in its GPIb- α -binding conformation (VWF-GPIb- α /BC) was higher than normal in all except the 16 cases without thrombocytopenia (values up to 6-fold higher than controls). The risk of bleeding was higher

in patients with thrombocytopenia (adjusted hazard ratio = 4.57; 95% confidence interval, 1.17-17.90) and in those with the highest tertile of bleeding severity score (5.66; 95% confidence interval, 1.03-31.07). Prediction of possible thrombocytopenia in VWD2B by measuring VWF-GPIb- α /BC is important because a low platelet count is an independent risk factor for bleeding. (Blood. 2009;113:526-534)

Introduction

Type 2B von Willebrand disease (VWD2B) is a rare autosomally inherited bleeding disorder (5%-8% of all VWD types), typically characterized by enhanced ristocetin-induced platelet aggregation (RIPA) in platelet-rich plasma (PRP).¹ In VWD2B, von Willebrand factor (VWF) is normally synthesized and assembled by endothelial cells but has heightened platelet affinity when released into the circulation.² Mutations, clustered in exon 28 of the *VWF* gene encoding the VWF A1 domain involved in VWF binding to the platelet receptor glycoprotein Ib alpha (GpIb- α), are responsible for this VWD2B defect.³⁻⁵ The gain of function of the VWF A1 domain creates the following conditions that may explain the bleeding tendency in these patients: (1) the high molecular weight (HMW), hemostatically most effective, VWF multimers are decreased in a patient's plasma; (2) thrombocytopenia may occur; and (3) platelets' GpIb- α receptors may be occupied by soluble VWF and become unable to interact with immobilized VWF exposed at the sites of injury.⁶ The plasma defect of the largest VWF multimers is thought to be the result of their removal from the circulation, consistent with the normal pattern in platelets and their transient appearance in plasma after 1-8-deamino-D-arginine vasopressin (DDAVP) administration.^{7,8} A wide degree of heterogeneity has

been found in the VWF multimeric patterns in VWD2B,⁹ ranging from the complete loss of the largest multimers to a normal pattern in the VWD2B variant previously known as type 1 Malmö or New York.¹⁰⁻¹²

VWD2B patients who lack the HMW multimers may also have moderate to severe thrombocytopenia,^{13,14} sometimes associated with giant platelets and spontaneous platelet aggregates,¹⁵⁻¹⁷ and this can predispose to more severe bleeding. Thrombocytopenia in VWD2B can be overlooked, especially in children¹⁸⁻²⁰; it can be transient or persistent with variable platelet count on different occasions but usually worsens after conditions characterized by a long- or short-term rise in plasma VWF such as pregnancy,^{21,22} infections, surgeries,²³ or DDAVP administration.²⁴ The thrombocytopenia phenotype is mainly caused by the enhanced 2BVWF-GpIb- α interactions resulting from gain of function of the A1 domain, leading to spontaneous platelet aggregation *in vitro* and *in vivo*.^{15,16} More recently, abnormal megakaryocytopoiesis with severe thrombocytopenia was described in a family with VWD2B characterized by the R1308P substitution, suggesting a major role of the enhanced 2BVWF-GpIb- α interactions in platelet formation and release.²⁵

No large studies on the effect of thrombocytopenia on the clinical manifestations of VWD2B patients have been performed

Submitted April 17, 2008; accepted August 23, 2008. Prepublished online as *Blood* First Edition paper, September 19, 2008; DOI 10.1182/blood-2008-04-152280.

An Inside *Blood* analysis of this article appears at the front of this issue.

Preliminary data were presented in part at the annual meeting of the American Society of Hematology, Orlando, FL, December 9-16, 2006, and discussed at

the International Society of Thrombosis and Hemostasis-Scientific and Standardization Committee Subcommittee on von Willebrand Factor, Vienna, Austria, July 4, 2008.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2009 by The American Society of Hematology

so far. To determine the prevalence of thrombocytopenia, the associated clinical and molecular markers, and to assess the risk of bleeding associated with this complication, we have followed for 2 years 67 patients with identified VWF2B mutations. Genotypes and VWF measurements were related to platelet count and morphology. In these patients, thrombocytopenia was also evaluated after the occurrence of stress conditions, such as pregnancy, infection, and surgery, and was correlated with the frequency of bleeding episodes. Another aim of this study was to confirm in a large number of well-characterized patients whether or not VWF in its GPIb- α -binding conformation (VWF-GPIb- α /BC) measured by a nanobody²⁶ helps to predict possible thrombocytopenia in VWD2B.

Methods

Clinical assessment of VWD2B patients

A cohort of 67 VWD2B patients regularly followed up by Hemophilia Centers in Milan and Vicenza (Italy) and Utrecht (The Netherlands) were evaluated at baseline and then prospectively at 3-month intervals during a follow-up period of 24 months (between October 1, 2005, and October 1, 2007).^{27,28} At baseline, a questionnaire originally devised to determine a bleeding severity score in VWD type 1 was administered to all patients.²⁹ During follow-up, all patients were investigated during physiologic and pathologic stress conditions that might trigger thrombocytopenia by increasing plasma levels of VWF2B. Bleeding episodes were also recorded. A study of family members was also performed to evaluate phenotype heterogeneity within the same pedigree in 4 families characterized by the following VWD2B mutations: P1266L, R1306W, R1308L, and I1309V (data not shown).

Blood collection and laboratory methods

Patients who were informed about the anonymous uses of their data and on the purpose of the study gave informed consent in accordance with the Declaration of Helsinki. This study received Institutional Review Board approval from all 3 participating centers. Venous blood was obtained at each Hemophilia Center at baseline and at the time of visits during follow-up. Plasma samples were prepared and kept frozen at -40°C . The skin bleeding time was measured by the Simplate-II (Organon Teknika, Durham, NC), factor VIII (FVIII) by one-stage coagulometric assay, VWF: antigen (VWF:Ag) by commercial kits, or homemade enzyme-linked immunosorbent assay using anti-VWF polyclonal antibodies. VWF Ristocetin cofactor activity (VWF:RCo) was measured by aggregometry using formalin-fixed platelets from normal donors; when plasma VWF:RCo was less than 15 U/dL, VWD2B patients were further investigated using the more sensitive enzyme-linked immunosorbent assay VWF:RCo.³⁰ RIPA was carried out in PRP by aggregometry, and changes in light transmission were inscribed in a potentiometric chart recorder. RIPA values were expressed as the minimal concentration of ristocetin giving 30% increase in transmission (normal range, 0.8-1.2 mg/mL ristocetin).^{1,28} The multimeric pattern of plasma VWF was evaluated centrally (Milan) by low-resolution gel electrophoresis using 0.9% low gelling temperature agarose.³¹ Four different patterns were identified in VWD2B patients (Figure 1): pattern A, all multimers present, as in normal persons; pattern B, partial loss of the HMW multimers; pattern C, complete loss of HMW multimers; pattern D, loss also of intermediate multimers. All the patients carried a mutation within the A1 domain of VWF. All identified mutations had been previously included in the International Society of Thrombosis and Hemostasis VWF database.³²

Platelet count and morphology

The number of single platelets and their mean platelet volumes were measured by SYSMEX XE-2100-Hemocouler (Sysmex, Kobe, Japan) immediately after blood withdrawal. VWD2B patients were considered as thrombocytopenic if their platelet counts were less than $140 \times 10^3/\mu\text{L}$




Figure 1. The 4 multimeric patterns found in the entire cohort of VWD2B patients. Pattern A indicates all HMW multimers present, as in plasma of normal persons; pattern B, partial loss of the HMW multimers; pattern C, complete loss of HMW multimers; pattern D, loss of high and intermediate multimers. Pattern A is identical to that of normal plasma (NP).

(corresponding to the lower limit of normal range of the method) on at least 3 different measurements. To prevent bias in the morphologic evaluation, blood smears were evaluated blindly (by A.P. who was not aware of the VWD2B mutations). Four different morphologic patterns were identified in the VWD2B patients (Figure 2): pattern A, no abnormalities of platelet size or distribution; pattern B, enlarged platelets, no platelet aggregates; pattern C, platelets forming small aggregates; pattern D, platelets forming large aggregates.

Measurement of VWF in its binding conformation to GPIb- α

The VWF in its binding conformation to GPIb- α was measured in plasma by immunoadsorbent assay based on the nanobody AU/VWFA-11.²⁶ This assay allows quantification of the abnormal VWF that circulates in plasma in a GPIb- α -binding conformation *in vivo*. In brief, AU/VWFA-11-coated microtiter wells were incubated with plasma samples diluted in phosphate-buffered saline. The amount of bound VWF was then measured by horseradish peroxidase-conjugated polyclonal antibodies against VWF. The slope of the initial linear part of the binding curve was calculated for each sample and for reference plasma. The ratio of the slope for the different plasma samples and the slope for reference plasma was subsequently calculated and designated as VWF in its GPIb- α -binding conformation (VWF-GPIb- α /BC). Normal values obtained on plasma samples from




Figure 2. The 4 platelet morphology patterns identified in the VWD2B patients. (A) No abnormalities of platelet size or distribution. (B) Enlarged platelets, no platelet aggregates. (C) Some platelets are associated to form small platelet aggregates. (D) Most platelets are associated to form large platelet aggregates. Microscopy was performed through the Axioscope 2 Plus microscope (Carl Zeiss, Gottingen, Germany), using a $20\times/0.50$ Plan Neofluar or a $63\times/1.25$ Plan Neofluar oil-immersion objective (Carl Zeiss), and the Eukitt mounting medium (Bio-Optica, Milan, Italy). Images were acquired by the AxioCam MRc camera (Carl Zeiss) and the Axiovision 4.5 software (Carl Zeiss).

Table 1. Demographic parameters and VWF measurements of the cohort of VWD 2B patients enrolled in the study and classified by genotype

Mutation	No. of cases (no. of families)	Age, y	Sex, F-M	Bleeding severity score	BT, min	RIPA, mg/mL	VWF:RCo, U/dL	VWF:RCo/Ag, ratio	FVIII, U/dL	Multimeric pattern
P1266Q	4 (2)	34 (19-52)	1-3	0.5 (0-2)	NT	0.6 (0.3-0.6)	32 (21-101)	0.7 (0.7-0.8)	58 (39-96)	A
P1266L	6 (2)	47 (11-84)	3-3	2 (0-6)	5 (3-9)	0.5 (0.4-0.6)	60 (43-105)	0.9 (0.7-1.1)	78 (56-125)	A
H1268D	1 (1)	36	1-0	7	5.5	0.4	24	0.5	62	B
R1306W	15 (9)	43 (14-77)	9-6	8 (3-24)	19 (6-35)	0.7 (0.5-0.8)	12 (7-36)	0.5 (0.25-0.8)	36 (27-75)	C
R1308C	13 (11)	35 (26-87)	10-3	9 (1-18)	15 (10-27)	0.7 (0.5-0.8)	19 (12-39)	0.5 (0.18-0.7)	43 (24-72)	D
R1308L	6 (1)	41 (18-50)	4-2	3 (0-16)	7 (5-8)	0.5 (0.3-0.6)	38 (23-49)	0.8 (0.65-1.0)	47 (35-57)	A
I1309V	9 (2)	55 (10-60)	4-5	4 (2-11)	9 (5-19)	0.4 (0.3-0.7)	65 (23-125)	0.7 (0.6-1.1)	91 (57-140)	B
V1316M	4 (4)	32 (22-62)	2-2	13 (12-16)	31 (28-35)	0.5 (0.5-0.6)	7 (6-20)	0.3 (0.2-0.7)	57	D
P1337L	2 (1)	27 (22-31)	2-0	3 (2-3)	8 (4-11)	0.6 (0.5-0.7)	70 (35-104)	1.0 (0.9-1.1)	75 (74-76)	B
R1341Q	4 (3)	48 (41-75)	3-1	5 (4-7)	8 (6-14)	0.5 (0.4-0.7)	36 (24-59)	0.6 (0.4-0.7)	64 (39-80)	C
R1341W	3 (2)	36 (6-62)	1-2	7 (0-9)	7 (4-9)	0.6 (0.4-0.7)	54 (42-58)	0.6 (0.6-1.0)	73 (59-86)	C
Total	67/38	42 (6-87)	40-27	5 (0-24)	10 (3-35)	0.6 (0.3-0.8)	32 (6-125)	0.7 (0.18-1.1)	57 (24-140)	A-D

Continuous variables are expressed as median (range). NT indicates not tested; and BT, bleeding time.

40 normal persons were 0.70 plus or minus 0.13 (mean ratio \pm SD). Data were also obtained from additional 36 patients with VWD types 1, 2A, or 2M all characterized by VWF mutations outside (VWD types 1 and 2A) or within (VWD type 2M) VWF A1 domain. VWF-GPIb- α /BC values were as follows: 0.30 plus or minus 0.25 in VWD1 ($n = 12$), 0.38 plus or minus 0.31 in VWD2A ($n = 14$), and 2.93 plus or minus 1.75 in VWD2M ($n = 10$). These data are used for comparison with results obtained in VWD2B patients enrolled in the study (for details, see Figure 5A).

Statistical analysis

Continuous variables were expressed as median and range. Comparison between groups was made by χ^2 test, Mann-Whitney U test, or one-way analysis of variance, where appropriate. The monthly incidence of bleeding during follow-up (and its 95% confidence interval [CI]) in the whole cohort and in different subgroups was calculated by dividing the total number of first bleeding episodes by the total number of patient-months. For each patient, the observation period started at October 1, 2005, and ended at October 1, 2007 (if no bleeding episodes occurred) or at the time of the first bleeding episode. Bleeding-free survival according to platelets count at baseline ($\geq 140 \times 10^3/\mu\text{L}$, $< 140 \times 10^3/\mu\text{L}$) and to different tertiles of bleeding severity scores (< 4 , 4-8, > 8) was calculated by the Kaplan-Meier method. The Cox proportional hazard model was performed, which allowed for adjustment for bleeding severity score and presence or absence of thrombocytopenia (1 for each other) and for age, sex, VWF:RCo divided in tertiles (> 40 , 20-40, < 20 IU/dL) and VWF-GPIb- α /BC (< 1 , 1-4, > 4). The final hazard ratio expresses the risk of bleeding of one group compared with the reference, adjusted for the other variables in the model. $P \leq .05$ was taken as cutoff for statistical significance. Statistical analysis was performed using SPSS software package (release 15.0; SPSS, Chicago, IL).

Results

Demographic, genotypic, and phenotypic data of VWD2B patients

In the cohort of 67 VWD2B patients (38 families), 40 (60%) were females and 27 (40%) males. Results of bleeding severity score and VWF measurements are summarized in Table 1 according to the specific VWF gene mutations, all located in exon 28 encoding the VWF A1 domain. The bleeding severity score was higher than 1 (normal values from -1 to 0) in 60 patients (89%). Patients with the P1266Q mutation had the lowest scores (median, 0.5; range, 0-2), those with V1316M the highest (median, 13; range, 12-16). The bleeding time was prolonged (normal value, < 7 minutes) in 54 patients (79%), the lowest values being in P1266L (median, 5; range, 3-9), and the highest in V1316M (median, 31; range, 28-35). RIPA was enhanced in all the patients by diagnostic definition,

whereas levels of VWF:RCo were only moderately reduced and were normal in 12 of 67 (18%) cases. VWF:Ag levels, normal in the majority of the cases, were used to calculate the VWF:RCo/Ag ratios, which were lower in the presence of defects of the HMW multimers. Results of multimeric analyses could be assembled into 4 different patterns (Figure 1). A normal multimeric pattern (pattern A) was found in the 16 patients with mutations P1266L/Q and R1308L. Partial loss of HMW multimers (pattern B) was found in the 12 patients with mutations H1268D, I1309V, and P1337L. Patterns C (complete loss of HMW multimers) and D (loss also of intermediate multimers) were found in 22 (R1306W, R1341Q/W) and 17 (R1308C, V1316M) cases, respectively. Plasma levels of factor VIII were normal or slightly reduced in the great majority of patients.

Platelet count at baseline and during physiologic or pathologic stress situations

At baseline, 20 of 67 patients (30%) had platelet count less than $140 \times 10^3/\mu\text{L}$ on at least 3 different measurements. An inverse relationship between platelet count at baseline and bleeding severity score was observed: the median platelet count was $249 \times 10^3/\mu\text{L}$ (range, 82-402; $n = 21$) in patients with bleeding severity score less than 4, $177 \times 10^3/\mu\text{L}$ (range, 66-251; $n = 24$) in those with bleeding severity score 4 to 8, and $144 \times 10^3/\mu\text{L}$ (range, 19-322; $n = 22$) in those with bleeding severity score more than 8 (Figure 3; 1-way analysis of variance: $P < .001$). During the 24-month follow-up, 31 patients were exposed to stress situations, including pregnancy ($n = 6$), infection ($n = 12$), and surgery




Figure 3. Distribution of platelet counts according to bleeding severity score in the 67 patients enrolled in the study. Bleeding severity score was obtained at enrollment, and platelet counts were measured at baseline. Each box plot represents interquartile range with median value (white line in the middle) and 95% intervals. \circ indicates an outlier.

Table 2. Platelet counts at baseline and after stress situations, platelet morphology, and VWF–GPIb- α /BC (nanobody) of the 67 VWD2B patients classified by genotype

Mutation	No. of cases (no. of families)	Platelet count at baseline ($\times 10^3/\mu\text{L}$)	Trough platelet counts after stress				MPV, fl	Platelet morphology	VWF–GPIb- α :BC
			Pregnancy, n	Infections, n	Surgery, n	DDAVP,* n			
P1266Q	4 (2)	302 (269-353)	199 (1)				8 (8-9)	A	0.7 (0.6-0.8)
P1266L	6 (2)	238 (172-302)			185 (1)	199.0 (2)	8 (8-9)	A	0.5 (0.5-1.1)
H1268D	1 (1)	223		117 (1)	123 (1)		10	B	2.2
R1306W	15 (9)	165 (54-214)	56 (4)	85 (2)	73 (2)		10 (4-11)	B	3.7 (1.9-5.5)
R1308C	13 (11)	165 (93-279)	64 (1)	84 (2)	25 (2)		11 (9-13)	D	2.6 (2-3)
R1308L	6 (1)	248 (193-402)		194 (2)	204 (1)	208.0 (1)	9 (8-10)	A	0.4 (0.3-0.7)
I1309V	9 (2)	111 (66-222)		76 (1)	41 (2)		10 (9-13)	C	2.4 (2.1-3.4)
V1316M	4 (4)	97 (19-149)		21 (1)	27 (2)		10 (10-10)	C	4.3 (4.1-4.4)
P1337L	2 (1)	247 (221-273)			141 (1)		9 (8-10)	A	1.2 (1.1-1.3)
R1341Q	4 (3)	199 (174-229)		105 (2)	73 (1)		10 (8-11)	B	3.0
R1341W	3 (2)	185 (171-319)		131 (1)			9 (8-10)	C	3.3
Total	67 (38)	184 (19-402)	56 (56-199)	85 (21-194)	73 (25-204)	202 (164-208)	10 (8-13)	AtoD	2.2 (0.3-5.5)

Continuous variables are expressed as median (range). MPV indicates mean platelet volume.

*DDAVP was used only in 3 patients with mutations P1266L, previously diagnosed as type 1 Malmö/New York¹⁰ and R1308L.³²

($n = 13$) (Table 2). During the last 3 months of pregnancy, 5 patients with mutations R1306W and R1308C had median values of 56 and 64 $\times 10^3/\mu\text{L}$, respectively, whereas a normal platelet count was found in a pregnant woman with P1266Q. Infectious episodes included viral hepatitis C ($n = 7$), gastroenteritis ($n = 3$), and bacterial infection complicating local hematomas ($n = 2$). Ten patients with mutations H1268D, R1306W, R1308C, I1309V, V1316M, and R1341Q/W developed thrombocytopenia during these episodes with trough values ranging from 21 to 131 $\times 10^3/\mu\text{L}$, whereas 2 patients with the R1308L mutation had no fall in platelet count. Six major surgical procedures were performed in the gastrointestinal tract for cancer or recurrent bleeding for angiodysplasia; caesarian sections, hysterectomy, and tonsillectomy were performed in 4, 2, and 1 case, respectively. Postsurgical thrombocytopenia was observed in 11 of 13 cases; however, platelet count remained normal in 2 patients with the P1266L and R1308L mutations. DDAVP is not recommended in VWD2B,²⁸ and it was therefore given only to 3 patients with P1266L (formerly diagnosed as type 1 Malmö or New York) or R1308L mutations^{10,33}; none of them developed thrombocytopenia after administration of the drug. To sum up, 26 of 34 patients who were exposed to stress situations developed transient thrombocytopenia. None of the 8 patients with P1266Q/L or R1308L mutations exposed to stress situations, including DDAVP infusion, developed thrombocytopenia.

Platelet volume and morphology in relation to VWF–GPIb- α /BC values

Morphologic analysis of peripheral blood smears identified 4 different patterns (Figure 2). Normal pattern (A) was found in the 18 patients with the P1266Q/L, R1308L, or P1337L mutations; pattern B (enlarged platelets without aggregates) was present in the 20 patients with H1268D, R1306W, and R1341Q; patterns C and D, with evidence of small or large platelet aggregates, were found in the 29 patients with R1308C, I1309V, V1316M, and R1341W (Table 2). Platelet aggregates correlated with a lower platelet count at baseline. Enlarged platelets at morphologic analysis were globally present in 24 patients, who corresponded to those who had increased mean platelet volume as measured by automated counters. The most severe thrombocytopenia at baseline and after stress conditions was observed in VWD2B patients carrying V1316M mutation.

VWF–GPIb- α /BC was measured in the VWD2B cases enrolled in the study, all characterized by mutations within VWF A1

domain. The localization of the gain-of-function mutations responsible for the 67 cases of VWD2B are shown in Figure 4: for comparison, also the sites of the loss-of-function mutations responsible for VWD2M are reported. Values of VWF–GPIb- α /BC were high in all but the 16 cases with mutations P1266L/Q, R1308L, with values up to 6 times higher than those of controls (Figure 5A). Interestingly, high VWF–GPIb- α /BC values were also found in 10 VWD2M patients with mutations (delD1277, R1315C/L, Y1321C, R1374H) within VWF A1 domain; on the other hand, values lower than normal were found in VWD types 1 and 2A, characterized by mutations outside VWF A1 domain.

In patients with loss of HMW multimers, values greater than 1.0 were inversely related with thrombocytopenia at baseline (Figure 5B, $R^2 = 0.537$) and predicted the occurrence of thrombocytopenia after exposure to stress. All the 16 patients with the P1266Q/L and R1308L mutations had the same phenotype, ie, normal levels of VWF–GPIb- α /BC, normal platelet count before and after stress, normal platelet morphology, and a normal multimeric pattern.




Figure 4. Localization of the 8 point mutations found in the 67 VWD2B patients within the VWFA1 domain. We are using the modeling derived by the crystal structure of the VWF A1 domain.³⁸ Seven gain-of-function VWD2B point mutations (H1268D, R1306Q, R1308C/L, I309V, V1316H, P1337L, R1341Q/W) are indicated in red, whereas the P1266L/Q is shown in yellow. Note that point mutation R1308 is indicated with 2 different colors according to the different amino acid substitutions: R1308C (red) and R1308L (yellow). For comparison, 4 loss-of-function mutations (delD1277, R1315C/L, Y1321C, R1374H) that were detected in patients with VWD2M are indicated in green. Note that both gain-of-function (VWD2B) and loss-of-function (VWD2M) mutations were characterized by VWF–GPIb- α /BC > 1. Only 2 point mutations, 1 with 2 different amino acid substitutions (P1266L/Q) and the other with 1 specific amino acid (R1308L), are indicated in yellow because they showed VWF–GPIb- α /BC values similar to normal controls (Table 2; Figure 5A).




Figure 5. Values of VWF-GPIIb- α /BC and correlation with platelet count. (A) Distribution of the values of VWF-GPIIb- α /BC as detected by nanobody in 6 different groups of persons (from the left to the right): 40 normal subjects (indicated as normal controls); 12 VWD1 patients with multimeric pattern A (VWD1 HMWM [A]); 14 VWD2A patients with multimeric patterns C or D (VWD2A no HMWM [C-D]); 16 VWD2B patients with P1266Q/L and R1308L mutations with multimeric pattern A (VWD2B HMWM [A]); 51 VWD2B with multimeric patterns B or C or D (VWD2B no HMWM [B, C, D]); 10 VWD2M patients with multimeric pattern B (VWD2M no HMWM [B]). Note that VWF-GPIIb- α /BC is lower than normal in VWD1 and VWD2A with mutations outside VWF A1 domain and is higher than normal in VWD2B and VWD2M with mutations within VWF A1 domain. Interestingly, the gain-of-function P1266Q/L and R1308L mutations characterized by the presence of all the multimers (pattern A) show VWF-GPIIb- α /BC values similar to controls. (B) Relationship between VWF-GPIIb- α /BC values by nanobody (vertical axis) and platelet count at baseline (horizontal axis) in the 67 patients with VWD2B. Patients with normal HMW multimers and P1266Q/L and R1308L mutations have normal values of VWF-GPIIb- α /BC (○), whereas VWD2B patients with loss of HMW multimers and H1268D, R1306Q, R1308C/L, I309V, V1316H, P1337L, and R1341Q/W mutations have increased values of VWF-GPIIb- α /BC (●). Note that an inverse relationship is present mainly in the second group of patients.

The variability of laboratory parameters within family members was assessed by providing details on 4 large VWD2B families characterized by 4 different mutations (P1226L, R1306W, R1308L, and I1309V): some phenotypic heterogeneity was found within the affected members, but the plasma multimeric pattern, occurrence of thrombocytopenia, and results of VWF-GPIIb- α /BC were consistent in these VWD2B pedigrees (data not shown).

VWD2B patients with normal multimeric pattern and platelet count

As mentioned above, 16 patients (24%) from 5 unrelated families had a normal multimeric pattern and no thrombocytopenia before and after stress conditions. Besides mutation R1308L, found in one family (6 patients),³³ more than 1 mutation was found in the remaining 4 families (10 patients), but all shared a substitution of

Table 3. Demographic data and laboratory measurements in the 16 VWD2B patients who never had thrombocytopenia

Mutation	Initials (no. of families)	Age, y	Sex, F-M	Bleeding severity score	BT, minutes	RIPA, mg/mL	VWF:RCo, U/dL	VWF:RCo/A _g , ratio	FVIII, U/dL	Platelet count at baseline, $\times 10^9/\mu$ L	MPV, fl	VWF-GPIIb- α /BC
N1231T/P1266Q	NP (1)	46	F	2	NT	0.70	25	0.66	62	353	NT	0.62
	PV (1)	21	M	0	NT	0.60	21	0.70	54	217	8.4	0.84
V1229G-N1231T-P1266L	GM (2)	19	M	0	NT	0.80	39	0.97	39	308	NT	0.65
	GF (2)	52	M	1	NT	0.40	101	0.89	96	269	7.8	0.75
P1266L	BS (3)	32	F	4	9.30	0.60	54	0.84	56	172	8.5	0.55
	BR (3)	49	M	0	6.00	0.50	76	1.03	78	216	8.1	NT
	BA (3)	59	M	2	4.00	0.50	66	1.12	76	225	7.9	1.09
R1308L	BR (3)	84	M	0	NT	0.60	105	0.98	125	302	8.3	NT
	TE (4)	46	F	6	3.0	0.40	45	0.76	57	251	7.6	0.40
V1229G-N1231T-P1266L	SL (4)	11	F	2	NT	0.50	53	0.99	60	285	8.3	0.49
	TA (5)	18	F	3	8.00	0.40	49	0.96	57	401	8.1	0.70
V1229G-N1231T-P1266L	CR (5)	46	F	16	8.00	0.50	38	1.01	48	193	8.6	NT
	CD (5)	49	F	2	7.00	0.80	39	1.08	35	401	8.1	0.32
V1229G-N1231T-P1266L	CI (5)	36	M	3	5.30	0.30	25	1.12	47	236	8.7	NT
	CMG (5)	50	F	0	NT	0.50	48	1.10	76	261	8.5	0.5
V1229G-N1231T-P1266L	CP (5)	30	M	3	5.00	0.50	25	1.12	44	275	8.2	0.40
	CP (5)	46 (11-84)	8-8	2 (0-16)	6.00 (3.00-9.30)	0.50 (0.30-0.80)	46.5 (21-105)	0.99 (0.66-1.12)	57 (35-125)	265 (172-401)	8.2 (7.6-8.7)	0.59 (0.32-1.09)

Continuous variables are expressed as median (range). MPV indicates mean platelet volume; and NT, not tested. Families are denoted by initials.

Table 4. Clinical and laboratory features of the 67 VWD2B patients, according to platelet levels

	Platelet count ≥140 × 10 ³ /μL (n = 47)	Platelet count <140 × 10 ³ /μL (n = 20)
Sex (M/F)	20/27	7/13
Median age, y* (range)	39 (6-87)	55 (8-75)
Median bleeding severity score† (range)	4 (0-24)	9 (2-19)
VWF:RCo, IU/dL Median (range)	24 (6-105)	24 (6-125)
Patient-months	917	162
No. of bleeding episodes‡	12	17
Incidence of bleeding, % patient-months (95% CI)	1.3 (0.7-2.2)	10.4 (6.0-16.1)

*Mann-Whitney U test: P = .047.

†Mann-Whitney U test: P = .010.

‡In the calculation of bleeding incidence, only the first bleeding episode after the start of follow-up is counted.

proline 1266. The following nucleotide changes were identified: 3692A>C (N1231T), 3789G>A, 3797C>A (P1266Q) in families 1 and 2, 3686T>G (V1229G), 3692A>C (N1231T), 3735G>A, 3789G>A, 3797C>T (P1266L) in family 3, and 3789G>A, 3797C>T (P1266L) in family 4, indicating gene conversion with pseudogene.³⁴ The data of these 16 patients are given in detail in Table 3: none of these patients characterized by enhanced RIPA, normal multimers, normal baseline platelet count, and morphology had increased values of VWF-GPIb-α/BC (median ratio, 0.59; range, 0.32-1.09), as shown in Figure 5A,B. Eight of these patients underwent a stress situation during follow-up without developing thrombocytopenia. The bleeding severity score was extremely heterogeneous (median, 2; range, 0-16), being abnormal in 11 of 16 (69%).

Thrombocytopenia and risk of bleeding and in VWD2B

During the 24-month follow-up, 29 of 67 VWD2B patients (43%) experienced at least 1 bleeding episode for a total of 73, mainly mucosal: 53% epistaxis, 32% gastrointestinal bleeding, 10% menorrhagia, 3% gum bleeds. Muscle hematomas were very rare (2%). The overall incidence of bleeding was 2.7% patient months (95% CI, 1.7%-3.7%). The main clinical and laboratory features of the 2 groups of patients with platelet count higher or lower than 140 × 10³/μL are summarized in Table 4. Patients with thrombocytopenia at baseline showed an approximately 8-fold higher monthly incidence of bleeding than those with normal platelet count. Bleeding-free survival (Kaplan-Meier method) was different in the 2 groups of patients with platelet count higher or lower than 140 × 10³/μL (Figure 6A; log-rank test: P < .001) and in those with bleeding severity score less than 4, 4 to 8, and more than 8 (Figure 6B; log-rank test: P = .003). The probability to be free of bleeding at 12 months was 79% for patients without thrombocytopenia and 20% for those with thrombocytopenia.

The adjusted hazard ratio for bleeding was approximately 6-fold higher in patients who had a bleeding severity score higher than 8 and 5 times higher in those with platelet count less than 140 × 10³/μL, compared with the corresponding reference groups (Table 5). VWF:RCo levels were milder predictors of bleeding in these patients on multivariate analysis. Crude hazard ratio for bleeding in the group of patients with VWF-GPIb-α/BC > 4.0 was increased compared with the group with values less than 1.0 (4.35; 95% CI, 1.41-13.42), but this difference disappeared after adjustment, perhaps because of the strong correlation between VWF-GPIb-α/BC and thrombocytopenia. The median number of individual bleeding episodes was significantly higher

Table 5. Risk of bleeding associated with bleeding severity score, platelet count, and VWF:RCo levels in the 67 patients with VWD2B

	n	Hazard ratio (95% CI)	Hazard ratio (95% CI)
Bleeding severity score			
< 4	21	1*	1*
4-8	24	3.69 (1.02-13.43)	4.97 (1.00-24.90)
> 8	22	6.52 (1.86-22.85)	5.66 (1.03-31.07)
Platelet count			
≥ 140 × 10 ³ /μL	47	1*	1*
< 140 × 10 ³ /μL	20	5.73 (2.69-12.20)	4.57 (1.17-17.90)
VWF:RCo (IU/dL)			
> 40	22	1*	1*
20-40	23	1.98 (0.73-5.35)	1.58 (0.47-5.34)
< 20	22	2.61 (1.00-7.00)	3.26 (0.92-11.56)

Values indicate an adjusted hazard ratio; all variables have been adjusted each one for the other one, and for age and sex.

*Reference group.

in patients with thrombocytopenia than in those without (2; range, 0-7 vs 0; range, 0-4; P < .001).

Discussion

This study investigated the prevalence of thrombocytopenia, its clinical and molecular predictors, and the risk of bleeding in a cohort of VWD2B patients characterized in terms of VWF mutations. Patients were investigated at the time of enrollment using the same questionnaire developed to calculate the bleeding severity score in VWD type 1.^{27,29} The bleeding severity score was higher than normal in the majority of patients (89%). Patients with the V1316M mutation had remarkably high scores, which correlated with the abnormal laboratory phenotype: the same patients had also very prolonged bleeding times (> 30 minutes). Although RIPA was by definition enhanced in all patients, VWF:RCo levels were only moderately reduced and normal in a substantial number of cases and were milder predictors of the risk of bleeding. Moreover, the VWF:RCo/VWF:Ag ratios were not invariably lower than 0.7 as typically observed in VWD types 2A and 2M resulting from dysfunctional VWF^{13,14,28}; normal or reduced ratios were determined by the presence or the absence of HMW multimers, respectively. In agreement with previous studies,⁹ the multimeric pattern of plasma VWF was highly heterogeneous and cannot be used for the diagnosis of VWD2B. Therefore, the only specific assay remains RIPA in patients' PRP.²⁸ Factor VIII levels were normal in most VWD2B, consistently with the very low frequency of soft tissue bleeding (hematomas and hemarthroses) in these patients.

Both the prevalence of thrombocytopenia and its role on the bleeding tendency of VWD2B patients were not well established so far. We found that bleeding severity score was inversely related to platelet count at baseline. During the 24-month follow-up, platelet count were obtained in all the patients both at baseline and during physiologic or pathologic stress conditions all associated with a short- or long-term rise of plasma VWF. Consistent with previous observations,^{9,13,14,33} platelet count were highly variable in VWD2B patients. The prevalence of patients with thrombocytopenia was only 30% at baseline but became almost double (58%) when patients were exposed to stress conditions. Among the 34 exposed patients, 26 (76%) developed transient thrombocytopenia. However, there were 16 patients from 5 unrelated families who always had a normal platelet count, both before and during stress conditions, including DDAVP. This therapeutic approach is usually not recommended in VWD2B²⁸ but has been used in patients with




Figure 6. Cumulative survival according to platelet count and bleeding score. Bleeding-free survival curves calculated according to platelet count assessed at baseline (A) and to bleeding severity score (B) of the 67 patients with VWD2B. In panel A, 2 groups of patients are represented according to absence ($\geq 140 \times 10^3/\mu\text{L}$; dashed line) or presence ($< 140 \times 10^3/\mu\text{L}$; solid line) of thrombocytopenia. In panel B, 3 groups of patients are represented according to tertiles of bleeding severity score, ie, < 4 (small dashed line), 4-8 (big dashed line), more than 8 (solid line). In all curves, each step represents a bleeding event.

P1266L or R1308L mutations^{10,33}: we confirmed that none of them developed thrombocytopenia after administration of the drug. This behavior was strictly correlated to genotype because these patients carried 3 mutations only (P1266Q/L or R1308L). In addition, they all had a normal multimeric pattern in plasma, like the patients previously identified as VWD type 1 Malmö/New York variants.¹⁰⁻¹² These variants were considered very rare so far: this study indicates that patients with VWD2B Malmö/New York are not so rare, they only need to be identified by a RIPA assay. Moreover, we could recently show that both mutations P1266L/Q are associated with VWD2B Malmö/New York phenotype and that both mutations are the result of gene conversion between the VWF gene and its pseudogene.³⁴ The finding of a bleeding score higher than 4 in several members of families with mutations localized at position 1266 as well as the demonstration of enhanced interaction with GPIb- α of this P1266L VWF mutant³⁵ makes the hypothesis of a polymorphism improbable. Specific amino acid substitutions are probably critical for some VWD2B phenotypes: 12 of 16 patients with normal multimers, no thrombocytopenia but increased RIPA have substitutions with leucine at positions P1266 and R1308. In this cohort of patients, platelet morphology was extremely heterogeneous, spanning from normal to large platelets with or without aggregate formation. As previously observed, platelet aggregates in the blood smears correlated with a lower platelet count at baseline.^{15,16} Nurden et al also evaluated platelet morphology by electron microscopy in VWD2B patients and found some ultrastructural abnormalities.³⁶

Another aim of this study was to confirm in a large series of patients the clinical significance of the measurement of VWF-GPIb- α /BC. In addition to the few cases with VWD2B and thrombotic thrombocytopenic purpura originally reported,²⁶ the VWF-GPIb- α /BC was investigated only in another study on the effect of exercise on VWF. This investigation was carried out in

patients with VWD type 1 and in 9 VWD2B patients with 4 different mutations (H1268N, R1306W, V1316M, R1341L)³⁷: there was a significant postexercise increase in VWF-GPIb- α /BC in both the control group and VWD2B but not in VWD type 1. In our study, VWF-GPIb- α /BC levels were increased in all but 16 cases with mutations P1266L/Q and R1308L, with values ranging from 2 to 6 times higher than normal. Indeed, assessment of VWF-GPIb- α /BC values was important for further characterization of VWD2B patients. In patients with loss of HMW multimers, values greater than 1 correlated with thrombocytopenia at baseline and predicted the development of thrombocytopenia after stress conditions. All the patients who had normal results on VWF-GPIb- α /BC carried the P1266Q/L or R1308L mutations, and, more importantly, the same phenotype, ie, normal platelet count both before and after stress, normal platelet morphology, and an intact multimeric pattern in plasma. Therefore, VWF-GPIb- α /BC, as tested by nanobody, is useful to predict thrombocytopenia in VWD2B patients after finding an enhanced RIPA test.

Moreover, in this study, we could confirm that VWF-GPIb- α /BC values were lower than normal not only in VWD types 1 but also in VWD2A, at least in those cases who are characterized by mutations outside VWF A1 domain. Interestingly, we could find, for the first time, values higher than normal in 10 VWD2M patients with mutations within VWF A1 domain (deI1277, R1315C/L, Y1321C, R1374H). These findings point out a major role for the nanobody assay in the diagnosis of VWD types because VWF-GPIb- α /BC values are always high in case of both gain-of-function (VWD2B) and loss-of-function (VWD2M) mutations within VWF A1 domain. Therefore, high values of VWF-GPIb- α /BC together with enhanced RIPA suggests VWD2B with possible thrombocytopenia; high values of VWF-GPIb- α /BC plus impaired RIPA suggests VWD2M; normal values of VWF-GPIb- α /BC plus enhanced RIPA, such as observed in certain P1266L/Q and R1308L mutations, suggests VWD2B without thrombocytopenia. We do not have definite explanations for the normal values of VWF-GPIb- α /BC observed in the 16 VWD2B patients with mutations P1266L/Q and R1308L. According to the observations found in several studies on the crystal structure of this complex,³⁸ we can only postulate that specific amino acid substitutions (ie, leucine) might induce significant changes of the electrostatic interactions between mutated VWF A1 domain and GPIb- α .

Several authors reported a wide degree of heterogeneity in the clinical and laboratory features of affected members within the same VWD2B families.³⁹⁻⁴¹ In this study, this heterogeneity was evaluated in 4 families with mutations P1226L, R1306W, R1308L, or I1309V. In these families, the VWF defects could be investigated not only in plasma^{1,9-11,33} but also by expression studies *in vitro*^{3,9,12,33} that gave homogeneous results. Despite some phenotypic heterogeneity observed within the affected members, the plasma multimeric pattern, occurrence of thrombocytopenia, and results of VWF-GPIb- α /BC were consistent in these VWD2B mutations.

Finally, we investigated whether or not thrombocytopenia associated with VWD2B was from VWD2B patients only when plasma levels of mutant VWF are raised by physiologic and independent risk factor for bleeding in these patients. The bleeding-free survival curves differed significantly in the group of VWD2B with or without thrombocytopenia, and the risk of bleeding in patients with thrombocytopenia was almost 5 times higher than that of patients with a normal platelet count, even

after correction for confounders. Perhaps these patients might deserve a trial with drugs inhibiting the heightened interaction between their abnormal VWF and platelet GPIIb-IIIa,^{42,43} to evaluate whether or not an increase in platelet count can be obtained during bleeding episodes or when they need invasive procedures.

In conclusion, thrombocytopenia is triggered in the majority pathologic stress situations. However, there is a group of patients with normal multimers who never develop thrombocytopenia. VWF-GPIIb-IIIa/BC as tested by nanobody can be a useful marker to further characterize VWD2B patients diagnosed by enhanced RIPA because it is always associated with loss of HMW multimers, large platelet volumes, and development of thrombocytopenia either in the absence or during stress situations. Thrombocytopenia is an independent risk factor for bleeding in patients with VWD2B.

Acknowledgments

The authors thank Rosanna Garavaglia who followed up the patients; biologists Giovanna Cozzi, Silvia La Marca, Margherita Punzo, and Valentina Rubini for the assays performed to diagnose VWD2B; Evelyn Groot who performed in Utrecht most of the assays to test VWF-GPIIb-IIIa/BC in additional patients with VWD; Zaverio M. Ruggeri for his helpful thoughts and suggestions on discussion of the data about molecular mechanisms of VWD2B;

Carlo Balduini for the critical revision of the manuscript; and Luigi Flamini Ghilardini for illustration work.

This work was supported in part by a grant from the Italian Ministry of Health (A.B.F., P.M.M.).

Authorship

Contribution: A.B.F. and P.G.D.G. were responsible for study initiation and coordination; A.B.F., P.M.M., G.C., L.B., P.B., M.T.C., A.P., P.J.L., and P.G.D.G. were involved in study design, data collection, and performing laboratory analyses; G.C., M.T.C., and L.B. were responsible for laboratory data of phenotype and genotype; A.P. was responsible for morphology of platelets; P.J.L. was responsible for setting up the assay of nanobody; P.B. was responsible for statistical analyses; A.B.F. and P.M.M. wrote the manuscript; G.C., L.B., P.B., A.P., P.J.L., and P.G.D.G. were responsible for revisions of draft manuscripts; and all authors were responsible for approval of the final manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Augusto B. Federici, Angelo Bianchi Bonomi Hemophilia Thrombosis Center, Department of Medicine and Medical Specialties, Istituti di Ricovero e Cura a Carattere Scientifico Maggiore Hospital, Mangiagalli, Regina Elena, Foundation and University of Milan, Via Pace, 9, 20122 Milan, Italy; e-mail: augusto.federici@unimi.it.

References

- Ruggeri ZM, Pareti FI, Mannucci PM, Ciavarella N, Zimmerman TS. Heightened interaction between platelets and factor VIII/von Willebrand factor in a new subtype of von Willebrand disease. *N Engl J Med*. 1980;302:1047-1051.
- De Groot PG, Federici AB, De Boer HC, et al. Von Willebrand factor synthesized by endothelial cells from a patient with type II B von Willebrand disease supports platelet adhesion normally but has an increased affinity for platelets. *Proc Natl Acad Sci U S A*. 1989;86:3793-3797.
- Randi AM, Rabinowitz I, Mancuso DJ, et al. Molecular basis of von Willebrand disease type IIB: candidate mutations cluster in one disulfide loop between proposed platelet glycoprotein Ib binding sequences. *J Clin Invest*. 1991;87:1220-1226.
- Cooney KA, Nichols WC, Bruck ME, et al. The molecular defect in type IIB von Willebrand disease: identification of four potential missense mutations within the putative GpIb binding domain. *J Clin Invest*. 1991;87:1227-1233.
- Cooney KA, Ginsburg D. Comparative analysis of type 2b von Willebrand disease mutations: implications for the mechanism of von Willebrand factor binding to platelets. *Blood*. 1996;87:2322-2328.
- Ruggeri ZM. Type IIB von Willebrand disease: a paradox explains how von Willebrand factor works. *J Thromb Haemost*. 2004;2:2-6.
- Ruggeri ZM, Mannucci PM, Lombardi R, Federici AB, Zimmerman TS. Multimeric composition of factor VIII/von Willebrand factor following administration of DDAVP: implications for pathophysiology and therapy of von Willebrand's disease subtypes. *Blood*. 1982;59:1272-1278.
- Ruggeri ZM, Lombardi R, Gatti L, et al. Type IIB von Willebrand disease: differential clearance of endogenous versus transfused large multimers von Willebrand factor. *Blood*. 1982;60:1453-1456.
- Federici AB, Mannucci PM, Stabile F, et al. A type 2b von Willebrand disease mutation (Ile546→Val) associated with an unusual phenotype. *Thromb Haemost*. 1997;78:1132-1137.
- Holmberg L, Berntorp E, Donner M, Nilsson IM. von Willebrand's disease characterized by increased ristocetin sensitivity and the presence of all von Willebrand factor multimers in plasma. *Blood*. 1986;68:668-672.
- Weiss HJ, Sussman II. A new von Willebrand variant (type I, New York): increased ristocetin-induced platelet aggregation and plasma von Willebrand factor containing the full range of multimers. *Blood*. 1986;68:149-156.
- Holmberg L, Dent JA, Schneppenheim, et al. von Willebrand factor mutation enhancing interaction with platelets in patients with normal multimeric structure. *J Clin Invest*. 1993;91:2169-2177.
- Castaman G, Federici AB, Rodeghiero F, Mannucci PM. von Willebrand's disease in the year 2003: towards the complete identification of gene defects for correct diagnosis and treatment. *Haematologica*. 2003;88:94-108.
- Sadler JE, Budde U, Eikenboom JCJ, et al. Update on the pathophysiology and classification of von Willebrand disease: a report of the Subcommittee on von Willebrand factor. *J Thromb Haemost*. 2006;4:2103-2114.
- Saba HI, Saba SR, Dent J, et al. Type IIB Tampa: a variant of von Willebrand disease with chronic thrombocytopenia, circulating platelet aggregates, and spontaneous platelet aggregation. *Blood*. 1985;66:282-286.
- Gralnick HR, Williams SB, McKeown LP, et al. von Willebrand's disease with spontaneous platelet aggregation induced by an abnormal plasma von Willebrand factor. *J Clin Invest*. 1985;76:1522-1529.
- Loffredo G, Baronciani L, Noris P, et al. Von Willebrand disease type 2B must be always considered in the differential diagnosis of genetic thrombocytopenias with giant platelets. *Platelets*. 2006;17:149-152.
- Bader-Meunier B, Proulle V, Trichet C, et al. Misdiagnosis of chronic thrombocytopenia in childhood. *J Pediatr Hematol Oncol*. 2003;25:548-552.
- Geddis AE, Balduini CL. Diagnosis of immune thrombocytopenic purpura in children. *Curr Opin Hematol*. 2007;14:520-525.
- Noris P, Pecci A, Di Bari F, et al. Application of a diagnostic algorithm for inherited thrombocytopenias to 46 consecutive patients. *Haematologica*. 2004;89:1219-1225.
- Rick ME, Williams SB, Sacher RA, McKeown LP. Thrombocytopenia associated with pregnancy in a patient with type IIB von Willebrand disease. *Blood*. 1987;69:786-789.
- Pareti FI, Federici AB, Cattaneo M, Mannucci PM. Spontaneous platelet aggregation during pregnancy in a patient with von Willebrand disease type II B can be blocked by monoclonal antibodies to both platelet glycoproteins Ib and IIb/IIIa. *Br J Haematol*. 1990;75:86-91.
- Hultin MB, Sussman II. Postoperative thrombocytopenia in type IIB von Willebrand disease. *Am J Hematol*. 1990;75:86-91.
- Holmberg L, Nilsson IM, Borge L, Gunnarsson M, Sjorin E. Platelet aggregation induced by 1-desamino-8-D-arginine vasopressin (DDAVP) in type IIB von Willebrand disease. *N Engl J Med*. 1983;309:816-821.
- Nurden P, Debili N, Vainchenker W, et al. Impaired megakaryocytopoiesis in type 2B von Willebrand disease with severe thrombocytopenia. *Blood*. 2006;108:2587-2595.
- Hulstein JJ, de Groot PG, Silence K, et al. A novel nanobody that detects the gain-of-function phenotype of von Willebrand factor in ADAMTS 13 deficiency and von Willebrand disease type 2B. *Blood*. 2005;106:3035-3042.
- Federici AB, Bucciarelli P, Castaman G, et al. Prevalence and determinants of bleeding in different types of von Willebrand disease: results of the first prospective multicenter study on 814 Italian patients. *Blood*. 2007; Abstract 713.
- Federici AB, Castaman G, Mannucci PM. Guidelines for the diagnosis and management of VWD in Italy. *Haemophilia*. 2002;8:607-621.

29. Tosetto A, Rodeghiero F, Castaman G, et al. A quantitative analysis of bleeding symptoms in type 1 von Willebrand disease: results from a multicenter European study (MCM-1 VWD). *J Thromb Haemost*. 2006;4:766-773.
30. Federici AB, Canciani MT, Forza I, et al. A sensitive ristocetin co-factor activity assay with recombinant glycoprotein Ib alpha for diagnosis of patients with low von Willebrand factor levels. *Haematologica*. 2004;89:77-85.
31. Ruggeri ZM, Zimmerman TS. The complex multimeric composition of factor VIII/von Willebrand factor. *Blood*. 1981;57:1140-1143.
32. Hampshire D. International Society of Thrombosis and Hemostasis von Willebrand factor database. www.VWF.group.shef.ac.uk. Accessed December 27, 2007.
33. Baronciani L, Federici AB, Beretta M, et al. Expression studies on a novel type 2B variant of the von Willebrand factor gene (R1308L) characterized by defective collagen binding. *J Thromb Haemost*. 2005;3:2689-2694.
34. Baronciani L, Federici AB, Castaman G, et al. Prevalence of type 2B "Malmoe/New York" von Willebrand disease among patients enrolled into the Italian registry: the role of von Willebrand factor gene conversion. *J Thromb Haemost*. 2008;6:887-890.
35. Gupta PK, Adamtziki E, Budde U, et al. Gene conversions are a common cause of von Willebrand disease. *Br J Haematol*. 2005;130:752-758.
36. Nurden P, Chretien F, Poujol C, et al. Platelet ultrastructural abnormalities in three patients with type 2B von Willebrand disease. *Br J Haematol*. 2000;110:704-714.
37. Stakiw J, Bowman M, Hegadorn C, et al. The effect of exercise on von Willebrand factor and ADAMTS-13 in individuals with type 1 and 2B von Willebrand disease. *J Thromb Haemost*. 2008;6:90-96.
38. Huizinga EG, Shizuko T, Romijn RAP, et al. Structure of glycoprotein Ib α and its complex with von Willebrand factor A1 domain. *Science*. 2002;297:1176-1179.
39. Federici AB, Mannucci PM, Bader R, et al. Heterogeneity in type IIB von Willebrand disease: two unrelated cases with no family history and mild abnormalities of ristocetin-induced interaction between von Willebrand factor and platelets. *Am J Hematol*. 1986;23:381-390.
40. Rendal E, Penas N, Larrabeiti B, et al. Type 2B von Willebrand's disease due to Va11316Met mutation: heterogeneity in the same sibship. *Ann Hematol*. 2001;80:345-360.
41. Mathew P, Greist A, Maahs JA, et al. Type 2B VWD: the varied clinical manifestations in two kindred. *Hemophilia*. 2003;9:137-144.
42. Gilbert JC, Feo-Fraulini T, Hutabarat RM, et al. First-in-human evaluation of anti von Willebrand factor therapeutic aptamer ARC1779 in healthy volunteers. *Circulation*. 2007;116:2678-2686.
43. Hennan JK, Swillo RE, Morgan GA, et al. Pharmacologic inhibition of platelet VWF-GpIb α interaction prevents coronary artery thrombosis. *Thromb Haemost*. 2006;95:469-475.