

Thrombophilic abnormalities, oral contraceptives, and risk of cerebral vein thrombosis: a meta-analysis

Francesco Dentali, Mark Crowther, and Walter Ageno

Recent studies suggest that thrombophilic abnormalities and the use of oral contraceptives (OCs) are the leading causes of cerebral vein thrombosis (CVT). The purpose of this study was to assess the association between CVT and thrombophilic states, OCs, and their interaction. For data sources, we used the MEDLINE, EMBASE, and Cochrane Library databases (January 1994 to March 2005), reference lists of retrieved articles, and contact with content experts. We selected studies comparing the prevalence of OC use and the

prevalence of prothrombotic abnormalities in patients with CVT compared with healthy controls. Two reviewers independently selected studies and extracted study characteristics, quality, and outcomes. Odds ratios (ORs) were calculated for each trial and pooled using the Mantel-Haenszel method. Seventeen studies were included. There was an increased risk of CVT in patients using OCs (OR 5.59; 95% confidence interval [CI] 3.95 to 7.91; $P < .001$), and in patients with factor V Leiden (OR 3.38; 95% CI 2.27

to 5.05; $P < .001$), with mutation G20 210A of prothrombin (OR 9.27; 95% CI 5.85 to 14.67; $P < .001$) and with hyperhomocysteinemia (OR 4.07; 95% CI 2.54 to 6.52; $P < .001$). We concluded that OC users, and patients with factor V Leiden, the prothrombin G20 120A mutation, and hyperhomocysteinemia are at a significantly increased risk of CVT. (Blood. 2006; 107:2766-2773)

© 2006 by The American Society of Hematology

Introduction

Cerebral vein thrombosis (CVT) is a relatively uncommon but potentially life-threatening disease. Recent series showed that as many as 8% of patients with CVT die.¹ In previous studies, the majority of CVT cases were found to be secondary to local or systemic infections and more than 30% of CVT cases were considered idiopathic.^{2,3} However, more recent trials reported other risk factors, such as thrombophilia or the use of oral contraceptives (OCs) to be associated with CVT.^{1,4} In the last decade several inherited or acquired factors causing hypercoagulable state have been studied in patients with deep vein thrombosis (DVT) and pulmonary embolism (PE). Resistance to activated protein C, the most common cause of inherited thrombophilia, was discovered in 1993.⁵ One year later, factor V Leiden was identified as the most frequent cause of this resistance.⁶ Finally, in 1996, a mutation in the prothrombin regulatory sequence was found to be a common prothrombotic factor.⁷ Several large studies and meta-analyses have confirmed that factor V Leiden and G20 210A of mutation of prothrombin are associated with an increased risk of DVT and PE.⁸⁻¹⁰ Large epidemiologic studies have also confirmed that OC users, particularly users of third-generation OCs, are at increased risk of venous thromboembolism (VTE).¹¹ Furthermore, the absolute risk of venous thrombosis in women who use OCs and have a thrombophilic state is higher than expected from the addition of these risks, suggesting a possible enhancement in their individual procoagulant effect.¹²

Unfortunately, due to the rarity of this pathology, the prevalence of these factors in patients with CVT has been evaluated only in small studies and the results are often conflicting or not conclusive.

The aims of our systematic review and meta-analysis are therefore to assess the prevalence and the risk of CVT associated with different inherited and acquired thrombophilic states, to evaluate the prevalence and the risk of CVT associated with use of OCs, and to evaluate the possible interaction between these different prothrombotic factors.

In particular, we considered the following factors: use of OCs, factor V Leiden, prothrombin mutation G20 210, hyperhomocysteinemia, anti-thrombin III, protein C, or protein S deficiency and antiphospholipid syndrome, and either anticardiolipin antibodies and/or lupus anticoagulants.

Methods

Study identification

We tried to identify all published studies that evaluated the presence of inherited or acquired risk factors for CVT using the MEDLINE (1994 to March 2005 Week 5), EMBASE (1994 to 2005 Week 15) and The Cochrane Library (2005, Issue 2) electronic databases. We excluded articles published before 1994 since that was the year that substantial evidence for the importance of activated protein C resistance/factor V Leiden began to become available. The search strategy was developed in collaboration with a professional librarian, had no language restrictions, and used the keywords and subject headings presented in Appendix 1. We supplemented our search by manually reviewing the reference list of all retrieved articles and contacted content experts for additional published or unpublished trials.

From the Department of Medicine, McMaster University, Hamilton, Ontario, Canada; and the Department of Clinical Medicine, University of Insubria, Varese, Italy.

Submitted September 6, 2005; accepted November 28, 2005. Prepublished online as *Blood* First Edition Paper, January 5, 2006; DOI 10.1182/blood-2005-09-3578.

M.C. is a Career Investigator of the Heart and Stroke Foundation of Canada.

Reprints: Mark A. Crowther, Rm L208, St Joseph's Hospital, 50 Charlton Ave East, Hamilton, Ontario, L8N 4A6, Canada; e-mail: crowthrm@mcmaster.ca.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 U.S.C. section 1734.

© 2006 by The American Society of Hematology

Study selection

Study selection was performed independently by 2 reviewers with disagreements resolved through discussion and by opinion of a third reviewer, if necessary. Studies were included if they met the following criteria: (1) diagnosis of CVT was objectively confirmed; (2) patients were 18 years or older; (3) patients were compared with a control group of healthy subjects without a history of thromboembolic disease or genetic relationship with the patients; (4) prevalence of at least one of the following risk factors was measured in patients and in control group: use of OCs, factor V Leiden, prothrombin mutation G20 210, hyperhomocysteinemia, anti-thrombin III, protein C, or protein S deficiency and antiphospholipid syndrome (anticardiolipin antibodies and/or lupus anticoagulant positive); (5) inherited or acquired factors causing hypercoagulable state were measured in an objectively and commonly accepted manner. We excluded case series of patients and all the studies in which CVT was diagnosed only on the basis of clinical symptoms and not confirmed by an objective imaging. When multiple papers for a single study had been published, we decided to use the latest publication and to supplement it, if necessary, with data from the earlier publications.

To assess the agreement between reviewers for study selection, we used the kappa (k) statistic, which measures agreement beyond chance.¹³ According to Maclure and Willett, k values higher than 0.6 are considered to represent a substantial agreement and values higher than 0.8 an almost perfect agreement.¹⁴

Study validity assessment

Two unmasked investigators independently completed the assessment of study validity (F.D., W.A.). Because the use of quality scoring systems or quality scales in observational studies is controversial,¹⁵ the internal validity of each study was evaluated considering 2 potential sources of bias of case-control studies.¹⁶ Studies were considered of low quality when subjects were arbitrarily excluded from either the case or control groups, and when baseline characteristics of the control group (age, sex) were not matched with characteristics of the patient group. Otherwise studies were considered of higher quality.

Data extraction

Two reviewers independently completed data extraction. Disagreement was resolved by consensus and by the opinion of a third reviewer, if necessary. The following baseline characteristics for cases and control groups were collected: number of subjects studied, mean age, variation in age, sex, and race. One or more of the following risk factors was collected in each study: (1) number and proportion of patients and controls with factor V Leiden, prothrombin mutation G20 210, hyperhomocysteinemia, anti-thrombin III, protein C, or protein S deficiency and antiphospholipid syndrome, either anticardiolipin antibodies and lupus anticoagulant positive; (2) number and proportion of women of reproductive age using OCs (pregnant, postpartum, and postmenopausal women were excluded). If the required data could not be located in the published report, we contacted the corresponding author by mail, with a reminder e-mail sent in 15 days.

Statistical analysis

We used Review Manager (RevMan; version 4.2 for Windows; Oxford, England; The Cochrane Collaboration, 2003) to pool data for each risk factor by using the Mantel-Haenszel method and a fixed-effects model.¹⁷ Pooled results are reported as odds ratio (OR) and are presented with 95% confidence interval (CI) and with 2-sided P values. A P value of .05 or less was considered statistically significant. Statistical heterogeneity was evaluated using the I² statistic, which assesses the appropriateness of pooling the individual study results.¹⁸ The I² value provides an estimate of the amount of variance across studies due to heterogeneity rather than chance. When heterogeneity was found, the analysis was repeated using the random-effects model, which includes a measure of variance between studies.¹⁹ To further clarify the role of thrombophilic states and OCs as risk factors for CVT, a priori secondary analysis was planned. We decided to assess, whenever possible, the prevalence and the risk of CVT associated with

different inherited and acquired thrombophilic states and associated with the use of OCs, in patients without any other risk factor known to predispose to thrombosis. Finally, funnel plots of effect size against standard error were completed, whenever possible, to assess for the presence of publication bias.²⁰

Results

Study identification and selection

We identified 436 studies using our search strategy: 187 from MEDLINE, 346 from EMBASE, and 20 from The Cochrane Library (Figure 1). There were 97 studies identified in duplicate. We could exclude 410 studies after title and abstract screening using the predefined inclusion and exclusion criteria, and 26 studies were retrieved for more detailed evaluation.²¹⁻⁴⁶ The interobserver agreement for the study selection was excellent, with a k of 0.99. Contact with the experts and manual review of references did not reveal any additional studies. Nine of the 26 studies were subsequently excluded for the following reasons: 8 did not meet inclusion criteria^{27,28,33,34,36,40,42,44} and 1 contained duplicate data.²⁵ Seventeen studies were therefore included in our systematic review.^{21-24,26,29-32,35,37-39,41,43,45,46}

Study characteristics

Of the 17 studies included, 16 were written in the English language^{21-24,26,29-31,35,37-39,41,43,45,46} and 1 in the Chinese language.³² All included studies were case-control studies. The number of subjects studied ranged from 48 to 2285. Baseline characteristics of included studies are summarized in Table 1. Eight studies evaluated the frequency of OC use in patients with CVT and in a control population.^{21-23,29,30,39,41,45} Sixteen studies evaluated the role of different thrombophilic factors in the CVT pathogenesis^{21-23,25,29-32,35,37-39,43,45,46}; 13 studies considered Leiden mutation of factor V^{22-24,26,29-32,34,38,43,45,46}; 9 mutation G20 210 of factor II^{22-24,29-31,35,39,46}; 4 hyperhomocysteinemia^{21-23,31}; 2 anti-thrombin III, protein C, and protein S deficiency^{23,24}; 1 antiphospholipid syndrome²³; and 1 anticardiolipin antibodies.³⁷ No studies specifically studied patients with more than 1 hereditary or acquired prothrombotic defect, except as discussed in "OCs and thrombophilic factor interactions."

Of the 4 studies that evaluated the presence of hyperhomocysteinemia, 2 measured only fasting levels of homocysteine^{22,31} and 2 also postmethionine load levels.^{21,23} Therefore, we decided to pool only fasting levels of homocysteine in our meta-analysis.




Figure 1. Study selection progression.

Table 1. Baseline characteristics

Study, year (reference no.)	Risk factors	Cases, n	Mean or median age for patients (range), y	Cases, description	Controls, n	Mean or median age for controls (range), y	Controls, description	Excluded patients
Gadelha et al, 2005 (29)	FV, FII, OCs	26	28.5 (3-46)	Patients < 50 aa with CVT confirmed by DA or MRA	217	29 (15-62)	Age- and racially matched healthy individuals with no history of thrombosis or genetic relationship brought by patients and volunteering physicians and health care workers	Patients with major systemic diseases known to predispose to thrombosis or with any positive antiphospholipid test
Boncoraglio et al, 2004 (31)	FV, FII, Hyper-Hcy	26	43 (21-73)	Consecutive patients with first episode of CVT confirmed by DA or MRA	100	42 (21-72)	Healthy hospital employees	NA
Ventura et al, 2004 (22)	FV, FII, Hyper-Hcy, OCs	30	35 (16-49)	Patients with CVT confirmed by CT DA or MRA	40	34 (18-51)	Age- and sex-matched healthy individuals with no history of thrombosis or vascular disease	CVT related to other risk factors
Rodrigues et al, 2004 (30)	FV, FII, OCs	42	28 (2-68)	Patients with objectively confirmed CVT	134	34 (-)	Healthy subjects with no history of thrombosis or genetic relationship with patients	NA
Cantu et al, 2004 (21)	Hyper-Hcy, OCs	45	28 (14-55)	Patients with CVT confirmed by DA or MRA	90	28 (16-53)	Age- and sex-matched healthy individuals with no history of thrombosis or vascular disease	Patients dead or lost to follow up
Martinelli et al, 2003 (23)	FV, FII, OCs, Hyper-Hcy, AT III, PC, PS, APL	121	33 (12-64)	Patients with first episode of CVT confirmed by CT DA or MRA	242	36 (13-62)	Age- and sex-matched healthy individuals with no history of thrombosis	Patients with incomplete thrombophilia screening, uncertainty in diagnosis, or previous thrombotic episode
Meng, 2002 (32)	FV	20	31 (20-48)	NA	50	38 (18-58)	NA	NA
Bombeli et al, 2002 (24)	FV, FII, AT III, PC, PS	51	36.7 (17-61)	Patients with CVT confirmed by DA or MRA	120	37.4 (19-62)	Healthy individuals with no history of thrombosis or vascular disease and without any medication	Patients with unclear diagnosis of acquired or hereditary protein C or S deficiency
Margaglione et al, 2001 (46)	FV, FII	28	NA	Patients with CVT confirmed by DA or MRA	1304	NA (22-66)	Healthy hospital employees	Patients < 14 aa or with previous thromboembolic episodes
Voetsch et al, 2000 (35)	FV, FII	14	24.8 (16-31)	Patients with CVT confirmed by DA or MRA considered idiopathic	225	34.1 (16-50)	Age- and sex-matched volunteer physicians, students, and laboratory staff	Patients with deficiency of natural anticoagulants or antiphospholipid antibodies: underlying venous malformation

Table 1. Baseline characteristics (continued)

Study, year (reference no.)	Risk factors	Cases, n	Mean or median age for patients (range), y	Cases, description	Controls, n	Mean or median age for controls (range), y	Controls, description	Excluded patients
Reuner et al, 1998 (39)	FII, OC	45	37 (3-69)	Patients with CVT confirmed by CT, DA, or MRA	354	39 (18-65)	Healthy blood donors	NA
De Bruijn et al, 1998 (41)	OC	37	NA (18-49)	Women with CVT confirmed by DA or MRA	2248	NA (18-49)	Age-matched randomly selected women	Women < 18 aa or pregnant
Weih et al, 1998 (26)	FV	12	33.8 (21-60)	Consecutive patients with CVT confirmed by DA or MRA	36	34.2 (NA)	Age- and sex-matched subjects with no history of thrombosis	Patients with septic venous thrombosis, neoplasm or non-German descent
Cristopher et al, 1998 (37)	ACL	31	27.6 (NA)	Patients with CVT confirmed by CT or DA	31	26.6 (NA)	Age- and sex-matched asymptomatic subjects	Patients with CVT related to infection or trauma
Ludemann et al, 1998 (38)	FV	55	40 (11-83)	Patients with CVT confirmed by DA or MR imaging	272	NA (18-55)	Healthy adult subjects	NA
Zuber et al, 1998 (43)	FV	19	38 (20-72)	Patients with CVT confirmed by DA or MR imaging	57	NA	Age- and sex-matched subjects with no history of cerebrovascular disease	NA
Martinelli et al, 1996 (45)	FV, OC	25	32 (21-64)	Patients with CVT confirmed by CT, DA, or MR imaging	75	NA	Age- and sex-matched subjects with no history of thrombosis	CVT secondary to a local infection

ACL indicates anticardiolipin antibodies; ATIII, deficiency of anti-thrombin III; CT, computerized tomography; DA, digital angiography; FII, mutation 20210 factor II; FV, mutation 1691 factor V (Leiden); Hyper-Hcy, hyperhomocysteinemia; MR, magnetic resonance; MRA, magnetic resonance angiography; OCs, oral contraceptives; PC, protein C deficiency; PS, protein S deficiency; and NA, not applicable.

Study quality

The quality of included studies was generally high. Only a few patients were arbitrarily excluded from either the case or control groups. Furthermore, in 10 of 17 studies, control subjects were matched with patients according to age and sex.^{21-23,26,29,35,37,41,43,45}

Oral contraceptives

There were 8 studies that evaluated the role of OCs in CVT.^{21-23,29,30,39,41,45} The pooled analysis included 263 women with CVT and 2862 women without CVT. In women using OCs, the summary OR for developing CVT was 5.59 in comparison to

controls (95% CI 3.95 to 7.91; *P* < .001), Figure 2. Heterogeneity between the studies was significant (*I*² = 68.4%; *P* = .002) and was maintained after the exclusion all the possible outliers from the analysis (data not shown). However, a similar OR was obtained using the random-effects model (OR 4.79; 95% CI 2.40 to 9.58; *P* < .001). This result probably reflects the difference among the studies in the selection of the population analyzed. Funnel plot of OR versus standard error appeared asymmetric with an absence of studies in the bottom right hand corner, suggesting that smaller, unpublished studies that demonstrate an increased OR of CVT in patients taking OCs were not included in our meta-analysis.




Figure 2. Odds ratio for cerebral vein thrombosis for oral contraceptive users. n indicates the number positive; N, total number.




Figure 3. Odds ratio for cerebral vein thrombosis in factor V Leiden carriers. n indicates the number positive; N, total number.

Factor V Leiden

Thirteen studies included 469 case and 3023 control subjects.^{22-24,26,29-32,35,37,43,45,46} In 12 studies the method used to determine the presence of the factor V Leiden mutation was described^{22-24,26,29,31,32,35,38,43,45,46}; in all studies detection of Leiden mutation was carried out by amplification of a fragment of exon 10 of the factor V and posterior digestion with an endonuclease, according to the method used by Bertina et al.⁶ Compared with controls, the pooled OR for CVT in patients with factor V Leiden mutation was 3.38 (95% CI 2.27 to 5.05; $P < .001$), Figure 3. Heterogeneity between the studies was extremely low ($I^2 = 27.7\%$; $P = .18$).

Funnel plot of OR versus standard error evaluated 12 of 13 studies included in our meta-analysis since in 1 study there were neither patients nor controls with Leiden mutation of factor V.³² The plot appeared symmetric, suggesting absence of publication bias.

Mutation G20 210A of factor II

Nine studies evaluated the role of mutation G20 210A in the risk of CVT.^{22-24,29-31,35,39,46} In the pooled analysis 360 patients and 2688 controls were studied. In 8 of 9 studies, the method to detect the G20 210A mutation of factor II was described.^{22-24,29,31,35,39,46} All the studies used the method described by Poort et al.⁷ The pooled OR of developing CVT was 9.27 (95% CI 5.85 to 14.67; $P < .001$) in patients with mutation G20 210A of factor II compared with controls, Figure 4. There was no significant between-study heterogeneity ($I^2 = 0\%$; $P = .49$). The funnel plot appeared symmetric, suggesting absence of publication bias.

Hyperhomocysteinemia

Four case-control studies evaluated hyperhomocysteinemia in patients with CVT.^{21-23,31} Fasting hyperhomocysteinemia was defined as above the 95th percentile of normal population value in 2 studies,^{23,31} as above the 90th percentile of normal population value in 1 study,²¹ and as higher than 12 μM in 1 study.²² Compared with controls, the pooled OR for CVT in patients with hyperhomocysteinemia was 4.07 (95% CI 2.54 to 6.52; $P < .001$), Figure 5. There was no heterogeneity between the studies ($I^2 = 0\%$; $P = .48$). In the 2 studies that also evaluated postmethionine load levels of homocysteine, patients with fasting or postload hyperhomocysteinemia had an OR of developing CVT of 4.3 and 4.2, respectively, compared with controls.^{21,23} Due to the low number of studies, funnel-plot analysis could not be done. Therefore, the presence of publication bias could not be excluded.

Other risk factors

Two studies analyzed the role of the deficiency of anti-thrombin III, protein C, and protein S^{23,24} as risk factors for CVT. Only 1 study considered antiphospholipid syndrome²³ and 1 anticardiolipin antibodies.³⁷ Antithrombin levels were measured using functional assay and then confirmed by an immunologic assay. The combined OR of the 2 studies was 2.69 (95% CI 0.66 to 10.96; $P = .19$). Protein C and protein S were measured by functional and immunologic assays. The combined OR of the 2 studies was 11.10 (95% CI 1.87 to 66.05; $P = .009$) for protein C and 12.49 for protein S (95% CI 1.45 to 107.29; $P = .03$). Due to the low number of eligible patients, the confidence interval is very wide and it is impossible to draw any major conclusion from these results.





Figure 4. Odds ratio for cerebral vein thrombosis in G20 210A factor II mutation carriers. n indicates the number positive; N, total number.

Figure 5. Odds ratio for cerebral vein thrombosis in patients with hyperhomocysteinemia. n indicates the number positive; N, total number.



The only study that considered the antiphospholipid antibodies syndrome²³ found a higher incidence of antiphospholipid antibodies in patients with CVT (9/121) compared with controls (0/242).

Finally, Christopher et al³⁷ evaluated the role of anticardiolipin antibodies in CVT. They found a significantly higher incidence of anticardiolipin antibodies in patients (7/31) in comparison to controls (1/31) (OR 8.75; 95% CI 1.01 to 75.64).

OCs and thrombophilic factor interactions

Few studies provided separate analyses of OR of CVT in thrombophilic women taking OCs.^{23,29} Therefore, it was not possible to pool these data. In their recent study, Martinelli et al²³ stratified patients for the presence of hyperhomocysteinemia, factor V Leiden, or prothrombin mutation and the intake of OCs. They found that the presence of both risk factors gave an OR of 19.5 (95% CI 5.7 to 67.3) for hyperhomocysteinemia, 30.0 (95% CI 3.4 to 263.0) for factor V Leiden, and 79.3 (95% CI 10.0 to 629.4) for prothrombin mutation in comparison to controls.

The multivariate analysis performed in the study conducted by Gadelha et al²⁹ confirmed the independent association between CVT, prothrombin mutation, and OC use.

Discussion

In the first meta-analysis published on this particular topic, we have reviewed 17 publications that analyzed the association between CVT and the most frequent prothrombotic states. We found a strong association between CVT and use of OCs, the factor V Leiden mutation, the G20 210A mutation of prothrombin, and hyperhomocysteinemia. Our conclusions are strengthened by the uniform nature of our results and the narrow confidence interval about the resulting ORs. Only 1 study failed to detect factor V Leiden mutation in both patient and control groups,³² but this study was carried out in Chinese patients within whom the prevalence of factor V Leiden is far lower than is seen in white patients.⁴⁷ Data on antiphospholipid antibodies syndrome, deficiencies of anti-thrombin III, protein C, and protein S deficiency were inadequate to allow a reliable statistical analysis. However, a non-statistically significant trend toward an association with the disease for antithrombin, protein C, and protein S was observed. Finally, only 2 studies have evaluated the effect of a concomitant thrombophilia in women of reproductive age using OCs. These data suggest a possible interaction between OCs and some prothrombotic states (hyperhomocysteinemia, factor V Leiden, prothrombin G20 210A). In particular, the risk of CVT in women on OCs who are carriers of the mutation G20 210A of prothrombin seemed to be much higher than in controls.

Several well-designed large trials, reviews, and meta-analyses have demonstrated that use of OCs and the presence of 1 or more

thrombophilic factors is associated with an increased risk of DVT and PE.^{8-12,48-50} Testing for most common thrombophilic factors in young patients presenting with unprovoked serious thrombotic events is currently recommended,⁵¹ a recommendation supported by our observation of a strong association between thrombophilic states and CVT.

Our meta-analysis has limitations. First, our systematic review was restricted to case-control studies, and the application of formal meta-analytic methods to observational studies is controversial, since bias implicit in the study design may misrepresent the strength of associations within the data.¹⁵ To minimize this potential bias, we selected only studies in which the diagnosis of CVT was objectively confirmed and in which prothrombotic factors were measured with objectively and widely accepted methods. Second, studies included in our meta-analysis have different inclusion and exclusion criteria, and to combine results across studies may be inappropriate. However, the heterogeneity between the studies, calculated using the I2 statistic, was generally low. Only when we pooled studies that evaluated OCs in patients with CVT was the heterogeneity between studies remarkable, and it remained high after adjustment for possible outliers. These results suggest that baseline characteristics of patients included in the analysis may be different and that results should be interpreted cautiously. However, after repeating the analysis using a random-effects model, an approach that accounts for some of the variance between studies, we found similar results. Third, despite a careful review of references and contact with content experts, we failed to identify any published or unpublished study not found in our initial literature search. Finally, despite several efforts, it was not possible to have information about the methods used to diagnose CVT in one study in which the authors stated that CVT was objectively confirmed in all patients.³⁰ However, since only 42 patients were included in that study, it is very unlikely that even in the worst-case scenario (the diagnosis of CVT was wrong in all patients) this could influence the results of our meta-analysis. Because it is recognized that publication bias can affect the results of meta-analyses, we attempted to assess this potential bias using a funnel plot. However, funnel plots appeared symmetric, suggesting the absence of publication bias in the 3 analyses that considered factor V Leiden, mutation G20 210A of the prothrombin, and hyperhomocysteinemia. In the funnel plot analysis of the association of OCs with CVT we observed asymmetry, with an absence of studies in the bottom right hand corner of the plot. This suggests that smaller, unpublished studies likely to demonstrate an increased risk of CVT with use of OCs were not included in our meta-analysis. However, inclusion of these studies and elimination of this bias, if it really existed, would increase the observed association between OCs and CVT.⁵²

In conclusion, our meta-analysis shows that CVT is strongly associated with factor V Leiden mutation, with mutation G20 210A of prothrombin, and with hyperhomocysteinemia.

The role of the other thrombophilic abnormalities in the pathogenesis of CVT is less clear. The use of OCs significantly increases the risk of CVT in women and preliminary data suggest that such risk is further increased in OC-using women with prothrombotic abnormalities. Therefore, our observations suggest that women with CVT should avoid use of OCs, and that the use of OCs in women with known thrombophilic states should be evaluated on an individual basis assessing the risk of CVT for different thrombophilic abnormalities. Future studies, including careful prospective studies of patients with CVT, are now warranted to better delineate the risk of recurrent thrombosis in patients with and without thrombophilic defects. However, due to the rarity of the disease, it is very unlikely that these studies will be performed.

Appendix: MEDLINE search strategy

1. (Intracranial embolism and thrombosis).mp. [mp = title, original title, abstract, name of substance word, subject heading word]/ (7436)
2. Exp cerebral veins/ (1672)
3. Exp intracranial thrombosis/ or exp sinus thrombosis, intracranial/ (2032)
4. 1 or 2 or 3/ (10 728)
5. Exp factor V/ (4134)
6. Anti-thrombin III/ (5505)
7. Protein C/ (3904)
8. Protein S/ (1375)
9. Homocysteine/ or 5-methyltetrahydrofolate-homocysteine S-methyltransferase/ (6860)
10. Exp antibodies, antiphospholipid/ (5050)
11. Exp contraceptives, oral/ (33 169)
12. Prothrombin/ (6075)
13. Cerebral vein thrombosis.tw./ (79)
14. Cerebral venous thrombosis.tw./ (548)
15. Hyperhomocysteinemia/ (1470)
16. 4 or 13 or 14/ (10 833)
17. 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 15/ (61 761)
18. 16 and 17/ (495)
19. Limit 18 to year = 1994-2005/ (236)
20. Limit 19 to "review articles"/ (28)
21. 19 not 20/ (208)
22. Limit 21 to "all infant (birth to 23 months)"/ (21)
23. 21 not 22/ (187)

References

1. Ferro JM, Canhao P, Stam J, Bousser MG, Barinagarrementeria F; ISCVT Investigators. Prognosis of cerebral vein and dural sinus thrombosis: results of the International Study on Cerebral Vein and Dural Sinus Thrombosis (ISCVT). *Stroke*. 2004;35:664-670.
2. Ameri A, Bousser MG. Cerebral venous thrombosis. *Neurol Clin*. 1992;10:87-111.
3. Gates PC. Cerebral venous thrombosis: a retrospective review. *Aust N Z J Med*. 1986;16:766-770.
4. De Bruijn SF, Stam J. Randomized, placebo-controlled trial of anticoagulant treatment with low-molecular-weight heparin for cerebral sinus thrombosis. *Stroke*. 1999;30:484-488.
5. Dahlback B, Carlsson M, Svensson PJ. Familial thrombophilia due to a previously unrecognized mechanism characterized by poor anticoagulant response to activated protein C: prediction of a cofactor to activated protein C. *Proc Natl Acad Sci U S A*. 1993;90:1004-1008.
6. Bertina RM, Koeleman BP, Koster T, et al. Mutation in blood coagulation factor V associated with resistance to activated protein C. *Nature*. 1994;369:64-67.
7. Poort SR, Rosendaal FR, Reitsma PH, Bertina RM. A common genetic variation in the 3'-untranslated region of the prothrombin gene is associated with elevated plasma prothrombin levels and an increase in venous thrombosis. *Blood*. 1996;88:3698-3703.
8. Rodeghiero F, Tosetto A. Activated protein C resistance and factor V Leiden mutation are independent risk factors for venous thromboembolism. *Ann Intern Med*. 1999;130:643-650.
9. Salomon O, Steinberg DM, Zivelin A, et al. Single and combined prothrombotic factors in patients with idiopathic venous thromboembolism: prevalence and risk assessment. *Arterioscler Thromb Vasc Biol*. 1999;19:511-518.
10. Emmerich J, Rosendaal FR, Cattaneo M, et al. Combined effect of factor V Leiden and prothrombin 20210A on the risk of venous thromboembolism—pooled analysis of 8 case-control studies including 2310 cases and 3204 controls. Study Group for Pooled-Analysis in Venous Thromboembolism. *Thromb Haemost*. 2001;86:809-816.
11. World Health Organization Collaborative Study of Cardiovascular Disease and Steroid Hormone Contraception. Venous thromboembolic disease and combined oral contraceptives: results of international multicentre case-control study. World Health Organization Collaborative Study of Cardiovascular Disease and Steroid Hormone Contraception. *Lancet*. 1995;346:1575-1582.
12. Vandenbroucke JP, Koster T, Briet E, Reitsma PH, Bertina RM, Rosendaal FR. Increased risk of venous thrombosis in oral-contraceptive users who are carriers of factor V Leiden mutation. *Lancet*. 1994;344:1453-1457.
13. McGinn T, Wyer PC, Newman TB, Keitz S, Leipzig R, Guyatt G, for Evidence-Based Medicine Teaching Tips Working Group. Tips for learners of evidence-based medicine, 3: measures of observer variability (kappa statistic). *CMAJ*. 2004;171:1369-1373.
14. Maclure M, Willett WC. Misinterpretation and misuse of the kappa statistic. *Am J Epidemiol*. 1987;126:161-169.
15. Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis of Observational Studies in Epidemiology (MOOSE) group. *JAMA*. 2000;283:2008-2012.
16. Hayden GF, Kramer MS, Horwitz RI. The case-control study: a practical review for the clinician. *JAMA*. 1982;247:326-331.
17. Mantel N, Haenszel W. Statistical aspects of the analysis of data from retrospective studies of disease. *J Natl Cancer Inst*. 1959;22:719-748.
18. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003;327:557-560.
19. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials*. 1986;7:177-188.
20. Sterne JA, Egger M, Smith GD. Systematic reviews in health care: investigating and dealing with publication and other biases in meta-analysis. *BMJ*. 2001;323:101-105.
21. Cantu C, Alonso E, Jara A, et al. Hyperhomocysteinemia, low folate and vitamin B12 concentrations, and methylene tetrahydrofolate reductase mutation in cerebral venous thrombosis. *Stroke*. 2004;35:1790-1794.
22. Ventura P, Cobelli M, Marietta M, Panini R, Rosa MC, Salvio G. Hyperhomocysteinemia and other newly recognized inherited coagulation disorders (factor V Leiden and prothrombin gene mutation) in patients with idiopathic cerebral vein thrombosis. *Cerebrovasc Dis*. 2004;17:153-159.
23. Martinelli I, Battaglioli T, Pedotti P, Cattaneo M, Mannucci PM. Hyperhomocysteinemia in cerebral vein thrombosis. *Blood*. 2003;102:1363-1366.
24. Bombeli T, Basic A, Fehr J. Prevalence of hereditary thrombophilia in patients with thrombosis in different venous systems. *Am J Hematol*. 2002;70:126-132.
25. Martinelli I, Sacchi E, Landi G, Taioli E, Duca F, Mannucci PM. High risk of cerebral-vein thrombosis in carriers of a prothrombin-gene mutation and in users of oral contraceptives. *N Engl J Med*. 1998;338:1793-1797.
26. Weih M, Vetter B, Ziemer S, et al. Increased rate of factor V Leiden mutation in patients with cerebral venous thrombosis. *J Neurol*. 1998;245:149-152.
27. Delport R, Ubbink JB, Vermaak WJ, Rossouw H, Becker PJ, Joubert J. Hyperhomocysteinemia in black patients with cerebral thrombosis. *QJM*. 1997;90:635-639.
28. Martinelli I, Rosendaal FR, Vandenbroucke JP, Mannucci PM. Oral contraceptives are a risk factor for cerebral vein thrombosis. *Thromb Haemost*. 1996;76:477-478.
29. Gadelha T, Andre C, Juca AA, Nucci M. Prothrombin 20210A and oral contraceptive use as risk factors for cerebral venous thrombosis. *Cerebrovasc Dis*. 2005;19:49-52.
30. Rodrigues CA, Rocha LK, Morelli VM, Franco RF, Lourenco DM. Prothrombin G20210A mutation, and not factor V Leiden mutation, is a risk factor for cerebral venous thrombosis in Brazilian patients. *J Thromb Haemost*. 2004;2:1211-1212.
31. Boncoraglio G, Carriero MR, Chiapparini L, et al. Hyperhomocysteinemia and other thrombophilic risk factors in 26 patients with cerebral venous thrombosis. *Eur J Neurol*. 2004;11:405-409.
32. Meng Q, Pu C. [Cerebral venous thrombosis and factor V Leiden mutation.] *Zhonghua Yi Xue Za Zhi*. 2002;82:47-49.
33. Marietta M, Bertesi M, Simoni L, Castelli I, Cappi C, Torelli G. Cerebral vein thrombosis and lupus anticoagulant antibodies. *Clin Appl Thromb Hemost*. 2001;7:238.
34. Madonna P, De Stefano V, Coppola A, Albinini R, Cerbone AM. G20210A PRTH gene mutation and other thrombophilic polymorphisms in patients with cerebral vein thrombosis. *Stroke*. 2000;31:1787-1788.

35. Voetsch B, Damasceno BP, Camargo EC, et al. Inherited thrombophilia as a risk factor for the development of ischemic stroke in young adults. *Thromb Haemost*. 2000;83:229-233.
36. Weih M, Vetter B, Castell S, Ziemer S, Kulozik AE, Einhaupl KM. Hereditary thrombophilia in cerebral venous thrombosis. *Cerebrovasc Dis*. 2000;10:161-162.
37. Christopher R, Nagaraja D, Dixit NS, Narayanan CP. Anticardiolipin antibodies: a study in cerebral venous thrombosis. *Acta Neurol Scand*. 1999;99:121-124.
38. Ludemann P, Nabavi DG, Junker R, et al. Factor V Leiden mutation is a risk factor for cerebral venous thrombosis: a case-control study of 55 patients. *Stroke*. 1998;29:2507-2510.
39. Reuner KH, Ruf A, Grau A, et al. Prothrombin gene G20210—A transition is a risk factor for cerebral venous thrombosis. *Stroke*. 1998;29:1765-1769.
40. de Bruijn SF, Stam J, Vandenbroucke JP. Increased risk of cerebral venous sinus thrombosis with third-generation oral contraceptives. Cerebral Venous Sinus Thrombosis Study Group. *Lancet*. 1998;351:1404.
41. de Bruijn SF, Stam J, Koopman MM, Vandenbroucke JP. Case-control study of risk of cerebral sinus thrombosis in oral contraceptive users and in [correction of who are] carriers of hereditary prothrombotic conditions. The Cerebral Venous Sinus Thrombosis Study Group. *BMJ*. 1998;316:589-592.
42. Scoditti U, Buccino GP, Pini M, Pattacini C, Mancina D. Risk of acute cerebrovascular events related to low oestrogen oral contraceptive treatment. *Ital J Neurol Sci*. 1998;19:15-19.
43. Zuber M, Toulon P, Marnet L, Mas JL. Factor V Leiden mutation in cerebral venous thrombosis. *Stroke*. 1996;27:1721-1723.
44. Kimber TE, Lloyd JV, Brophy BP, Thompson PD. Cerebral venous sinus thrombosis associated with factor V gene mutation. *J Neurol Neurosurg Psychiatry*. 1996;61:204-205.
45. Martinelli I, Landi G, Merati G, Cella R, Tosi A, Mannucci PM. Factor V gene mutation is a risk factor for cerebral venous thrombosis. *Thromb Haemost*. 1996;75:393-394.
46. Margaglione M, Brancaccio V, Ciampa A, Papa ML, Grandone E, Di Minno G. Inherited thrombophilic risk factors in a large cohort of individuals referred to Italian thrombophilia centers: distinct roles in different clinical settings. *Haematologica*. 2001;86:634-639.
47. Rees DC, Cox M, Clegg JB. World distribution of factor V Leiden. *Lancet*. 1995;346:1133-1134.
48. Price DT, Ridker PM. Factor V Leiden mutation and the risks for thromboembolic disease: a clinical perspective. *Ann Intern Med*. 1997;127:895-903.
49. Den Heijer M, Rosendaal FR, Blom HJ, Gerrits WB, Bos GM. Hyperhomocysteinemia and venous thrombosis: a meta-analysis. *Thromb Haemost*. 1998;80:874-847.
50. College of American Pathologists Consensus Conference XXXVI: diagnostic issues in thrombophilia. *Arch Pathol Lab Med*. 2002;126:1277-1433.
51. Seligsohn U, Lubetsky A. Genetic susceptibility to venous thrombosis. *N Engl J Med*. 2001;344:1222-1231.
52. Sutton AJ, Duval SJ, Tweedie RL, Abrams KR, Jones DR. Empirical assessment of effect of publication bias on meta-analyses. *BMJ*. 2000;320:1574-1577.