

TP53 alterations in acute myeloid leukemia with complex karyotype correlate with specific copy number alterations, monosomal karyotype, and dismal outcome

Frank G. Rücker,¹ Richard F. Schlenk,¹ Lars Bullinger,¹ Sabine Kayser,¹ Veronica Teleanu,¹ Helena Kett,¹ Marianne Habdank,¹ Carla-Maria Kugler,¹ Karlheinz Holzmann,¹ Verena I. Gaidzik,¹ Peter Paschka,¹ Gerhard Held,² Marie von Lilienfeld-Toal,³ Michael Lübbert,⁴ Stefan Fröhling,¹ Thorsten Zenz,^{1,5,6} Jürgen Krauter,⁷ Brigitte Schlegelberger,⁸ Arnold Ganser,⁷ Peter Lichter,⁹ Konstanze Döhner,¹ and Hartmut Döhner¹

¹Department of Internal Medicine III, University Hospital of Ulm, Ulm, Germany; ²Department of Internal Medicine I, University of Saarland, Homburg, Germany; ³Department of Internal Medicine III, University Hospital of Bonn, Bonn, Germany; ⁴Department of Hematology-Oncology, University of Freiburg Medical Center, Freiburg, Germany; ⁵Department of Translational Oncology, National Center for Tumor Diseases and German Cancer Research Center, Heidelberg, Germany; ⁶Department of Medicine V, University of Heidelberg, Heidelberg, Germany; ⁷Department of Hematology, Hemostasis, Oncology, and Stem Cell Transplantation, Hannover Medical School, Hannover, Germany; ⁸Institute of Cell and Molecular Pathology, Hannover Medical School, Hannover, Germany; and ⁹Division of Molecular Genetics, German Cancer Research Center, Heidelberg, Germany

To assess the frequency of *TP53* alterations and their correlation with other genetic changes and outcome in acute myeloid leukemia with complex karyotype (CK-AML), we performed integrative analysis using *TP53* mutational screening and array-based genomic profiling in 234 CK-AMLs. *TP53* mutations were found in 141 of 234 (60%) and *TP53* losses were identified in 94 of 234 (40%) CK-AMLs; in total, 164 of 234 (70%) cases had *TP53* alterations. *TP53*-altered CK-AML were characterized by a higher degree of

genomic complexity (aberrations per case, 14.30 vs 6.16; $P < .0001$) and by a higher frequency of specific copy number alterations, such as $-5/5q-$, $-7/7q-$, $-16/16q-$, $-18/18q-$, $+1/+1p$, and $+11/+11q/amp11q13-25$; among CK-AMLs, *TP53*-altered more frequently exhibited a monosomal karyotype (MK). Patients with *TP53* alterations were older and had significantly lower complete remission rates, inferior event-free, relapse-free, and overall survival. In multivariable analysis for overall survival, *TP53* alterations, white

blood cell counts, and age were the only significant factors. In conclusion, *TP53* is the most frequently known altered gene in CK-AML. *TP53* alterations are associated with older age, genomic complexity, specific DNA copy number alterations, MK, and dismal outcome. In multivariable analysis, *TP53* alteration is the most important prognostic factor in CK-AML, outweighing all other variables, including the MK category. (*Blood*. 2012;119(9):2114-2121)

Introduction

Chromosomal abnormalities are found in approximately 55% of adult patients with acute myeloid leukemia (AML) and are among the most important independent prognostic factors.¹⁻³ AMLs exhibiting 3 or more acquired chromosome aberrations in the absence of chromosomal rearrangements listed in the World Health Organization (WHO) 2008 category “AML with recurrent genetic abnormalities” are now defined as AML with complex karyotype (CK-AML).^{1,2} CK-AMLs account for 10% to 15% of adult AMLs, and the frequency increases with age. CK-AMLs belong to the cytogenetic adverse-risk group because they are associated with very poor outcome when treated with intensive or nonintensive conventional chemotherapy.^{1,3,4} Recently, a new cytogenetic category was introduced, that is, the monosomal karyotype (MK) defined by the presence of one single autosomal monosomy in association with at least one additional autosomal monosomy or one structural chromosomal abnormality (in the absence of core binding factor AML and acute promyelocytic leukemia).⁵ This MK category was reported to be associated with a dismal prognosis and to add prognostic information, even in CK-AML.

Complex karyotypes often contain numerous chromosome aberrations that can only be partially or not at all interpreted using standard cytogenetic techniques. Such aberrations include unbalanced translocations with chromosomal material of unknown origin, marker or ring chromosomes, homogeneously staining regions, or double minutes, the latter representing cytogenetic equivalents of high-level DNA amplifications. In general, CK-AMLs are characterized by chromosomal gains and losses, rather than balanced translocations, suggesting distinct mechanisms in leukemogenesis.⁶

In recent years, molecular cytogenetic and array-based techniques have enabled a more precise characterization of these complex genetic changes. The imbalances most frequently found are losses affecting chromosome 5 or 5q ($-5/5q-$), $-17/17p-$, $-7/7q-$, $-18/18q-$, $-16/16q-$, $-12/12p-$, and gains affecting chromosome 8 or 8q ($+8/+8q$), $+11/+11q$, $+21/+21q$, $+22/+22q$, and $+1/+1p$.⁷ Furthermore, novel potential target genes have been delineated based on the observation that they are located in critical regions of deletions⁸ or contained in amplicons, such as *MYC* in

Submitted August 24, 2011; accepted November 8, 2011. Prepublished online as *Blood* First Edition paper, December 20, 2011; DOI 10.1182/blood-2011-08-375758.

The online version of this article contains a data supplement.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked “advertisement” in accordance with 18 USC section 1734.

There is an Inside *Blood* commentary on this article in this issue.

© 2012 by The American Society of Hematology

8q24, *ETS1* and *FLII* in 11q24, *CDX2* in 13q12, and *ETS2* and *ERG* in 21q22.^{7,9-12}

One target located in the commonly deleted region of 17p13 is the tumor suppressor gene *TP53*. In AML, *TP53* alterations (mutations and/or losses, *TP53*^{altered}) are rare and have been closely associated with CK-AML.¹³⁻¹⁶ Clinically, *TP53* alterations appear to be associated with inferior outcome.¹⁵⁻¹⁷ However, these data are based on a small number of studies, and only one of these addressed both *TP53* losses and mutations but not the prognostic significance.¹⁵

The objectives of our study were: (1) to study a large cohort of CK-AMLs (n = 234) for *TP53* mutations; (2) to analyze these cases for DNA copy number alterations (CNAs) using array-based techniques; and (3) to correlate *TP53* alterations with specific chromosome abnormalities, CNAs, MK, and clinical outcome.

Methods

Patients

Peripheral blood (PB) and/or bone marrow (BM) samples from 234 adult patients with CK-AML were analyzed. The definition of CK-AML followed recommended criteria.^{1,2} The diagnosis of AML was based on French-American-British Cooperative Group criteria,¹⁸ and after 2004 on WHO criteria.¹⁹ A total of 133 patients had de novo AML, 31 secondary AML (s-AML) after myelodysplastic syndrome or myeloproliferative neoplasms, 30 therapy-related (t-AML), and in 40 patients it was unknown. Of these 234 patients, 155 (66%) were treated on consecutive multicenter treatment trials of the German-Austrian AML Study Group (AMLSG) applying age-adjusted intensive chemotherapy: AML HD93 (n = 1),²⁰ AML HD98A (n = 30),²¹ and AMLSG 07-04 (n = 54; NCT00151242) for younger patients (16-60 years); and AML HD98B (n = 27)²² and AMLSG 06-04 (n = 43; NCT00151255) for elderly patients (> 60 years). All trials were approved by the local ethics committees of all participating institutions; all patients gave informed consent for treatment, cryopreservation of samples, and molecular analyses according to the Declaration of Helsinki. Samples were primarily selected based on availability of sufficient material for genomic profiling and mutational analysis.

Cytogenetic and molecular genetic analysis

For cytogenetic classification, metaphases of sufficient quality could be studied by chromosome banding analysis in 219 patients; karyotypes were described according to the International System for Human Cytogenetic Nomenclature.²³

Because of evolving technology that occurred in the course of the study, we switched from array comparative genomic hybridization (CGH) to single nucleotide polymorphism (SNP) array-based genomic profiling. Array CGH (n = 131) using the 2.8k platform and/or the 8.0k platform and unpaired SNP analyses using Affymetrix GeneChip Human Mapping 250K Array (n = 61) were performed as previously described^{7,24}; Genome-Wide Human SNP Array 6.0 profiling (n = 42) was performed according to the manufacturer's protocols (Affymetrix). Genotyping Version 2.0 console (Affymetrix) was used for analysis of 6.0 arrays. Microarray data will be available at gene expression omnibus at <http://www.ncbi.nlm.nih.gov/geo/> (GEO accession number GSE34542).

TP53 sequence analysis

To identify mutations in exons 4 to 10 of *TP53*, denaturing high-performance liquid chromatography was performed as previously described.²⁵ Aberrant profiles were verified by bidirectional sequencing and compared with wild-type sequence (GenBank; X54156). Mutations were described using 2 different databases (IARC *TP53* Database; www.p53.iarc.fr and The *TP53* Web site; www.p53.free.fr).^{26,27} Subcloning analyses using the TOPO TA Cloning Kit and resequencing were performed according to manufacturer's protocols (Invitrogen).

Statistical analyses

The section on statistical analyses is provided in supplemental Methods (available on the *Blood* Web site; see the Supplemental Materials link at the top of the online article).

Results

TP53 mutation analysis

In 141 of the 234 (60%) patients with CK-AML, a total of 168 mutations were identified; 161 (96%) were located in the sequence-specific DNA-binding domain of p53 (residues 102-292). The majority were missense mutations (n = 130), followed by deletions/insertions (n = 21); 17 resulted in premature stop; 4 preserved the open reading frame (g.13149del6, g.13149del9, g.14001del21, and g.13162ins3), nonsense mutations (n = 9), and splice site mutations (n = 8; Figure 1). Seventeen of 25 patients harboring 2 or more *TP53* mutations exhibited no *TP53* loss; 5 of these showed a homozygous *TP53* mutation; and in all remaining patients with available DNA for subcloning (n = 8), compound heterozygous mutations were confirmed. Hemizygous mutations (loss of 1 allele and at least 1 mutation in the remaining allele; 47% [79 of 168]) were more frequent than homozygous mutations (26% [43 of 168]), heterozygous (18% [30 of 168]), including possible compound heterozygous mutations among the 4 patients with more than 1 heterozygous mutation, and compound heterozygous mutations (10% [16 of 168]); 65 (39%) mutations affected common hot spots, such as codons 175, 245, 248, 273, and 275. Mutations affecting codons 175, 248, and 273 were associated with biallelic *TP53* alteration compared with all other *TP53* mutations (100% [27 of 27] vs 79% [90 of 114], *P* = .008).

CNAs and copy number neutral loss of heterozygosity UPD

In the entire cohort of 234 CK-AML, genomic losses (n = 1.845) were more frequent than gains (n = 778) or high-level DNA amplifications (n = 153). The median number of aberrations per case was 10 (range, 0-51); median numbers of losses, gains, and amplifications per case were 6 (range, 0-43), 2 (range, 0-28), and 0 (range, 0-7), respectively.

Recurrent losses were identified for the following chromosomes: monosomy 5 or losses of 5q (-5/5q-) (n = 147; 63%), -7/7q- (n = 123; 53%), -17/17p- (n = 106; 45%), -16/16q- (n = 66; 28%), -18/18q-, -12/12p- (n = 65 each; 28%), -20/20q- (n = 55; 24%), -3/3p- (n = 54; 23%), and -11/11q- (n = 35; 15%). Most frequent gains were +8/+8q (n = 67; 29%), +11/+11q (n = 61; 26%), +21/+21q (n = 39; 17%), +1/+1p (n = 37; 16%), +22/+22q (n = 33; 14%), +13/+13q (n = 29; 12%), +9/+9p (n = 28; 12%), and +19/+19p (n = 25; 11%). Most frequent high-level DNA amplifications mapped to 21q22, 11q13~25 (n = 22 each; 9%), and 8q24 (n = 8; 4%).

Usually large, but also submicroscopic, losses (down to 800 kb in size) affecting the *TP53* locus on 17p13 were identified in 94 of 234 (40%) cases. Uniparental disomy of 17p [UPD(17p); 8.14-22.50 Mb in size] encompassing the *TP53* locus was detected in 15 of 103 (15%) cases analyzed by SNP arrays.

Biallelic TP53 alteration

Combining mutational and microarray findings, 164 of 234 (70%) CK-AMLs exhibited *TP53* alteration (mutation and/or loss of *TP53*); 71 of 164 (43%) cases had biallelic *TP53* alteration by hemizygous mutation pattern, and 38 of 70 (54%) cases without




Figure 1. Mapping of 168 *TP53* mutations in 141 CK-AMLs. Hemizygous mutations are indicated in the bottom panel, heterozygous, and/or homozygous mutations are marked in the top panel. Exons 4 to 10 are drawn to relative scale; missense mutations (green), nonsense mutations (red), and insertion/deletion mutations (blue) are shown at their approximate location along the exons. Bold represents homozygous mutations, and blue italics, frameshift mutations leading to a premature stop codon.

TP53 loss exhibited homozygous *TP53* mutation caused by UPD(17p) in 15 of 19 SNP-analyzed cases. In the 4 cases lacking evidence for UPD(17p) in SNP profiling, homozygous *TP53* mutations possibly resulted from intragenic loss of heterozygosity. Furthermore, 19 of 131 (15%) cases analyzed by array CGH on DNA sequence analysis exhibited homozygous *TP53* mutation that are probably caused by UPD(17p) considering the frequency of UPD(17p) found by SNP array analysis. In addition, subcloning analysis confirmed biallelic *TP53* alteration by compound heterozygous mutations in 8 patients. Together, at least 117 of 164 (71%) *TP53*^{altered} CK-AMLs had biallelic *TP53* inactivation (not taken into account the 4 patients with potentially compound heterozygous *TP53* mutations). Patient 96 exhibited a homozygous missense mutation in exon 6 (p.R213L) and an additional heterozygous frameshift mutation in exon 4 (p.A74fs), suggesting that these had occurred sequentially, with p.R213L being the primary event followed by UPD(17p) resulting in the homozygous mutation pattern, whereas the p.A74fs mutation followed the recombination event (supplemental Figure 1).

Correlation of *TP53* alteration with pattern of chromosome abnormalities and CNAs

We correlated *TP53* alterations, as assessed by DNA sequence analysis and array profiling, with the pattern of chromosome abnormalities identified by conventional cytogenetics and with the pattern of CNAs detected by array-based analyses (Table 1).

Correlation with chromosome abnormalities. *TP53* alterations were identified in 157 of 219 (72%) CK-AMLs that could be analyzed by conventional cytogenetics. *TP53*^{altered} CK-AMLs had a higher degree of genomic complexity as measured by total number of aberrations (≥ 5 aberrations, $P < .0001$) and the presence of marker chromosomes ($P = .0005$). *TP53* alterations were correlated with the presence of specific cytogenetic abnormalities, such as $-5/5q-$ ($P < .0001$), concomitant $-5/5q-$ and $-7/7q-$

($P = .0006$), and $20q-$ ($P = .02$); we found no correlation with $-7/7q-$ ($P = .14$; Table 1).

Correlation with CNAs. *TP53* alterations were correlated with the total number of losses (mean \pm SD; 9.54 ± 7.49 vs 4.00 ± 4.88 , $P < .0001$), gains (3.91 ± 3.80 vs 1.94 ± 1.92 , $P < .0001$), high-level DNA amplifications (0.84 ± 1.31 vs 0.21 ± 0.83 , $P = .0002$), and genomic complexity as measured by total number of aberrations per case (14.30 ± 9.41 vs 6.16 ± 5.53 , $P < .0001$). Moreover, *TP53* alterations were positively correlated with specific genomic aberrations, such as $-5/5q-$ ($P < .0001$), $-7/7q-$ ($P = .003$), concomitant $-5/5q-$ and $-7/7q-$ ($P < .0001$), and also $-3/3p-$ ($P = .002$), $-16/16q-$ ($P < .0001$), $-18/18q-$ ($P = .0008$), and $-20/20q-$ ($P = .004$); further correlations were identified for $+1/+1p$ ($P = .001$), $+11/+11q$ ($P = .0002$), $+13/+13q$ ($P = .02$), $+19/+19p$ ($P = .04$), and amplifications in $11q13\sim 25$ [$\text{amp}(11)(q13\sim 25)$] ($P = .0004$; Table 1; Figure 2).

Correlation of *TP53* alterations with MK

By conventional cytogenetics, 171 of 219 (78%) CK-AMLs fulfilled the MK criteria (CK⁺/MK⁺ AML) as previously defined. *TP53* alterations were found in 137 of 171 (80%) CK⁺/MK⁺ AMLs and in only 20 of 48 (42%) CK⁺/MK⁻ AMLs ($P < .0001$; Table 1). Compared with CK⁺/MK⁻ AMLs, CK⁺/MK⁺ AMLs were characterized by a higher degree of genomic complexity determined by cytogenetics: more than or equal to 5 aberrations, 88% (151 of 171) versus 54% (26 of 48), $P < .0001$; and by genomic profiling as measured by total number of losses (mean \pm SD; 9.29 ± 7.40 vs 3.67 ± 5.72 ; $P < .0001$) and aberrations per case (13.59 ± 9.61 vs 6.81 ± 6.11 , $P < .0001$).

We subsequently determined MK⁺ AML based on array data (molMK). The frequency of CK⁺/molMK⁺ AML was much lower

Table 1. Genetic and clinical characteristics according to TP53 alteration

	TP53 ^{unaltered}	TP53 ^{altered}	P
Cytogenetics	n = 62	n = 157	
≥ 5 aberrations	38 (61%)	139 (89%)	< .0001
Marker chromosomes	29 (47%)	114 (73%)	.0005
–5/5q–	20 (32%)	124 (79%)	< .0001
–7/7q–	25 (40%)	81 (52%)	.14
–5/5q– and –7/7q–	12 (19%)	70 (45%)	.0006
–20/20q–	8 (13%)	44 (28%)	.02
MK	34 (55%)	137 (87%)	< .0001
Array-based genomics	n = 70	n = 164	
Total no. of losses (mean ± SD)	4.00 ± 4.88	9.54 ± 7.49	< .0001
Total no. of gains (mean ± SD)	1.94 ± 1.92	3.91 ± 3.80	< .0001
Total no. of amplifications (mean ± SD)	0.21 ± 0.83	0.84 ± 1.31	.0002
Total no. of genomic aberrations (mean ± SD)	6.16 ± 5.53	14.30 ± 9.41	< .0001
–3/3p–	7 (10%)	47 (29%)	.002
–5/5q–	20 (29%)	127 (77%)	< .0001
–7/7q–	26 (37%)	97 (59%)	.003
–5/5q– and –7/7q–	13 (19%)	87 (53%)	< .0001
–11/11q–	13 (19%)	22 (13%)	.32
–12/12p–	13 (19%)	52 (32%)	.06
–16/16q–	5 (7%)	61 (37%)	< .0001
–18/18q–	9 (13%)	56 (34%)	.0008
–20/20q–	8 (11%)	47 (29%)	.004
+1/+1p	3 (4%)	34 (21%)	.001
+8/+8q	21 (30%)	46 (28%)	.75
+9/+9p	11 (16%)	17 (10%)	.27
+11/+11q	7 (10%)	54 (33%)	.0002
+13/+13q	3 (4%)	26 (16%)	.02
+19/+19p	3 (4%)	22 (13%)	.04
+21/+21q	7 (10%)	32 (20%)	.09
+22/+22q	6 (9%)	27 (16%)	.15
amp(8)(q24)	4 (6%)	4 (2%)	.24
amp(11)(q13~25)	0 (0%)	22 (13%)	.0004
amp(21)(q22)	3 (4%)	19 (12%)	.09
Molecular MK	16 (23%)	59 (36%)	.07
Molecular genetics	n = 50	n = 99	
FLT3-ITD positive	3 (6%)	1 (1%)	.11
FLT3-TKD mutation	3 (6%)	1 (1%)	.11
NPM1 mutation	3 (6%)	0 (0%)	.04
Clinical data	n = 52	n = 103	
Sex (male/female)	26 (50%)/26 (50%)	54 (52%)/49 (48%)	.87
Median age, y	54	61	.002
AML history			
De novo	36 (69%)	76 (74%)	.57
Secondary	6 (12%)	9 (9%)	.58
Therapy-related	9 (17%)	16 (16%)	.82
Median WBC, × 10 ⁶ /L	12.9	6.5	.18
Median platelet count, × 10 ⁶ /L	48	41	.46
Median hemoglobin, g/dL	9.1	8.9	.38
Median BM blast count, %	78	65	.04
Median PB blast count, %	45	30	.18
Median LDH serum level, U/L	391	438	.25
Response	n = 52	n = 103	
CR after induction therapy	26 (50%)	29 (28%)	.01
RD after induction therapy	18 (35%)	53 (51%)	.06
Outcome	n = 52	n = 103	
OS			
Median, mo	10.97	4.14	< .0001
3-y survival rate, %	28	3	
EFS			
Median, mo	1.94	1.12	.0007
3-y survival rate, %	13	1	
RFS			
Median, mo	12.16	6.51	.01
3-y survival rate, %	30	7	

ITD indicates internal tandem duplication; and TKD, tyrosine kinase domain.




Figure 2. Relative frequencies and pairwise co-occurrences of *TP53* alteration-associated genomic aberrations illustrated using Circos Table Viewer Version 0.52.²⁸ The percentages indicate the proportion of each aberration associated with (A) *TP53*^{altered} CK-AMLs and (B) *TP53*^{unaltered} CK-AMLs. Unaltered *TP53* and amp(11)(q13~q25) were mutually exclusive. MK is based on cytogenetics analysis.

(75 of 234; 32%) because many monosomies described in chromosome banding analysis were not real monosomies but part of chromosomal material hidden in unbalanced translocation or marker chromosomes. *TP53* alterations were found in 59 of 75 (79%) CK⁺/molMK⁺ AMLs and in 105 of 159 (66%) CK⁺/molMK⁻ AMLs ($P = .07$; Table 1).

Correlation of *TP53* alterations with clinical characteristics, response to therapy, and survival

Analyses were restricted to patients enrolled into AMLSG multicenter treatment trials applying age-adjusted intensive chemotherapy ($n = 155$, median age, 59 years; range, 18-81 years). Because there were no significant differences regarding clinical characteristics, response to therapy, and survival for *TP53*^{monoallelic altered} and *TP53*^{biallelic altered} CK-AMLs (supplemental Table 1; supplemental Figure 2), these genotypes were grouped as *TP53*^{altered} CK-AML for further analyses.

Clinical characteristics. *TP53*^{altered} CK-AML patients were older (median 61 vs 54 years, $P = .002$) and had lower BM blast counts (median 65% vs 78%, $P = .04$; Table 1).

Response to therapy. *TP53* alterations were associated with resistance to chemotherapy. Response to induction therapy was as follows: complete remission (CR) 28% and 50% ($P = .01$), refractory disease (RD) 51% and 35% ($P = .06$), and early/hypoplastic death 21% and 15% ($P = .52$) for CK⁺/*TP53*^{altered} and CK⁺/*TP53*^{unaltered} AML, respectively (Table 1). Other variables predicting for poor response to induction therapy were age ($P < .0001$) and genomic losses affecting 5q ($P = .02$), 7q ($P = .03$), and 16q ($P = .04$). Lactate dehydrogenase (LDH) serum levels, white blood cell count (WBC), *s/t*-AML, and cytogenetic MK did not impact CR achievement.

For multivariable analysis, a conditional model was used with an age cut-point at 60 years to address the different treatment

intensities applied in the different age cohorts. This model revealed as significant factors *TP53*^{altered} (odds ratio [OR] = 0.55; 95% confidence interval [CI], 0.30-1.00; $P = .05$) and age (OR for a 10-year difference, 0.67; 95% CI, 0.52-0.87; $P = .003$). No significant impact on CR achievement was found for the variables WBC, platelet counts, cytogenetic MK, and *s/t*-AML (Table 2).

Survival analysis. The median follow-up time for survival in the 155 CK-AML was 36.6 months (95% CI, 29.9-51.4 months); the estimated 3-year event-free survival (EFS), relapse-free survival (RFS), and overall survival (OS) of the entire cohort were 5% (95% CI, 2%-10%), 17% (95% CI, 9%-31%), and 12% (95% CI, 7%-19%), respectively.

TP53 alterations were associated with inferior survival; the 3-year estimated survival rates for CK⁺/*TP53*^{altered} and CK⁺/*TP53*^{unaltered} patients were as follows: EFS, 1% versus 13% (log-rank, $P = .0007$); RFS, 7% versus 30% ($P = .01$); and OS, 3% versus 28% ($P < .0001$), respectively (Table 1; Figure 3). Other variables predicting for inferior OS in univariable analysis were age ($P < .0001$), cytogenetic MK ($P = .03$), and genomic losses of 5q ($P = .03$), 7q ($P = .003$), 16q ($P = .0004$), and gains of 1p ($P = .04$), and amp(11)(q13~25; $P = .05$). LDH and WBC did not impact OS. Among CK⁺/MK⁺ AMLs, those with *TP53* alterations had significantly worse OS ($P = .0004$; Figure 4).

Multivariable analysis stratified again for age at a cut-point of 60 years revealed *TP53*^{altered} (hazard ratio [HR] = 2.43; 95% CI, 1.56-3.77; $P = .0001$), logarithm of WBC (HR = 1.62; 95% CI, 1.17-2.26; $P = .004$), and age (HR for 10-year difference, 1.26; 95% CI, 1.01-1.56, $P = .04$) as significant variables; not significant for OS were platelet counts, cytogenetic MK, and *s/t*-AML (Table 2).

Allogeneic hematopoietic stem cell transplantation in first CR was performed in 30 CK-AML patients. Of those, 14 of 15 *TP53*^{altered} CK-AML relapsed and died, whereas in *TP53*^{unaltered} CK-AML 9 of 15 relapsed and died ($P = .04$). This translated into significantly worse OS for *TP53*^{altered} CK-AML ($P = .04$; supplemental Figure 3).

Table 2. Multivariate analyses of outcome

CK-AML	Response		OS	
	OR	P	HR	P
<i>TP53</i> alteration	0.55	.05	2.43	.0001
Age (difference of 10 y)	0.67	.003	1.26	.04
<i>s/t</i> -AML	0.67	.24	1.05	.81
Logarithm of WBC	0.74	.19	1.62	.004
Logarithm of platelets	0.76	.39	1.13	.62
MK*	0.75	.43	0.87	.57

*Determined by chromosome banding analysis.

Discussion

In our series of 234 CK-AMLs, *TP53* was deleted and/or mutated in 70% of cases, thus representing the most frequently known altered gene in this AML subgroup. *TP53* alterations were associated with older age, genomic complexity, specific chromosome abnormalities, monosomal karyotype, specific CNAs, and predicted for dismal outcome.




Figure 3. Kaplan-Meier survival estimates according to the *TP53* status. Data are shown for *TP53*^{unaltered} CK-AMLs and *TP53*^{altered} CK-AMLs. (A) OS. (B) EFS. (C) RFS.

Loss of *TP53* was found in approximately 40% of CK-AMLs by array-based techniques, a value that corresponded well with that of 17p abnormalities found on chromosome banding analysis. By DNA sequence analysis, 60% of cases exhibited *TP53* mutations, consistent with previous reports.^{6,15,16} Of note, at least two-thirds of mutated cases had biallelic *TP53* alteration resulting from hemizygous, compound heterozygous, and homozygous mutations commonly as a result of homologous recombination leading to UPD. Thus, when assessing for *TP53* mutational status in CK-AML, it will be necessary to include DNA sequence analysis.

TP53^{altered} CK-AMLs were characterized by a significantly higher degree of genomic complexity, as assessed by total number of genomic losses and gains, as well as the frequency of high-level DNA amplifications. This observation fits well into the p53 pathomechanism of genomic instability.²⁹⁻³⁵ *TP53*^{altered} CK-AMLs were also associated with specific abnormalities. As previously reported,^{15,16,36,37} $-5/5q-$ and/or $-7/7q-$ were significantly more frequent among *TP53*^{altered} CK-AMLs. Because we also applied array-based techniques, we identified additional CNAs associated with *TP53*^{altered} CK-AML, that is, $-3/3p-$, $-16/16q-$, $-18/18q-$, $-20/20q-$, and gains or amplifications of 1p, 11q, 13q, and 19p. Such genomic pattern associated with *TP53* alterations may pinpoint to candidates cooperating in p53-dependent leukemogenesis.

Recently, the cytogenetic category of “monosomal karyotype” was described, allowing further risk stratification of CK-AML patients.⁵ Of note, in our study, CK⁺/MK⁺ AMLs were significantly associated with *TP53* alterations, found in 80% of CK⁺/MK⁺ AML compared with only 42% in CK⁺/MK⁻ AML. Thus, *TP53* alterations appear to be one molecular basis for this purely descriptive cytogenetic subset. The association of *TP53* alterations with CK⁺/MK⁺ AML was lost when *TP53* alterations were correlated with CNAs identified by array-based assays. Not unexpectedly, many monosomies described in chromosome banding analysis were not real monosomies but were part of chromosomal material hidden in unbalanced translocations or marker chromosomes.




Figure 4. Kaplan-Meier survival estimates according to the cytogenetic status. (A) Data are shown for OS for CK⁺/MK⁻ AML and CK⁺/MK⁺ AML. (B) Data are shown for OS for the subgroups CK⁺/MK⁻/*TP53*^{unaltered}, CK⁺/MK⁻/*TP53*^{altered}, CK⁺/MK⁺/*TP53*^{unaltered}, and CK⁺/MK⁺/*TP53*^{altered}.

Little is known about the pathogenesis of CK-AML, but the high frequency of *TP53* alteration, and in particular biallelic alteration, suggests an important role of p53 in leukemogenesis. Evidence for this hypothesis comes from several observations in mice and human disease: (1) mouse studies requiring biallelic *TP53* inactivation and a concomitant “second hit” for myeloid leukemogenesis^{38,39} demonstrated that p53^{lost} myeloid progenitors exhibit aberrant self-renewal, thereby promoting AML⁴⁰; (2) in high-risk MDS and/or AML evolving from a 5q- syndrome, the expansion of preexisting *TP53* mutated subclones was observed^{41,42}; and (3) recently, next-generation sequencing of a therapy-related CK-AML genome identified several acquired genetic lesions and a heterozygous intragenic germline *TP53* deletion, becoming homozygous in AML as a result of acquired UPD(17p),⁴³ a mechanism possibly underlying the sequential *TP53* inactivation in patient 96 (supplemental Figure 1).

Besides being older and having lower BM blasts, *TP53*^{altered} CK-AML had no distinct clinical phenotype, possibly because of the complexity of concurrent genetic events and different consequences of *TP53* alterations. *TP53* losses or mutations entail various tumor phenotypes,⁴⁴ and mouse models investigating *TP53* inactivation identified gain of function for hot spot mutations, such as R175H, R248W, and R273H, as well as increased proliferation related to accelerated tumorigenesis and leukemogenesis, resulting in a more aggressive AML.^{35,44-46}

p53 loss of function has been shown to be related to resistance to chemotherapy, also to cytarabine.^{46,47} Consistent with this finding, *TP53* alterations in our study were associated with resistance to “3 + 7”-based induction chemotherapy (Tables 1 and 2). Refractory disease was observed in 51% of CK⁺/*TP53*^{altered} compared with 35% of CK⁺/*TP53*^{unaltered} AMLs. In univariable analysis, *TP53* alteration also predicted for inferior OS; median survival times for CK⁺/*TP53*^{altered} and CK⁺/*TP53*^{unaltered} patients were 4.14 and 10.97 months, respectively. In multivariable analysis, *TP53* alteration was by far the strongest prognostic factor for OS, followed by logarithm of WBC and age; of note, the cytogenetic category MK completely lost its prognostic impact. Explorative subset analysis suggested that allogeneic hematopoietic stem cell transplantation had no favorable impact on outcome in *TP53*^{altered} CK-AML.

TP53 alterations are the most common molecular lesions in CK-AML and predict for resistance to conventional chemotherapy and dismal outcome. *TP53* alterations correlate with specific CNAs and with the MK category. In CK-AML, *TP53* alteration represents the most important prognostic marker, even outweighing the MK category in multivariable analysis. Therefore, *TP53* mutational status should be assessed in clinical trials investigating novel agents to identify compounds that may be effective in this subset of patients.

Acknowledgments

The authors thank all members of the AMLSG for their participation in this study and for providing patient samples. The following AML Study Group institutions and investigators participated in this study: Daniel Oruzio, MD, Klinikum Augsburg, Augsburg, Germany; Dietmar Reichert, MD, Ubbo-Emmius-Klinik Aurich, Aurich, Germany; Jörg Westermann, MD, Charité Berlin, Berlin, Germany; Christian Teschendorf, MD, Klinikum Bochum, Bochum, Germany; Ingo Schmidt-Wolf, MD, Universitätsklinikum Bonn, Bonn, Germany; Florian Lordick, MD, Städtisches Klinikum Braunschweig, Braunschweig, Germany; Bernd Hertenstein, MD, Klinikum Bremen-Mitte, Bremen, Germany; Helga

Bernhard, MD, Klinikum Darmstadt, Darmstadt, Germany; Ulrich Germing, MD, Universitätsklinikum Düsseldorf, Düsseldorf, Germany; Carsten Schwänen, MD, Klinikum Esslingen, Esslingen, Germany; Mohammed Wattad, MD, Kliniken Essen Süd, Ev. Krankenhaus Essen-Werden gGmbH, Essen, Germany; Elke Jäger, MD, Krankenhaus Nordwest GmbH, Frankfurt, Germany; Michael Lübbert, MD, Universitätsklinikum Freiburg, Freiburg, Germany; Andrea Distelrath, MD, Klinikum Fulda, Fulda, Germany; Volker Runde, MD, Wilhelm-Anton-Hospital, Goch, Germany; Detlef Haase, MD, Universitätsklinikum Göttingen, Göttingen, Germany; Walter Fiedler, Universität Hamburg, Hamburg, Germany; Hans Salwender, MD, Allgemeines Krankenhaus Altona, Hamburg, Germany; Elisabeth Lange, MD, Klinikum Hamm, Hamm, Germany; Andrea Sendler, MD, Klinikum Hanau, Hanau, Germany; Hartmut Kirchner, MD, Krankenhaus Siloah, Hannover, Germany; Uwe Martens, SLK-Kliniken GmbH Heilbronn, Heilbronn, Germany; Ulrich Kaiser, MD, Bernward-Krankenhaus Hildesheim, Hildesheim, Germany; Michael Pfreundschuh, MD, Universitätsklinikum Homburg/Saar, Homburg, Germany; David Nachbaur, Universitätsklinikum Innsbruck, Austria; Martin Bentz, MD, Städtisches Klinikum Karlsruhe, Karlsruhe, Germany; Heinz A. Horst, MD, Universitätsklinikum Kiel, Kiel, Germany; Stefan Kremers, MD, Caritas-Krankenhaus Lebach, Lebach, Germany; Frank Hartmann, MD, Klinikum Lemgo, Lemgo, Germany; Andreas Petzer, MD, Krankenhaus der Barmherzigen Schwestern Linz, Linz, Austria; Gerhard Heil, MD, Klinikum Lüdenschheid, Lüdenschheid, Germany; Thomas Fischer, MD, Klinikum Magdeburg, Magdeburg, Germany; Thomas Kindler, MD, Universitätsklinikum Mainz, Mainz, Germany; Martin Griebhammer, MD, Klinikum Minden, Minden, Germany; Katharina Götze, MD, Technische Universität München, München, Germany; Ali-Nuri Hünerlitürkoglu, Lukaskrankenhaus Neuss, Neuss, Germany; Claus-Henning Köhne, MD, Klinikum Oldenburg, Oldenburg, Germany; Thomas Südhoff, MD, Klinikum Passau, Passau, Germany; Karin Corduan, MD, Elisabeth Krankenhaus Recklinghausen, Recklinghausen, Germany; Michael Schenk, MD, Krankenhaus der Barmherzigen Brüder, Regensburg, Germany; Artur Wehmeier, MD, Klinikum Remscheid, Remscheid, Germany; Axel Matzdorff, MD, Caritas-Klinik St Theresia, Saarbrücken, Germany; Richard Greil, MD, Salzburger Landeskliniken, Salzburg, Austria; Wolfgang Grimminger, MD, Klinikum Schwäbisch-Gmünd, Mutlangen, Germany; Hans-Günther Mergenthaler, MD, Klinikum Stuttgart, Stuttgart, Germany; Else Heidemann, MD, Diakonie-Klinikum Stuttgart, Stuttgart, Germany; Heinz Kirchen, MD, Krankenhaus der Barmherzigen Brüder, Trier, Germany; Helmut Salih, MD, Universitätsklinikum Tübingen, Tübingen, Germany; Wolfgang Brugger, MD, Klinikum Villingen-Schwenningen, Villingen-Schwenningen, Germany; Elisabeth Koller, MD, Hanuschkrankenhaus, Vienna, Austria; and Aruna Raghavachar, MD, Helios Klinikum Wuppertal, Wuppertal, Germany.

This study was supported in part by the Bundesministerium für Bildung und Forschung (grants 01GS0439 NGFN2 and 01GS0871 NGFNplus).

Authorship

Contribution: F.G.R. designed and performed research, analyzed and interpreted data, performed statistical analysis, and wrote the manuscript; R.F.S. provided study materials or patients, collected data, and analyzed and interpreted data; L.B. collected, analyzed, and interpreted data; S.K., V.T., V.I.G., P.P., G.H., M.v.L.-T., M.L., J.K., B.S., and A.G. provided study materials or patients and collected data; H.K., M.H., and C.-M.K. performed research and collected data; K.H. designed research and collected data; S.F., T.Z., and P.L. analyzed and interpreted data; and

K.D. and H.D. designed research, provided study materials or patients, collected data, analyzed and interpreted data, and wrote the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Hartmut Döhner, Department of Internal Medicine III, University Hospital of Ulm, Albert-Einstein-Allee 23, 89081 Ulm, Germany; e-mail: hartmut.doechner@uniklinik-ulm.de.

References

- Döhner H, Estey EH, Amadori S, et al. Diagnosis and management of acute myeloid leukemia in adults: recommendations from an international expert panel, on behalf of the European LeukemiaNet. *Blood*. 2010;115(3):453-474.
- Swerdlow SH, Campo E, Harris NL, et al. *WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues*. Lyon, France: IARC Press; 2008
- Grimwade D, Hills RK, Moorman AV, et al. Refinement of cytogenetic classification in acute myeloid leukemia: determination of prognostic significance of rare recurring chromosomal abnormalities among 5876 younger adult patients treated in the United Kingdom Medical Research Council trials. *Blood*. 2010;116(3):354-365.
- Schoch C, Haferlach T, Haase D, et al. Patients with de novo acute myeloid leukaemia and complex karyotype aberrations show a poor prognosis despite intensive treatment: a study of 90 patients. *Br J Haematol*. 2001;112(1):118-126.
- Breems DA, Van Putten WL, De Greef GE, et al. Monosomal karyotype in acute myeloid leukemia: a better indicator of poor prognosis than a complex karyotype. *J Clin Oncol*. 2008;26(29):4791-4797.
- Mrózek K. Cytogenetic, molecular genetic, and clinical characteristics of acute myeloid leukemia with a complex karyotype. *Semin Oncol*. 2008;35(4):365-377.
- Rücker FG, Bullinger L, Schwaenen C, et al. Disclosure of candidate genes in acute myeloid leukemia with complex karyotypes using microarray-based molecular characterization. *J Clin Oncol*. 2006;24(24):3887-3894.
- Delhommeau F, Dupont S, Della Valle V, et al. Mutation in TET2 in myeloid cancers. *N Engl J Med*. 2009;360(22):2289-2301.
- Baldus CD, Liyanarachchi S, Mrózek K, et al. Acute myeloid leukemia with complex karyotypes and abnormal chromosome 21: amplification discloses overexpression of APP, ETS2, and ERG genes. *Proc Natl Acad Sci U S A*. 2004;101(11):3915-3920.
- Poppe B, Vandesompele J, Schoch C, et al. Expression analyses identify MLL as a prominent target of 11q23 amplification and support an etiologic role for MLL gain of function in myeloid malignancies. *Blood*. 2004;103(1):229-235.
- Storlazzi CT, Fioretos T, Surace C, et al. MYC-containing double minutes in hematologic malignancies: evidence in favor of the episome model and exclusion of MYC as the target gene. *Hum Mol Genet*. 2006;15(6):933-942.
- Zatkova A, Schoch C, Speleman F, et al. GAB2 is a novel target of 11q amplification in AML/MDS. *Genes Chromosomes Cancer*. 2006;45(9):798-807.
- Fenaux P, Preudhomme C, Quiquandon I, et al. Mutations of the P53 gene in acute myeloid leukaemia. *Br J Haematol*. 1992;80(2):178-183.
- Stirewalt DL, Kopecky KJ, Meshinchi S, et al. FLT3, RAS, and TP53 mutations in elderly patients with acute myeloid leukemia. *Blood*. 2001;97(11):3589-3595. Erratum in: *Blood*. 2001;98(4):924.
- Haferlach C, Dicker F, Herholz H, et al. Mutations of the TP53 gene in acute myeloid leukemia are strongly associated with a complex aberrant karyotype. *Leukemia*. 2008;22(8):1539-1541.
- Bowen D, Groves MJ, Burnett AK, et al. TP53 gene mutation is frequent in patients with acute myeloid leukemia and complex karyotype, and is associated with very poor prognosis. *Leukemia*. 2009;23(1):203-206.
- Seifert H, Mohr B, Thiede C, et al. Study Alliance Leukemia (SAL): the prognostic impact of 17p (p53) deletion in 2272 adults with acute myeloid leukemia. *Leukemia*. 2009;23(4):656-663.
- Bennett JM, Catovsky D, Daniel MT, et al. Proposed revised criteria for the classification of acute myeloid leukemia: a report of the French-American-British Cooperative Group. *Ann Intern Med*. 1985;103(4):620-625.
- Jaffe ES, Harris NL, Stein H, et al. *Pathology and Genetics of Tumours of Haematopoietic and Lymphoid Tissues* (3rd Ed). Lyon, France: IARC Press; 2001.
- Schlenk RF, Benner A, Hartmann F, et al. Risk-adapted postremission therapy in acute myeloid leukemia: results of the German multicenter AML HD93 treatment trial. *Leukemia*. 2003;17(8):1521-1528.
- Schlenk RF, Döhner K, Mack S, et al. Prospective evaluation of allogeneic hematopoietic stem-cell transplantation from matched related and matched unrelated donors in younger adults with high-risk acute myeloid leukemia: German-Austrian trial AMLHD98A. *J Clin Oncol*. 2010;28(30):4642-4648.
- Schlenk RF, Frohling S, Hartmann F, et al. Phase III study of all-trans retinoic acid in previously untreated patients 61 years or older with acute myeloid leukemia. *Leukemia*. 2004;18(11):1798-1803.
- Shaffer LG, Tommerup N. *An International System for Human Cytogenetic Nomenclature*. Basel, Switzerland: S. Karger; 2005.
- Bullinger L, Krönke J, Schön C, et al. Identification of acquired copy number alterations and uniparental disomies in cytogenetically normal acute myeloid leukemia using high-resolution single-nucleotide polymorphism analysis. *Leukemia*. 2010;24(2):438-449.
- Zenz T, Kröber A, Scherer K, et al. Monoallelic TP53 inactivation is associated with poor prognosis in chronic lymphocytic leukemia: results from a detailed genetic characterization with long-term follow-up. *Blood*. 2008;112(8):3322-3329.
- Olivier M, Petitjean A, Marcel V, et al. Recent advances in p53 research: an interdisciplinary perspective. *Cancer Gene Ther*. 2009;16(1):1-12.
- Hamroun D, Kato S, Ishioka C, et al. The UMD TP53 database and website: update and revisions. *Hum Mutat*. 2006;(1):27:14-20.
- Krzywinski M, Schein J, Birol I, et al. Circos: an information aesthetic for comparative genomics. *Genome Res*. 2009;19(9):1639-1645.
- Kirsch DG, Kastan MB. Tumor-suppressor p53: implications for tumor development and prognosis. *J Clin Oncol*. 1998;16(9):3158-3168.
- Bouffler SD, Kemp CJ, Balmain A, et al. Spontaneous and ionizing radiation-induced chromosomal abnormalities in p53-deficient mice. *Cancer Res*. 1995;55(17):3883-3889.
- Jacks T, Remington L, Williams BO, et al. Tumor spectrum analysis in p53-mutant mice. *Curr Biol*. 1994;4(1):1-7.
- Liu PK, Kraus E, Wu TA, et al. Analysis of genomic instability in Li-Fraumeni fibroblasts with germline p53 mutations. *Oncogene*. 1996;12(11):2267-2278.
- Mekeel KL, Tang W, Kachnic LA, et al. Inactivation of p53 results in high rates of homologous recombination. *Oncogene*. 1997;14(15):1847-1857.
- Xu Y. A new role for p53 in maintaining genetic stability in embryonic stem cells. *Cell Cycle*. 2005;4(3):363-364.
- Song H, Hollstein M, Xu Y. p53 gain-of-function cancer mutants induce genetic instability by inactivating ATM. *Nat Cell Biol*. 2007;9(5):573-580.
- Christiansen DH, Andersen MK, Pedersen-Bjergaard J. Mutations with loss of heterozygosity of p53 are common in therapy-related myelodysplasia and acute myeloid leukemia after exposure to alkylating agents and significantly associated with deletion or loss of 5q, a complex karyotype, and a poor prognosis. *J Clin Oncol*. 2001;19(5):1405-1413.
- Jasek M, Gondek LP, Bejanyan N, et al. TP53 mutations in myeloid malignancies are either homozygous or hemizygous due to copy number-neutral loss of heterozygosity or deletion of 17p. *Leukemia*. 2010;24(1):216-219.
- Shing DC, Trubia M, Marchesi F, et al. Overexpression of sPRDM16 coupled with loss of p53 induces myeloid leukemias in mice. *J Clin Invest*. 2007;117(12):3696-3707.
- Kawasaki Y, Hirabayashi Y, Kaneko T, et al. Benzene-induced hematopoietic neoplasms including myeloid leukemia in Trp53-deficient C57BL/6 and C3H/He mice. *Toxicol Sci*. 2009;110(2):293-306.
- Zhao Z, Zuber J, Diaz-Flores E, et al. p53 loss promotes acute myeloid leukemia by enabling aberrant self-renewal. *Genes Dev*. 2010;24(13):1389-1402.
- Jädersten M, Saft L, Pellagatti A, et al. Clonal heterogeneity in the 5q- syndrome: p53 expressing progenitors prevail during lenalidomide treatment and expand at disease progression. *Haematologica*. 2009;94(12):1762-1766.
- Jädersten M, Saft L, Smith A, et al. TP53 mutations in low-risk myelodysplastic syndromes with del(5q) predict disease progression. *J Clin Oncol*. 2011;29(15):1971-1979.
- Link DC, Schuettpeiz LG, Shen D, et al. Identification of a novel TP53 cancer susceptibility mutation through whole-genome sequencing of a patient with therapy-related AML. *JAMA*. 2011;305(15):1568-1576.
- Donehower LA, Lozano G. 20 years studying p53 functions in genetically engineered mice. *Nat Rev Cancer*. 2009;9(11):831-841.
- Sigal A, Rotter V. Oncogenic mutations of the p53 tumor suppressor: the demons of the guardian of the genome. *Cancer Res*. 2000;60(24):6788-6793.
- Zuber J, Radtke I, Pardee TS, et al. Mouse models of human AML accurately predict chemotherapy response. *Genes Dev*. 2009;23(7):877-889.
- Yin B, Kogan SC, Dickins RA, et al. Trp53 loss during in vitro selection contributes to acquired Ara-C resistance in acute myeloid leukemia. *Exp Hematol*. 2006;34(5):631-641.