

Interferons limit inflammatory responses by induction of tristetraprolin

Ines Sauer, Barbara Schaljo, Claus Vogl, Irene Gattermeier, Thomas Kolbe, Mathias Müller, Perry J. Blackshear, and Pavel Kovarik

Interferons (IFNs) are cytokines with pronounced proinflammatory properties. Here we provide evidence that IFNs also play a key role in decline of inflammation by inducing expression of tristetraprolin (*Ttp*). TTP is an RNA-binding protein that destabilizes several AU-rich element-containing mRNAs including $TNF\alpha$. By promoting mRNA decay, TTP significantly contributes to cytokine homeostasis. Now we report that IFNs strongly stimulate expression of TTP if a costimulatory

stress signal is provided. IFN-induced expression of *Ttp* depends on the IFN-activated transcription factor STAT1, and the costimulatory stress signal requires p38 MAPK. Within the *Ttp* promoter we have identified a functional gamma interferon-activated sequence that recruits STAT1. Consistently, STAT1 is required for full expression of *Ttp* in response to LPS that stimulates both p38 MAPK and, indirectly, interferon signaling. We demonstrate that in macrophages IFN-induced

TTP protein limits LPS-stimulated expression of several proinflammatory genes, such as $TNF\alpha$, *IL-6*, *Ccl2*, and *Ccl3*. Thus, our findings establish a link between interferon responses and TTP-mediated mRNA decay during inflammation, and propose a novel immunomodulatory role of IFNs. (Blood. 2006;107:4790-4797)

© 2006 by The American Society of Hematology

Introduction

Interferons (IFNs) are cytokines that were originally described as polypeptides causing activation of antiviral responses.¹ Over the years it became clear that interferons play a major role in immune responses to essentially all infections.^{2,3} IFNs exert their biological effects by regulating gene transcription. Type II IFN (IFN- γ) induces phosphorylation of the transcription factor STAT1 at Tyr701 that causes STAT1:1 dimer formation, nuclear translocation, and DNA binding.^{4,5} Type I IFNs (eg, IFN- α and IFN- β) induce tyrosine phosphorylation of STAT1 and STAT2 causing the formation of STAT1:2 heterodimers and STAT1:1 homodimers.^{4,5} STAT1:2 heterodimers acquire their DNA binding ability by associating with the transcription factor IRF9 to form the transcription complex ISGF3. ISGF3 binds to ISRE elements, whereas STAT1 dimers target GAS elements in the promoters of interferon-regulated genes. For full biological activity, the transactivation domain of STAT1 requires phosphorylation at Ser727.^{6,7} IFNs cause phosphorylation of both Tyr701 and Ser727 residues. p38 MAPK induces Ser727 phosphorylation only, but can enhance STAT1-driven transcription independently of Ser727 phosphorylation.⁸⁻¹¹ Thus, STAT1 function and IFN-stimulated gene expression are modulated by p38 MAPK, one of the key inflammation-activated serine kinases. In fact, many inflammation-related genes require both IFNs and p38 MAPK for their maximal stimulation.^{3,12}

One of the major proinflammatory mediators is tumor necrosis factor alpha ($TNF\alpha$). An acute overproduction or chronically high levels of $TNF\alpha$ cause pathologic conditions such as septic shock or inflammatory arthritis, respectively.^{13,14} The immune system has developed several mechanisms to limit the production of this potentially harmful cytokine. Regulation of mRNA stability plays a major role in preventing overproduction of $TNF\alpha$.¹⁵ The 3' end of $TNF\alpha$ mRNA contains AU-rich elements (AREs) that are recognized by a group of ARE-binding proteins regulating mRNA stability.¹⁶⁻¹⁸ Gene-targeting experiments provided evidence that tristetraprolin (*Ttp*) is an anti-inflammatory gene that suppresses $TNF\alpha$ production by stimulating ARE-mediated $TNF\alpha$ mRNA decay.^{19,20} Mice deficient in TTP display elevated $TNF\alpha$ serum levels resulting in cachexia and inflammatory polyarthritis. *Ttp* is an immediate-early gene induced by inflammatory stimuli like LPS or $TNF\alpha$ with p38 MAPK playing a key role herein.^{20,21} TTP is also induced by IL-4 and TGF β through the activation of the transcription factors STAT6 and SMAD3/SMAD4, respectively.^{22,23} TTP regulates stability of mRNAs other than $TNF\alpha$ as well, most notably granulocyte-macrophage colony-stimulating factor (GM-CSF) and TTP itself.²⁴⁻²⁶ The biological activity of TTP is regulated by phosphorylation that plays a role in association of TTP with stress granules, the sites of ARE-mediated mRNA decay.^{21,27-29}

From the Max F. Perutz Laboratories, Department of Microbiology and Immunobiology, University of Vienna; the Austrian Center of Biomodelling and Transgenetics and the Institute of Animal Breeding and Genetics, Vienna University of Veterinary Medicine; the Department of Agrobiotechnology, IFA (Interuniversity Research Center)-Tulln, University of Natural Resources and Applied Life Sciences, Vienna, Austria; and the Laboratory of Neurobiology, National Institute of Environmental Health Science, Research Triangle Park, NC.

Submitted July 29, 2005; accepted February 2, 2006. Prepublished online as *Blood* First Edition Paper, March 2, 2006; DOI 10.1182/blood-2005-07-3058.

Supported by the Austrian Research Foundation (FWF) and the European Science Foundation (ESF) through grants P16 726-B14 and I27-B03 (P.K.) and FWF grant SFB F28 (P.K., M.M.); the Viennese Foundation for Science, Research, and Technology (WWTF; project LS133; M.M.); and the Austrian

Ministry of Education and Science (BMBWK; project BM:BWK OEZBT GZ200.074/I-VI/1a/2002; M.M.).

I.S. designed and performed research and analyzed data; P.K. designed research, analyzed data, and wrote the paper; B.S. and I.G. performed research and analyzed data; C.V. analyzed data; and T.K., M.M., and P.J.B. contributed new reagents.

Reprints: Pavel Kovarik, Max F. Perutz Laboratories, Department of Microbiology and Immunobiology, University of Vienna, Dr Bohr-Gasse 9, A-1030 Vienna, Austria; e-mail: pavel.kovarik@univie.ac.at.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 U.S.C. section 1734.

© 2006 by The American Society of Hematology

Previously we have shown that transcription of *Irf1*, a known IFN target gene, is synergistically induced by IFN and p38 MAPK signaling.¹⁰ In a DNA microarray-based screen we have identified *Ttp* as another gene synergistically induced by IFN- γ and p38 MAPK. We have characterized a functional GAS element in the *Ttp* promoter that is capable of recruiting STAT1. Our study shows that STAT1 is required for maximal expression of *Ttp* after stimulation of macrophages with LPS. LPS-induced expression of *Ttp* can be further strongly enhanced by cotreatment with IFNs. We further demonstrate that IFN-induced *Ttp* expression restrains the induction of the proinflammatory genes *TNF α* , *IL-6*, *Ccl2*, and *Ccl3*. Thus, by amplifying TTP expression IFNs generate a negative feedback that limits inflammatory signals elicited by both IFNs and microbial products.

Materials and methods

Cell culture

p38 α ^{-/-}, *p38 α ^{+/+}*, *STAT1^{-/-}*, and *STAT1^{+/+}* immortalized fibroblasts have been described recently.^{10,30} Cells were maintained in Dulbecco modified Eagle medium (DMEM) supplemented with 10% fetal calf serum (FCS). Primary bone marrow macrophages (BMMs) were obtained from 8- to 12-week-old mice and cultivated in L cell-derived CSF-1 as reported previously.³¹ *STAT1*-deficient mice³² were of mixed 129Sv/CD1 background, and *Ttp*-deficient¹⁹ as well as *IFN- β* -deficient mice were of C57Bl/6 background. Mice were housed under specific pathogen-free conditions. Mouse macrophage line Bac 1.2F5 was grown as described.³³ Mouse macrophage line J774 was grown in DMEM supplemented with 10% FCS.

Cytokines, reagents, and antibodies

Recombinant IFN- γ (kindly provided by G. Adolf, Boehringer Ingelheim, Austria) was used at 10 ng/mL for the times indicated in figure legends. Recombinant IFN- β was purchased from Calbiochem (Darmstadt, Germany) and used at 200 U/mL. LPS from *Salmonella minnesota* (Alexis, Lausen, Switzerland) and anisomycin (Sigma, Vienna, Austria) were used at 100 ng/mL for the times specified in figure legends. Antiserum to STAT1 C-terminus (S1) has been described.³⁴ Antibodies to Tyr701-phosphorylated STAT1 (pY701-S1) and phosphorylated p38 (pp38) were from Cell Signaling (NEB, Frankfurt/Main, Germany). Monoclonal antibodies to STAT1 N-terminus (S1-N) and to extracellular signal regulated kinases (panERK) were from BD Transduction Laboratories (BD Biosciences, Erembodegem, Belgium). p38 antibody was obtained from Santa Cruz Biotechnology (Santa Cruz, CA). Antibody to the C-terminus of TTP was kindly provided by A. R. Clark (Imperial College School of Medicine London, United Kingdom).

Transient transfection and luciferase assays

pGL2 luciferase reporter vectors, pGL2basic and pGL2promoter, were from Promega (Mannheim, Germany). Fragments of the mouse *Ttp* promoter, size 2 kb (-2025 to +25) and size 194 bp (-2085 to -1891), were amplified by polymerase chain reaction (PCR) from genomic mouse fibroblast DNA, and cloned into pGL2basic and pGL2promoter to obtain the plasmids pGL-TTP and pGL-TTP-GAS, respectively. Both constructs were verified by sequencing. The 2-kb fragment was amplified using the forward primer 5'-ATTCGAGGTTACAAGAGAAGGAACCCAA-3' and reverse primer 5'-ATAAGCTTTGTCGGTTCCGACAGAAGTCAGG-3', and inserted into pGL2basic using *XhoI* and *HindIII* restriction sites. The 194-bp fragment was amplified using 5'-ACGACGCGTCCGCCCTCTCTCCCTTCTTTCC-3' as forward and 5'-CCCTCGAGGGCAGTAGGCTGCTGTGGCACGG-3' as reverse primer and cloned into the pGL2promoter plasmid using *MluI* and *XhoI* to obtain the plasmid pGL2-TTP-GAS. Transfections were performed using ExGen500 (Fermentas, St Leon-Rot, Germany). pGL2-TTP together with pEF-Zeo³⁰ was cotransfected into

p38 α ^{+/+} cells and a bulk culture was established by a 2-week-long selection with 100 μ g/mL zeocin. pGL-TTP-GAS was transiently transfected into *STAT1^{+/+}* and *STAT1^{-/-}* fibroblasts. Transfection efficiency was controlled using ecdysone inducible system (Invitrogen, Lofer, Austria) as previously described.¹⁰

Luciferase assays were performed in triplicate, according to standard protocols.³⁵

ELISAs

For enzyme-linked immunosorbent assays (ELISAs), BMMs were seeded the day before use at 5×10^4 cells per well in 96-well tissue plates. Supernatants were diluted 1:7 in DMEM, and TNF α and IL-6 were assayed in triplicate using Quantikine kits (R&D Systems, Minneapolis, MN).

Western blot and electrophoretic mobility-shift assay (EMSA)

Western blot analysis was performed using whole-cell extracts from 2×10^6 BMMs as previously described.³⁴ For EMSA, whole-cell extracts from 5×10^6 Bac 1.2F5 cells were prepared and processed as described.³⁴ Oligonucleotides used in EMSA were as follows: TTP-GAS (fwd 5'-CTAGCGGCTTCCAGGAAGCCCG-3', rev 5'-GCCGAAGGTCCTTCGGGCGATC-3') and mutTTP-GAS (fwd 5'-CTAGCGGCTTAAAGGAAGCCCG-3', rev 5'-GCCGAATTTCTTCGGGCGATC-3').

Quantitation of gene expression by quantitative reverse transcription-PCR (qRT-PCR)

Total RNA was isolated using nucleospin reagent kit (Clontech, Mountain View, CA). Reverse transcription was performed with the Mu-MLV reverse transcriptase (Fermentas). The following primers were used: for *Hprt*, the housekeeping gene used for normalization, HPRT-fwd 5'-GGATTTGAATCAGCTTTGTGTCAT-3' and HPRT-rev 5'-ACACCTGCTAATTTTACTGCAA-3'; for TTP, TTP-fwd 5'-CTCTGCCATCTACGAGAGCC-3' and TTP-rev 5'-GATGGAGTCCGAGTTTATGTTCC-3'; for luciferase, Luc-fwd 5'-GCAGGTCTTCCCGACGATGA-3' and Luc-rev 5'-GTACTTCGTCCACAAACACAACACTC-3'. Amplification of DNA was monitored by SYBR Green (Molecular Probes, Eugene, OR).³⁶ For detection of TNF α , *Ccl2*, *Ccl3*, *Gbp2*, and *Cxcl10* Taqman assays from Applied Biosystems (Weiterstadt, Germany) were used.

Chromatin immunoprecipitation assay (ChIP)

STAT1-WT fibroblasts (2×10^6) were treated for 30 minutes with 10 ng/mL IFN- γ or left untreated. ChIP was performed using the Upstate Biotechnology protocol with slight modifications.⁷ Analysis of the immunoprecipitated DNA was performed by PCR using primers for the *Ttp* promoter (fwd 5'-ATTGGCTGGCTCAGGGATTTGT-3' and rev 5'-CATAGGCTGCTGTGGCACGG-3').

Nuclear run-on

Nuclear run-on reactions were performed as previously described,³⁷ with some modifications. Briefly, nuclei were prepared from 8×10^6 cells by lysis in hypotonic buffer (10 mM HEPES, pH 7.9, 1.5 mM MgCl₂, 10 mM KCl, 0.1% Nonidet P-40, 1 mM DTT), followed by 6 up and down strokes through a 25 G syringe. Nuclei were isolated by centrifugation (500g), and pellets were resuspended in 100 μ L glycerol buffer (40% glycerol, 50 mM Tris-HCl, pH 8.8, 5 mM MgCl₂, 0.1 mM EDTA). 2x reaction buffer (100 μ L containing 10 mM Tris-HCl, pH 8, 300 mM KCl, 5 mM MgCl₂, 5 mM DTT, and 1 mM each of ATP, CTP, and GTP), 100 U/ μ L ribonuclease inhibitor (Sigma), and 3.7×10^6 Bq (100 μ Ci) ³²P-labeled UTP (Perkin Elmer, Wellesley, MA) were added to the nuclei. After incubation for 30 minutes at room temperature and addition of 6 μ g DNase I (Sigma) for 10 minutes (at room temperature), RNA was isolated using Trizol reagent (Invitrogen) and ammonium acetate precipitation. Pellets were resuspended in 1 mL ETS hybridization buffer (10 mM Tris-HCl, pH 7.4, 10 mM EDTA, 0.2% SDS, 2x Denhardt, 400 mM NaCl, 200 μ g/mL yeast tRNA). cDNA (400 ng per slot) was slot-blotted on a nylon membrane (Perkin Elmer) and UV cross-linked. Membranes were prehybridized for 1 hour in TESS buffer

(20 mM TES, pH 7.4, 400 mM NaCl, 10 mM EDTA, 0.2% SDS, 2x Denhardt, 200 μ g/mL yeast tRNA) at 65°C. The prehybridization solution was removed and 1 mL of the labeled RNA was incubated with the membranes at 65°C for 18 hours. Membranes were washed twice in 2x SSC/1% SDS at 60°C and once in 0.1x SSC at room temperature.

Statistical analysis

Data from independent experiments were analyzed using univariate linear regression models and the SPSS program (SPSS, Chicago, IL). For qRT-PCR, untransformed normalized Ct-values, for ELISA, pg/mL were used as raw data. Residuals were plotted, visually inspected, and tested for normality. Design matrices were specified such that the coefficients for the relevant comparisons could be calculated, for example, between the baseline and induced states and between genotypes. Only the significance levels are reported.

Results

Interferons require STAT1 and activation of p38 MAPK to induce transcription of *Ttp*

STAT1 is activated by IFNs independently of p38 MAPK, which is, however, able to further enhance STAT1 transcriptional activity.^{10,11} We employed mouse 16K cDNA microarrays to find genes that are strongly activated only if both STAT1 (by IFN- γ) and p38 MAPK (by anisomycin) are stimulated (data not shown). We identified tristetraprolin (*Ttp*) to be regulated in the desired way and validated the DNA-microarray data by quantitative RT-PCR (qRT-PCR). Immortalized p38 $\alpha^{-/-}$ mouse embryonic fibroblasts (MEFs) and control MEFs were stimulated with IFN- γ , anisomycin, or both. p38 $\alpha^{-/-}$ cells are lacking the p38 α isoform, the most abundant of the known p38 α , β , γ , and δ isoforms. p38 $\alpha^{-/-}$ cells display a very weak p38 activity, which is due to the low expression of the remaining p38 genes.¹⁰ The p38 $\alpha^{-/-}$ cells have therefore been widely used to evaluate the contribution of p38 MAPK to various biological phenomena.^{10,11,38} A 1-hour treatment with anisomycin (a p38 MAPK agonist³⁹) induced *Ttp* mRNA levels in p38 $\alpha^{+/+}$ but not in p38 $\alpha^{-/-}$ cells (Figure 1A). The requirement for p38 MAPK in expression of *Ttp* is in agreement with previous studies employing the p38 MAPK inhibitor SB203 580.²¹ Treatment with IFN- γ for 1 hour weakly increased *Ttp* mRNA in p38 $\alpha^{+/+}$ or p38 $\alpha^{-/-}$ fibroblasts. However, a combined treatment with IFN- γ and anisomycin caused a strong synergistic induction of *Ttp* mRNA in p38 $\alpha^{+/+}$ but not in p38 $\alpha^{-/-}$ cells. The level of *Ttp* mRNA after the double treatment was 3 times higher than that

induced by anisomycin alone. The effect was more than additive: the cumulative induction by the single treatments with IFN- γ and anisomycin was 14-fold, whereas the induction by the double treatment was 33-fold. The synergistic induction by the double treatment as well as the effect of p38 MAPK were highly significant ($P < .01$). Similar results were obtained when IFN- β was used instead of IFN- γ (Figure 1B).

The transcription factor STAT1 is indispensable for the majority of responses to both types of IFNs.^{32,40} To find out whether STAT1 was involved in the IFN-mediated *Ttp* expression we employed immortalized STAT1 $^{-/-}$ MEFs and STAT1 $^{-/-}$ MEFs reconstituted with STAT1 cDNA (STAT1 $^{+/+}$).³⁰ The enhancing effect of IFN- γ (Figure 1C) or IFN- β (Figure 1D) on p38-stimulated *Ttp* expression was abolished in STAT1 $^{-/-}$ cells.

These data demonstrate that IFNs strongly induce *Ttp* expression if a p38 MAPK stimulus is provided simultaneously.

Ttp promoter contains a functional GAS element that binds STAT1

Ttp contains within the 3' UTR an ARE that regulates the mRNA stability.^{25,26} The low induction of *Ttp* mRNA levels by IFNs alone (without the p38 MAPK stimulus) could be explained by constitutive decay of *Ttp* mRNA due to the ARE. The destabilizing activity of AREs is known to be released by activation of p38 MAPK.^{18,41} Thus, the requirement of p38 MAPK in IFN-induced *Ttp* expression might reflect the need to stabilize the newly synthesized mRNA by p38 rather than a p38-derived input in the transcription machinery. To address this issue, we cloned a 2-kb fragment of the *Ttp* promoter into the pGL2basic vector to obtain the plasmid pGL-TTP. The cloned fragment spans the region from base pair -2025 to +25 and contains a GAS element (ie, STAT1 binding site) as described in Figure 2A. The plasmid pGL-TTP together with the pEF-Zeo vector plasmid (to allow for selection with zeocin)³⁰ were cotransfected into p38 $\alpha^{+/+}$ cells and a bulk culture was established. These cells were assayed using qRT-PCR for induction of luciferase mRNA by IFN- γ , anisomycin, or both (Figure 2B). The experiment revealed that both IFN- γ and p38 MAPK are needed to induce transcription of the reporter gene. Since the reporter gene does not contain ARE sequences we conclude that IFN- γ requires the activity of p38 MAPK for stimulating *Ttp* transcription rather than for inhibiting ARE-mediated decay of *Ttp* mRNA. To substantiate this finding and because of the lower induction of the reporter construct compared with endogenous *Ttp* we conducted nuclear run-on assays. Nuclear




Figure 1. p38 MAPK and STAT1 synergistically increase *Ttp* mRNA expression. p38 $\alpha^{+/+}$ and p38 $\alpha^{-/-}$ MEFs were left untreated (w/o) or treated for 1 hour with IFN- γ (g), anisomycin (a) or both (a/g) (A), or treated with IFN- β (b), anisomycin (a) or both (a/b) (B). STAT1 $^{+/+}$ and STAT1 $^{-/-}$ fibroblasts were stimulated as described for panels A and B, respectively (C,D). Total RNA was isolated and *Ttp* mRNA levels were determined using qRT-PCR. To obtain *Ttp* mRNA induction, values were normalized to those of untreated cells. Error bars indicate SD. * $P < .01$ (a/g) or (a/b) versus (a) treatment in $+/+$ cells; † $P < .01$ $+/+$ versus $-/-$ MEFs by univariate linear regression models; n = 3 experiments.




Figure 2. *Ttp* promoter contains a functional GAS element that binds STAT1. (A) Schematic representation of the *Ttp* promoter as cloned into the reporter construct pGL2-TTP, comprising the region from -2025 to $+25$. The *Ttp* promoter contains a GAS element at the position -1976 to -1964 . The reporter plasmid pGL2-TTP-GAS contains the TTP-GAS with flanking sequences (-2085 to -1891). The consensus for STAT1-binding GAS⁴² is shown (according to IUPAC nomenclature, H stands for A, C, or T; S stands for G or C; D stands for G, A, or T). Mutated TTP-GAS (TTP-GASMut) was obtained by introducing 2 point mutations. (B) p38^{+/+} cells stably transfected with a reporter plasmid containing the luciferase gene under the control of a 2-kb fragment of the *Ttp* promoter (pGL2-TTP) were left untreated (w/o) or treated for 1 hour with IFN- γ (g), anisomycin (a), or both (a/g). Total RNA was isolated and luciferase expression was assayed by qRT-PCR. Error bars indicate SD. * $P < .01$ (a/g) versus (a) treatment by univariate linear regression models; $n = 3$ experiments. (C) For nuclear run-on assay p38^{+/+} cells were left untreated (w/o) or treated for 25 minutes with IFN- γ (g), anisomycin (a), or both (a/g), nuclei were prepared and run-on reaction was performed. Nuclear RNA was isolated and hybridized to membranes containing cDNA of *Gapdh* (as a control) and *Ttp*. (D) Bac 1.2F5 mouse macrophages were stimulated for 30 minutes with IFN- γ (g) or left untreated (–), and whole-cell extracts were assayed for binding to a radioactively labeled TTP-GAS probe using EMSA. The results demonstrate IFN- γ -inducible binding of STAT1 to TTP-GAS but not to mutated TTP-GAS (TTP-GASMut). The presence of STAT1 in the complexes was confirmed by super-shift using STAT1 antibodies (lanes marked with +S1). (E) A luciferase reporter containing TTP-GAS (pGL-TTPGAS) was transfected into STAT1^{-/-} MEFs reconstituted with STAT1 cDNA (STAT1^{+/+}) and for control into the parental STAT1^{-/-} cells. After the transfection, cells were divided in 2 halves and 24 hours later treated for 6 hours with IFN- γ (g) or left untreated. Error bars indicate SD. † $P < .01$ +/+ versus –/– MEFs by univariate linear regression models; $n = 3$ experiments. (F) For ChIP, STAT1^{+/+} MEFs were treated for 30 minutes with IFN- γ (g) or left untreated (–). Recruitment of STAT1 to the TTP promoter was assayed using STAT1 antibodies (+S1). A control ChIP was performed using rabbit preimmune serum (C). Equal amount of material used in the ChIP experiments was confirmed by PCR-amplification of the input DNA (input).

run-ons assess transcription by measuring density of engaged RNA polymerases on specific promoters.³⁷ They provide a direct measure of transcriptional activity by eliminating any posttranscriptional effects. The nuclear run-on revealed that treatment with both anisomycin and IFN- γ strongly enhanced the transcriptional activity on the *Ttp* promoter, confirming the role of both stimuli in *Ttp* transcription (Figure 2C).

Activation of transcription by STAT1 is mediated by GAS elements, 13-bp-long stretches of DNA that recruit STAT1 dimers (Figure 2A).⁴² We have identified a potential GAS at the position -1976 to -1964 in the mouse *Ttp* promoter (Figure 2A). The TTP-GAS was investigated for its STAT1 binding activity using EMSA experiments with TTP-GAS and a mutated version (mutTTP-GAS) that contained 2 point mutations within the GAS core sequence (Figure 2A). We detected STAT1 binding to the TTP-GAS that was abolished by the point mutations (Figure 2D). The presence of STAT1 was confirmed by super-shift using a STAT1 antibody.

To investigate whether the TTP-GAS is a functional transcription-enhancing element, the TTP-GAS together with its flanking sequences (-2085 to -1891) was cloned into the pGLpromoter vector, and the resulting plasmid pGL-TTP-GAS was transfected into STAT1-WT cells and STAT1^{-/-} cells. Luciferase was induced in IFN- γ -stimulated STAT1-WT cells but not in STAT1^{-/-} cells (Figure 2E). The luciferase induction was reproducibly 2- to 3-fold, which is consistent with reported constructs containing one GAS element only.⁴³ Thus, we identified a functional GAS within the promoter of the mouse *Ttp* gene. Inspection of the human *Ttp* gene uncovered the presence of a GAS at a similar position, implicating a conserved transcriptional regulation of both human and mouse *Ttp*.

To strengthen the evidence for STAT1 binding to the *Ttp* promoter, chromatin immunoprecipitations (ChIPs) were performed. Chromatin was precipitated from IFN- γ -treated or untreated MEFs using a STAT1 antibody. DNA was recovered from the immunocomplexes and a fragment of the TTP promoter comprising the TTP-GAS was PCR amplified. The promoter fragment was amplified from the IFN- γ -treated sample, proving IFN- γ -inducible binding of STAT1 to the *Ttp* promoter (Figure 2F).

These data establish STAT1 to bind to the *Ttp* promoter, thereby facilitating transcription of the *Ttp* gene.

Maximal LPS-induced *Ttp* expression requires STAT1

To find out whether the increase of *Ttp* mRNA by the synergy of IFN and p38 MAPK pathways causes an equivalent rise in protein levels, Western blotting together with qRT-PCR analyses were performed. To activate p38 MAPK, LPS was applied instead of anisomycin since LPS is a physiological p38 agonist known to strongly induce TTP.^{20,21,27} Primary bone marrow-derived macrophages (BMMs) were stimulated for 1 hour (for qRT-PCR experiments; Figure 3A) or 3 hours (for Western blot analysis; Figure 3B) with IFN- γ , LPS, or both. The experiments revealed that IFN- γ caused only a weak increase of *Ttp* mRNA (Figure 3A) or TTP protein levels (Figure 3B). Stimulation of cells with LPS resulted in a significant induction of both mRNA and protein. However, a combined treatment of macrophages with LPS and IFN- γ caused a robust amplification of *Ttp* mRNA and protein that by far exceeded that induced by LPS alone. Thus, the induction of TTP protein by IFN- γ and/or LPS virtually mirrored the increase of *Ttp* mRNA that was mediated by the same stimuli. The qRT-PCR experiments were also performed in STAT1 knockout (KO) macrophages to confirm




Figure 3. STAT1 is required for full expression of *Ttp* in LPS-treated primary macrophages. (A) Macrophages derived from bone marrow (BMM) of STAT1WT and STAT1KO mice were left untreated or treated with IFN- γ (g), LPS (L), or both (L/g) and *Ttp* mRNA induction was measured by qRT-PCR and normalized to samples from untreated cells. Error bars indicate SD, $n = 3$ experiments. (B) BMMs isolated from STAT1WT mice were treated as explained for panel A, except the time of the treatment was 3 hours instead of 1 hour. TTP protein levels were analyzed by Western blotting of whole-cell extracts using a TTP antibody. The blot was reprobed with a panERK antibody to control for equal protein loading. (C) BMMs from STAT1WT and STAT1KO mice were treated for 1 hour and 4 hours with LPS or left untreated, and mRNA induction was analyzed as described in panel A. Error bars indicate SD, $n = 3$ experiments. (D) Same cells as those used in panel C were treated for 2 and 4 hours with LPS or left untreated. TTP protein was detected by Western blotting of whole-cell extracts using a TTP antibody. Activation of IFN signaling by endogenous production of type I interferon in LPS-treated macrophages was demonstrated using antibody to tyrosine-phosphorylated STAT1 (pY701-S1). Equal protein loading was confirmed using a STAT1 antibody and a panERK antibody. (E) BMMs from *IFN* β knockout (IFN β KO) and wild-type controls (IFN β WT) were stimulated with LPS for the times indicated. Total RNA was isolated and qRT PCR was performed. Error bars indicate SD, $n = 3$ experiments. (F) Whole-cell extracts of the same cells as used in panel E were stimulated with LPS as indicated. TTP protein was detected by Western blotting using a TTP antibody.

the STAT1 dependency (Figure 3A). Interestingly, LPS induced *Ttp* mRNA 50-fold in wild-type (WT) macrophages but only 25-fold in STAT1KO cells, indicating that STAT1 was also involved in LPS-induced *Ttp* expression. It is known that in macrophages LPS causes activation of MAPKs, NF κ B, and IRF3, resulting in rapid production of a range of cytokines including IFN- β .⁴⁴⁻⁴⁷ IFN- β in turn activates STAT1 and STAT2, and consequently the expression of GAS- and ISRE-containing genes, resulting in maximal activation of macrophages. Hence, LPS causes activation of both p38 MAPK and IFN pathways, thereby providing a system to examine their influence on gene expression under physiologic conditions. We used this system to investigate the biological significance of the synergistic effects of IFNs and p38 MAPK on *Ttp* expression. BMMs isolated from WT and STAT1KO mice were treated with LPS for 1 hour and 4 hours (in case of qRT-PCR; Figure 3C), and for 2 hours and 4 hours (in case of Western blots; Figure 3D). The experiments demonstrated that both TTP mRNA and protein are rapidly and strongly induced in WT macrophages whereas the induction in STAT1KO macrophages is weak and does not reach, at any time point, the levels detected in WT cells. LPS treatment causes phosphorylation of TTP, resulting in the appearance of slower migrating bands.²¹ Activation of STAT1 by LPS-induced endogenous production of IFN- β was revealed using phosphoTyr701-STAT1 antibody. To confirm the role of IFN- β in the full induction of TTP by LPS, we examined *Ttp* expression in BMM from IFN β KO mice (Figure 3E-F). The data show that initially (at 1 hour), LPS induced similar *Ttp* mRNA (Figure 3E) and protein (Figure 3F) levels in both control and IFN β KO cells. However, in control cells *Ttp* mRNA levels remained detectable and protein levels continued to rise whereas in the IFN β KO cells *Ttp* mRNA dropped close to basal levels at the 6-hour time point, and protein expression was declining. The *Ttp* expression profile, which is initially dependent mostly on the p38 MAPK and in the second phase also on IFN, is consistent with the biphasic expression of *Ttp* observed in RAW macrophages.²⁵


Our data proved that activation of STAT1 by the endogenous IFN- β is required for the maximal expression of *Ttp* in LPS-treated macrophages. The differences in *Ttp* expression between control and STAT1KO cells (Figure 3C-D) were more dramatic than those between control and IFN β KO (Figure 3E-F), indicating that other IFNs or STAT1-activating cytokines also play a role.

We conclude that *Ttp* mRNA and protein are induced by synergistic function of p38 MAPK and STAT1, and IFN/STAT1 signaling is required for full induction of *Ttp* expression by LPS.

IFN-stimulated *Ttp* expression limits induction of TNF α and IL-6 in activated macrophages

We asked whether the increased *Ttp* expression in macrophages treated with both IFNs and LPS compared with single treatments alone might affect the expression of TNF α , a known TTP target. BMMs from TTPKO and littermate TTPWT mice were treated with LPS, IFN- γ , or both, and the *TNF* α mRNA and protein levels were monitored (Figure 4A-B). IFN- γ did not induce *TNF* α mRNA in TTPWT cells, whereas it caused a weak but consistent 6- to 8-fold induction in TTPKO cells, indicating that IFN- γ can activate *TNF* α expression in the absence of TTP. LPS-induced *TNF* α mRNA levels were approximately 3-fold higher in TTPKO cells than in TTPWT cells, thus in agreement with published data on TTP-mediated destabilization of TNF α transcripts.^{19,48} Importantly, *TNF* α mRNA was even more induced in TTPKO macrophages treated with both IFN- γ and LPS, whereas it slightly decreased in double-treated TTPWT macrophages (Figure 4A). This effect resulted in 10-fold higher TNF α mRNA levels in double-treated TTPKO cells compared with TTPWT cells. Quantification of secreted TNF α revealed that during the treatment (2-4 hours) with both IFN- γ and LPS, the TNF α levels increased by 80% in supernatants of TTPKO but only by 5% in the supernatants of TTPWT macrophages (Figure 4B). This continuous production caused 3-fold higher

Figure 4. Interferon-stimulated *Ttp* expression limits induction of TNF α and IL-6. (A) BMMs derived from *Ttp* wild-type (TTPWT) and deficient (TTPKO) mice were left untreated or treated with IFN- γ (g), LPS (L), or both (L/g) for 3 hours and TNF α mRNA induction was determined by qRT-PCR and normalized to untreated samples. * $P < .05$ treated versus untreated cells by univariate linear regression models. (B) BMMs from TTPWT and TTPKO were stimulated for the times indicated with IFN- γ (G), LPS, or both (L/G), or left untreated (0). Supernatants were collected and analyzed for TNF α cytokine by ELISA. * $P < .01$, 4 hours versus 2 hours in L/G-treated KO cells; † $P < .01$, KO versus WT BMMs by univariate linear regression models, $n = 3$ experiments. (C) TTPWT BMMs were treated for 30 minutes (30') with IFN- γ and LPS, and activation of p38 was demonstrated using an antibody against phosphorylated p38 MAPK (pp38). Equal protein loading was confirmed using a p38 antibody. (D) BMMs from TTPWT and TTPKO mice were treated as described in panel A. Absence of TTP protein in TTPKO cells was confirmed by Western blotting of whole-cell extracts using a TTP antibody. Activation of IFN signaling by endogenous production of type I interferon in LPS-treated macrophages was demonstrated using an antibody to tyrosine-phosphorylated STAT1 (pY701-S1). Equal protein loading was confirmed using a STAT1 antibody and a panERK antibody. (E) BMMs from TTPWT and TTPKO were stimulated for the times indicated with IFN- β (B), LPS, or both (L/B) or left untreated (0). Supernatants were collected and IL-6 was measured by ELISA. * $P < .01$, 6 hours versus 4 hours in L/B-treated KO cells; † $P < .01$, KO versus WT BMMs by univariate linear regression models, $n = 3$ experiments.



TNF α levels after 4 hours in supernatants of TTPKO compared with control cells. Importantly, although LPS caused higher production of TNF α in TTPKO cells compared with TTPWT cells, the total TNF α levels remained below those of double-treated cells and the increase over time remained low (15% in TTPKO versus no increase in TTPWT cells). IFN- γ alone did not induce TNF α secretion in either cell type, although it caused a modest increase of *TNF α* mRNA in TTPKO cells. This result is consistent with the inability of IFN- γ to activate p38 MAPK, which is required for translation of *TNF α* mRNA and secretion of the cytokine.¹⁷ The lack of p38 MAPK activation by IFN- γ alone is shown in Figure 4C. To rule out that an increased IFN signaling in TTPKO cells was responsible for the higher IFN- γ -dependent *TNF α* expression in TTPKO macrophages, we analyzed the activation of STAT1. Figure 4D demonstrates that activation of STAT1 by IFN- γ was not augmented in TTPKO cells. In fact, we observed an approximately 2-fold lower STAT1 activation at very early time points (data not shown).

One of the other major cytokines produced by activated macrophages is IL-6. To investigate whether TTP plays a similar role in IL-6 expression as in TNF α expression, we performed ELISA to detect secreted IL-6 in TTPKO and control BMMs stimulated with IFN- β , LPS, or both (Figure 4E). Similar to TNF α , the production of IL-6 was strongly enhanced by double treatment (LPS and IFN- β) in TTPKO cells and raised by 60% between 4 hours and 6 hours of treatment, whereas in TTPWT cells IL-6 was 2-fold lower after 4 hours and increased only by 5% after 6 hours. Importantly, IFN- β alone did not induce IL-6 in either cell type, and LPS caused only a weak IL-6 expression that was 20% higher in TTPKO cells.

These results demonstrate that IFNs have the potential to strongly augment LPS-induced TNF α and IL-6 expression, and this effect is counter-balanced by the simultaneous induction of TTP.

IFNs exhibit a TTP-dependent suppressive effect on expression of *Ccl2* and *Ccl3*

We reasoned that the IFN-induced TTP protein might act on other proinflammatory molecules such as chemokines. According to the ARE database, human *Ccl3* contains AREs in its 3' UTRs.⁴⁹ Similar sequences are found in the mouse chemokine genes *Ccl2* (MCP1) and *Ccl3* (MIP-1 α). We performed qRT-PCR analysis of the *Ccl2* and *Ccl3* mRNA isolated from TTPKO and littermate TTPWT BMMs treated with IFN- γ , LPS, or both. Figure 5A shows that *Ccl2* mRNA was induced 3- to 4-fold in TTPWT cells treated with either stimulus. In TTPKO cells the induction was 8-fold by IFN- γ , 12-fold by LPS, and augmented to 16-fold by the double treatment. These data suggest that *Ccl2* is a TTP target and that the synergistic induction of *Ccl2* by LPS and IFN- γ is suppressed by the simultaneous synergistic induction of TTP. Similar analysis of *Ccl3* expression revealed that in TTPWT cells *Ccl3* mRNA levels were 60% lower upon combined LPS and IFN- γ treatment as compared with LPS alone (Figure 5B). In contrast, in TTPKO cells treated with both LPS and IFN- γ slightly increased *Ccl3* mRNA if compared with LPS treatment alone. The expression of the *Gbp2* or *Cxcl10* genes, which are both stimulated by IFN- γ and do not contain an obvious ARE in their 3' UTRs, was not (in the case of *Cxcl10*) or only modestly (in the case of *Gbp2*) affected by the TTP deficiency (Figure 5C-D). Thus, IFN-induced TTP can specifically reduce expression of a subset of inflammation-related genes that contain AREs in their mRNAs.

Discussion

This work emphasizes the role of IFNs in the negative control of inflammatory responses. The negative control mechanism is based on the ability of IFNs to induce the expression of *Ttp*, a key




Figure 5. Interferon- and LPS-induced *Ttp* expression has a suppressive effect on *Ccl2* and *Ccl3* mRNA production. BMMs derived from TTP wild-type (TTPWT) and deficient (TTPKO) mice were left untreated or treated with IFN- γ (g), LPS (L), or both (L/g) for 3 hours. Total RNA was extracted and the induction of *Ccl2* (A), *Ccl3* (B), *Gbp2* (C), and *Cxcl10* (D) genes was detected by real-time RT-PCR using Taqman assays from Applied Biosystems. To obtain mRNA induction, qRT-PCR values were normalized to those of untreated cells. * $P < .05$, treated versus untreated cells by univariate linear regression models.

mediator of the decay of several inflammatory mRNAs that contain AU-rich elements in their 3' UTRs.

Both IFN and p38 MAPK pathways are stimulated in macrophages upon challenge with microbes or microbial products. These signaling cascades cause, either individually or in synergy with each other, vast changes in gene expression. The reprogramming of gene expression is essential for activation of immune responses but also for balancing them. For the maintenance of immune homeostasis both signaling pathways elicit various negative regulatory mechanisms. The most prominent inhibitory molecules induced by IFNs are SOCS1 and SOCS3.⁵⁰ Feedback inhibition caused by p38 MAPK is brought about by transcriptional induction of diverse phosphatases acting either directly on p38 MAPK or on upstream activating enzymes such as MKK3.⁵¹ In addition, p38 MAPK plays a key role in anti-inflammatory responses by increasing levels of TTP that possesses strong anti-inflammatory properties and destabilizes the mRNA of the most potent immunostimulating cytokine, TNF α .^{19-21,25,26} p38 MAPK also negatively interferes with responses to IFNs by stimulating the expression of SOCS1.⁵⁰ On the other hand, there is only a limited evidence for IFNs playing a role in suppressing the proinflammatory function of p38 MAPK that is elicited mainly by induction of TNF α . In this work, we show for the first time that IFNs and the IFN-activated transcription factor STAT1 play a fundamental role in expression of *Ttp*, an important factor in negative regulation of cytokine expression. We demonstrate that for the full induction of TTP by LPS the autocrine, IFN- β -mediated activation of STAT1 is required. These data suggest that IFNs can augment the feedback inhibition exerted by TTP on cytokine (eg, TNF α) production. We confirmed this hypothesis by employing bone marrow-derived macrophages from TTPKO mice. The expression of TNF α , IL-6, *Ccl2*, and *Ccl3* was augmented in TTPKO cells treated with both LPS and IFNs, as compared with single treatments alone. In contrast, the expression of these genes in wild-type cells was only moderately increased or even reduced by the combined treatment as compared with single treatments. Thus, the strong proinflammatory activity of IFNs is limited by their ability to induce expression of *Ttp*. Since *Ttp* has to be transcribed, translated, and posttranscriptionally modified before engagement in the regulatory loop, it is likely that the *Ttp*-mediated negative feedback becomes apparent at later phases of inflammatory responses. In this scenario, IFN and LPS first act synergistically on activation of proinflammatory molecules that are, however, down-regulated by the synergistic increase of TTP levels later on. We have employed IFN- β -deficient cells to prove

the role of the autocrine production of IFN- β in LPS-mediated induction of *Ttp*. Interestingly, the *Ttp* expression was more efficiently affected by the lesion in STAT1 than in the IFN- β gene. Since mouse macrophages do not produce IFN- γ in response to LPS,⁵² the members of the IFN- α gene family^{2,53} might be responsible for the lower reduction of *Ttp* expression in the IFN- β -deficient cells. However, in macrophages the IFN- α production has been reported to depend entirely on IFN- β synthesis.⁵⁴ Thus, the more likely STAT1 agonists in IFN- β knockout cells are cytokines, such as IL-6, that are released from LPS-treated macrophages and known to activate (albeit weakly) STAT1.

In this study we have characterized a GAS element within the *Ttp* promoter that is conserved in mice and men. The TTP-GAS binds STAT1, yet STAT1 needs, in addition to IFNs, a p38-derived stimulus to activate *Ttp* transcription. This finding is consistent with reports on the role of p38 MAPK in promoting STAT1-driven transcription.^{10,11} While the molecular mechanism of the synergy is currently not known, our findings indicate that the number of IFN-stimulated genes (currently more than 300⁵⁵), will grow considerably if the IFN stimulus is supported by a p38 MAPK agonist.

In conclusion, we establish IFNs and the IFN-activated transcription factor STAT1 to play an important role in the transcription of *Ttp*, a major ARE-directed TNF α mRNA destabilizing factor. The IFN signal must be accompanied by a p38 MAPK stimulus for STAT1 to become transcriptionally active at the *Ttp* promoter. Under physiological conditions, both signaling pathways (IFN and p38 MAPK) are activated when macrophages encounter bacteria or bacteria-derived products such as LPS. Consistently, STAT1 is required for complete induction of *Ttp* in LPS-treated macrophages. The IFN-dependent *Ttp* induction exerts a negative effect on the expression of several ARE-containing proinflammatory molecules, thereby limiting potentially harmful consequences of uncontrolled cytokine or chemokine production. These findings reveal a previously unrecognized immunomodulatory mechanism by which IFNs contribute to immune system homeostasis.

Acknowledgments

We wish to thank Andy Clark for kindly providing us with TTP antibodies, and Matthias Gaestel and Alexey Kotlyarov for sharing TTP mice. Birgit Strobl, Silvia Stockinger, and Heather Zwaferink are gratefully acknowledged for critically reading the manuscript.

References

- Sen GC. Viruses and interferons. *Annu Rev Microbiol.* 2001;55:255-281.
- Bogdan C. The function of type I interferons in antimicrobial immunity. *Curr Opin Immunol.* 2000;12:419-424.
- Schroder K, Hertzog PJ, Ravasi T, Hume DA. Interferon-gamma: an overview of signals, mechanisms and functions. *J Leukoc Biol.* 2004;75:163-189.
- Schindler C, Brutsaert S. Interferons as a paradigm for cytokine signal transduction. *Cell Mol Life Sci.* 1999;55:1509-1522.
- Levy DE, Darnell JE Jr. Stats: transcriptional control and biological impact. *Nat Rev Mol Cell Biol.* 2002;3:651-662.
- Wen Z, Zhong Z, Darnell JE Jr. Maximal activation of transcription by Stat1 and Stat3 requires both tyrosine and serine phosphorylation. *Cell.* 1995;82:241-250.
- Varinou L, Ramsauer K, Karaghiosoff M, et al. Phosphorylation of the Stat1 transactivation domain is required for full-fledged IFN-gamma-dependent innate immunity. *Immunity.* 2003;19:793-802.
- Kovarik P, Stoiber D, Eysers PA, et al. Stress-induced phosphorylation of STAT1 at Ser727 requires p38 mitogen-activated protein kinase whereas IFN-gamma uses a different signaling pathway. *Proc Natl Acad Sci U S A.* 1999;96:13956-13961.
- Gollob JA, Schnipper CP, Murphy EA, Ritz J, Frank DA. The functional synergy between IL-12 and IL-2 involves p38 mitogen-activated protein kinase and is associated with the augmentation of STAT serine phosphorylation. *J Immunol.* 1999;162:4472-4481.
- Ramsauer K, Sadzak I, Porras A, et al. p38 MAPK enhances STAT1-dependent transcription independently of Ser-727 phosphorylation. *Proc Natl Acad Sci U S A.* 2002;99:12859-12864.
- Li Y, Sassano A, Majchrzak B, et al. Role of p38alpha Map kinase in type I interferon signaling. *J Biol Chem.* 2004;279:970-979.
- Iwasaki A, Medzhitov R. Toll-like receptor control of the adaptive immune responses. *Nat Immunol.* 2004;5:987-995.
- Tracey KJ, Cerami A. Tumor necrosis factor: a pleiotropic cytokine and therapeutic target. *Annu Rev Med.* 1994;45:491-503.
- Feldmann M, Brennan FM, Maini RN. Rheumatoid arthritis. *Cell.* 1996;85:307-310.
- Anderson P, Phillips K, Stoecklin G, Kedersha N. Post-transcriptional regulation of proinflammatory proteins. *J Leukoc Biol.* 2004;76:42-47.
- Kontoyiannis D, Pasparakis M, Pizarro TT, Cominelli F, Kollias G. Impaired on/off regulation of TNF biosynthesis in mice lacking TNF AU-rich elements: implications for joint and gut-associated immunopathologies. *Immunity.* 1999;10:387-398.
- Kotlyarov A, Neiningner A, Schubert C, et al. MAPKAP kinase 2 is essential for LPS-induced TNF-alpha biosynthesis. *Nat Cell Biol.* 1999;1:94-97.
- Kontoyiannis D, Kotlyarov A, Carballo E, et al. Interleukin-10 targets p38 MAPK to modulate ARE-dependent TNF mRNA translation and limit intestinal pathology. *EMBO J.* 2001;20:3760-3770.
- Taylor GA, Carballo E, Lee DM, et al. A pathogenic role for TNF alpha in the syndrome of cachexia, arthritis, and autoimmunity resulting from tristetraprolin (TTP) deficiency. *Immunity.* 1996;4:445-454.
- Carballo E, Lai WS, Blackshear PJ. Feedback inhibition of macrophage tumor necrosis factor-alpha production by tristetraprolin. *Science.* 1998;281:1001-1005.
- Mahtani KR, Brook M, Dean JL, Sully G, Saklatvala J, Clark AR. Mitogen-activated protein kinase p38 controls the expression and posttranslational modification of tristetraprolin, a regulator of tumor necrosis factor alpha mRNA stability. *Mol Cell Biol.* 2001;21:6461-6469.
- Suzuki K, Nakajima H, Ikeda K, et al. IL-4-Stat6 signaling induces tristetraprolin expression and inhibits TNF-alpha production in mast cells. *J Exp Med.* 2003;198:1717-1727.
- Ogawa K, Chen F, Kim YJ, Chen Y. Transcriptional regulation of tristetraprolin by transforming growth factor-beta in human T cells. *J Biol Chem.* 2003;278:30373-30381.
- Carballo E, Lai WS, Blackshear PJ. Evidence that tristetraprolin is a physiological regulator of granulocyte-macrophage colony-stimulating factor messenger RNA deadenylation and stability. *Blood.* 2000;95:1891-1899.
- Tchen CR, Brook M, Saklatvala J, Clark AR. The stability of tristetraprolin mRNA is regulated by mitogen activated protein kinase p38 and by tristetraprolin itself. *J Biol Chem.* 2004;279:32393-32400.
- Brooks SA, Connolly JE, Rigby WF. The role of mRNA turnover in the regulation of tristetraprolin expression: evidence for an extracellular signal-regulated kinase-specific, AU-rich element-dependent, autoregulatory pathway. *J Immunol.* 2004;172:7263-7271.
- Stoecklin G, Stubbs T, Kedersha N, et al. MK2-induced tristetraprolin:14-3-3 complexes prevent stress granule association and ARE-mRNA decay. *EMBO J.* 2004;23:1313-1324.
- Chrestensen CA, Schroeder MJ, Shabanowitz J, et al. MAPKAP kinase 2 phosphorylates tristetraprolin on in vivo sites including Ser178, a site required for 14-3-3 binding. *J Biol Chem.* 2004;279:10176-10184.
- Rigby WF, Roy K, Collins J, et al. Structure/function analysis of tristetraprolin (TTP): p38 stress-activated protein kinase and lipopolysaccharide stimulation do not alter TTP function. *J Immunol.* 2005;174:7883-7893.
- Kovarik P, Mangold M, Ramsauer K, et al. Specificity of signaling by STAT1 depends on SH2 and C-terminal domains that regulate Ser727 phosphorylation, differentially affecting specific target gene expression. *EMBO J.* 2001;20:91-100.
- Baccarini M, Bistoni F, Lohmann-Matthes ML. In vitro natural cell-mediated cytotoxicity against *Candida albicans*: macrophage precursors as effector cells. *J Immunol.* 1985;134:2658-2665.
- Durbin JE, Hackenmiller R, Simon MC, Levy DE. Targeted disruption of the mouse Stat1 gene results in compromised innate immunity to viral disease. *Cell.* 1996;84:443-450.
- Morgan C, Pollard JW, Stanley ER. Isolation and characterization of a cloned growth factor dependent macrophage cell line, BAC1.2F5. *J Cell Physiol.* 1987;130:420-427.
- Kovarik P, Stoiber D, Novy M, Decker T. Stat1 combines signals derived from IFN-gamma and LPS receptors during macrophage activation [published erratum appears in *EMBO J* 1998 Jul 15;17:4210]. *EMBO J.* 1998;17:3660-3668.
- Sambrook J, Fritsch EF, Maniatis T. *Molecular Cloning: A Laboratory Manual.* Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press; 1989.
- Morrison TB, Weis JJ, Wittwer CT. Quantification of low-copy transcripts by continuous SYBR Green I monitoring during amplification. *Biotechniques.* 1998;24:954-958,960,962.
- Groudine M, Peretz M, Weintraub H. Transcriptional regulation of hemoglobin switching in chicken embryos. *Mol Cell Biol.* 1981;1:281-288.
- Porras A, Zuluaga S, Black E, et al. P38 alpha mitogen-activated protein kinase sensitizes cells to apoptosis induced by different stimuli. *Mol Biol Cell.* 2004;15:922-933.
- Fukunaga R, Hunter T. MNK1, a new MAP kinase-activated protein kinase, isolated by a novel expression screening method for identifying protein kinase substrates. *EMBO J.* 1997;16:1921-1933.
- Meraz MA, White JM, Sheehan KC, et al. Targeted disruption of the Stat1 gene in mice reveals unexpected physiologic specificity in the JAK-STAT signaling pathway. *Cell.* 1996;84:431-442.
- Rousseau S, Morrice N, Pegg M, Campbell DG, Gaestel M, Cohen P. Inhibition of SAPK2a/p38 prevents hnRNP A0 phosphorylation by MAPKAP-K2 and its interaction with cytokine mRNAs. *EMBO J.* 2002;21:6505-6514.
- Decker T, Kovarik P, Meinke A. GAS elements: a few nucleotides with a major impact on cytokine-induced gene expression. *J Interferon Cytokine Res.* 1997;17:121-134.
- Chen ZY, Shie JL, Tseng CC. STAT1 is required for IFN-gamma-mediated gut-enriched Kruppel-like factor expression. *Exp Cell Res.* 2002;281:19-27.
- Gao J, Morrison DC, Parmely TJ, Russell SW, Murphy WJ. An interferon-gamma-activated site (GAS) is necessary for full expression of the mouse iNOS gene in response to interferon-gamma and lipopolysaccharide. *J Biol Chem.* 1997;272:1226-1230.
- Gao JJ, Filla MB, Fultz MJ, Vogel SN, Russell SW, Murphy WJ. Autocrine/paracrine IFN-alpha-beta mediates the lipopolysaccharide-induced activation of transcription factor Stat1alpha in mouse macrophages: pivotal role of Stat1alpha in induction of the inducible nitric oxide synthase gene. *J Immunol.* 1998;161:4803-4810.
- Sing A, Merlin T, Knopf HP, et al. Bacterial induction of beta interferon in mice is a function of the lipopolysaccharide component. *Infect Immun.* 2000;68:1600-1607.
- Toshchakov V, Jones BW, Perera PY, et al. TLR4, but not TLR2, mediates IFN-beta-induced STAT1alpha/beta-dependent gene expression in macrophages. *Nat Immunol.* 2002;3:392-398.
- Lai WS, Carballo E, Strum JR, Kennington EA, Phillips RS, Blackshear PJ. Evidence that tristetraprolin binds to AU-rich elements and promotes the deadenylation and destabilization of tumor necrosis factor alpha mRNA. *Mol Cell Biol.* 1999;19:4311-4323.
- Bakheet T, Williams BR, Khabar KS. ARE2.0: an update of AU-rich element mRNA database. *Nucleic Acids Res.* 2003;31:421-423.
- Alexander WS, Hilton DJ. The role of suppressors of cytokine signaling (SOCS) proteins in regulation of the immune response. *Annu Rev Immunol.* 2004;22:503-529.
- English J, Pearson G, Wilsbacher J, et al. New insights into the control of MAP kinase pathways. *Exp Cell Res.* 1999;253:255-270.
- Schleicher U, Hesse A, Bogdan C. Minute numbers of contaminant CD8⁺ T cells or CD11b⁺-CD11c⁺ NK cells are the source of IFN-gamma in IL-12/IL-18-stimulated mouse macrophage populations. *Blood.* 2005;105:1319-1328.
- Marie I, Durbin JE, Levy DE. Differential viral induction of distinct interferon-alpha genes by positive feedback through interferon regulatory factor-7. *EMBO J.* 1998;17:6660-6669.
- Stockinger S, Reutterer B, Schaljo B, et al. IFN regulatory factor 3-dependent induction of type I IFNs by intracellular bacteria is mediated by a TLR- and Nod2-independent mechanism. *J Immunol.* 2004;173:7416-7425.
- de Veer MJ, Holko M, Frevel M, et al. Functional classification of interferon-stimulated genes identified using microarrays. *J Leukoc Biol.* 2001;69:912-920.