

Identification of an active, well-tolerated dose of pralatrexate in patients with relapsed or refractory cutaneous T-cell lymphoma

Steven M. Horwitz,¹ Youn H. Kim,² Francine Foss,³ Jasmine M. Zain,⁴ Patricia L. Myskowski,¹ Mary Jo Lechowicz,⁵ David C. Fisher,⁶ Andrei R. Shustov,⁷ Nancy L. Bartlett,⁸ Maria L. Delioukina,⁹ Tony Koutsoukos,⁹ Michael E. Saunders,¹⁰ Owen A. O'Connor,¹¹ and Madeleine Duvic¹²

¹Department of Medicine, Memorial Sloan-Kettering Cancer Center, New York, NY; ²Stanford Cancer Center, Stanford University, Stanford, CA; ³Yale Medical Oncology, Yale Cancer Center, New Haven, CT; ⁴Hematology/Oncology, New York University Langone Medical Center, New York, NY; ⁵Hematology and Medical Oncology, Winship Cancer Institute of Emory University School of Medicine, Atlanta, GA; ⁶Department of Medicine, Dana-Farber Cancer Institute, Boston, MA; ⁷University of Washington School of Medicine, Seattle, WA; ⁸Chair in Medical Oncology, Washington University School of Medicine, St Louis, MO; ⁹Blood and Marrow Transplant Program, Cedars-Sinai Samuel Oschin Cancer Center, Los Angeles, CA; ¹⁰Allos Therapeutics Inc, Princeton, NJ; ¹¹Hematology/Oncology, Irving Cancer Research Center, The New York Presbyterian Hospital-Columbia University Medical Center, New York, NY; and ¹²Department of Dermatology, Division of Internal Medicine, University of Texas MD Anderson Cancer Center, Houston, TX

Systemic treatment for cutaneous T-cell lymphoma (CTCL) involves the use of less aggressive, well-tolerated therapies. Pralatrexate is a novel antifolate with high affinity for reduced folate carrier-1. A dose de-escalation strategy identified recommended pralatrexate dosing for patients with CTCL that demonstrated high activity, good rates of disease control, and an acceptable toxicity profile for continuous long-term dosing. Eligibility included mycosis fungoides, Sézary syndrome, or

primary cutaneous anaplastic large cell lymphoma, with disease progression after ≥ 1 prior systemic therapy. The starting dose and schedule was 30 mg/m²/wk intravenously for 3 of 4 (3/4) weeks. Subsequent starting doses were 20, 15, and 10 mg/m²/wk for 3/4 or 2 of 3 (2/3) weeks. Response was evaluated by the modified severity-weighted adjustment tool. Fifty-four patients were treated. The recommended regimen was identified as 15 mg/m²/wk for 3/4 weeks and was explored in

the expansion cohort. In 29 patients treated overall with the recommended dosing regimen, the median number of prior systemic therapies was 4. Pralatrexate was administered for a median of 4 cycles; response rate was 45%. The most common grade 3 adverse event (AE) was mucositis (17%); the only grade 4 AE was leukopenia (3%). Pralatrexate 15 mg/m²/wk for 3/4 weeks shows high activity with acceptable toxicity in patients with relapsed/refractory CTCL. (*Blood*. 2012;119(18):4115-4122)

Introduction

Cutaneous T-cell lymphomas (CTCLs) are typically indolent T-cell lymphomas that present primarily or exclusively in the skin. The term CTCL comprises several distinct clinical entities including mycosis fungoides (MF), the Sézary syndrome (SS), and primary cutaneous anaplastic large cell lymphoma (ALCL), which are all characterized by infiltration of the skin by malignant T cells.¹

Therapeutic approaches for patients with CTCL depend on the disease stage. For early stage disease, skin-directed therapies are generally used, with systemic treatments reserved for patients with relapsed or more extensive disease.^{2,3} Durable remissions off-therapy are uncommon in MF/SS, and, historically, more aggressive first-line approaches have not resulted in improved outcomes, but are associated with increased toxicity.⁴ In pivotal studies of approved systemic therapies for advanced relapsed/refractory CTCLs, response rates were reported for vorinostat (30%), denileukin diftitox (30%), oral bexarotene (45%-54%), and romidepsin (34%-35%), with median response durations of 6 to 15 months.⁵⁻¹⁰ Recently, denileukin diftitox demonstrated a 44% response rate versus placebo (15.9% response rate) in a randomized study, with a median progression-free survival (PFS) of > 2 years compared with 124 days in the placebo group.¹¹ Several other agents are used

in CTCL treatment, although their use is generally derived from smaller studies.¹²⁻¹⁵ Traditional cytotoxic chemotherapy agents are used later in the disease course because of risks of myelosuppression associated with poor skin integrity and the underlying immunosuppression associated with MF/SS.^{14,16-19} The moderate response rates and frequent lack of durable responses to current therapies for CTCL underline the need for additional effective and tolerable treatments.

Pralatrexate (FOLOTYN; Allos Therapeutics Inc), an antineoplastic folate analog, has high affinity for the reduced folate carrier type-1 oncoprotein and is an efficient substrate for polyglutamyl-ation by the enzyme folylpolyglutamyl synthetase, resulting in extensive internalization and accumulation within tumor cells. Pralatrexate inhibits dihydrofolate reductase, resulting in disruption of DNA synthesis and subsequent tumor cell death. This agent has been extensively studied in preclinical and clinical settings, alone and in combination,²⁰ and has been shown to have superior activity compared with methotrexate against human non-Hodgkin lymphoma in both in vitro and in vivo models.^{21,22} The pharmacokinetics and toxicology of pralatrexate, as well as its efficacy and safety in peripheral T-cell lymphoma (PTCL), are previously well

Submitted November 30, 2011; accepted February 12, 2012. Prepublished online as *Blood* First Edition paper, March 6, 2012; DOI 10.1182/blood-2011-11-390211.

Presented in poster form (Poster 2800) at the American Society of Hematology 52nd Annual Meeting, Orlando, FL, December 4, 2010.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2012 by The American Society of Hematology

described.²³⁻³¹ Pralatrexate received accelerated approval from the US Food and Drug Administration for patients with relapsed or refractory PTCL at a dose of 30 mg/m² weekly by IV push for 6 of 7 (6/7) weeks, based on the results of the PROPEL study.²⁹ To reduce the risk of mucositis, the most common toxicity encountered in the PROPEL study, vitamin B₁₂ and folate supplementation, are used with treatment.^{23,26,28,32}

CTCL is approached as a more indolent disease compared with PTCL, and combination chemotherapies are rarely used. Because of the activity in several patients with CTCL in prior phase 1-2 pralatrexate studies at various doses, this dose de-escalation study (PDX-010) was designed to try to identify an effective dose with acceptable toxicity for patients with CTCL.^{29,32}

Methods

This study was approved by the institutional review boards of all participating institutions and was conducted in accordance with the principles set forth in the Declaration of Helsinki.

Patients

Eligible patients included those with histologically confirmed CTCL subtypes of MF (stage \geq IB), SS, or primary cutaneous ALCL with measurable disease at the time of enrollment. Patients must have progressed after \geq 1 prior systemic therapy, had an Eastern Cooperative Oncology Group performance status of 0-2, adequate hematologic, hepatic, and renal function, and provided written informed consent before study entry.

Study design

This was a dose-finding study to determine an effective and well-tolerated pralatrexate dosing regimen with vitamin B₁₂ and folic acid supplementation in patients with relapsed/refractory CTCLs. Patients in the dose-finding cohorts were treated at sequentially decreasing pralatrexate dose intensities to identify an optimal dosing regimen, and an expansion cohort further explored that identified regimen. The optimal regimen was defined as the dose and schedule with \geq 1 response in up to 9 patients, an incidence of dose-limiting toxicities (DLTs) of $<$ 33%, and no patients with grade 4 hematologic toxicity, grade 3-4 infection, or febrile neutropenia.

The pralatrexate starting dose and schedule was 30 mg/m²/wk by IV push for 3 consecutive weeks followed by 1 rest week in a 4-week cycle. This regimen was based on a previous phase 1-2 dose-escalation study in patients with relapsed/refractory lymphoid malignancies and was the lowest dose explored in that study.³² On occurrence of predefined DLTs, which included many grade 2 toxicities in an effort to find a minimally toxic dose, each subsequent cohort was treated at the next lower pralatrexate dose of 20, 15, or 10 mg/m², or the next less frequent dosing schedule of 3 of 4 (3/4) or 2 of 3 (2/3) weeks.

Protocol-defined DLTs that would trigger dose and/or schedule de-escalation included grade 3-4 neutropenia, grade 2-4 thrombocytopenia or any thrombocytopenia with clinically significant bleeding (excluding epistaxis), any febrile neutropenia, grade 2-4 nonhematologic adverse event (AE), dose reduction/omission in cycle 1 for treatment-related AE, or initiation of cycle 2 delayed more than 1 week for treatment-related AE.

Pralatrexate was continued until progressive disease (PD) or 1 of the following criteria for treatment discontinuation was met: initiation of subsequent lymphoma therapy; intolerance of pralatrexate treatment; lapse of 2 weeks between pralatrexate doses within a cycle or 3 weeks between cycles; withdrawal of consent; or investigator/sponsor decision. Patients received treatment for a maximum of 12 months; however, on clinical or radiologic evidence of benefit in the investigator's opinion, patients could continue to receive pralatrexate. All patients received supplementation with vitamin B₁₂ (1 mg intramuscularly every 8-10 weeks) and folic acid (1 mg orally once daily) before study treatment initiation (at least 10 days prior for folic acid and within 10 weeks prior for vitamin B₁₂) through

30 days after study treatment discontinuation. Patients were not permitted to initiate treatment with systemic or topical corticosteroids while on study; however, patients could continue to receive topical corticosteroids or a dose of no more than 10 mg/d of prednisone (or equivalent) provided the corticosteroids were ongoing for at least 1 month before study entry.

Pralatrexate dose modifications for mucositis were: (1) for grade 2, treatment was held until recovery to grade 0-1, with the following dose administered without reduction; and (2) for recurrent grade 2 or grade 3-4, treatment was held until recovery to grade 0-1, followed by reduction to the next lower dose level, with discontinuation for recurrence. For other nonhematologic toxicities, absolute neutrophil counts, and platelet counts, doses were omitted for grade 3-4 until recovery to grade 0-2, followed by reduction to the next lower dose, with discontinuation for recurrence.

Assessments

Disease staging was performed using the tumor, node, metastasis (TNM) classification system.³³ During the course of the study, the TNM system was modified to account for blood involvement (TNMB).³⁴ Disease staging was conducted using the TNMB for patients enrolled in this study after the modified criteria were published; however, no retrospective staging was performed for previously enrolled patients.

Response to treatment was evaluated by the investigator every 2 cycles for 6 months and every 4 cycles thereafter using a standard response assessment for CTCL studies: the modified severity-weighted adjustment tool.⁵ For patients with lymph node involvement, computed tomography scans were obtained at baseline and on clinical response or end of treatment, whichever occurred first. Clinical photographs were not required for response assessment. Response rate, PFS, and response duration were determined. Patients were evaluable for response if they received \geq 1 pralatrexate dose.

Safety was evaluated by physical examinations, clinical laboratory evaluations, and treatment-emergent AEs. The incidence and severity of AEs were graded using the National Cancer Institute Common Terminology Criteria for AEs Version 3.0, and were coded using the Medical Dictionary for Regulatory Activities Version 11.0.

Statistical analysis

Safety and efficacy data, baseline values, and demographics were analyzed for all patients who received \geq 1 pralatrexate dose. Data were collected during treatment and for 35 (\pm 5) days after the last pralatrexate dose. Comparative statistical testing was not performed in this single-arm study.

Investigator assessment of response was defined as either complete response (CR) or partial response (PR). Duration of response was calculated from the day of first response (per investigator) until PD or death. Responding patients who had not progressed were censored at the last response assessment before study end. Progression was defined as either PD or death, and PFS was calculated from the first pralatrexate dose until PD or death. Patients who came off therapy without PD or death were censored for PFS at the last response assessment before study end.³⁵ Kaplan-Meier estimates of the median duration of response and PFS were generated.

Results

Overall patient characteristics

Between August 2007 and October 2010, 54 patients were treated at 8 US sites (31 in the dose-finding cohorts and 23 in the expansion cohort). Patient characteristics are shown in Table 1. This was a heavily pretreated population, with a median of 6.5 prior regimens (range, 1-25) and 4 prior systemic regimens (range, 1-11), given either sequentially or in combination. Overall, 69% of patients had received prior oral bexarotene, 83% had received other prior systemic chemotherapy, and 70% had received prior immunotherapy (mostly interferon and denileukin diftitox); most had received both systemic chemotherapy and immunotherapy.

Table 1. Baseline patient characteristics

Parameter	Patients (N = 54)
Sex, n (%)	
Male	32 (59)
Female	22 (41)
Race/ethnicity, n (%)	
White	39 (72)
Black	12 (22)
Hispanic	2 (4)
Other	1 (2)
Median age (range), y	61.5 (30-81)
CTCL subtype (per investigator)	
Mycosis fungoides	38 (70)
Sézary syndrome	15 (28)
Primary cutaneous anaplastic large cell lymphoma	1 (2)
Median number of all prior therapies (range)	6.5 (1-25)
Median number of systemic prior therapies (range)	4.0 (1-11)
Nonsystemic therapies, n (%)	
Phototherapy	27 (50)
Topical corticosteroids	27 (50)
Radiation therapy	26 (48)
Topical chemotherapy	24 (44)
Topical bexarotene	10 (19)
Other nonsystemic therapy	8 (15)
Systemic therapies, n (%)	
Oral bexarotene	37 (69)
Interferon	28 (52)
HDAC inhibitors	27 (50)
Denileukin diftitox	14 (26)
Methotrexate	13 (24)
mAb*	12 (22)
Gemcitabine	10 (19)
Other single-agent chemotherapy	29 (54)
CHOP or CHOP-like regimens	7 (13)
Other multiagent chemotherapy	8 (15)
Extracorporeal photopheresis	19 (35)
Corticosteroids	13 (24)
Other systemic therapy	8 (15)

CHOP indicates cyclophosphamide/doxorubicin/vincristine/prednisone; CTCL, cutaneous T-cell lymphoma; and HDAC, histone deacetylase.

*Includes alemtuzumab and anti-CD4 Ab.

Dose-finding cohorts

In the dose-finding cohorts, 31 patients were enrolled sequentially and treated at de-escalating doses and schedules until an optimal dosing regimen was determined. The regimens evaluated, number of patients treated, and observed DLTs and responses are summarized in Table 2. The optimal dosing regimen was originally defined as the dose and schedule with ≥ 1 response in up to 9 patients and a DLT incidence rate of < 33%. The only dosing regimen to demonstrate this DLT rate was pralatrexate 10 mg/m² weekly for

3/4 weeks (cohort 6: 3/10 patients [30%] with DLTs), but this cohort did not demonstrate sufficient activity, with only one response in 10 patients. As no dosing regimen met the predefined criteria for expansion, a review of toxicity and efficacy data was performed across the overall pralatrexate safety profile to assess the risk:benefit profiles of each cohort.

Table 2 presents responses observed in the dose-finding cohorts. High response rates (≥ 50%) were seen at 30 mg/m² for 3/4 weeks, 20 mg/m² for 3/4 and 2/3 weeks, and 15 mg/m² for 3/4 weeks, with a combined response rate at these doses (cohorts 1-4) of 11/18 (61%), and an apparent threshold for activity was observed with a response rate of 1/13 (8%) in patients treated at lower dose intensities (cohorts 5-6). While an active and essentially nontoxic dose was not identified, cohorts 3 (20 mg/m²/wk for 2/3 weeks) and 4 (15 mg/m²/wk for 3/4 weeks) had the greatest risk:benefit profiles and both were considered for the recommended dose. After review among the investigators, DLTs in cohort 4 were considered to be less clinically significant with an absence of grade 3 toxicities compared with those in cohort 3. Therefore, the dose and schedule evaluated in cohort 4 (15 mg/m²/wk for 3/4 weeks) was chosen as the recommended dosing regimen to be explored in the expansion cohort, and the protocol was amended to reflect this change.

Expansion cohort

The cohort that received 15 mg/m² weekly for 3/4 weeks (n = 6) was further expanded to an additional 23 patients, for a total of 29 patients treated with this recommended dosing regimen. At the time of this analysis, 6 of these patients remained on pralatrexate therapy and 23 patients had discontinued therapy. Reasons for discontinuation were: 6 patients (21%) experienced PD; 7 patients (24%) per investigator decision because of lack of response (n = 5), intolerance of therapy (n = 1), and surgery for preexisting condition (n = 1); 3 patients (10%) for AEs; 3 patients (10%) for receiving subsequent therapy (includes one patient who went on to a transplantation); and 4 patients (14%) per patient decision (because of withdrawal of consent [n = 2] and patient relocation [n = 2]).

All 29 patients in the expansion cohort were evaluable. Pralatrexate was administered for a median of 4 cycles (range, 1-23 cycles), and the median treatment duration was 99 days. The response rate among patients treated with a dose of 15 mg/m² weekly for 3/4 weeks was 45% (13/29; 1 CR, 12 PRs; 95% confidence interval [CI]: 26.4%-64.3%), as displayed in Table 3 by CTCL stage and subtype shown. The median time to best response was 57 days (range, 44-168 days). The median response duration was not reached because of censoring (range, 1*-372* days; * = censored for subsequent therapy, study termination, in response at time of analysis, or 1 day if no further response

Table 2. DLT and response in the dose-finding cohorts

Cohort	Pralatrexate dose, mg/m ² /wk, schedule	n	Patients with DLTs,* N (toxicity, grade)	Response rate, % (n/N)	Response type
1	30, 3/4 wks	2	2 (anorexia, grade 2; weakness, grade 3)	100 (2/2)	2 PR
2	20, 3/4 wks	3	2 (stomatitis, grade 2)	67 (2/3)	2 PR
3	20, 2/3 wks	7	3 (stomatitis, grade 2-3; elevated liver function tests, grade 3)	57 (4/7)	1 CR/3 PR
4	15, 3/4 wks	6	3 (stomatitis, grade 2; fatigue, grade 2)	50 (3/6)	3 PR
5	15, 2/3 wks	3	2 (stomatitis, grade 2; dehydration, grade 2)	0 (0/3)	—
6	10, 3/4 wks	10	3 (thrombocytopenia/neutropenia, grade 3; skin lesion, grade 3; zoster, grade 3)	10 (1/10)	1 CR

2/3 indicates 2 of 3 weeks; 3/4, 3 of 4 weeks; DLT, dose limiting toxicity; CR, complete response; PR, partial response; and —, no response noted.

*Some patients experienced more than 1 DLT.

Table 3. Response by CTCL subtype and stage for all patients (N = 54)

CTCL subtype (per investigator)	Stage	Best response and response rate						Overall, % (n/N)	Best response
		15 mg/m ² , 3/4 wk % (n/N)	Best response	15 mg/m ² or more, 3/4 wk % (n/N)	Best response	< 15 mg/m ² , 3/4 wk % (n/N)	Best response		
Mycosis fungoides	IB	60 (3/5)	3 PR	63 (5/8)	5 PR	0 (0/2)	—	50 (5/10)	5 PR
	IIA	0 (0/2)	—	0 (0/2)	—	—	—	0 (0/2)	—
	IIB	67 (4/6)	4 PR	67 (8/12)	8 PR	20 (1/5)	1 CR	53 (9/17)	1 CR; 8 PR
	III	50 (1/2)	1 PR	50 (1/2)	1 PR	—	—	50 (1/2)	1 PR
	IVA	60 (3/5)	3 PR	60 (3/5)	3 PR	0 (0/1)	—	50 (3/6)	3 PR
Sézary syndrome	IVB	0 (0/1)	—	0 (0/1)	—	—	—	0 (0/1)	—
	IIB*	50 (1/2)	1 PR	50 (1/2)	1 PR	—	—	50 (1/2)	1 PR
	III	33 (1/3)	1 CRu	25 (1/4)	1 CRu	—	—	25 (1/4)	1 CRu
	IVA	0 (0/2)	—	0 (0/2)	—	0 (0/5)	—	0 (0/7)	—
Primary cutaneous anaplastic large cell lymphoma	IVB	0 (0/1)	—	50 (1/2)	1 PR	—	—	50 (1/2)	1 PR
	IIB	—	—	100 (1/1)	—	—	—	100 (1/1)	1 CR
All patients		45 (13/29) (95% CI: 26.4-64.3)		51 (21/41) (95% CI: 35.1-67.1)		8 (1/13) (95% CI: 0.2-36.0)		41 (22/54) (95% CI: 27.6-55.0)	

3/4 indicates 3 of 4 weeks; CTCL, cutaneous T-cell lymphoma; CI, confidence interval; CR, complete response; CRu, CR unconfirmed; TNM, tumor, node, metastasis; PR, partial response; and —, no response noted.

*Patient was assessed using the original TNM criteria, which did not account for blood involvement.

assessments were performed after response was observed), although the Kaplan-Meier estimate shows 73% of responses continuing at 6 months. Among the 13 patients who responded at the pralatrexate dose of 15 mg/m² weekly for 3/4 weeks, the median number of cycles administered was 7 (range, 4-23 cycles) and the median duration of treatment was 159 days.

PFS was calculated for the 29 patients treated with 15 mg/m² weekly for 3/4 weeks (Figure 1A) and for all patients treated at a dose intensity of ≥ 15 mg/m² weekly for 3/4 weeks (Figure 1B). Median PFS was not reached for the 15 mg/m² weekly for 3/4 weeks regimen (range, 1*-429* days; * = censored for subsequent therapy, study termination, continuation of follow-up, or censored to treatment day 1). Median PFS for the 41 patients who received ≥ 15 mg/m² weekly for 3/4 weeks was 388 days (range, 1*-722* days).

Two patients in the 20 mg/m² for 2/3 weeks dose-finding cohort stopped therapy per protocol while still in response to pralatrexate (1 PR [response duration at study end = 218 days] and 1 CR [response duration at study end = 94 days]). On disease recurrence or progression, both patients were reentered into the study as permitted by the protocol. The patient with a PR on pralatrexate was retreated at 20 mg/m² weekly for 2/3 weeks and again achieved a PR that was continuing as of the data collection end point. The patient with a CR on pralatrexate was retreated at a lower dose (15 mg/m²/wk for 2/3 weeks) and again achieved a CR on pralatrexate that lasted 154 days; this patient was still in response when he discontinued pralatrexate treatment because of a nontreatment-related AE.

Dose escalation to 20 mg/m² was permitted per investigator discretion for patients in the expansion cohort who did not experience a CR and tolerated treatment for 2 complete cycles. Two of the 23 patients treated in the expansion cohort met the criteria for dose escalation to 20 mg/m² weekly for 3/4 weeks. One of these patients achieved a PR at the assessment directly following escalation, and received a total of 5 doses at 20 mg/m² weekly for 3/4 weeks. The other patient maintained stable disease and received 13 doses at 20 mg/m² weekly for 3/4 weeks.

The most common AEs, regardless of causality, reported in patients treated with pralatrexate 15 mg/m² weekly for 3/4 weeks

included mucositis (48%), fatigue (41%), nausea (31%), edema (28%), epistaxis (24%), pyrexia (21%), anorexia (21%), and skin toxicity (21%). The only grade 3 AEs that occurred in > 1 patient in this dosing regimen were mucositis (17%) and skin toxicity (7%),




Figure 1. Kaplan-Meier estimates of progression-free survival based on investigator assessment of response. (A) PFS for 29 patients treated at the optimal dosing regimen of pralatrexate 15 mg/m² weekly for 3 of 4 (3/4) weeks and (B) for 41 patients treated at pralatrexate ≥ 15 mg/m² weekly for 3/4 weeks.

Table 4. Treatment-related AEs in $\geq 5\%$ of patients in the overall population (N = 54)

AE, n (%)	15 mg/m ² /wk for 3/4 wk, n = 29				15 mg/m ² /wk or more for 3/4 wk, n = 41				< 15 mg/m ² /wk for 3/4 wk, n = 13			
	All	Grade 1-2	Grade 3	Grade 4+	All	Grade 1-2	Grade 3	Grade 4+	All	Grade 1-2	Grade 3	Grade 4+
Any treatment-related AE	20 (69)	12 (41)	8 (28)	0 (0)	32 (78)	16 (39)	16 (39)	0 (0)	12 (92)	7 (54)	4 (31)	1 (8)
Mucosal inflammation (stomatitis)*	14 (48)	9 (31)	5 (17)	0 (0)	23 (56)	16 (39)	7 (17)	0 (0)	6 (46)	5 (38)	1 (8)	0 (0)
Fatigue	11 (38)	10 (34)	1 (3)	0 (0)	17 (41)	16 (39)	1 (2)	0 (0)	5 (38)	5 (38)	0 (0)	0 (0)
Nausea	9 (31)	9 (31)	0 (0)	0 (0)	16 (39)	16 (39)	0 (0)	0 (0)	5 (38)	5 (38)	0 (0)	0 (0)
Skin toxicity*	6 (21)	4 (14)	2 (7)	0 (0)	7 (17)	5 (12)	2 (5)	0 (0)	3 (23)	2 (15)	1 (8)	0 (0)
Vomiting	4 (14)	4 (14)	0 (0)	0 (0)	6 (15)	6 (15)	0 (0)	0 (0)	2 (15)	2 (15)	0 (0)	0 (0)
Pyrexia	2 (7)	2 (7)	0 (0)	0 (0)	5 (12)	5 (12)	0 (0)	0 (0)	3 (23)	3 (23)	0 (0)	0 (0)
Epistaxis	7 (24)	7 (24)	0 (0)	0 (0)	7 (17)	7 (17)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Edema*	4 (14)	4 (14)	0 (0)	0 (0)	4 (10)	4 (10)	0 (0)	0 (0)	3 (23)	3 (23)	0 (0)	0 (0)
Anemia*	1 (3)	1 (3)	0 (0)	0 (0)	4 (10)	2 (5)	2 (5)	0 (0)	3 (23)	3 (23)	0 (0)	0 (0)
Anorexia*	3 (10)	3 (10)	0 (0)	0 (0)	4 (10)	4 (10)	0 (0)	0 (0)	1 (8)	1 (8)	0 (0)	0 (0)
Diarrhea	2 (7)	2 (7)	0 (0)	0 (0)	3 (7)	3 (7)	0 (0)	0 (0)	2 (15)	2 (15)	0 (0)	0 (0)
Upper respiratory tract infection	2 (7)	2 (7)	0 (0)	0 (0)	3 (7)	3 (7)	0 (0)	0 (0)	1 (8)	1 (8)	0 (0)	0 (0)
Pruritus*	2 (7)	2 (7)	0 (0)	0 (0)	2 (5)	2 (5)	0 (0)	0 (0)	1 (8)	1 (8)	0 (0)	0 (0)
Thrombocytopenia*	2 (7)	1 (3)	1 (3)	0 (0)	2 (5)	1 (2)	1 (2)	0 (0)	1 (8)	0 (0)	1 (8)	0 (0)
Burning sensation	1 (3)	1 (3)	0 (0)	0 (0)	1 (2)	1 (2)	0 (0)	0 (0)	2 (15)	2 (15)	0 (0)	0 (0)
Dyspnea	1 (3)	1 (3)	0 (0)	0 (0)	2 (5)	2 (5)	0 (0)	0 (0)	1 (8)	1 (8)	0 (0)	0 (0)
Liver function test abnormal*	0 (0)	0 (0)	0 (0)	0 (0)	2 (5)	1 (2)	1 (2)	0 (0)	2 (15)	2 (15)	0 (0)	0 (0)
Rash*	1 (3)	1 (3)	0 (0)	0 (0)	2 (5)	2 (5)	0 (0)	0 (0)	1 (8)	1 (8)	0 (0)	0 (0)
Staphylococcal infection	1 (3)	1 (3)	0 (0)	0 (0)	2 (5)	2 (5)	0 (0)	0 (0)	1 (8)	0 (0)	1 (8)	0 (0)
Headache	0 (0)	0 (0)	0 (0)	0 (0)	2 (5)	2 (5)	0 (0)	0 (0)	1 (8)	1 (8)	0 (0)	0 (0)
Lymphopenia	0 (0)	0 (0)	0 (0)	0 (0)	1 (2)	0 (0)	1 (2)	0 (0)	2 (15)	1 (8)	1 (8)	0 (0)

3/4 indicates 3 of 4; and AE, adverse event.

*Includes closely related Medical Dictionary for Regulatory Activities preferred terms that have been grouped as a single term.

and the only grade 4 AE was leukopenia, which was reported in 1 patient ~ 2 weeks after discontinuing pralatrexate treatment. Treatment-related AEs in $\geq 5\%$ of patients are detailed in Table 4. At a dose of 15 mg/m² weekly for 3/4 weeks, no treatment-related AEs of neutropenia were reported, the only treatment-related AE of anemia was grade 2, and no grade 4 treatment-related AEs were reported.

All patients

For the entire study, the overall response rate was 41% (22/54; 95% CI: 27.6%-55.0%), including 3 CRs and 19 PRs. Response is presented by CTCL subtype and disease stage in Table 3 for the overall patient population (N = 54), for patients treated with 15 mg/m² weekly for 3/4 weeks (n = 29), for patients treated with ≥ 15 mg/m² weekly for 3/4 weeks (n = 41), and for patients treated with < 15 mg/m² weekly for 3/4 weeks (n = 13). Patients with both advanced MF and SS achieved responses with pralatrexate; Figure 2 shows a patient with MF stage II B who experienced a PR. There was only one patient with ALCL in this study, and this patient achieved a CR on pralatrexate, as illustrated in Figure 3. Logistic regression analyses revealed no correlations between response and disease stage or number of prior therapies (data not shown).

The type of prior therapy received did not predict responses to pralatrexate. Response rates in patients who were previously treated with and progressed following bexarotene (n = 37), histone deacetylase (HDAC) inhibitors (n = 27; mostly vorinostat), interferon (n = 28), and methotrexate (n = 13) were 46%, 41%, 36%, and 46%, respectively.

Among the entire population, 50 patients (93%) experienced an AE on study or within 30 days of the last dose. The most frequently reported AEs regardless of causality were mucositis (54%), fatigue (43%), nausea (39%), skin toxicity (28%), edema (26%), anemia

(22%), and pyrexia (22%). Grade 3 treatment-related AEs reported in > 1 patient included mucositis (n = 8 [15%]), skin toxicity (n = 3 [6%]), and anemia, lymphopenia, and thrombocytopenia (each in 2 patients [4%]). Only one grade 4 treatment-related AE was reported (pulmonary embolism in a patient treated in the 15 mg/m²/wk for 2/3 weeks cohort). The only serious AE reported in > 2 patients was skin toxicity (n = 3 [6%]); serious AEs reported in 2 patients (4%) included failure to thrive, hypotension, mucositis, acute renal failure, and sepsis. When the incidence of \geq grade 2 mucositis was compared with baseline methylmalonic acid levels (above and below 200 nmol/L), no definitive correlation was observed.

Discussion

None of the current therapeutic options available for patients with CTCLs are curative. Many patients are treated sequentially and suffer from frequent morbidity because of the burden of their disease and the cumulative toxicity of therapy.

The present study identified an effective CTCL dosing regimen for pralatrexate with an AE profile that is acceptable for continuous long-term use. For patients treated with 15 mg/m² weekly for 3/4 weeks and ≥ 15 mg/m² weekly for 3/4 weeks, the response rates were 45% and 51%, respectively. These results were obtained in heavily pretreated patients with relapsed/refractory CTCLs with a median of 4 prior systemic therapies. These results compare with the single agents commonly used for the treatment of CTCLs (vorinostat, denileukin diftitox, oral bexarotene, and romidepsin) for which these patients had been previously treated and have reported response rates between 30% and 54% with response durations of 6 to 15 months.⁵⁻¹⁰ The median time to response for patients in the expansion cohort (57 days) is also comparable with

A

Patient with Stage IIB MF Pretreatment

B

Partial Response Post Cycle 3

Figure 2. Response to pralatrexate in patients with cutaneous T-cell lymphoma. Skin photographs documenting disease in a patient with mycosis fungoides (MF) stage IIB (A) before the start of treatment with pralatrexate and (B) after 4 cycles of treatment with pralatrexate.

or shorter than those reported for these other agents, which range from 42 to 180 days.⁵⁻¹⁰ While most of the responses observed in the expansion cohort were PRs (12 of 13; 92%), this proportion falls within the range of other published data, where 67% to 96% of all responses were PRs.⁵⁻¹⁰ The most frequent AEs experienced with this effective pralatrexate regimen were stomatitis (48%; 14 of 29) and fatigue (38%; 11 of 29). Similar incidence and grades of AEs have been reported for these other agents.⁵⁻¹⁰

In patients receiving pralatrexate 15 mg/m² weekly for 3/4 weeks, the median response duration was not reached (1*-372* days [* = censored]), and the Kaplan-Meier estimate for duration of response at 6 months was 73%. Many of the single agents used for CTCL treatment are biologic agents or non-traditional cytotoxic chemotherapies such as retinoids, interferons, and HDAC inhibitors. These agents are preferred because they can be administered long-term without cumulative toxicities, particularly immunosuppression. At the pralatrexate dosing regimens explored in this study, patients were treated for up to ~ 2 years without cumulative myelosuppression or an increased rate of infection. Among the 54 treated patients, there was only one patient (2%) each with treatment-related grade 3 neutropenia and grade 3-4 staphylococcal infection, which is one of the more significant and dangerous toxicities commonly associated with chemotherapy for patients with CTCLs. Furthermore, the rate of AEs reported across all doses was significantly lower than that seen in the pivotal study in PTCL, where the administered dose intensity was 30 mg/m² weekly for 6/7 weeks.

Interestingly, pralatrexate had a 46% response rate in patients who progressed following prior methotrexate, reinforcing that pralatrexate may exhibit potentially non-cross-resistant mechanisms of action compared with methotrexate.

In conclusion, the starting dose of pralatrexate 15 mg/m² weekly for 3/4 weeks demonstrated high activity with acceptable toxicity in a relapsed/refractory CTCL population. While this 54-patient study is reasonably sized for this uncommon illness, a confirmatory study is warranted. Pralatrexate represents a viable treatment option with the potential to provide durable benefit for patients with advanced-stage CTCLs who have been previously treated with many of the currently available agents. Moreover, the relatively low toxicity profile coupled with the described activity suggests that pralatrexate could form part of future combination studies with some of the active and nonmyelosuppressive agents used in CTCLs. Currently, phase 1 studies exploring pralatrexate in combination with HDAC inhibitors and retinoids are under way.

Acknowledgments

Monica Ramchandani, PhD, and Lindsey Lozano of Allos Therapeutics Inc provided medical writing support with direction and guidance from authors. Adrienne Schreiber of *inScience* Communications provided editing support, which was funded by Allos Therapeutics Inc.


This study was sponsored by Allos Therapeutics Inc.

The opinions expressed are those of the authors, who received no honoraria for manuscript development.

Authorship

Contribution: S.M.H. contributed to conceiving and designing the study, collecting, analyzing, and interpreting the data, and writing and approving the manuscript; Y.H.K. contributed to designing, collecting, and analyzing the data, interpreting the data, writing/commenting and approving the manuscript, and providing study materials or patients; F.F. contributed to writing/commenting and approving the manuscript; J.M.Z. contributed to collecting data, providing study material or patients, writing/commenting, and final approval of the manuscript; P.L.M. contributed to collecting and

Figure 3. Response to pralatrexate in patients with cutaneous T-cell lymphoma. Skin photographs documenting disease in a patient with anaplastic large cell lymphoma before the start of treatment with pralatrexate and after 4 cycles of treatment.



assembling data, writing/commenting, and approving the manuscript; M.J.L. contributed to analyzing and interpreting data, writing/commenting, and approving the manuscript; D.C.F. contributed to collecting, assembling, and interpreting data, writing/commenting on the manuscript, and providing final approval; A.R.S. contributed to collecting, analyzing, and interpreting the data, writing/commenting and approving the manuscript, and providing study material and patients; N.L.B. contributed to performing research, collecting and analyzing the data, writing/commenting, and approving the manuscript; M.L.D. contributed to collecting data and writing/commenting and approving the manuscript; T.K. contributed to concept, design, and analysis of the data, interpreting data and performing statistical analysis, writing/commenting, and approving the manuscript; M.E.S. contributed to the conception and design of the study, the collection and assembly of data, analysis and interpretation of the data, writing/commenting, and approving the final manuscript; O.A.O. contributed to designing research, analyzing and interpreting data, and writing/commenting on the manuscript; and M.D. contributed to the concept and design of the study, the collection, analysis, and interpretation of the data, writing/commenting, and approving the manuscript.

Conflict-of-interest disclosure: S.M.H. has served as a consultant for and received research funding from Allos Therapeutics Inc, Celgene, and Seattle Genetics, and has also received honorarium from Merck. Y.H.K. has served as a compensated consultant or advisor for Allos Therapeutics Inc, Gloucester, Celgene, Kyowa

Kirin, Seattle Genetics, Millennium, Therakos, Merck, and Eisai Co Ltd, and has received research funding from Kyowa Kirin, Merck, Allos Therapeutics Inc, Yaupon, and Seattle Genetics. F.F. has served as a consultant for Celgene, Eisai Co Ltd, and Merck, and received honoraria from Celgene, Cephalon, Eisai Co Ltd, and Merck. J.M.Z. has received honoraria for a speaking engagement for Allos Therapeutic Inc. M.J.L. has consulted with Allos Therapeutics Inc and was compensated. D.C.F. has served on an advisory board for Allos Therapeutics Inc for which he received honoraria. A.R.S. has received research funding and honorarium from Allos Therapeutics Inc. N.L.B. has received research funding from Allos Therapeutics Inc. M.L.D. has served as a compensated consultant and received honoraria for a speaking engagement for Allos Therapeutics Inc. T.K. is the Vice President of Biometrics for Allos Therapeutics Inc. M.E.S. is the Vice President of Clinical Drug Safety and Pharmacovigilance for Allos Therapeutics Inc, and has stock or other ownership interest in Allos Therapeutics Inc. O.A.O. has received grant funding from the US Food and Drug Administration (1 RO1 FD0003498-01), and has received funding for clinical trial support from Allos Therapeutics Inc. M.D. has received funding for clinical trial support from Allos Therapeutics Inc, and has served as a compensated advisor for Allos. P.L.M. declares no competing financial interests.

Correspondence: Steven M. Horwitz, MD, Dept of Medicine, Memorial Sloan-Kettering Cancer Center, 1275 York Ave, New York, NY 10065; e-mail: horwitzs@mskcc.org.

References

- Willemze R, Meijer CJ. Primary cutaneous CD30-positive lymphoproliferative disorders. *Hematol Oncol Clin North Am*. 2003;17(6):1319-1332.
- Horwitz SM, Olsen EA, Duvic M, Porcu P, Kim YH. Review of the treatment of mycosis fungoides and Sezary syndrome: a stage-based approach. *J Natl Compr Canc Netw*. 2008;6(4):436-442.
- Kim YH, Liu HL, Mraz-Gernhard S, Varghese A, Hoppe RT. Long-term outcome of 525 patients with mycosis fungoides and Sezary syndrome: clinical prognostic factors and risk for disease progression. *Arch Dermatol*. 2003;139(7):857-866.
- Kaye FJ, Bunn PA Jr, Steinberg SM, et al. A randomized trial comparing combination electron-beam radiation and chemotherapy with topical therapy in the initial treatment of mycosis fungoides. *N Engl J Med*. 1989;321(26):1784-1790.
- Olsen EA, Kim YH, Kuzel TM, et al. Phase IIb multicenter trial of vorinostat in patients with persistent, progressive, or treatment refractory cutaneous T-cell lymphoma. *J Clin Oncol*. 2007;25(21):3109-3115.
- Olsen E, Duvic M, Frankel A, et al. Pivotal phase III trial of two dose levels of denileukin diftitox for the treatment of cutaneous T-cell lymphoma. *J Clin Oncol*. 2001;19(2):376-388.
- Duvic M, Hymes K, Heald P, et al. Bexarotene is effective and safe for treatment of refractory advanced-stage cutaneous T-cell lymphoma: multinational phase II-III trial results. *J Clin Oncol*. 2001;19(9):2456-2471.
- Duvic M, Martin AG, Kim Y, et al. Phase 2 and 3 clinical trial of oral bexarotene (Targretin capsules) for the treatment of refractory or persistent early-stage cutaneous T-cell lymphoma. *Arch Dermatol*. 2001;137(5):581-593.
- Whittaker SJ, Demierre MF, Kim EJ, et al. Final results from a multicenter, international, pivotal study of romidepsin in refractory cutaneous T-cell lymphoma. *J Clin Oncol*. 2010;28(29):4485-4491.
- Piekarz RL, Frye R, Turner M, et al. Phase II multi-institutional trial of the histone deacetylase inhibitor romidepsin as monotherapy for patients with cutaneous T-cell lymphoma. *J Clin Oncol*. 2009;27(32):5410-5417.
- Prince HM, Duvic M, Martin A, et al. Phase III placebo-controlled trial of denileukin diftitox for patients with cutaneous T-cell lymphoma. *J Clin Oncol*. 2010;28(11):1870-1877.
- Zinzani PL, Musuraca G, Tani M, et al. Phase II trial of proteasome inhibitor bortezomib in patients with relapsed or refractory cutaneous T-cell lymphoma. *J Clin Oncol*. 2007;25(27):4293-4297.
- Lundin J, Hagberg H, Repp R, et al. Phase 2 study of alemtuzumab (anti-CD52 monoclonal antibody) in patients with advanced mycosis fungoides/Sezary syndrome. *Blood*. 2003;101(11):4267-4272.
- Zackheim HS, Kashani-Sabet M, McMillan A. Low-dose methotrexate to treat mycosis fungoides: a retrospective study in 69 patients. *J Am Acad Dermatol*. 2003;49(5):873-878.
- Olsen EA. Interferon in the treatment of cutaneous T-cell lymphoma. *Dermatol Ther*. 2003;16(4):311-321.
- Duvic M, Talpur R, Wen S, Kurzrock R, David CL, Apisarnthanarax N. Phase II evaluation of gemcitabine monotherapy for cutaneous T-cell lymphoma. *Clin Lymphoma Myeloma*. 2006;7(1):51-58.

17. Zinzani PL, Baliva G, Magagnoli M, et al. Gemcitabine treatment in pretreated cutaneous T-cell lymphoma: experience in 44 patients. *J Clin Oncol*. 2000;18(13):2603-2606.
18. Kurzrock R, Pilat S, Duvic M. Pentostatin therapy of T-cell lymphomas with cutaneous manifestations. *J Clin Oncol*. 1999;17(10):3117-3121.
19. Wollina U, Dummer R, Brockmeyer NH, et al. Multicenter study of pegylated liposomal doxorubicin in patients with cutaneous T-cell lymphoma. *Cancer*. 2003;98(5):993-1001.
20. Foss FM. Evaluation of the pharmacokinetics, preclinical and clinical efficacy of pralatrexate for the treatment of T-cell lymphoma. *Expert Opin Drug Metab Toxicol*. 2011;7(9):1141-1152.
21. Wang ES, O'Connor O, She Y, Zelenetz AD, Sirotiak FM, Moore MA. Activity of a novel antifolate (PDX, 10-propargyl 10-deazaaminopterin) against human lymphoma is superior to methotrexate and correlates with tumor RFC-1 gene expression. *Leuk Lymphoma*. 2003;44(6):1027-1035.
22. Sirotiak FM, DeGraw JI, Colwell WT, Piper JR. A new analogue of 10-deazaaminopterin with markedly enhanced curative effects against human tumor xenografts in mice. *Cancer Chemother Pharmacol*. 1998;42(4):313-318.
23. Krug LM, Ng KK, Kris mg, et al. Phase I and pharmacokinetic study of 10-propargyl-10-deazaaminopterin, a new antifolate. *Clin Cancer Res*. 2000;6(9):3493-3498.
24. Azzoli CG, Patel JD, Krug LM, et al. Pralatrexate with vitamin supplementation in patients with previously treated, advanced non-small cell lung cancer: safety and efficacy in a phase 1 trial. *J Thorac Oncol*. 2011;6(11):1915-1922.
25. Fury mg, Krug LM, Azzoli CG, et al. A phase I clinical pharmacologic study of pralatrexate in combination with probenecid in adults with advanced solid tumors. *Cancer Chemother Pharmacol*. 2006;57(5):671-677.
26. Krug LM, Azzoli CG, Kris mg, et al. 10-propargyl-10-deazaaminopterin: an antifolate with activity in patients with previously treated non-small cell lung cancer. *Clin Cancer Res*. 2003;9(6):2072-2078.
27. Krug LM, Heelan RT, Kris MG, Venkatraman E, Sirotiak FM. Phase II trial of pralatrexate (10-propargyl-10-deazaaminopterin, PDX) in patients with unresectable malignant pleural mesothelioma. *J Thorac Oncol*. 2007;2(4):317-320.
28. Azzoli CG, Krug LM, Gomez J, et al. A phase 1 study of pralatrexate in combination with paclitaxel or docetaxel in patients with advanced solid tumors. *Clin Cancer Res*. 2007;13(9):2692-2698.
29. O'Connor OA, Pro B, Pinter-Brown L, et al. Pralatrexate in patients with relapsed or refractory peripheral T-cell lymphoma: results from the pivotal PROPEL study. *J Clin Oncol*. 2011;29(9):1182-1189.
30. Mould DR, Sweeney K, Duffull SB, et al. A population pharmacokinetic and pharmacodynamic evaluation of pralatrexate in patients with relapsed or refractory non-Hodgkin's or Hodgkin's lymphoma. *Clin Pharmacol Ther*. 2009;86(2):190-196.
31. Horwitz SM, Vose JM, Advani R, et al. Pralatrexate and gemcitabine in patients with relapsed or refractory lymphoproliferative malignancies: phase 1 results [abstract]. *Blood (ASH Annual Meeting Abstracts)*. 2009:Abstract 1674.
32. O'Connor OA, Horwitz S, Hamlin P, et al. Phase II-I-II study of two different doses and schedules of pralatrexate, a high-affinity substrate for the reduced folate carrier, in patients with relapsed or refractory lymphoma reveals marked activity in T-cell malignancies. *J Clin Oncol*. 2009;27(26):4357-4364.
33. AJCC Cancer Staging Handbook. *American Joint Committee on Cancer*. 6th Ed. New York, NY: Springer; 2002.
34. Olsen E, Vonderheid E, Pimpinelli N, et al. Revisions to the staging and classification of mycosis fungoides and Sezary syndrome: a proposal of the International Society for Cutaneous Lymphomas (ISCL) and the cutaneous lymphoma task force of the European Organization of Research and Treatment of Cancer (EORTC). *Blood*. 2007;110(6):1713-1722.
35. Cheson BD, Pfistner B, Juweid ME, et al. Revised response criteria for malignant lymphoma. *J Clin Oncol*. 2007;25(5):579-586.